

Ruelle zeta at zero for nearly hyperbolic 3-manifolds

Semyon Dyatlov (MIT)

Feb 23, 2021

Overview

- Studying $m_R(0)$: order of vanishing at 0 of the **Ruelle zeta function** for the geodesic flow on a negatively curved 3-manifold (Σ, g)
- $g = g_H$ hyperbolic $\implies m_R(0) = 4 - 2b_1(\Sigma)$ [Fried '86]
- $g =$ generic perturbation of $g_H \implies m_R(0) = 4 - b_1(\Sigma)$ [Cekić–D–Küster–Paternain '20]
- This is in contrast with the case $\dim \Sigma = 2$ where $m_R(0) = b_1(\Sigma) - 2$ for all negatively curved (Σ, g) [D–Zworski '17]
- Motivated by **Fried's conjecture** '87 relating the values at 0 of twisted dynamical zeta functions to analytic torsion

Overview

- Studying $m_{\mathbb{R}}(0)$: order of vanishing at 0 of the **Ruelle zeta function** for the geodesic flow on a negatively curved 3-manifold (Σ, g)
- $g = g_H$ hyperbolic $\implies m_{\mathbb{R}}(0) = 4 - 2b_1(\Sigma)$ [Fried '86]
- $g =$ generic perturbation of $g_H \implies m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$ [Cekić–D–Küster–Paternain '20]
- This is in contrast with the case $\dim \Sigma = 2$ where $m_{\mathbb{R}}(0) = b_1(\Sigma) - 2$ for all negatively curved (Σ, g) [D–Zworski '17]
- Motivated by **Fried's conjecture** '87 relating the values at 0 of twisted dynamical zeta functions to analytic torsion

Overview

- Studying $m_{\mathbb{R}}(0)$: order of vanishing at 0 of the **Ruelle zeta function** for the geodesic flow on a negatively curved 3-manifold (Σ, g)
- $g = g_H$ hyperbolic $\implies m_{\mathbb{R}}(0) = 4 - 2b_1(\Sigma)$ [Fried '86]
- $g =$ generic perturbation of $g_H \implies m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$ [Cekić–D–Küster–Paternain '20]
- This is in contrast with the case $\dim \Sigma = 2$ where $m_{\mathbb{R}}(0) = b_1(\Sigma) - 2$ for all negatively curved (Σ, g) [D–Zworski '17]
- Motivated by **Fried's conjecture '87** relating the values at 0 of twisted dynamical zeta functions to analytic torsion

Overview

- Studying $m_{\mathbb{R}}(0)$: order of vanishing at 0 of the **Ruelle zeta function** for the geodesic flow on a negatively curved 3-manifold (Σ, g)
- $g = g_H$ hyperbolic $\implies m_{\mathbb{R}}(0) = 4 - 2b_1(\Sigma)$ [Fried '86]
- $g =$ generic perturbation of $g_H \implies m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$ [Cekić–D–Küster–Paternain '20]
- This is in contrast with the case $\dim \Sigma = 2$ where $m_{\mathbb{R}}(0) = b_1(\Sigma) - 2$ for all negatively curved (Σ, g) [D–Zworski '17]
- Motivated by **Fried's conjecture** '87 relating the values at 0 of twisted dynamical zeta functions to analytic torsion

Geodesic and contact flows

- (Σ, g) a compact connected oriented Riemannian n -dim manifold
- $M = S\Sigma$ the sphere bundle of (Σ, g) , $\pi_\Sigma : M \rightarrow \Sigma$ projection map
- $\alpha_{(x,v)}(\xi) = \langle v, d\pi_\Sigma(x, v)\xi \rangle_g$ canonical 1-form on M
- α is a contact form: $d\text{vol}_\alpha := \alpha \wedge (d\alpha)^{n-1}$ is nonvanishing
- Geodesic flow: $\varphi_t = e^{tX} : M \rightarrow M$ where $X \in C^\infty(M; TM)$ given by

$$\iota_X \alpha = 1, \quad \iota_X d\alpha = 0$$

- g has negative sectional curvature $\implies \varphi_t$ is Anosov:

$$TM = E_0 \oplus E_u \oplus E_s, \quad E_0 = \mathbb{R}X,$$

$$\exists C, \theta > 0 : \quad \|d\varphi_{-t}|_{E_u}\|, \|d\varphi_t|_{E_s}\| \leq Ce^{-\theta t}, \quad t \geq 0$$

- Define $E_u^* := (E_0 \oplus E_u)^\perp$, $E_s^* := (E_0 \oplus E_s)^\perp$ subsets of T^*M

Geodesic and contact flows

- (Σ, g) a compact connected oriented Riemannian n -dim manifold
- $M = S\Sigma$ the sphere bundle of (Σ, g) , $\pi_\Sigma : M \rightarrow \Sigma$ projection map
- $\alpha_{(x,v)}(\xi) = \langle v, d\pi_\Sigma(x, v)\xi \rangle_g$ canonical 1-form on M
- α is a contact form: $d\text{vol}_\alpha := \alpha \wedge (d\alpha)^{n-1}$ is nonvanishing
- Geodesic flow: $\varphi_t = e^{tX} : M \rightarrow M$ where $X \in C^\infty(M; TM)$ given by

$$\iota_X \alpha = 1, \quad \iota_X d\alpha = 0$$

- g has negative sectional curvature $\implies \varphi_t$ is **Anosov**:

$$TM = E_0 \oplus E_u \oplus E_s, \quad E_0 = \mathbb{R}X,$$

$$\exists C, \theta > 0 : \quad \|d\varphi_{-t}|_{E_u}\|, \|d\varphi_t|_{E_s}\| \leq Ce^{-\theta t}, \quad t \geq 0$$

- Define $E_u^* := (E_0 \oplus E_u)^\perp$, $E_s^* := (E_0 \oplus E_s)^\perp$ subsets of T^*M

Geodesic and contact flows

- (Σ, g) a compact connected oriented Riemannian n -dim manifold
- $M = S\Sigma$ the sphere bundle of (Σ, g) , $\pi_\Sigma : M \rightarrow \Sigma$ projection map
- $\alpha_{(x,v)}(\xi) = \langle v, d\pi_\Sigma(x, v)\xi \rangle_g$ canonical 1-form on M
- α is a contact form: $d\text{vol}_\alpha := \alpha \wedge (d\alpha)^{n-1}$ is nonvanishing
- Geodesic flow: $\varphi_t = e^{tX} : M \rightarrow M$ where $X \in C^\infty(M; TM)$ given by

$$\iota_X \alpha = 1, \quad \iota_X d\alpha = 0$$

- g has negative sectional curvature $\implies \varphi_t$ is **Anosov**:

$$TM = E_0 \oplus E_u \oplus E_s, \quad E_0 = \mathbb{R}X,$$

$$\exists C, \theta > 0 : \quad \|d\varphi_{-t}|_{E_u}\|, \|d\varphi_t|_{E_s}\| \leq Ce^{-\theta t}, \quad t \geq 0$$

- Define $E_u^* := (E_0 \oplus E_u)^\perp$, $E_s^* := (E_0 \oplus E_s)^\perp$ subsets of T^*M

Ruelle zeta function

Define the **Ruelle zeta function**

$$\zeta_{\mathbb{R}}(\lambda) = \prod_{\gamma} (1 - e^{-\lambda T_{\gamma}}), \quad \operatorname{Re} \lambda \gg 1$$

where the product is over all primitive closed geodesics γ of periods T_{γ}

- The function $\zeta_{\mathbb{R}}(\lambda)$ continues meromorphically to $\lambda \in \mathbb{C}$
[Giulietti–Liverani–Pollicott '13, D–Zworski '16]
Conjectured by Smale '67; partial progress by
Ruelle '76, Parry–Pollicott '90, Rugh '96, Fried '95
- Define the vanishing order $m_{\mathbb{R}}(0) \in \mathbb{Z}$:

$$\lambda^{-m_{\mathbb{R}}(0)} \zeta_{\mathbb{R}}(\lambda) \text{ holomorphic and nonvanishing at } \lambda = 0$$

Question

Can we describe $m_{\mathbb{R}}(0)$ in terms of topological invariants of Σ , such as the Betti numbers $b_k(\Sigma) = \dim H^k(\Sigma; \mathbb{R})$?

Ruelle zeta function

Define the **Ruelle zeta function**

$$\zeta_{\mathbb{R}}(\lambda) = \prod_{\gamma} (1 - e^{-\lambda T_{\gamma}}), \quad \operatorname{Re} \lambda \gg 1$$

where the product is over all primitive closed geodesics γ of periods T_{γ}

- The function $\zeta_{\mathbb{R}}(\lambda)$ **continues meromorphically** to $\lambda \in \mathbb{C}$
 [Giulietti–Liverani–Pollicott '13, D–Zworski '16]
 Conjectured by Smale '67; partial progress by
 Ruelle '76, Parry–Pollicott '90, Rugh '96, Fried '95
- Define the vanishing order $m_{\mathbb{R}}(0) \in \mathbb{Z}$:

$$\lambda^{-m_{\mathbb{R}}(0)} \zeta_{\mathbb{R}}(\lambda) \text{ holomorphic and nonvanishing at } \lambda = 0$$

Question

Can we describe $m_{\mathbb{R}}(0)$ in terms of topological invariants of Σ , such as the Betti numbers $b_k(\Sigma) = \dim H^k(\Sigma; \mathbb{R})$?

Ruelle zeta function

Define the **Ruelle zeta function**

$$\zeta_{\mathbb{R}}(\lambda) = \prod_{\gamma} (1 - e^{-\lambda T_{\gamma}}), \quad \operatorname{Re} \lambda \gg 1$$

where the product is over all primitive closed geodesics γ of periods T_{γ}

- The function $\zeta_{\mathbb{R}}(\lambda)$ **continues meromorphically** to $\lambda \in \mathbb{C}$
 [Giulietti–Liverani–Pollicott '13, D–Zworski '16]
 Conjectured by Smale '67; partial progress by
 Ruelle '76, Parry–Pollicott '90, Rugh '96, Fried '95
- Define the vanishing order $m_{\mathbb{R}}(0) \in \mathbb{Z}$:

$$\lambda^{-m_{\mathbb{R}}(0)} \zeta_{\mathbb{R}}(\lambda) \quad \text{holomorphic and nonvanishing at } \lambda = 0$$

Question

Can we describe $m_{\mathbb{R}}(0)$ in terms of topological invariants of Σ , such as the Betti numbers $b_k(\Sigma) = \dim H^k(\Sigma; \mathbb{R})$?

Previous work I

- More general zeta functions $\zeta_\rho(\lambda)$ **twisted** by a representation $\rho : \pi_1(\Sigma) \rightarrow U(m)$; $\zeta_{\mathbb{R}}$ corresponds to the trivial $\rho : \pi_1(\Sigma) \rightarrow U(1)$
- ρ is called **acyclic** if $H_\rho^k(\Sigma; \mathbb{R}) = 0$ for all k
- Fried '86 studied the **hyperbolic** case (curvature = -1):

$$m_{\mathbb{R}}(0) = \begin{cases} b_1(\Sigma) - 2, & \dim \Sigma = 2 \\ 4 - 2b_1(\Sigma), & \dim \Sigma = 3 \end{cases}$$

For ρ acyclic, he computed $m_\rho(0) = 0$ and $\zeta_\rho(0) = T_\rho^2$ where T_ρ is the **analytic torsion**. **Fried's conjecture**: same formula for $\zeta_\rho(0)$ holds for general locally homogeneous (Σ, g)

- Fried's conjecture proved for **locally symmetric** spaces by Shen '16, following Moscovici–Stanton '91, Bismut '11
- All the above use **Selberg trace formulas + representation theory**

Previous work I

- More general zeta functions $\zeta_\rho(\lambda)$ **twisted** by a representation $\rho : \pi_1(\Sigma) \rightarrow U(m)$; $\zeta_{\mathbb{R}}$ corresponds to the trivial $\rho : \pi_1(\Sigma) \rightarrow U(1)$
- ρ is called **acyclic** if $H_\rho^k(\Sigma; \mathbb{R}) = 0$ for all k
- **Fried** '86 studied the **hyperbolic** case (curvature = -1):

$$m_{\mathbb{R}}(0) = \begin{cases} b_1(\Sigma) - 2, & \dim \Sigma = 2 \\ 4 - 2b_1(\Sigma), & \dim \Sigma = 3 \end{cases}$$

For ρ acyclic, he computed $m_\rho(0) = 0$ and $\zeta_\rho(0) = T_\rho^2$ where T_ρ is the **analytic torsion**. **Fried's conjecture**: same formula for $\zeta_\rho(0)$ holds for general locally homogeneous (Σ, g)

- Fried's conjecture proved for **locally symmetric** spaces by **Shen** '16, following **Moscovici–Stanton** '91, **Bismut** '11
- All the above use **Selberg trace formulas + representation theory**

Previous work I

- More general zeta functions $\zeta_\rho(\lambda)$ **twisted** by a representation $\rho : \pi_1(\Sigma) \rightarrow U(m)$; $\zeta_{\mathbb{R}}$ corresponds to the trivial $\rho : \pi_1(\Sigma) \rightarrow U(1)$
- ρ is called **acyclic** if $H_\rho^k(\Sigma; \mathbb{R}) = 0$ for all k
- **Fried** '86 studied the **hyperbolic** case (curvature = -1):

$$m_{\mathbb{R}}(0) = \begin{cases} b_1(\Sigma) - 2, & \dim \Sigma = 2 \\ 4 - 2b_1(\Sigma), & \dim \Sigma = 3 \end{cases}$$

For ρ acyclic, he computed $m_\rho(0) = 0$ and $\zeta_\rho(0) = T_\rho^2$ where T_ρ is the **analytic torsion**. **Fried's conjecture**: same formula for $\zeta_\rho(0)$ holds for general locally homogeneous (Σ, g)

- Fried's conjecture proved for **locally symmetric** spaces by **Shen** '16, following **Moscovici–Stanton** '91, **Bismut** '11
- All the above use **Selberg trace formulas + representation theory**

Previous work II

What happens for general (not locally symmetric) negatively curved Σ ?

- [D-Zworski '17](#): $m_{\mathbb{R}}(0) = b_1(\Sigma) - 2$ when $\dim \Sigma = 2$;
applies to general contact Anosov flows in dimension 3
- Extended to surfaces with boundary by [Hadfield '18](#),
to the nonorientable case by [Borns-Weil-Shen '20](#)
- [Cekić-Paternain '19](#): studied $m_{\mathbb{R}}(0)$ for general volume preserving
Anosov flows on a 3-manifold M and showed it depends on the
properties of the flow, not just on the topology of M
- [Dang-Guillarmou-Rivière-Shen '20](#) proved Fried's conjecture on $\zeta_{\rho}(0)$
when Σ is any nearly hyperbolic 3-manifold
- Related works: [Dang-Rivière '17](#), [Chaubet-Dang '19](#),
[Küster-Weich '20](#)

Previous work II

What happens for general (not locally symmetric) negatively curved Σ ?

- [D-Zworski '17](#): $m_{\mathbb{R}}(0) = b_1(\Sigma) - 2$ when $\dim \Sigma = 2$;
applies to general contact Anosov flows in dimension 3
- Extended to surfaces with boundary by [Hadfield '18](#),
to the nonorientable case by [Borns-Weil-Shen '20](#)
- [Cekić-Paternain '19](#): studied $m_{\mathbb{R}}(0)$ for general volume preserving Anosov flows on a 3-manifold M and showed it depends on the properties of the flow, not just on the topology of M
- [Dang-Guillarmou-Rivière-Shen '20](#) proved Fried's conjecture on $\zeta_{\rho}(0)$ when Σ is any nearly hyperbolic 3-manifold
- Related works: [Dang-Rivière '17](#), [Chaubet-Dang '19](#),
[Küster-Weich '20](#)

Previous work II

What happens for general (not locally symmetric) negatively curved Σ ?

- [D-Zworski '17](#): $m_{\mathbb{R}}(0) = b_1(\Sigma) - 2$ when $\dim \Sigma = 2$;
applies to general contact Anosov flows in dimension 3
- Extended to surfaces with boundary by [Hadfield '18](#),
to the nonorientable case by [Borns-Weil-Shen '20](#)
- [Cekić-Paternain '19](#): studied $m_{\mathbb{R}}(0)$ for general volume preserving
Anosov flows on a 3-manifold M and showed it depends on the
properties of the flow, not just on the topology of M
- [Dang-Guillarmou-Rivière-Shen '20](#) proved Fried's conjecture on $\zeta_{\rho}(0)$
when Σ is any nearly hyperbolic 3-manifold
- Related works: [Dang-Rivière '17](#), [Chaubet-Dang '19](#),
[Küster-Weich '20](#)

Previous work II

What happens for general (not locally symmetric) negatively curved Σ ?

- [D-Zworski '17](#): $m_{\mathbb{R}}(0) = b_1(\Sigma) - 2$ when $\dim \Sigma = 2$;
applies to general contact Anosov flows in dimension 3
- Extended to surfaces with boundary by [Hadfield '18](#),
to the nonorientable case by [Borns-Weil-Shen '20](#)
- [Cekić-Paternain '19](#): studied $m_{\mathbb{R}}(0)$ for general volume preserving
Anosov flows on a 3-manifold M and showed it depends on the
properties of the flow, not just on the topology of M
- [Dang-Guillarmou-Rivière-Shen '20](#) proved Fried's conjecture on $\zeta_{\rho}(0)$
when Σ is any nearly hyperbolic 3-manifold
- Related works: [Dang-Rivière '17](#), [Chaubet-Dang '19](#),
[Küster-Weich '20](#)

Statement of the result

Theorem 1 [Cekić–D–Küster–Paternain '20]

Let (Σ, g_H) be a compact connected oriented hyperbolic 3-manifold. Then:

1. If $g = g_H$ then $m_{\mathbb{R}}(0) = 4 - 2b_1(\Sigma)$
2. If g is a generic conformal perturbation of g_H then $m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$

Here **generic conformal perturbation** is understood as follows:

there exists an open dense $\mathcal{O} \subset C^\infty(\Sigma; \mathbb{R})$ such that for any $a \in \mathcal{O}$ there exists $\varepsilon > 0$ such that for all $\tau \in (-\varepsilon, \varepsilon) \setminus \{0\}$ the metric $g = e^{\tau a} g_H$ has $m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$

- First result on instability of $m_{\mathbb{R}}(0)$ under metric perturbations
- Our proof of part 1 is different from [Fried '86], using geometric rather than algebraic techniques

Statement of the result

Theorem 1 [Cekić–D–Küster–Paternain '20]

Let (Σ, g_H) be a compact connected oriented hyperbolic 3-manifold. Then:

1. If $g = g_H$ then $m_{\mathbb{R}}(0) = 4 - 2b_1(\Sigma)$
2. If g is a generic conformal perturbation of g_H then $m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$

Here **generic conformal perturbation** is understood as follows:

there exists an open dense $\mathcal{O} \subset C^\infty(\Sigma; \mathbb{R})$ such that
 for any $a \in \mathcal{O}$ there exists $\varepsilon > 0$ such that for all $\tau \in (-\varepsilon, \varepsilon) \setminus \{0\}$
 the metric $g = e^{\tau a} g_H$ has $m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$

- First result on instability of $m_{\mathbb{R}}(0)$ under metric perturbations
- Our proof of part 1 is different from [Fried '86], using geometric rather than algebraic techniques

Spectral interpretation of zeta functions I

- General idea: “ $\zeta(\lambda) = \det(\lambda - P)$ ” for some operator P
- This should be understood as $\partial_\lambda \log \zeta(\lambda) = \text{tr}(\lambda - P)^{-1}$ with the right definition of trace
- Vanishing order of ζ at 0 = dimension of the space of generalized eigenstates at 0 $\{u \mid \exists \ell : P^\ell u = 0\}$
- One can write the vanishing order $m_{\mathbb{R}}(0)$ of $\zeta_{\mathbb{R}}$ using the dimensions of certain spaces of Pollicott–Ruelle generalized resonant forms $\text{Res}_0^{k,\infty}$
- Our strategy is to describe $\text{Res}_0^{k,\infty}$ in terms of the de Rham cohomology of Σ
- In this talk we will focus on the case $k = 1$

Spectral interpretation of zeta functions I

- General idea: “ $\zeta(\lambda) = \det(\lambda - P)$ ” for some operator P
- This should be understood as $\partial_\lambda \log \zeta(\lambda) = \text{tr}(\lambda - P)^{-1}$ with the right definition of trace
- Vanishing order of ζ at 0 = dimension of the space of **generalized eigenstates at 0** $\{u \mid \exists \ell : P^\ell u = 0\}$
- One can write the vanishing order $m_{\mathbb{R}}(0)$ of $\zeta_{\mathbb{R}}$ using the dimensions of certain spaces of Pollicott–Ruelle generalized resonant forms $\text{Res}_0^{k,\infty}$
- Our strategy is to describe $\text{Res}_0^{k,\infty}$ in terms of the de Rham cohomology of Σ
- In this talk we will focus on the case $k = 1$

Spectral interpretation of zeta functions I

- General idea: “ $\zeta(\lambda) = \det(\lambda - P)$ ” for some operator P
- This should be understood as $\partial_\lambda \log \zeta(\lambda) = \text{tr}(\lambda - P)^{-1}$ with the right definition of trace
- Vanishing order of ζ at 0 = dimension of the space of **generalized eigenstates at 0** $\{u \mid \exists \ell : P^\ell u = 0\}$
- One can write the vanishing order $m_{\mathbb{R}}(0)$ of $\zeta_{\mathbb{R}}$ using the dimensions of certain spaces of Pollicott–Ruelle generalized resonant forms $\text{Res}_0^{k,\infty}$
- Our strategy is to describe $\text{Res}_0^{k,\infty}$ in terms of the de Rham cohomology of Σ
- In this talk we will focus on the case $k = 1$

Spectral interpretation of zeta functions I

- General idea: “ $\zeta(\lambda) = \det(\lambda - P)$ ” for some operator P
- This should be understood as $\partial_\lambda \log \zeta(\lambda) = \text{tr}(\lambda - P)^{-1}$ with the right definition of trace
- Vanishing order of ζ at 0 = dimension of the space of **generalized eigenstates at 0** $\{u \mid \exists \ell : P^\ell u = 0\}$
- One can write the vanishing order $m_{\mathbb{R}}(0)$ of $\zeta_{\mathbb{R}}$ using the dimensions of certain spaces of Pollicott–Ruelle generalized resonant forms $\text{Res}_0^{k,\infty}$
- Our strategy is to describe $\text{Res}_0^{k,\infty}$ in terms of the de Rham cohomology of Σ
- In this talk we will focus on the case $k = 1$

Spectral interpretation of zeta functions II

- $M = S\Sigma$, $\dim \Sigma = 3$, $X \in C^\infty(M; TM)$ generates the geodesic flow
- Our operators: $P_{k,0} = \mathcal{L}_X$ acting on $\Omega_0^k := \{\omega \in \wedge^k T^*M \mid \iota_X \omega = 0\}$
- For certain anisotropic Sobolev spaces $\mathcal{H}, \mathcal{D}_P$ the operator $P_{k,0} - \lambda : \mathcal{D}_P(M; \Omega_0^k) \rightarrow \mathcal{H}(M; \Omega_0^k)$ is Fredholm of index 0
Blank–Keller–Liverani '02, Liverani '04, '05, Baladi '05,
Gouëzel–Liverani '06, Baladi–Tsujii '07, Butterley–Liverani '07
- We will use the microlocal/scattering theory approach:
Faure–Roy–Sjöstrand '08, Faure–Sjöstrand '11, D–Zworski '16

Spectral interpretation of zeta functions II

- $M = S\Sigma$, $\dim \Sigma = 3$, $X \in C^\infty(M; TM)$ generates the geodesic flow
- Our operators: $P_{k,0} = \mathcal{L}_X$ acting on $\Omega_0^k := \{\omega \in \wedge^k T^*M \mid \iota_X \omega = 0\}$
- For certain **anisotropic Sobolev spaces** $\mathcal{H}, \mathcal{D}_P$ the operator $P_{k,0} - \lambda : \mathcal{D}_P(M; \Omega_0^k) \rightarrow \mathcal{H}(M; \Omega_0^k)$ is Fredholm of index 0
 Blank–Keller–Liverani '02, Liverani '04,'05, Baladi '05,
 Gouëzel–Liverani '06, Baladi–Tsuji '07, Butterley–Liverani '07
- We will use the microlocal/scattering theory approach:
 Faure–Roy–Sjöstrand '08, Faure–Sjöstrand '11, D–Zworski '16

Spectral interpretation of zeta functions II

- $M = S\Sigma$, $\dim \Sigma = 3$, $X \in C^\infty(M; TM)$ generates the geodesic flow
- Our operators: $P_{k,0} = \mathcal{L}_X$ acting on $\Omega_0^k := \{\omega \in \wedge^k T^*M \mid \iota_X \omega = 0\}$
- For certain **anisotropic Sobolev spaces** $\mathcal{H}, \mathcal{D}_P$ the operator $P_{k,0} - \lambda : \mathcal{D}_P(M; \Omega_0^k) \rightarrow \mathcal{H}(M; \Omega_0^k)$ is Fredholm of index 0
 Blank–Keller–Liverani '02, Liverani '04,'05, Baladi '05,
 Gouëzel–Liverani '06, Baladi–Tsujii '07, Butterley–Liverani '07
- We will use the microlocal/scattering theory approach:
 Faure–Roy–Sjöstrand '08, Faure–Sjöstrand '11, D–Zworski '16

Spectral interpretation of zeta functions II

- $M = S\Sigma$, $\dim \Sigma = 3$, $X \in C^\infty(M; TM)$ generates the geodesic flow
- Our operators: $P_{k,0} = \mathcal{L}_X$ acting on $\Omega_0^k := \{\omega \in \wedge^k T^*M \mid \iota_X \omega = 0\}$
- For certain **anisotropic Sobolev spaces** \mathcal{H} , \mathcal{D}_P the operator $P_{k,0} - \lambda : \mathcal{D}_P(M; \Omega_0^k) \rightarrow \mathcal{H}(M; \Omega_0^k)$ is Fredholm of index 0
- The poles of $(P_{k,0} - \lambda)^{-1}$ are called **Pollicott–Ruelle resonances**
- Generalized resonant states at $\lambda = 0$:

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}_P(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X^\ell u = 0\}$$

- D–Zworski '16, using Hörmander's propagation of singularities, Melrose's radial estimates, and Atiyah–Bott–Guillemin trace formula:

$$m_{\mathbb{R}}(0) = \sum_{k=0}^4 (-1)^k \dim \text{Res}_0^{k,\infty}$$

Spectral interpretation of zeta functions II

- $M = S\Sigma$, $\dim \Sigma = 3$, $X \in C^\infty(M; TM)$ generates the geodesic flow
- Our operators: $P_{k,0} = \mathcal{L}_X$ acting on $\Omega_0^k := \{\omega \in \wedge^k T^*M \mid \iota_X \omega = 0\}$
- For certain **anisotropic Sobolev spaces** $\mathcal{H}, \mathcal{D}_P$ the operator $P_{k,0} - \lambda : \mathcal{D}_P(M; \Omega_0^k) \rightarrow \mathcal{H}(M; \Omega_0^k)$ is Fredholm of index 0
- The poles of $(P_{k,0} - \lambda)^{-1}$ are called **Pollicott–Ruelle resonances**
- Generalized resonant states at $\lambda = 0$:

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}_P(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X^\ell u = 0\}$$

- **D–Zworski** '16, using **Hörmander**'s propagation of singularities, **Melrose**'s radial estimates, and **Atiyah–Bott–Guillemin** trace formula:

$$m_{\text{R}}(0) = \sum_{k=0}^4 (-1)^k \dim \text{Res}_0^{k,\infty}$$

Resonance multiplicities

Theorem 1 follows from $m_{\mathbb{R}}(0) = \sum_{k=0}^4 (-1)^k \dim \text{Res}_0^{k,\infty}$ and

Theorem 2 [Cekić–D–Küster–Paternain '20]

Let (Σ, g_H) be a compact connected oriented hyperbolic 3-manifold. Then the dimensions of $\text{Res}_0^{k,\infty}$ are:

k	Hyperbolic	Perturbation
0	1	1
1	$2b_1(\Sigma)$	$b_1(\Sigma)$
2	$2b_1(\Sigma) + 2$	$b_1(\Sigma) + 2$
3	$2b_1(\Sigma)$	$b_1(\Sigma)$
4	1	1

$(d\alpha)^j \wedge : \text{Res}_0^{2-j,\infty} \rightarrow \text{Res}_0^{2+j,\infty}$ isomorphisms \Rightarrow study $k = 0, 1, 2$

Resonance multiplicities

Theorem 1 follows from $m_{\mathbb{R}}(0) = \sum_{k=0}^4 (-1)^k \dim \text{Res}_0^{k,\infty}$ and

Theorem 2 [Cekić–D–Küster–Paternain '20]

Let (Σ, g_H) be a compact connected oriented hyperbolic 3-manifold. Then the dimensions of $\text{Res}_0^{k,\infty}$ are:

k	Hyperbolic	Perturbation
0	1	1
1	$2b_1(\Sigma)$	$b_1(\Sigma)$
2	$2b_1(\Sigma) + 2$	$b_1(\Sigma) + 2$
3	$2b_1(\Sigma)$	$b_1(\Sigma)$
4	1	1

$(d\alpha)^j \wedge : \text{Res}_0^{2-j,\infty} \rightarrow \text{Res}_0^{2+j,\infty}$ isomorphisms \Rightarrow study $k = 0, 1, 2$

Resonant and coresonant states

- **Generalized resonant states:**

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}_P(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X^\ell u = 0\}$$

- $\mathcal{D}'_{E_u^*}(M; \Omega_0^k) = \{u \in \mathcal{D}'(M; \Omega_0^k) \mid \text{WF}(u) \subset E_u^*\}$

defined using **wavefront set** $\text{WF}(u) \subset T^*M \setminus 0$

- **Resonant states:** $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$

- **Coresonant states:** $\text{Res}_{0*}^k = \{u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^k) \mid \mathcal{L}_X u_* = 0\}$

$\text{Res}_{0*}^k = \mathcal{J}^* \text{Res}_0^k$ where $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$

- **Pairing:** $u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k)$, $u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^{4-k}) \mapsto \int_M \alpha \wedge u \wedge u_*$

- **Semisimplicity:** $\text{Res}_0^{k,\infty} = \text{Res}_0^k$, equivalent to the pairing being nondegenerate on $\text{Res}_0^k \times \text{Res}_{0*}^{4-k}$

- The case $k = 0$ is simple: $\text{Res}_0^{0,\infty} = \text{Res}_0^0 = \mathbb{R}1$

Resonant and coresonant states

- **Generalized resonant states:**

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X^\ell u = 0\}$$

- $\mathcal{D}'_{E_u^*}(M; \Omega_0^k) = \{u \in \mathcal{D}'(M; \Omega_0^k) \mid \text{WF}(u) \subset E_u^*\}$

defined using **wavefront set** $\text{WF}(u) \subset T^*M \setminus 0$

- **Resonant states:** $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$

- **Coresonant states:** $\text{Res}_{0*}^k = \{u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^k) \mid \mathcal{L}_X u_* = 0\}$

$\text{Res}_{0*}^k = \mathcal{J}^* \text{Res}_0^k$ where $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$

- **Pairing:** $u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k)$, $u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^{4-k}) \mapsto \int_M \alpha \wedge u \wedge u_*$

- **Semisimplicity:** $\text{Res}_0^{k,\infty} = \text{Res}_0^k$, equivalent to the pairing being nondegenerate on $\text{Res}_0^k \times \text{Res}_{0*}^{4-k}$

- The case $k = 0$ is simple: $\text{Res}_0^{0,\infty} = \text{Res}_0^0 = \mathbb{R}1$

Resonant and coresonant states

- **Generalized resonant states:**

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X u = 0\}$$

- $\mathcal{D}'_{E_u^*}(M; \Omega_0^k) = \{u \in \mathcal{D}'(M; \Omega_0^k) \mid \text{WF}(u) \subset E_u^*\}$

defined using **wavefront set** $\text{WF}(u) \subset T^*M \setminus 0$

- **Resonant states:** $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$

- **Coresonant states:** $\text{Res}_{0*}^k = \{u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^k) \mid \mathcal{L}_X u_* = 0\}$

$\text{Res}_{0*}^k = \mathcal{J}^* \text{Res}_0^k$ where $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$

- **Pairing:** $u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k)$, $u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^{4-k}) \mapsto \int_M \alpha \wedge u \wedge u_*$

- **Semisimplicity:** $\text{Res}_0^{k,\infty} = \text{Res}_0^k$, equivalent to the pairing being nondegenerate on $\text{Res}_0^k \times \text{Res}_{0*}^{4-k}$

- The case $k = 0$ is simple: $\text{Res}_0^{0,\infty} = \text{Res}_0^0 = \mathbb{R}1$

Resonant and coresonant states

- **Generalized resonant states:**

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X u = 0\}$$

- $\mathcal{D}'_{E_u^*}(M; \Omega_0^k) = \{u \in \mathcal{D}'(M; \Omega_0^k) \mid \text{WF}(u) \subset E_u^*\}$

defined using **wavefront set** $\text{WF}(u) \subset T^*M \setminus 0$

- **Resonant states:** $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$

- **Coresonant states:** $\text{Res}_{0*}^k = \{u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^k) \mid \mathcal{L}_X u_* = 0\}$

$\text{Res}_{0*}^k = \mathcal{J}^* \text{Res}_0^k$ where $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$

- **Pairing:** $u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k)$, $u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^{4-k}) \mapsto \int_M \alpha \wedge u \wedge u_*$

- **Semisimplicity:** $\text{Res}_0^{k,\infty} = \text{Res}_0^k$, equivalent to the pairing being nondegenerate on $\text{Res}_0^k \times \text{Res}_{0*}^{4-k}$

- The case $k = 0$ is simple: $\text{Res}_0^{0,\infty} = \text{Res}_0^0 = \mathbb{R}1$

Resonant and coresonant states

- **Generalized resonant states:**

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X u = 0\}$$

- $\mathcal{D}'_{E_u^*}(M; \Omega_0^k) = \{u \in \mathcal{D}'(M; \Omega_0^k) \mid \text{WF}(u) \subset E_u^*\}$

defined using **wavefront set** $\text{WF}(u) \subset T^*M \setminus 0$

- **Resonant states:** $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$

- **Coresonant states:** $\text{Res}_{0*}^k = \{u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^k) \mid \mathcal{L}_X u_* = 0\}$

$\text{Res}_{0*}^k = \mathcal{J}^* \text{Res}_0^k$ where $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$

- **Pairing:** $u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k)$, $u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^{4-k}) \mapsto \int_M \alpha \wedge u \wedge u_*$

- **Semisimplicity:** $\text{Res}_0^{k,\infty} = \text{Res}_0^k$, equivalent to the pairing being nondegenerate on $\text{Res}_0^k \times \text{Res}_{0*}^{4-k}$

- The case $k = 0$ is simple: $\text{Res}_0^{0,\infty} = \text{Res}_0^0 = \mathbb{R}1$

Resonant and coresonant states

- **Generalized resonant states:**

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X u = 0\}$$

- $\mathcal{D}'_{E_u^*}(M; \Omega_0^k) = \{u \in \mathcal{D}'(M; \Omega_0^k) \mid \text{WF}(u) \subset E_u^*\}$

defined using **wavefront set** $\text{WF}(u) \subset T^*M \setminus 0$

- **Resonant states:** $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$

- **Coresonant states:** $\text{Res}_{0*}^k = \{u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^k) \mid \mathcal{L}_X u_* = 0\}$

$\text{Res}_{0*}^k = \mathcal{J}^* \text{Res}_0^k$ where $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$

- **Pairing:** $u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k)$, $u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^{4-k}) \mapsto \int_M \alpha \wedge u \wedge u_*$

- **Semisimplicity:** $\text{Res}_0^{k,\infty} = \text{Res}_0^k$, equivalent to the pairing being nondegenerate on $\text{Res}_0^k \times \text{Res}_{0*}^{4-k}$

- The case $k = 0$ is simple: $\text{Res}_0^{0,\infty} = \text{Res}_0^0 = \mathbb{R}1$

Resonant and coresonant states

- **Generalized resonant states:**

$$\text{Res}_0^{k,\infty} = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \exists \ell : \mathcal{L}_X u = 0\}$$

- $\mathcal{D}'_{E_u^*}(M; \Omega_0^k) = \{u \in \mathcal{D}'(M; \Omega_0^k) \mid \text{WF}(u) \subset E_u^*\}$

defined using **wavefront set** $\text{WF}(u) \subset T^*M \setminus 0$

- **Resonant states:** $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$

- **Coresonant states:** $\text{Res}_{0*}^k = \{u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^k) \mid \mathcal{L}_X u_* = 0\}$

$\text{Res}_{0*}^k = \mathcal{J}^* \text{Res}_0^k$ where $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$

- **Pairing:** $u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k)$, $u_* \in \mathcal{D}'_{E_s^*}(M; \Omega_0^{4-k}) \mapsto \int_M \alpha \wedge u \wedge u_*$

- **Semisimplicity:** $\text{Res}_0^{k,\infty} = \text{Res}_0^k$, equivalent to the pairing being nondegenerate on $\text{Res}_0^k \times \text{Res}_{0*}^{4-k}$

- The case $k = 0$ is simple: $\text{Res}_0^{0,\infty} = \text{Res}_0^0 = \mathbb{R}\mathbf{1}$

Closed resonant 1-forms

- $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^k) \mid \mathcal{L}_X u = 0\}$, $\mathcal{L}_X = d\iota_X + \iota_X d$
- Closed forms: $\text{Res}_0^k \cap \ker d = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, du = 0\}$
- Cohomology map: $\pi_k : \text{Res}_0^k \cap \ker d \rightarrow H^k(M; \mathbb{R})$, $\pi_k(u) = [u]_{H^k}$
- π_k can be defined because $\mathcal{D}'_{E_u^*}$ is closed under $(d\delta + \delta d + 1)^{-1}$:
 $u \in \mathcal{D}'_{E_u^*}, du \in C^\infty \implies u = v + dw$ for some $v \in C^\infty, w \in \mathcal{D}'_{E_u^*}$

Lemma: π_1 is an isomorphism

Injectivity: if $u \in \text{Res}_0^1$ and $u = df$, $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$, then $Xf = \iota_X u = 0$, so $f \in \text{Res}_0^0 = \mathbb{R}\mathbf{1}$ and $u = df = 0$

Surjectivity: if $v \in C^\infty(M; \Omega^1)$ and $dv = 0$, then $\int_M (\iota_X v) d\text{vol}_\alpha = 0$, so by the Fredholm property there exists $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$ with $Xf = \iota_X v$. Take $u := v - df \in \text{Res}_0^1 \cap \ker d$, then $\pi_1(u) = [v]_{H^1}$

Closed resonant 1-forms

- $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, \iota_X du = 0\}$, $\Omega^k = \wedge^k T^*M$
- **Closed forms:** $\text{Res}_0^k \cap \ker d = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, du = 0\}$
- **Cohomology map:** $\pi_k : \text{Res}_0^k \cap \ker d \rightarrow H^k(M; \mathbb{R})$, $\pi_k(u) = [u]_{H^k}$
- π_k can be defined because $\mathcal{D}'_{E_u^*}$ is closed under $(d\delta + \delta d + 1)^{-1}$:
 $u \in \mathcal{D}'_{E_u^*}, du \in C^\infty \implies u = v + dw$ for some $v \in C^\infty, w \in \mathcal{D}'_{E_u^*}$

Lemma: π_1 is an isomorphism

Injectivity: if $u \in \text{Res}_0^1$ and $u = df$, $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$, then $Xf = \iota_X u = 0$, so $f \in \text{Res}_0^0 = \mathbb{R}1$ and $u = df = 0$

Surjectivity: if $v \in C^\infty(M; \Omega^1)$ and $dv = 0$, then $\int_M (\iota_X v) d\text{vol}_\alpha = 0$, so by the Fredholm property there exists $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$ with $Xf = \iota_X v$. Take $u := v - df \in \text{Res}_0^1 \cap \ker d$, then $\pi_1(u) = [v]_{H^1}$

Closed resonant 1-forms

- $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, \iota_X du = 0\}$, $\Omega^k = \wedge^k T^*M$
- **Closed forms:** $\text{Res}_0^k \cap \ker d = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, du = 0\}$
- **Cohomology map:** $\pi_k : \text{Res}_0^k \cap \ker d \rightarrow H^k(M; \mathbb{R})$, $\pi_k(u) = [u]_{H^k}$
- π_k can be defined because $\mathcal{D}'_{E_u^*}$ is closed under $(d\delta + \delta d + 1)^{-1}$:
 $u \in \mathcal{D}'_{E_u^*}, du \in C^\infty \implies u = v + dw$ for some $v \in C^\infty, w \in \mathcal{D}'_{E_u^*}$

Lemma: π_1 is an isomorphism

Injectivity: if $u \in \text{Res}_0^1$ and $u = df$, $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$, then
 $Xf = \iota_X u = 0$, so $f \in \text{Res}_0^0 = \mathbb{R}1$ and $u = df = 0$

Surjectivity: if $v \in C^\infty(M; \Omega^1)$ and $dv = 0$, then $\int_M (\iota_X v) d\text{vol}_\alpha = 0$,
 so by the Fredholm property there exists $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$ with $Xf = \iota_X v$.
 Take $u := v - df \in \text{Res}_0^1 \cap \ker d$, then $\pi_1(u) = [v]_{H^1}$

Closed resonant 1-forms

- $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, \iota_X du = 0\}$, $\Omega^k = \wedge^k T^*M$
- **Closed forms:** $\text{Res}_0^k \cap \ker d = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, du = 0\}$
- **Cohomology map:** $\pi_k : \text{Res}_0^k \cap \ker d \rightarrow H^k(M; \mathbb{R})$, $\pi_k(u) = [u]_{H^k}$
- π_k can be defined because $\mathcal{D}'_{E_u^*}$ is closed under $(d\delta + \delta d + 1)^{-1}$:
 $u \in \mathcal{D}'_{E_u^*}, du \in C^\infty \implies u = v + dw$ for some $v \in C^\infty, w \in \mathcal{D}'_{E_u^*}$

Lemma: π_1 is an isomorphism

Injectivity: if $u \in \text{Res}_0^1$ and $u = df$, $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$, then $Xf = \iota_X u = 0$, so $f \in \text{Res}_0^0 = \mathbb{R}1$ and $u = df = 0$

Surjectivity: if $v \in C^\infty(M; \Omega^1)$ and $dv = 0$, then $\int_M (\iota_X v) d\text{vol}_\alpha = 0$, so by the Fredholm property there exists $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$ with $Xf = \iota_X v$. Take $u := v - df \in \text{Res}_0^1 \cap \ker d$, then $\pi_1(u) = [v]_{H^1}$

Closed resonant 1-forms

- $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, \iota_X du = 0\}$, $\Omega^k = \wedge^k T^*M$
- **Closed forms:** $\text{Res}_0^k \cap \ker d = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, du = 0\}$
- **Cohomology map:** $\pi_k : \text{Res}_0^k \cap \ker d \rightarrow H^k(M; \mathbb{R})$, $\pi_k(u) = [u]_{H^k}$
- π_k can be defined because $\mathcal{D}'_{E_u^*}$ is closed under $(d\delta + \delta d + 1)^{-1}$:
 $u \in \mathcal{D}'_{E_u^*}, du \in C^\infty \implies u = v + dw$ for some $v \in C^\infty, w \in \mathcal{D}'_{E_u^*}$

Lemma: π_1 is an isomorphism

Injectivity: if $u \in \text{Res}_0^1$ and $u = df$, $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$, then $Xf = \iota_X u = 0$, so $f \in \text{Res}_0^0 = \mathbb{R}\mathbf{1}$ and $u = df = 0$

Surjectivity: if $v \in C^\infty(M; \Omega^1)$ and $dv = 0$, then $\int_M (\iota_X v) d\text{vol}_\alpha = 0$, so by the Fredholm property there exists $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$ with $Xf = \iota_X v$. Take $u := v - df \in \text{Res}_0^1 \cap \ker d$, then $\pi_1(u) = [v]_{H^1}$

Closed resonant 1-forms

- $\text{Res}_0^k = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, \iota_X du = 0\}$, $\Omega^k = \wedge^k T^*M$
- **Closed forms:** $\text{Res}_0^k \cap \ker d = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega^k) \mid \iota_X u = 0, du = 0\}$
- **Cohomology map:** $\pi_k : \text{Res}_0^k \cap \ker d \rightarrow H^k(M; \mathbb{R})$, $\pi_k(u) = [u]_{H^k}$
- π_k can be defined because $\mathcal{D}'_{E_u^*}$ is closed under $(d\delta + \delta d + 1)^{-1}$:
 $u \in \mathcal{D}'_{E_u^*}, du \in C^\infty \implies u = v + dw$ for some $v \in C^\infty, w \in \mathcal{D}'_{E_u^*}$

Lemma: π_1 is an isomorphism

Injectivity: if $u \in \text{Res}_0^1$ and $u = df$, $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$, then $Xf = \iota_X u = 0$, so $f \in \text{Res}_0^0 = \mathbb{R}\mathbf{1}$ and $u = df = 0$

Surjectivity: if $v \in C^\infty(M; \Omega^1)$ and $dv = 0$, then $\int_M (\iota_X v) d\text{vol}_\alpha = 0$, so by the Fredholm property there exists $f \in \mathcal{D}'_{E_u^*}(M; \mathbb{R})$ with $Xf = \iota_X v$. Take $u := v - df \in \text{Res}_0^1 \cap \ker d$, then $\pi_1(u) = [v]_{H^1}$

Resonant forms, hyperbolic case

- We know that $\mathcal{C} := \text{Res}_0^1 \cap \ker d$ has dimension $b_1(M) = b_1(\Sigma)$
- We show every $u \in \text{Res}_0^1$ is a section of $E_u^* = (E_0 \oplus E_u)^\perp \subset \Omega_0^1$
- The $\frac{\pi}{2}$ -rotation $\mathcal{I} : E_u^* \rightarrow E_u^*$ commutes with \mathcal{L}_X because the flow $\varphi_t = e^{tX}$ is conformal on E_u^* : $|d\varphi_t(\rho)^{-T}\xi| = e^t|\xi|$, $\xi \in E_u^*(\rho)$
- Thus \mathcal{I} acts on $\text{Res}_0^1 = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^1) \mid \mathcal{L}_X u = 0\}$
- If $u \in \mathcal{C} \setminus \{0\}$ then $d\mathcal{I}(u) \neq 0$: express $[d\alpha \wedge \mathcal{I}(u)]_{H^3}$ via $\pi_1(u)$
- We show that $\text{Res}_0^1 = \mathcal{C} \oplus \mathcal{I}(\mathcal{C})$ is $2b_1(\Sigma)$ -dimensional and semisimplicity holds for $k = 1$, so $\dim \text{Res}_0^{1,\infty} = 2b_1(\Sigma)$
- We also show that $\text{Res}_0^2 = \text{Res}_0^2 \cap \ker d = \mathbb{R}d\alpha \oplus \mathbb{R}\psi \oplus d \text{Res}_0^1$ is $(b_1(\Sigma) + 2)$ -dimensional where ψ is an explicit smooth 2-form
- We finally show $\dim \text{Res}_0^{2,\infty} = 2b_1(\Sigma) + 2$: get $b_1(\Sigma)$ Jordan blocks

Resonant forms, hyperbolic case

- We know that $\mathcal{C} := \text{Res}_0^1 \cap \ker d$ has dimension $b_1(M) = b_1(\Sigma)$
- We show every $u \in \text{Res}_0^1$ is a section of $E_u^* = (E_0 \oplus E_u)^\perp \subset \Omega_0^1$
- The $\frac{\pi}{2}$ -rotation $\mathcal{I} : E_u^* \rightarrow E_u^*$ commutes with \mathcal{L}_X because the flow $\varphi_t = e^{tX}$ is conformal on E_u^* : $|d\varphi_t(\rho)^{-T}\xi| = e^t|\xi|$, $\xi \in E_u^*(\rho)$
- Thus \mathcal{I} acts on $\text{Res}_0^1 = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^1) \mid \mathcal{L}_X u = 0\}$
- If $u \in \mathcal{C} \setminus \{0\}$ then $d\mathcal{I}(u) \neq 0$: express $[d\alpha \wedge \mathcal{I}(u)]_{H^3}$ via $\pi_1(u)$
- We show that $\text{Res}_0^1 = \mathcal{C} \oplus \mathcal{I}(\mathcal{C})$ is $2b_1(\Sigma)$ -dimensional and semisimplicity holds for $k = 1$, so $\dim \text{Res}_0^{1,\infty} = 2b_1(\Sigma)$
- We also show that $\text{Res}_0^2 = \text{Res}_0^2 \cap \ker d = \mathbb{R}d\alpha \oplus \mathbb{R}\psi \oplus d \text{Res}_0^1$ is $(b_1(\Sigma) + 2)$ -dimensional where ψ is an explicit smooth 2-form
- We finally show $\dim \text{Res}_0^{2,\infty} = 2b_1(\Sigma) + 2$: get $b_1(\Sigma)$ Jordan blocks

Resonant forms, hyperbolic case

- We know that $\mathcal{C} := \text{Res}_0^1 \cap \ker d$ has dimension $b_1(M) = b_1(\Sigma)$
- We show every $u \in \text{Res}_0^1$ is a section of $E_u^* = (E_0 \oplus E_u)^\perp \subset \Omega_0^1$
- The $\frac{\pi}{2}$ -rotation $\mathcal{I} : E_u^* \rightarrow E_u^*$ commutes with \mathcal{L}_X because the flow $\varphi_t = e^{tX}$ is conformal on E_u^* : $|d\varphi_t(\rho)^{-T}\xi| = e^t|\xi|$, $\xi \in E_u^*(\rho)$
- Thus \mathcal{I} acts on $\text{Res}_0^1 = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^1) \mid \mathcal{L}_X u = 0\}$
- If $u \in \mathcal{C} \setminus \{0\}$ then $d\mathcal{I}(u) \neq 0$: express $[d\alpha \wedge \mathcal{I}(u)]_{H^3}$ via $\pi_1(u)$
- We show that $\text{Res}_0^1 = \mathcal{C} \oplus \mathcal{I}(\mathcal{C})$ is $2b_1(\Sigma)$ -dimensional and semisimplicity holds for $k = 1$, so $\dim \text{Res}_0^{1,\infty} = 2b_1(\Sigma)$
- We also show that $\text{Res}_0^2 = \text{Res}_0^2 \cap \ker d = \mathbb{R}d\alpha \oplus \mathbb{R}\psi \oplus d \text{Res}_0^1$ is $(b_1(\Sigma) + 2)$ -dimensional where ψ is an explicit smooth 2-form
- We finally show $\dim \text{Res}_0^{2,\infty} = 2b_1(\Sigma) + 2$: get $b_1(\Sigma)$ Jordan blocks

Resonant forms, hyperbolic case

- We know that $\mathcal{C} := \text{Res}_0^1 \cap \ker d$ has dimension $b_1(M) = b_1(\Sigma)$
- We show every $u \in \text{Res}_0^1$ is a section of $E_u^* = (E_0 \oplus E_u)^\perp \subset \Omega_0^1$
- The $\frac{\pi}{2}$ -rotation $\mathcal{I} : E_u^* \rightarrow E_u^*$ commutes with \mathcal{L}_X because the flow $\varphi_t = e^{tX}$ is conformal on E_u^* : $|d\varphi_t(\rho)^{-T}\xi| = e^t|\xi|$, $\xi \in E_u^*(\rho)$
- Thus \mathcal{I} acts on $\text{Res}_0^1 = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^1) \mid \mathcal{L}_X u = 0\}$
- If $u \in \mathcal{C} \setminus \{0\}$ then $d\mathcal{I}(u) \neq 0$: express $[d\alpha \wedge \mathcal{I}(u)]_{H^3}$ via $\pi_1(u)$
- We show that $\text{Res}_0^1 = \mathcal{C} \oplus \mathcal{I}(\mathcal{C})$ is $2b_1(\Sigma)$ -dimensional and semisimplicity holds for $k = 1$, so $\dim \text{Res}_0^{1,\infty} = 2b_1(\Sigma)$
- We also show that $\text{Res}_0^2 = \text{Res}_0^2 \cap \ker d = \mathbb{R}d\alpha \oplus \mathbb{R}\psi \oplus d \text{Res}_0^1$ is $(b_1(\Sigma) + 2)$ -dimensional where ψ is an explicit smooth 2-form
- We finally show $\dim \text{Res}_0^{2,\infty} = 2b_1(\Sigma) + 2$: get $b_1(\Sigma)$ Jordan blocks

Resonant forms, hyperbolic case

- We know that $\mathcal{C} := \text{Res}_0^1 \cap \ker d$ has dimension $b_1(M) = b_1(\Sigma)$
- We show every $u \in \text{Res}_0^1$ is a section of $E_u^* = (E_0 \oplus E_u)^\perp \subset \Omega_0^1$
- The $\frac{\pi}{2}$ -rotation $\mathcal{I} : E_u^* \rightarrow E_u^*$ commutes with \mathcal{L}_X because the flow $\varphi_t = e^{tX}$ is conformal on E_u^* : $|d\varphi_t(\rho)^{-T}\xi| = e^t|\xi|$, $\xi \in E_u^*(\rho)$
- Thus \mathcal{I} acts on $\text{Res}_0^1 = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^1) \mid \mathcal{L}_X u = 0\}$
- If $u \in \mathcal{C} \setminus \{0\}$ then $d\mathcal{I}(u) \neq 0$: express $[d\alpha \wedge \mathcal{I}(u)]_{H^3}$ via $\pi_1(u)$
- We show that $\text{Res}_0^1 = \mathcal{C} \oplus \mathcal{I}(\mathcal{C})$ is $2b_1(\Sigma)$ -dimensional and semisimplicity holds for $k = 1$, so $\dim \text{Res}_0^{1,\infty} = 2b_1(\Sigma)$
- We also show that $\text{Res}_0^2 = \text{Res}_0^2 \cap \ker d = \mathbb{R}d\alpha \oplus \mathbb{R}\psi \oplus d \text{Res}_0^1$ is $(b_1(\Sigma) + 2)$ -dimensional where ψ is an explicit smooth 2-form
- We finally show $\dim \text{Res}_0^{2,\infty} = 2b_1(\Sigma) + 2$: get $b_1(\Sigma)$ Jordan blocks

Resonant forms, hyperbolic case

- We know that $\mathcal{C} := \text{Res}_0^1 \cap \ker d$ has dimension $b_1(M) = b_1(\Sigma)$
- We show every $u \in \text{Res}_0^1$ is a section of $E_u^* = (E_0 \oplus E_u)^\perp \subset \Omega_0^1$
- The $\frac{\pi}{2}$ -rotation $\mathcal{I} : E_u^* \rightarrow E_u^*$ commutes with \mathcal{L}_X because the flow $\varphi_t = e^{tX}$ is conformal on E_u^* : $|d\varphi_t(\rho)^{-T}\xi| = e^t|\xi|$, $\xi \in E_u^*(\rho)$
- Thus \mathcal{I} acts on $\text{Res}_0^1 = \{u \in \mathcal{D}'_{E_u^*}(M; \Omega_0^1) \mid \mathcal{L}_X u = 0\}$
- If $u \in \mathcal{C} \setminus \{0\}$ then $d\mathcal{I}(u) \neq 0$: express $[d\alpha \wedge \mathcal{I}(u)]_{H^3}$ via $\pi_1(u)$
- We show that $\text{Res}_0^1 = \mathcal{C} \oplus \mathcal{I}(\mathcal{C})$ is $2b_1(\Sigma)$ -dimensional and semisimplicity holds for $k = 1$, so $\dim \text{Res}_0^{1,\infty} = 2b_1(\Sigma)$
- We also show that $\text{Res}_0^2 = \text{Res}_0^2 \cap \ker d = \mathbb{R}d\alpha \oplus \mathbb{R}\psi \oplus d \text{Res}_0^1$ is $(b_1(\Sigma) + 2)$ -dimensional where ψ is an explicit smooth 2-form
- We finally show $\dim \text{Res}_0^{2,\infty} = 2b_1(\Sigma) + 2$: get $b_1(\Sigma)$ Jordan blocks

Resonant forms for perturbations

- Consider now the **perturbed metric** $g_\tau = e^{\tau a} g_H$, $a \in C^\infty(\Sigma; \mathbb{R})$
- Define $\pi_\Sigma : M = S\Sigma \rightarrow \Sigma$; $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$
- We still have $\dim(\text{Res}_0^1 \cap \ker d) = b_1(\Sigma)$, need to show that all non-closed elements of Res_0^1 are moved by the perturbation
- A first variation calculation shows that we need nondegeneracy of

$$du \in d(\text{Res}_0^1), \quad du_* \in d(\text{Res}_{0*}^1) \quad \mapsto \quad \int_M (\pi_{\Sigma*}^* a) \alpha \wedge du \wedge du_*$$

- Take for simplicity $b_1(\Sigma) = 1$, then enough to show

$$u \in \text{Res}_0^1, \quad du \neq 0 \quad \implies \quad \int_M (\pi_{\Sigma*}^* a) \alpha \wedge du \wedge \mathcal{J}^*(du) \neq 0$$

- That's true for generic a as long as $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) \neq 0$ where $\pi_{\Sigma*} : \mathcal{D}'(M; \Omega^k) \rightarrow \mathcal{D}'(\Sigma; \Omega^{k-2})$ is the pushforward on forms

Resonant forms for perturbations

- Consider now the **perturbed metric** $g_\tau = e^{\tau a} g_H$, $a \in C^\infty(\Sigma; \mathbb{R})$
- Define $\pi_\Sigma : M = S\Sigma \rightarrow \Sigma$; $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$
- We still have $\dim(\text{Res}_0^1 \cap \ker d) = b_1(\Sigma)$, need to show that all non-closed elements of Res_0^1 are moved by the perturbation
- A first variation calculation shows that we need nondegeneracy of

$$du \in d(\text{Res}_0^1), \quad du_* \in d(\text{Res}_{0*}^1) \quad \mapsto \quad \int_M (\pi_{\Sigma*}^* a) \alpha \wedge du \wedge du_*$$

- Take for simplicity $b_1(\Sigma) = 1$, then enough to show

$$u \in \text{Res}_0^1, \quad du \neq 0 \quad \implies \quad \int_M (\pi_{\Sigma*}^* a) \alpha \wedge du \wedge \mathcal{J}^*(du) \neq 0$$

- That's true for generic a as long as $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) \neq 0$ where $\pi_{\Sigma*} : \mathcal{D}'(M; \Omega^k) \rightarrow \mathcal{D}'(\Sigma; \Omega^{k-2})$ is the pushforward on forms

Resonant forms for perturbations

- Consider now the **perturbed metric** $g_\tau = e^{\tau a} g_H$, $a \in C^\infty(\Sigma; \mathbb{R})$
- Define $\pi_\Sigma : M = S\Sigma \rightarrow \Sigma$; $\mathcal{J} : M \rightarrow M$, $\mathcal{J}(x, v) = (x, -v)$
- We still have $\dim(\text{Res}_0^1 \cap \ker d) = b_1(\Sigma)$, need to show that all non-closed elements of Res_0^1 are moved by the perturbation
- A first variation calculation shows that we need nondegeneracy of

$$du \in d(\text{Res}_0^1), \quad du_* \in d(\text{Res}_{0*}^1) \quad \mapsto \quad \int_M (\pi_{\Sigma^*}^* a) \alpha \wedge du \wedge du_*$$

- Take for simplicity $b_1(\Sigma) = 1$, then enough to show

$$u \in \text{Res}_0^1, \quad du \neq 0 \quad \implies \quad \int_M (\pi_{\Sigma^*}^* a) \alpha \wedge du \wedge \mathcal{J}^*(du) \neq 0$$

- That's true for generic a as long as $\pi_{\Sigma^*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) \neq 0$ where $\pi_{\Sigma^*} : \mathcal{D}'(M; \Omega^k) \rightarrow \mathcal{D}'(\Sigma; \Omega^{k-2})$ is the pushforward on forms

Nontriviality of first variation

- Working only with the hyperbolic metric now
- Given $u \in \text{Res}_0^1$, $du \neq 0$, need $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) \neq 0$
- Write $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) = F d \text{vol}_g$ for some $F \in \mathcal{D}'(\Sigma; \mathbb{R})$
- Difficult to show that $F \neq 0$ because cannot evaluate F at points

Main identity

We have $Q_4 F = -\frac{1}{6} \Delta_g |\sigma|_g^2$ where

- $\sigma = \pi_{\Sigma*}(\alpha \wedge du)$ is a nonzero harmonic 1-form on Σ
- $Q_4 f(x) = \int_{\mathbb{H}^3} \cosh^{-4} d_{\mathbb{H}^3}(x, y) f(y) d \text{vol}_g(y)$ descends to $Q_4 : \mathcal{D}'(\Sigma) \rightarrow C^\infty(\Sigma)$ where $\Sigma = \Gamma \backslash \mathbb{H}^3$
- If $F = 0$, then $\Delta_g |\sigma|_g^2 = 0$, so $|\sigma|_g$ is constant, but this is impossible!

Nontriviality of first variation

- Working only with the hyperbolic metric now
- Given $u \in \text{Res}_0^1$, $du \neq 0$, need $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) \neq 0$
- Write $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) = F d \text{vol}_g$ for some $F \in \mathcal{D}'(\Sigma; \mathbb{R})$
- Difficult to show that $F \neq 0$ because cannot evaluate F at points

Main identity

We have $Q_4 F = -\frac{1}{6} \Delta_g |\sigma|_g^2$ where

- $\sigma = \pi_{\Sigma*}(\alpha \wedge du)$ is a nonzero harmonic 1-form on Σ
- $Q_4 f(x) = \int_{\mathbb{H}^3} \cosh^{-4} d_{\mathbb{H}^3}(x, y) f(y) d \text{vol}_g(y)$ descends to $Q_4 : \mathcal{D}'(\Sigma) \rightarrow C^\infty(\Sigma)$ where $\Sigma = \Gamma \backslash \mathbb{H}^3$

- If $F = 0$, then $\Delta_g |\sigma|_g^2 = 0$, so $|\sigma|_g$ is constant, but this is impossible!

Nontriviality of first variation

- Working only with the hyperbolic metric now
- Given $u \in \text{Res}_0^1$, $du \neq 0$, need $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) \neq 0$
- Write $\pi_{\Sigma*}(\alpha \wedge du \wedge \mathcal{J}^*(du)) = F d \text{vol}_g$ for some $F \in \mathcal{D}'(\Sigma; \mathbb{R})$
- Difficult to show that $F \neq 0$ because cannot evaluate F at points

Main identity

We have $Q_4 F = -\frac{1}{6} \Delta_g |\sigma|_g^2$ where

- $\sigma = \pi_{\Sigma*}(\alpha \wedge du)$ is a nonzero harmonic 1-form on Σ
- $Q_4 f(x) = \int_{\mathbb{H}^3} \cosh^{-4} d_{\mathbb{H}^3}(x, y) f(y) d \text{vol}_g(y)$ descends to $Q_4 : \mathcal{D}'(\Sigma) \rightarrow C^\infty(\Sigma)$ where $\Sigma = \Gamma \backslash \mathbb{H}^3$
- If $F = 0$, then $\Delta_g |\sigma|_g^2 = 0$, so $|\sigma|_g$ is constant, but this is impossible!

Two conjectures

Conjecture 1

Let (Σ, g) be a **generic** negatively curved compact connected oriented 3-manifold. Then:

- semisimplicity holds and $d(\text{Res}_0^k) = 0$ for all $k = 0, \dots, 4$
- $\dim \text{Res}_0^0 = 1$, $\dim \text{Res}_0^1 = b_1(\Sigma)$, $\dim \text{Res}_0^2 = b_1(\Sigma) + 2$
- $m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$

The set of g satisfying Conjecture 1 is open:

$$\dim \text{Res}_0^{1,\infty} \leq b_1(\Sigma), \dim \text{Res}_0^{2,\infty} \leq b_1(\Sigma) + 2 \implies \text{Conjecture 1 holds}$$

Conjecture 2

Let $\rho : \pi_1(\Sigma) \rightarrow U(m)$ be acyclic: $H_\rho^*(\Sigma; \mathbb{R}) = 0$. Then $\text{Res}_0^k = 0$ for all k

Conjecture 2 + **DGRS** '20 $\implies \zeta_\rho(0)$ is locally constant under perturbations of ρ, g , which could lead to a solution of Fried's conjecture

Two conjectures

Conjecture 1

Let (Σ, g) be a **generic** negatively curved compact connected oriented 3-manifold. Then:

- semisimplicity holds and $d(\text{Res}_0^k) = 0$ for all $k = 0, \dots, 4$
- $\dim \text{Res}_0^0 = 1$, $\dim \text{Res}_0^1 = b_1(\Sigma)$, $\dim \text{Res}_0^2 = b_1(\Sigma) + 2$
- $m_{\mathbb{R}}(0) = 4 - b_1(\Sigma)$

The set of g satisfying Conjecture 1 is open:

$$\dim \text{Res}_0^{1,\infty} \leq b_1(\Sigma), \dim \text{Res}_0^{2,\infty} \leq b_1(\Sigma) + 2 \implies \text{Conjecture 1 holds}$$

Conjecture 2

Let $\rho : \pi_1(\Sigma) \rightarrow U(m)$ be acyclic: $H_\bullet^\rho(\Sigma; \mathbb{R}) = 0$. Then $\text{Res}_0^k = 0$ for all k

Conjecture 2 + **DGRS** '20 $\implies \zeta_\rho(0)$ is locally constant under perturbations of ρ, g , which could lead to a solution of Fried's conjecture

Thank you for your attention!