BOUNCING BALL MODES AND QUANTUM CHAOS

NICOLAS BURQ AND MACIEJ ZWORSKI

ABSTRACT. Quantum ergodicity for classically chaotic systems has been studied extensively both theoretically and experimentally in mathematics and in physics. Despite this long tradition we are able to prove a new rigorous result using only elementary calculus. In the case of the famous Bunimovich stadium shown in Fig.1 we prove that the wave functions have to spread to any neighbourhood of the wings.

The quantum/classical correspondence is a puzzling issue that has been with us since the advent of quantum mechanics hundred years ago. Many aspects of it go back to the Newton/Huyghens debate over the wave vs. corpuscular theories of light.

The simplest description of a wave comes from solving the Helmholtz equation:

$$(-\Delta - \lambda^2)u = 0\,, \quad \Delta = \partial_x^2 + \partial_y^2\,, \quad (x,y) \in \Omega\,, \quad u\restriction_{\partial\Omega} = 0\,.$$

Here we put our wave inside a two dimensional region Ω. In classical wave mechanics the limit $\lambda \to \infty$ is described using geometrical optics where the waves propagate along straight lines reflecting in the boundary $\partial\Omega$. Roughly speaking, we expect something similar in the classical/quantum correspondence with the Helmholtz equation replaced by its quantum mechanical version, the Schrödinger equation. For many fascinating illustrations of this we refer to the web art gallery of the Harvard physicist Eric Heller [10].



FIGURE 1. An experimental image of bouncing ball modes in a Bunimovich stadium cavity – see [5] and `http://www.bath.ac.uk/~pyscmd/acoustics`. The bouncing ball modes spread all the way to the boundaries of the rectangular part of the billiard which is consistent with the results of this paper.

Many researchers on different aspects of semiclassical analysis have been interested in the correspondence of solutions to the equation above and the classical geometry of balls bouncing from the walls of Ω: Bäcker, Cvitanović, Eckhardt, Gaspard, Heller, Sridhar, in physics, and

Colin de Verdière, Melrose, Sjöstrand, Zelditch, in mathematics, to mention some (see [1] for references to the physics literature, and [15] for mathematics).

Billiard tables for which the motion is chaotic are a particularly interesting model to study. One of the most famous is the Bunimovich billiard table shown in Fig.1. By adding two circular "wings" to a rectangular table the motion of a reflecting billiard ball becomes chaotic, or more precisely, hyperbolic, in the sense that changes in initial conditions lead to exponentially large changes in motion as time goes on. It is also ergodic in the sense that any flow invariant subset of the phase space has either full measure or zero measure.

As a model for studying quantum phenomena in chaotic systems this billiard table has become popular in experimental physics. A genuinely quantum example is shown in Fig.2 – it comes from the scanning tunnelling microscope work of Crommie, Lutz and Eigler [6].



FIGURE 2. Quantum corral in the shape of the Bunimovich stadium. Courtesy of IBM® Research.

The basic mathematical result in the theory of quantum chaos is the so called

Shnirelman Theorem. *Suppose that the billiard flow on a bounded domain, Ω, with boundary, $\partial\Omega$, is ergodic. Let u_j be the sequence of normalized eigenfunctions of the Dirichlet (or Neumann) Laplacian,*

$$-\Delta u_j = \lambda_j^2 u_j \,, \quad u_j\restriction_{\partial\Omega} = 0 \,, \quad \int_\Omega |u_j(x)|^2 dx = 1 \,. \tag{1}$$

Then there exists a sequence $\{j_k\}_{k=1}^\infty \subset \mathbb{N}$ of density one, that is, $\lim_{k\to\infty} j_k/k = 1$, such that for any open subset V, of Ω,

$$\lim_{k\to\infty} \int_V |u_{j_k}(x)|^2 dx = \frac{\text{Area}(V)}{\text{Area}(\Omega)} \,. \tag{2}$$

This means that for almost all eigenfunctions there cannot be any concentration: they have to be uniformly spread out in the billiard table. The integral of the square of the eigenfuction over V is interpreted as the probability of finding the quantum state in V. A stronger version of the theorem gives a phase space version of this statement.

The Shnirelman Theorem for billiards was first proved for convex billiards (in particular the Bunimovich billiard) by Gérard-Leichtnam [9], and for arbitrary manifolds with piecewise smooth boundaries by Zelditch-Zworski [16]. We refer to these papers and to [1],[8],[15] for history and pointers to the literature.

One question which is still mysterious to mathematicians and physicists alike is if the quantum states of a classically ergodic system (in our case, solutions of the Helmholtz equation for an ergodic billiard) can concentrate on the highly unstable closed orbits of the classical flow, or on some invariant tori formed by such orbits. The Shnirelman theorem allows the possibility of such concentration on sequences of density zero.

A system is called *quantum unique ergodic* if there is no such concentration – see [15] and references given there. In particular, quantum unique ergodicity means that (2) holds for the *full* sequence of eigenfuctions, that is

$$\lim_{j\to\infty}\int_V |u_j(x)|^2 dx = \frac{\text{Area}(V)}{\text{Area}(\Omega)}\,. \tag{3}$$

In the arithmetic† case investigated and popularized by Sarnak [14], spectacular advances have been recently achieved by Lindenstrauss [12]. On the other hand for the quantization of the *Arnold cat map* results showing the *lack* of quantum unique ergodicity were produced by Bonechi, De Bièvre, Faure, and Nonnenmacher [7].

Here we prove an elementary but striking result which shows that no sequence of eigenfunctions of the Bunimovitch stadium can concentrate on a single bouncing ball orbit, or indeed on any smaller phase space set than the union of the bouncing ball orbits of the entire the rectangle. It follows from adapting the first author's earlier work in control theory. Although motivated by the more general aspects of [2] we give a simple self-contained proof. Without estimates, which will be given in Theorem′ below, it can be stated as

Theorem. *Let Ω be the Bunimovich billiard table, $\Omega = R \cup W$ where R is the rectangular part, and W the circular wings. Let V be any open neighbourhood of the closure of W in Ω. With the notation of* (1) *we have*

$$\liminf_{j\to\infty}\int_V |u_j(x)|^2 dx > \frac{1}{C_V}\,, \tag{4}$$

where $C_V > 0$ depends only on V.

In particular, the result says that single bouncing ball orbits (that is orbits following an interval perpendicular to the horizontal straight boundaries) cannot produce localized waves. Our result allows concentration on the full invariant set of all vertical orbits over R – that is consistent with the existing physical literature – both numerical and experimental – see [1] and [5]. In fact, it is expected that there exists a density zero subsequence, $\{p_k\}_{k=1}^{\infty}$, for which

$$\lim_{k\to\infty}\int_R |u_{p_k}(x)|^2 dx = 1 > \frac{\text{Area}(R)}{\text{Area}(\Omega)}\,.$$

Establishing this would constitute a major breakthrough.

In [2] we show a stronger result than (4), namely that the neighbourhood of the wings can be replaced by any neighbourhood of the vertical intervals between the wings and the rectangular part. The proof of that predictable (to experts) improvement is however no longer elementary.

The proof of theorem depends on the following unpublished result of the first author (see [2] for detailed references and background material):

---

†That is, for billiards given by arithmetic surfaces with hyperbolic metrics: the motion is given by the geodesic flow and the quantum Hamiltonian is the Laplace-Beltrami operator.

Proposition. *Let $\Delta = \partial_x^2 + \partial_y^2$ be the Laplace operator on the rectangle $R = [0,1]_x \times [0,a]_y$. Then for any open $\omega \subset R$ of the form $\omega = \omega_x \times [0,a]_y$, there exists C such that for any solution of*

$$(-\Delta - \lambda^2)u = f + \partial_x g \quad \text{on } R , \quad u\restriction_{\partial R} = 0 \, ,$$

with an arbitrary $\lambda \geq 0$ we have

$$\int_R |u(x,y)|^2 dx dy \leq C \left(\int_R (|f(x,y)|^2 + |g(x,y)|^2) dx dy + \int_\omega |u(x,y)|^2 dx dy \right) . \tag{5}$$

Here, f and g are assumed to be square integrable.

Proof. We decompose u, and $f + \partial_x g$ in terms of the basis of $L^2([0,a])$ formed by the Dirichlet eigenfunctions $e_k(y) = \sqrt{2/a}\sin(2k\pi y/a)$,

$$u(x,y) = \sum_k e_k(y) u_k(x), \qquad f(x,y) + \partial_x g(x,y) = \sum_k e_k(y)(f_k(x) + \partial_x g_k(x)) \tag{6}$$

we get for u_k, f_k the equation

$$-\left(\partial_x^2 + z\right) u_k = f_k + \partial_x g_k, \qquad u_k(0) = u_k(1) = 0 \, , \quad z = \lambda^2 - (2k\pi/a)^2 \, .$$

It is now easy to see that

$$\int_0^1 |u_k(x)|^2 dx \leq C \left(\int_0^1 (|f_k(x)|^2 + |g_k(x)|^2) dx + \int_{\omega_x} |u_k(x)|^2 dx \right) , \tag{7}$$

where C is independent of z. In fact, let us first assume that $\omega_x = (0,\delta)$, $\delta > 0$, and $z = \lambda_1^2$, with $\mathrm{Im}\lambda_1 \leq 1$. We then choose $\chi \in \mathcal{C}_c^\infty([0,1])$ identically zero near 0 and identically one on $[\delta/2, 1]$. Then

$$\left(\partial_x^2 + \lambda_1^2\right)(\chi u_k) = F_k \, , \quad F_k = -\left(\chi(f_k + \partial_x g_k) + 2\partial_x \chi \partial_x u_k + \partial_x^2 \chi u_k\right) \, .$$

We can now use the explicit solution given by

$$\chi(x) u_k(x) = \frac{1}{\lambda_1} \int_0^x \sin(\lambda_1(x-y)) F_k(y) dy \, .$$

All the terms with $\partial_x g_k$ and $\partial_x u_k$ can be converted to g_k and u_k by integration by parts (with boundary terms 0 at both ends). Due to the λ_1^{-1} factor that produces no loss and the estimate follows. The argument is symmetric under the $x \mapsto -x$ change, so we can place our control interval anywhere.

It remains to discuss the case $z \leq -1 < 0$. Then the estimate (7) follows from integration by parts (where now we do not need ω_x):

$$\int_0^1 \left(f(x)\overline{u(x)} - g(x)\overline{\partial_x u(x)} \right) dx = \int_0^1 (f(x) + \partial_x g(x))\overline{u(x)} dx =$$

$$\int_0^1 (-\partial_x^2 - z) u(x)\overline{u(x)} dx = \int_0^1 \left(|\partial_x u(x)|^2 + |z||u(x)|^2 \right) dx \, .$$

But the Cauchy-Schwartz shows that the left hand side is bounded from above by

$$\left(\int_0^1 \left(|f(x)|^2 + |g(x)|^2 \right) dx \right)^{\frac{1}{2}} \left(\int_0^1 \left(|u(x)|^2 + |\partial_x u(x)|^2 \right) dx \right)^{\frac{1}{2}} .$$

Since $|z| > 1 > 0$, (7) follows from elementary inequalities (see [2, Lemma 4.1] for a general microlocal argument). We can now sum the estimate in k to obtain the proposition. $\square$

We can now present a more precise version of the theorem. For a yet finer version we refer the reader to [2, Theorem 3′] and [2, Fig.5].

Theorem′. *Consider Ω, the Bunimovich stadium constructed from a rectangle R and wings W. Let V be any open neighbourhood of $\overline{W}$ in Ω. There exists a constant C depending only on Ω and V such that for any solution of the equation*

$$(-\Delta-\lambda^2)v = F\,, \quad u\restriction_{\partial\Omega}= 0\,, \quad \lambda \geq 0\,,$$

we have

$$\int_\Omega |v(x,y)|^2 dxdy \leq C\left(\int_\Omega |F(x,y)|^2 dxdy + \int_V |v(x,y)|^2 dxdy\right).$$

We apply the second theorem with $F=0$ to obtain the first one.

Proof. Let us take x, y as the coordinates on the stadium, so that x is the horizontal direction, y the vertical direction, and the internal rectangle is $[0,1]_x \times [0,a]_y$. Let us then consider u and F satisfying $(-\Delta-\lambda^2)u = F$, $u=0$ on the boundary of the stadium, and $\chi(x) \in \mathcal{C}^\infty_{\mathrm{c}}(0,1)$ equal to 1 on $[\varepsilon, 1-\varepsilon]$. Then $\chi(x)u(x,y)$ is solution of

$$(-\Delta-\lambda^2)\chi u = \chi F + [\Delta,\chi]u \ \text{ in } R$$

with Dirichlet boundary conditions on ∂R. Since $[\Delta,\chi]u = 2\partial_x(\chi' u) - \chi'' u$ we can apply the proposition with $f = \chi F - \chi'' u$ and $g = 2\chi' u$ to obtain

$$\int_R |\chi(x)u(x,y)|^2 dxdy \leq C\left(\int_R |\chi(x)f(x,y)|^2 dxdy + \int_{\omega_\varepsilon} |u(x,y)|^2 dxdy\right)$$

where ω_ε is a neighbourhood of the support of $\partial_x\chi$. Since we can choose it to be contained in $R\cap V$, the theorem follows. □

The reader can notice that the structure of the wings did not play any rôle in the argument and hence the result holds for any billiard $\Omega = R \cup W$ as long $\partial\Omega$ contains a pair of parallel sides of R – see Fig.3.



FIGURE 3. A partially rectangular billiard

In any such billiard it is easy to construct a weak quasimode, that is an approximation to an eigenstate. Let again $R = [0,1]_x \times [0,a]_y$ with the sides parallel to the x axis contained in the boundary of $\Omega = R \cup W$. Consider $\phi \in \mathcal{C}^\infty_{\rm c}((0,1))$, a smooth one dimensional function vanishing near the boundaries of $[0,1]$, with the property that $\int_0^1 |\phi(x)|^2 dx = 1$. Then put

$$v_k(x,y) = e_k(y)\phi(x)\,, \quad e_k(y) = \sqrt{\frac{2}{a}} \sin\left(\frac{2k\pi y}{a}\right)\,,$$

so that

$$(-\Delta - (2\pi k/a)^2) v_k = -e_k(y)\phi''(x) = \mathcal{O}(1)\,, \quad k \to \infty\,. \tag{8}$$

A strong quasimode would be defined by (8) with $\mathcal{O}(1)$ replaced by $\mathcal{O}(k^{-N})$, $N \gg 1$. Theorem′ shows however that the trivial construction of v_k is the best possible: if $\mathcal{O}(1)$ in (8) were replaced by $o(1)$, as $k \to \infty$, we would obtain a contradiction with the estimate (5) once we took ω outside the support of ϕ. The same argument works in the settings of [8] and [15] where similar weak quasimodes were considered, since the rectangle can be replaced by a torus.

In [3] we show how similar methods imply control theory result of Jaffard [11] and some new results for Sinai billiards. They however require slightly more advanced tools.

Acknowledgments. The authors would like to thank the National Science Foundation for partial support under the grant DMS-0200732. They are also grateful to Arnd Bäcker, Stephan De Bièvre, Richard Melrose, and Steve Zelditch for helpful discussions and comments, and to Hilary Ockendon for her patient reading of the first version and numerous useful suggestions. The first author thanks the Mathematical Science Research Institute for its hospitality during spring 2003.

References

[1] A. Bäcker, R. Schubert, and P. Stifter. On the number of bouncing ball modes in billiards. *J. Phys. A: Math. Gen.* 30:6783-6795, 1997.
[2] N. Burq and M. Zworski. Control in the presence of a black box. `math.AP/0304184`, Journal of A.M.S., to appear.
[3] N. Burq and M. Zworski. Eigenfunctions for partially rectangular billiards, `math.SP/0312098` .
[4] C. Bardos, G. Lebeau and J. Rauch. Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary. *SIAM J. Control Optim.* 30:1024–1065, 1992.
[5] P.A. Chinnery and V.F. Humphrey. Experimental visualization of acoustic resonances within a stadium-shaped cavity. *Physical Review E*, 53, 1996, 272–276.
[6] M. F. Crommie, C. P. Lutz, and D. M. Eigler. Confinement of electrons to quantum corrals on a metal surface *Science* 262, 1993, 218–221.
[7] F. Faure, S. Nonnenmacher, and Stephan De Biévre. Scarred Eigenstates for Quantum Cat Maps of Minimal Periods, *Comm. Math. Phys.*, 239, 2003, 449–492.
[8] H. Donnelly. Quantum unique ergodicity *Proc. Amer. Math. Soc.* 131, 2945–2951, 2003.
[9] P. Gérard and E. Leichtnam. Ergodic Properties of Eigenfunctions for the Dirichlet Problem, *Duke Math. Jour.* 71, 559–607, 1993
[10] Eric J. Heller Gallery, `http://www.ericjhellergallery.com`.
[11] S. Jaffard. Contrôle interne exact des vibrations d'une plaque rectangulaire. *Portugal. Math.* 47 (1990), no. 4, 423-429.
[12] E. Lindenstrauss. Invariant measures and arithmetic quantum unique ergodicity, *preprint*, 2003
[13] R.B. Melrose and J. Sjöstrand. Singularities of Boundary Value Problems I & II, *Comm. in Pure Appl. Math.*, 31 & 35, 593- 617 & 129-168, 1978 & 1982.
[14] P. Sarnak. Arithmetic quantum chaos. In *The Schur lectures (1992) (Tel Aviv)*, volume 8 of *Israel Math. Conf. Proc.*, pages 183–236. Bar-Ilan Univ., Ramat Gan, 1995.
[15] S. Zelditch. A note on quantum unique ergodicity. *Proc. Amer. Math. Soc.* 2004, to appear.
[16] S. Zelditch and M. Zworski. Ergodicity of eigenfunctions for ergodic billiards. *Comm. Math. Phys.*, 175:673–682, 1996.

Université Paris Sud, Mathématiques, Bât 425, 91405 Orsay Cedex
E-mail address: Nicolas.burq@math.u-psud.fr

Mathematics Department, University of California, Evans Hall, Berkeley, CA 94720, USA
E-mail address: zworski@math.berkeley.edu