

# MATH 54 Lecture Notes 5

GSI Carter

June 30, 2006

## 1 Vector Space Axioms

Let  $V$  be a set on which addition and scalar multiplication are defined. Suppose all of the following conditions hold for any  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and all scalars  $r$  and  $s$ .

1.  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
2.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ .
3. There exists an element  $\mathbf{0} \in V$  such that  $\mathbf{u} + \mathbf{0} = \mathbf{u}$  for all  $\mathbf{u} \in V$ .
4. For every  $\mathbf{u} \in V$ , there exists  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
5.  $r(\mathbf{u} + \mathbf{v}) = r\mathbf{u} + r\mathbf{v}$ .
6.  $(r + s)\mathbf{u} = r\mathbf{u} + s\mathbf{u}$ .
7.  $(rs)\mathbf{u} = r(s\mathbf{u})$ .
8.  $1 \cdot \mathbf{u} = \mathbf{u}$ .

Then we say that  $V$  is a vector space. If the field of scalars is  $\mathbb{R}$ ,  $V$  is an  $\mathbb{R}$ -vector space. If the field of scalars is  $\mathbb{C}$ , then  $V$  is a  $\mathbb{C}$ -vector space. Every  $\mathbb{C}$ -vector space is also an  $\mathbb{R}$ -vector space.

## 2 Vector Space Examples

- $\mathbb{R}^n$  is an  $\mathbb{R}$ -vector space.  $\mathbb{C}^n$  is a  $\mathbb{C}$ -vector space.
- Spans. This includes lines, planes, and the zero vector space.
- Matrices,  $M_{mn}$ .
- Polynomials,  $P_n$ .
- Continuous functions,  $C[a, b]$ .

### 3 Vector Space Properties

- $0 \cdot \mathbf{u} = \mathbf{0}$ . Here  $\mathbf{0}$  means the additive identity.
- $r \cdot \mathbf{0} = \mathbf{0}$ .
- $(-1)\mathbf{u} = -\mathbf{u}$ . The thing on the left is scalar multiplication; the thing on the right is the additive inverse of  $\mathbf{u}$ .
- If  $r\mathbf{u} = \mathbf{0}$ , then either  $r = 0$  or  $\mathbf{u} = \mathbf{0}$ .