

YOUR NAME: Alexander Givental

Math 110. Midterm I. Feb. 9, 2009

1. Formulate and prove the “toy version” of the Orthogonal Diagonalization Theorem (about quadratic forms on the plane).

Solution. See page 11 of the text.

2. A regular triangle ABC is inscribed into a circle of radius R . Prove that for every point X of this circle, $XA^2 + XB^2 + XC^2 = 6R^2$.

Solution. Let O denote the center of the circle, Then $\vec{XA} = \vec{OA} - \vec{OX}$, and hence

$$XA^2 = \langle \vec{OA} - \vec{OX}, \vec{OA} - \vec{OX} \rangle = OA^2 + OX^2 - 2\langle \vec{OA}, \vec{OX} \rangle,$$

and similarly for XB and XC . Therefore

$$XA^2 + XB^2 + XC^2 = 3OX^2 + OA^2 + OB^2 + OC^2 - 2\langle \vec{OA} + \vec{OB} + \vec{OC}, \vec{OX} \rangle.$$

Since the triangle ABC is equilateral, we have $\vec{OA} + \vec{OB} = -\vec{OC}$. Since $OA = OB = OC = OX = R$, the result follows.

3. Find all roots of the polynomial $z^4 + 4z^2 + 4$.

Solution. $z^4 + 4z^2 + 4 = (z^2 + 2)^2$. Thus there are two roots, $z = \sqrt{-2} = \pm i\sqrt{2}$, of multiplicity 2 each.

4. Find the place of the curve $x^2 + 2xy + y^2 = 2x$ in classification of quadratic plane curves (i.e. find out if it is an ellipse, hyperbola, parabola, two crossing lines, two parallel lines, one line, one point, or the empty set).

Solution. We have: $x^2 + 2xy + y^2 = (x + y)^2$, and therefore in new coordinates $X = x + y, Y = 2x$, the curve is given by the equation $X^2 = Y$. Thus it is a parabola.

5. Classify quadratic forms $Q = ax^2$ in one variable with complex coefficients (i.e. if $a \in \mathbb{C}$) up to complex linear changes $x = cy, c \in \mathbb{C}, c \neq 0$.

Solution. When $a = 0$, the quadratic form Q is equal to zero identically, and remains zero under any change of the variable. When $a \neq 0$, take $c = a^{-1/2}$ to transform Q to the form $Q = y^2$. Thus, there are two equivalence classes, whose representatives are 0 and y^2 .