

Math 110, Spring 2014. Midterm Review

Things You Must Know:

Definitions:

- Ch. 1 - vector space over F , subspace, sums of subspaces, direct sums of subspace (in particular, they are special types of sums).
- Ch. 2 - finite dimensional spaces, spanning lists, linearly (in)dependent lists, basis, dimension.
- Ch. 3 - linear maps, $\text{null}(T)$, $\text{range}(T)$, $L(V, W)$, $L(V)$, operators, matrix of a linear map, injective, surjective, bijective, invertible linear map, isomorphic, dual space, dual basis.

Theorems:

- Ch. 1 -
- $U \subset V$ is a subspace if and only if $\forall \lambda, \mu \in F, u, v \in U$ we have $\lambda u + \mu v \in U$.
 - for $U, W \subset V$ subspaces, $U + W$ is a subspace.
 - $V = U \oplus W$ if and only if $V = U + W$ and $U \cap W = \{0\}$.
- Ch. 2 -
- if V is f.d and (v_1, \dots, v_m) is linearly dependent in V , and $v_1 \neq 0$ then there is $j \in \{2, \dots, m\}$ such that (v_1, \dots, v_j) is linearly independent, for $i = 1, \dots, j - 1$, and $v_j \in \text{span}(v_1, \dots, v_{j-1})$. Hence, $\text{span}(v_1, \dots, v_j) = \text{span}(v_1, \dots, v_{j-1})$.
 - if V is f.d. with $V = \text{span}(v_1, \dots, v_m)$, and (v_1, \dots, v_k) is linearly independent, then $k \leq m$.
 - if $\dim V = n$ and (v_1, \dots, v_k) is linearly independent then $k \leq n$. (*linearly independent lists are 'small'*)
 - if $\dim V = n$ and (v_1, \dots, v_m) is a spanning list then $m \geq n$. (*spanning lists are 'big'*)
 - bases exist!
 - every linearly independent list of length $\dim V$ is a basis.
 - every spanning list of length $\dim V$ is a basis.
 - every linearly independent list can be extended to a basis.
 - if (v_1, \dots, v_m) is a spanning list in V , $\dim V = n$, then there are $i_1, \dots, i_n \in \{1, \dots, m\}$ distinct, such that $(v_{i_1}, \dots, v_{i_n})$ is a basis.
 - if $U \subset V$ is a subspace then $\dim U \leq \dim V$. In particular, any basis of U can be extended to a basis of V .
 - (**Dimension Formula**) if $U, W \subset V$ are subspaces, V f.d., then
$$\dim(U + W) = \dim U + \dim W - \dim U \cap W.$$
 - $U + W$ is a direct sum if and only if $\dim(U + W) = \dim U + \dim W$.
- Ch. 3 -
- let $T \in L(V, W)$. Then, $\text{null}(T) \subset V$ is a subspace; $\text{range}(T) \subset W$ is a subspace.
 - $T \in L(V, W)$ is injective if and only if $\text{null}(T) = \{0\}$.
 - $T \in L(V, W)$ is surjective if and only if $\text{range}(T) = W$.
 - let V be f.d., $T \in L(V, W)$. Then, $\dim V = \dim \text{null}(T) + \dim \text{range}(T)$.
 - if $\dim V > \dim W$ then no linear map $T \in L(V, W)$ is injective.

- if $\dim V < \dim W$ then no linear map $T \in L(V, W)$ is surjective.
- if $T \in L(V, W)$ is injective and $(v_1, \dots, v_k) \subset V$ is linearly independent then $(T(v_1), \dots, T(v_k)) \subset W$ is linearly independent.
- if $T \in L(V, W)$ is surjective and (v_1, \dots, v_m) spans V then $(T(v_1), \dots, T(v_m))$ spans W .
- $L(V, W)$ is a vector space over F ; it has $\dim L(V, W) = \dim V \dim W$
- $L(V)$ is a vector space over F .
- $T \in L(V)$ is injective if and only if T is surjective if and only if T is invertible.
- V and W are isomorphic if and only if $\dim V = \dim W$.
- if $(v_1, \dots, v_n) \subset V$ is a basis of V and $w_1, \dots, w_n \in W$ arbitrary, then there is a unique linear map $T \in L(V, W)$ such that $T(v_i) = w_i$.
- to any basis of V there is a dual basis of the dual space V' .