

Uniqueness of Weak Solutions to the Ricci Flow and Topological Applications

Richard H Bamler
(joint work with Bruce Kleiner, NYU)

May 2020

Structure of Talk

- **Part I:** Topological Results
- **Part II:** Ricci flow, Weak solutions, Uniqueness, Continuous dependence
- **Part III:** Applications to Topology

Part I: Topological Results

Basic definitions

M (mostly) 3-dimensional, compact, orientable manifold

Recall: The topology of 3-manifolds is sufficiently well understood due to the resolution of the Poincaré and Geometrization Conjectures by Perelman, using Ricci flow.

Main objects of study:

- $\text{Met}(M)$: space of Riemannian metrics on M
- $\text{Met}_{PSC}(M) \subset \text{Met}(M)$: subset of metrics with positive scalar curvature
- $\text{Diff}(M)$: space of diffeomorphisms $\phi : M \rightarrow M$

... each equipped with the C^∞ -topology.

Goal: Classify these spaces up to homotopy (using Ricci flow)!

$\text{Met}(M)$ is contractible

Main Result 1:

Ba., Kleiner 2019

$\text{Met}_{PSC}(M)$ is either contractible or empty.

History:

- true in dimension 2 (via Uniformization Theorem or Ricci flow (see later))
- Hitchin 1974; Gromov, Lawson 1984; Botvinnik, Hanke, Schick, Walsh 2010:
Further examples with $\pi_i(\text{Met}_{PSC}(M^n)) \neq 1$ for certain (large) i, n .
- Marques 2011 (using Ricci flow with surgery):
 $\text{Met}_{PSC}(M^3)/\text{Diff}(M^3)$ is path-connected
 $\text{Met}_{PSC}(S^3)$ is path-connected,

Diffeomorphism groups

Smale 1958: $O(3) \simeq \text{Diff}(S^2)$

Smale Conjecture: $O(4) \simeq \text{Diff}(S^3)$

proven by Hatcher in 1983

For a general spherical space form $M = S^3/\Gamma$ consider the injection

$$\text{Isom}(M) \longrightarrow \text{Diff}(M)$$

Generalized Smale Conjecture

This map is a homotopy equivalence.

- Verified for a handful of other spherical space forms, but open e.g. for $\mathbb{R}P^3$.
- All proofs so far are purely topological and technical. No uniform treatment.

Main Result 2:

Theorem (Ba., Kleiner 2019)

The Generalized Smale Conjecture is true.

Remarks:

- Proof via Ricci flow (first purely topological application of Ricci flow since Perelman's work \sim 15 years ago).
- Uniform treatment of all cases.
- Alternative proof in the S^3 -case (Smale Conjecture).
- There are two proofs:
 - "Short" proof (Ba., Kleiner 2017): GSC if $M \not\approx S^3, \mathbb{R}P^3$, M hyperbolic
 - Long proof (Ba., Kleiner 2019): full GSC and $S^2 \times \mathbb{R}$ -cases

Similar techniques imply results in non-spherical case:

- If M is closed and hyperbolic, then $\text{Isom}(M) \simeq \text{Diff}(M)$.
(topological proof by Gabai 2001)
- If (M, g) is aspherical and geometric and g has maximal symmetry, then $\text{Isom}(M) \simeq \text{Diff}(M)$.
(new in non-Haken infranil case.)
- $\text{Diff}(S^2 \times S^1) \simeq O(2) \times O(3) \times \Omega O(3)$
(topological proof by Hatcher)
- $\text{Diff}(\mathbb{R}P^3 \# \mathbb{R}P^3) \simeq O(1) \times O(3)$
(topological proof by Hatcher)

Connection to Ricci flow

Lemma

For any $g \in \text{Met}_{K \equiv \pm 1}(M)$:

$$\text{Isom}(M, g) \simeq \text{Diff}(M) \iff \text{Met}_{K \equiv \pm 1}(M) \text{ contractible}$$

Proof: Fiber bundle

$$\begin{aligned} \text{Isom}(M, g) &\longrightarrow \text{Diff}(M) \longrightarrow \text{Met}_{K \equiv \pm 1}(M) \\ \phi &\longmapsto \phi^* g \end{aligned}$$

Apply long exact homotopy sequence.

This reduces both results to:

Theorem (Ba., Kleiner 2019)

$\text{Met}_{PSC}(M)$ and $\text{Met}_{K \equiv 1}(M)$ are each either contractible or empty.

or equivalently:

$$\pi_k(\text{Met}(M), \text{Met}_{PSC/K \equiv 1}(M)) = 1.$$

Part II: Ricci flow, Weak solutions, Uniqueness, Continuous dependence

Ricci flow

Ricci flow: $(M, g(t)), t \in [0, T)$

$$\partial_t g(t) = -2 \operatorname{Ric}_{g(t)}, \quad g(0) = g_0 \quad (*)$$


Short-time existence (Hamilton):

- For every initial condition g_0 the initial value problem $(*)$ has a unique solution for maximal $T \in (0, \infty]$.
- If $T < \infty$, then “singularity at time T ”. Curvature $|\operatorname{Rm}|$ blows up as $t \nearrow T$.

Example: Round shrinking sphere

$$M = S^n$$

$$g(t) = (1 - 2(n-1)t)g_{S^n}.$$




Ricci flow in 2D

Hamilton, Chow: On S^2 for any initial condition g_0 we have

$$T = \frac{\text{vol}(S^2, g_0)}{8\pi}, \quad (T - t)^{-1}g(t) \rightarrow g_{\text{round}}$$

Interpretation on the space of metrics:



- Preservation of positive scalar curvature (in all dimensions)
- \rightsquigarrow deformation retractions from $\text{Met}(S^2)$ and $\text{Met}_{PSC}(S^2)$ onto $\text{Met}_{K \equiv 1}(S^2)$

Theorem

$\text{Met}_{PSC}(S^2) \simeq \text{Met}_{K \equiv 1}(S^2) \simeq \text{Met}(S^2) \simeq *$
Therefore $\text{Diff}(S^2) \simeq O(3)$.


Difficulties:

- Flow may incur non-round and non-global singularities.
- Necessary to extend the flow past the first singular time (surgeries).
- Continuous dependence on initial data?

Results:

- Perelman: Qualitative classification of singularity models (κ -solutions)
- Brendle 2018 / Ba., Kleiner 2019: Further classification / rotational symmetry of κ -solutions

Example: rotationally symmetric dumbbell



Ricci flow with surgery

Given (M, g_0) construct
Ricci flow with surgery:


$$(M_1, g_1(t)), t \in [0, T_1],$$

$$(M_2, g_2(t)), t \in [T_1, T_2],$$

$$(M_3, g_3(t)), t \in [T_2, T_3], \dots$$

Observations:

- surgery scale $\approx \delta \ll 1$
- high curvature regions are ε -close to singularity models from before:
“ ε -canonical neighborhood assumption”




RF with surgery was used to prove Poincaré & Geometrization Conjectures

Drawback:

surgery process is not canonical (depends on surgery parameters)

Perelman:

- *It is likely that [...] one would get a canonically defined Ricci flow through singularities, but at the moment I don't have a proof of that.*
- *Our approach [...] is aimed at eventually constructing a canonical Ricci flow, [...] - a goal, that has not been achieved yet in the present work.*



Theorem (Ba., Kleiner, Lott)

Perelman's "conjecture" is true:

- There is a notion of a **weak Ricci flow** "through singularities" and we have **existence** and **uniqueness** within this class.
- This weak flow is a **limit** of Ricci flows with surgery, where surgery scale $\delta \rightarrow 0$.

Comparison with Mean Curvature Flow:

- Notions of weak flows: Level Set Flow, Brakke flow
- General case: fattening \cong non-uniqueness
- Mean convex case: non-fattening \cong uniqueness
- 2-convex case: uniqueness + weak flow is limit of MCF with surgery as surgery scale $\delta \rightarrow 0$

How to take limits of sequences of Ricci flows with surgery?

Space-time picture

Space-time 4-manifold:

$$\mathcal{M}^4 = (M_1 \times [0, T_1] \cup M_2 \times [T_1, T_2] \cup M_3 \times [T_2, T_3] \cup \dots) - \text{surgery points}$$

Time function: $t: \mathcal{M} \rightarrow [0, \infty)$

Time-slices: $\mathcal{M}_t = t^{-1}(t)$

Time vector field:

∂_t on \mathcal{M} (with $\partial_t \cdot t = 1$)


Metric g : on the distribution
 $\ker dt \subset T\mathcal{M}$

Ricci flow equation:

$$\mathcal{L}_{\partial_t} g = -2 \text{Ric}_g$$

$\mathcal{M} = (\mathcal{M}, t, \partial_t, g)$ is called a
Ricci flow spacetime.

Note: there are “holes” at scale $\approx \delta$
space-time is δ -complete




Kleiner, Lott 2014: Compactness theorem and $\delta_i \rightarrow 0$
 \implies existence of **singular Ricci flow** starting from any (M, g)

Singular Ricci flow: Ricci flow spacetime \mathcal{M} that:

- is **0-complete** (i.e. “surgery scale $\delta = 0$ ”)
- satisfies the **ε -canonical neighborhood assumption** for small ε .

Remarks:

- \mathcal{M} is smooth everywhere and not defined at singularities
- singular times may accumulate



Theorem (Ba., Kleiner 2016)

\mathcal{M} is uniquely determined by its initial time-slice (\mathcal{M}_0, g_0) up to isometry.


So for any (M, g) there is (up to isometry) a **canonical singular Ricci flow** \mathcal{M} with initial time-slice $(\mathcal{M}_0, g_0) \cong (M, g)$.

Write: \mathcal{M}^g .

Uniqueness \longrightarrow Continuous dependence

continuous family of metrics $(g^s)_{s \in [0,1]}$ on M

$\rightsquigarrow \{\mathcal{M}^s := \mathcal{M}^{g^s}\}_{s \in [0,1]}$ singular RFs



Continuity of singular RFs

\mathcal{M}^g depends continuously on its initial metric g .

Precise statement:

Theorem (Ba., Kleiner 2019)

Given a continuous family $(g^s)_{s \in X}$ of Riemannian metrics on M over some topological space X , there is a continuous family of singular RFs $(\mathcal{M}^s = \mathcal{M}^{g^s})_{s \in X}$. That is:

- A topology on $\sqcup_{s \in X} \mathcal{M}^s$ such that the projection

$$\bigsqcup_{s \in X} \mathcal{M}^{g^s} \longrightarrow X$$

is a topological submersion.

- A compatible lamination structure on $\sqcup_{s \in X} \mathcal{M}^s$ with leaves \mathcal{M}^s with respect to which all objects t^s, ∂_t^s, g^s are transversely continuous.

Part III: Applications to Topology

Theorem (Ba., Kleiner 2019)

$$\pi_k(\text{Met}(M), \text{Met}_{\text{PSC}/K \equiv 1}(M)) = 1$$

Meaning (PSC-case): For any family of metrics $(h_{s,0})_{s \in D^k}$ on M , where

$$h_{s,0} \text{ has PSC for } s \in \partial D^k,$$

there is a **homotopy** $(h_{s,t})_{s \in D^k \times [0,1]}$, s.t.

$$h_{s,1} \text{ has PSC for } s \in D^k$$

$$h_{s,t} \text{ has PSC for } s \in \partial D^k, \quad t \in [0, 1]$$

Previous results: $(h_{s,0})_{s \in D^k} \rightsquigarrow$ cont. family of singular RFs $(\mathcal{M}^s = \mathcal{M}^{g^s})_{s \in D^k}$

Remaining Conversion Problem: Convert continuous family of sing. RFs $(\mathcal{M}^s)_{s \in X}$ with initial time-slice $\mathcal{M}_0^s = M$ to $(h_{s,t})_{s \in X, t \in [0,1]}$ with:

- 1 $(M, h_{s,0}) \cong (\mathcal{M}_0^s, g_0^s)$.
- 2 $h_{s,1}$ has PSC.
- 3 If \mathcal{M}^s has PSC, then so does $h_{s,t}$ for all $t \in [0, 1]$.

Conversion Problem: Given $(\mathcal{M}^s)_{s \in X}$, find $(h_{s,t})_{s \in X, t \in [0,1]}$ s.t.:

- 1 $(M, h_{s,0}) \cong (\mathcal{M}_0^s, g_0^s)$
- 2 $h_{s,1}$ has PSC
- 3 If \mathcal{M}^s has PSC, then so does $h_{s,t}$ for all $t \in [0, 1]$.

- **Rounding procedure:** perturb metrics on $(\mathcal{M}^s)_{s \in X}$ so that they are round or rotationally symmetric in high curvature regions (this works because κ -solutions are round or rot. symmetric)
- **Strategy:** Construct $(h_{s,t})$ by backwards induction over time.
- **Problem:** For any fixed $T \geq 0$ the family $s \mapsto \mathcal{M}_T^s$ of time- T -slices is a “continuous family of Riemannian manifolds” whose topology may vary.
- **New notion:** “Partial homotopy at time T ”

Conversion Problem: Given $(\mathcal{M}^s)_{s \in X}$, find $(h_{s,t})_{s \in X, t \in [0,1]}$ s.t.:

- 1 $(M, h_{s,0}) \cong (\mathcal{M}_0^s, g_0^s)$
- 2 $h_{s,1}$ has PSC
- 3 If \mathcal{M}^s has PSC, then so does $h_{s,t}$ for all $t \in [0,1]$.

Partial homotopy at time T :

Notion involving families of metrics $(h_{s,t})$ as in 1-3, but with

$$“(M, h_{s,0}) \cong (\mathcal{M}_T^s, g_T^s)”$$

defined where $|\text{Rm}| \lesssim r^{-2}$ over a simplicial decomposition of X .

Lemma

- If $T \gg 0$, then there is an (empty) partial homotopy at time T for $(\mathcal{M}^s)_{s \in X}$.
- A partial homotopy at time T for $(\mathcal{M}^s)_{s \in X}$, can be transformed into a partial homotopy at time $T - \varepsilon$, where $\varepsilon > 0$ is uniform, through certain [modification moves](#).
- If there is a partial homotopy at time $T = 0$ for $(\mathcal{M}^s)_{s \in X}$, then there is a family $(h_{s,t})$ satisfying 1-3

Partial homotopy at time T :

Fix a simplicial decomposition of X . For each simplex $\sigma \subset X$ choose:

- a continuous family of compact domains $(Z_s^\sigma \subset \mathcal{M}_T^s)_{s \in \sigma}$.
(roughly: $Z_s^\sigma \approx \{|\text{Rm}| \lesssim r_{\dim \sigma}^{-2}\}$ for $r_0 \ll \dots \ll r_n$.)
- a continuous family of Riemannian metrics $(h_{s,t}^\sigma)_{s \in \sigma, t \in [0,1]}$ on (Z_s^σ) .

such that ①-③ hold at time T and:

- **Compatibility:** If $s \in \tau \subset \sigma$, then $Z_s^\sigma \subset Z_s^\tau$ and $h_{s,t}^\sigma = h_{s,t}^\tau|_{Z_s^\sigma}$.
- **Largeness of the domains:** $|\text{Rm}| \gtrsim r_{\dim \sigma}^{-2}$ on $\mathcal{M}_T^s \setminus Z_s^\sigma$ (\Rightarrow round or rot. symmetric)
- **“Contractible ambiguity”:** $h_{s,t}^\tau$ is round or rot. symmetric on any $Z_s^\tau \setminus Z_s^\sigma$.

Modification Moves:

- Passing to a simplicial refinement.
- Enlarging some (Z_s^σ) by a family of round or rot. symmetric subsets.
- Shrinking some (Z_s^σ) by removing a family of disks.
- Reducing T to $T - \varepsilon$ if (Z_s^σ) stay away from high curvature regions.