

# Uniqueness of Weak Solutions to the Ricci Flow and Topological Applications

Richard H Bamler  
(joint work with Bruce Kleiner, NYU)

June 2020



# Structure of Talk

- **Part I:** Topological Results
- **Part II:** Ricci flow, Weak solutions, Uniqueness, Continuous dependence
- **Part III:** Applications to Topology

# Part I: Topological Results

# Basic definitions

$M$  (mostly) 3-dimensional, compact, orientable manifold

**Recall:** The topology of 3-manifolds is sufficiently well understood due to the resolution of the Poincaré and Geometrization Conjectures by Perelman, using Ricci flow.

## Main objects of study:

- $\text{Met}(M)$ : space of Riemannian metrics on  $M$
- $\text{Met}_{PSC}(M) \subset \text{Met}(M)$ : subset of metrics with positive scalar curvature
- $\text{Diff}(M)$ : space of diffeomorphisms  $\phi : M \rightarrow M$

... each equipped with the  $C^\infty$ -topology.

**Goal:** Classify these spaces up to homotopy (using Ricci flow)!

$\text{Met}(M)$  is contractible

## Main Result 1:

Ba., Kleiner 2019

$\text{Met}_{PSC}(M)$  is either contractible or empty.

## History:

- true in dimension 2 (via Uniformization Theorem or Ricci flow (see later))
- Hitchin 1974; Gromov, Lawson 1984; Botvinnik, Hanke, Schick, Walsh 2010:  
Further examples with  $\pi_i(\text{Met}_{PSC}(M^n)) \neq 1$  for certain (large)  $i, n$ .
- Marques 2011 (using Ricci flow with surgery):  
 $\text{Met}_{PSC}(M^3)/\text{Diff}(M^3)$  is path-connected  
 $\text{Met}_{PSC}(S^3)$  is path-connected,

# Diffeomorphism groups

**Smale 1958:**  $O(3) \simeq \text{Diff}(S^2)$

**Smale Conjecture:**  $O(4) \simeq \text{Diff}(S^3)$

proven by Hatcher in 1983

For a general spherical space form  $M = S^3/\Gamma$  consider the injection

$$\text{Isom}(M) \longrightarrow \text{Diff}(M)$$

## Generalized Smale Conjecture

This map is a homotopy equivalence.

- Verified for a handful of other spherical space forms, but open e.g. for  $\mathbb{R}P^3$ .
- All proofs so far are purely topological and technical. No uniform treatment.

## Main Result 2:

Theorem (Ba., Kleiner 2019)

The Generalized Smale Conjecture is true.

### Remarks:

- Proof via Ricci flow (first purely topological application of Ricci flow since Perelman's work  $\sim$  15 years ago).
- Uniform treatment of all cases.
- Alternative proof in the  $S^3$ -case (Smale Conjecture).
- There are two proofs:
  - "Short" proof (Ba., Kleiner 2017): GSC if  $M \not\approx S^3, \mathbb{R}P^3$ ,  $M$  hyperbolic
  - Long proof (Ba., Kleiner 2019): full GSC and  $S^2 \times \mathbb{R}$ -cases



## Similar techniques imply results in non-spherical case:

- If  $M$  is closed and hyperbolic, then  $\text{Isom}(M) \simeq \text{Diff}(M)$ .  
(topological proof by Gabai 2001)
- If  $(M, g)$  is aspherical and geometric and  $g$  has maximal symmetry, then  $\text{Isom}(M) \simeq \text{Diff}(M)$ .  
(new in non-Haken infranil case.)
- $\text{Diff}(S^2 \times S^1) \simeq O(2) \times O(3) \times \Omega O(3)$   
(topological proof by Hatcher)
- $\text{Diff}(\mathbb{R}P^3 \# \mathbb{R}P^3) \simeq O(1) \times O(3)$   
(topological proof by Hatcher)

# Connection to Ricci flow

## Lemma

For any  $g \in \text{Met}_{K \equiv \pm 1}(M)$ :

$$\text{Isom}(M, g) \simeq \text{Diff}(M) \iff \text{Met}_{K \equiv \pm 1}(M) \text{ contractible}$$

**Proof:** Fiber bundle

$$\begin{aligned} \text{Isom}(M, g) &\longrightarrow \text{Diff}(M) \longrightarrow \text{Met}_{K \equiv \pm 1}(M) \\ &\phi \longmapsto \phi^* g \end{aligned}$$

Apply long exact homotopy sequence.

**This reduces both results to:**

**Theorem (Ba., Kleiner 2019)**

$\text{Met}_{PSC}(M)$  and  $\text{Met}_{K \equiv 1}(M)$  are each either contractible or empty.

## Part II: Ricci flow, Weak solutions, Uniqueness, Continuous dependence

# Ricci flow

**Ricci flow:**  $(M, g(t)), t \in [0, T)$

$$\partial_t g(t) = -2 \operatorname{Ric}_{g(t)}, \quad g(0) = g_0 \quad (*)$$


**Short-time existence (Hamilton):**

- For every initial condition  $g_0$  the initial value problem  $(*)$  has a unique solution for maximal  $T \in (0, \infty]$ .
- If  $T < \infty$ , then “singularity at time  $T$ ”. Curvature  $|\operatorname{Rm}|$  blows up as  $t \nearrow T$ .

**Example:** Round shrinking sphere

$$M = S^n$$

$$g(t) = (1 - 2(n-1)t)g_{S^n}.$$




# Ricci flow in 2D

**Hamilton, Chow:** On  $S^2$  for any initial condition  $g_0$  we have

$$T = \frac{\text{vol}(S^2, g_0)}{8\pi}, \quad (T - t)^{-1}g(t) \rightarrow g_{\text{round}}$$

**Interpretation on the space of metrics:**



- Preservation of positive scalar curvature (in all dimensions)
- $\rightsquigarrow$  deformation retractions from  $\text{Met}(S^2)$  and  $\text{Met}_{PSC}(S^2)$  onto  $\text{Met}_{K \equiv 1}(S^2)$

## Theorem

$\text{Met}_{PSC}(S^2) \simeq \text{Met}_{K \equiv 1}(S^2) \simeq \text{Met}(S^2) \simeq *$   
Therefore  $\text{Diff}(S^2) \simeq O(3)$ .


## Difficulties:

- Flow may incur non-round and non-global singularities.
- Necessary to extend the flow past the first singular time (surgeries).
- Continuous dependence on initial data?

## Results:

- Perelman: Qualitative classification of singularity models ( $\kappa$ -solutions)
- Brendle 2018 / Ba., Kleiner 2019: Further classification / rotational symmetry of  $\kappa$ -solutions

**Example:** rotationally symmetric dumbbell



# Ricci flow with surgery

Given  $(M, g_0)$  construct  
Ricci flow with surgery:


$$(M_1, g_1(t)), t \in [0, T_1],$$

$$(M_2, g_2(t)), t \in [T_1, T_2],$$

$$(M_3, g_3(t)), t \in [T_2, T_3], \dots$$

## Observations:

- surgery scale  $\approx \delta \ll 1$
- high curvature regions are  $\varepsilon$ -close to singularity models from before:  
“ $\varepsilon$ -canonical neighborhood assumption”






RF with surgery was used to prove  
Poincaré & Geometrization Conjectures

**Drawback:**

surgery process is not canonical  
(depends on surgery parameters)

**Perelman:**

- *It is likely that [...] one would get a canonically defined Ricci flow through singularities, but at the moment I don't have a proof of that.*
- *Our approach [...] is aimed at eventually constructing a canonical Ricci flow, [...] - a goal, that has not been achieved yet in the present work.*



## Theorem (Ba., Kleiner, Lott)

Perelman's "conjecture" is true:

- There is a notion of a **weak Ricci flow** "through singularities" and we have **existence** and **uniqueness** within this class.
- This weak flow is a **limit** of Ricci flows with surgery, where surgery scale  $\delta \rightarrow 0$ .

### Comparison with Mean Curvature Flow:

- Notions of weak flows: Level Set Flow, Brakke flow
- General case: fattening  $\cong$  non-uniqueness
- Mean convex case: non-fattening  $\cong$  uniqueness
- 2-convex case: uniqueness + weak flow is limit of MCF with surgery as surgery scale  $\delta \rightarrow 0$

### How to take limits of sequences of Ricci flows with surgery?

# Space-time picture

Space-time 4-manifold:

$$\mathcal{M}^4 = (M_1 \times [0, T_1] \cup M_2 \times [T_1, T_2] \cup M_3 \times [T_2, T_3] \cup \dots) - \text{surgery points}$$

Time function:  $t: \mathcal{M} \rightarrow [0, \infty)$

Time-slices:  $\mathcal{M}_t = t^{-1}(t)$

Time vector field:

$\partial_t$  on  $\mathcal{M}$  (with  $\partial_t \cdot t = 1$ )


Metric  $g$ : on the distribution  
 $\ker dt \subset T\mathcal{M}$

Ricci flow equation:

$$\mathcal{L}_{\partial_t} g = -2 \text{Ric}_g$$

$\mathcal{M} = (\mathcal{M}, t, \partial_t, g)$  is called a  
Ricci flow spacetime.

**Note:** there are “holes” at scale  $\approx \delta$   
space-time is  $\delta$ -complete




**Kleiner, Lott 2014:** Compactness theorem and  $\delta_i \rightarrow 0$   
 $\implies$  existence of (weak) singular Ricci flow starting from any  $(M, g)$

**Singular Ricci flow:** Ricci flow spacetime  $\mathcal{M}$  that:

- is 0-complete (i.e. “surgery scale  $\delta = 0$ ”)
- satisfies the  $\varepsilon$ -canonical neighborhood assumption for small  $\varepsilon$ .

**Remarks:**

- $\mathcal{M}$  is smooth everywhere and not defined at singularities
- singular times may accumulate



## Theorem (Ba., Kleiner 2016)

$\mathcal{M}$  is uniquely determined by its initial time-slice  $(\mathcal{M}_0, g_0)$  up to isometry.


So for any  $(M, g)$  there is (up to isometry) a **canonical singular Ricci flow**  $\mathcal{M}$  with initial time-slice  $(\mathcal{M}_0, g_0) \cong (M, g)$ .

Write:  $\mathcal{M}^g$ .

# Uniqueness $\longrightarrow$ Continuous dependence

continuous family of metrics  $(g^s)_{s \in [0,1]}$  on  $M$


$\rightsquigarrow \{\mathcal{M}^s := \mathcal{M}^{g^s}\}_{s \in [0,1]}$  singular RFs



# Uniqueness $\longrightarrow$ Continuous dependence

continuous family of metrics  $(g^s)_{s \in [0,1]}$  on  $M$

$\rightsquigarrow \{\mathcal{M}^s := \mathcal{M}^{g^s}\}_{s \in [0,1]}$  singular RFs



# Continuity of singular RFs

$\mathcal{M}^g$  depends continuously on its initial metric  $g$ .

## Precise statement:

### Theorem (Ba., Kleiner 2019)

Given a continuous family  $(g^s)_{s \in X}$  of Riemannian metrics on  $M$  over some topological space  $X$ , there is a continuous family of singular RFs  $(\mathcal{M}^s = \mathcal{M}^{g^s})_{s \in X}$ . That is:

- A topology on  $\sqcup_{s \in X} \mathcal{M}^s$  such that the projection

$$\bigsqcup_{s \in X} \mathcal{M}^{g^s} \longrightarrow X$$

is a topological submersion.

- A compatible lamination structure on  $\sqcup_{s \in X} \mathcal{M}^s$  with leaves  $\mathcal{M}^s$  with respect to which all objects  $t^s, \partial_t^s, g^s$  are transversely continuous.



## Part III: Applications to Topology

## Generalized Smale Conjecture

Let  $M \approx S^3/\Gamma$ , then  $\text{Diff}(M) \simeq \text{Isom}(M)$ .

**Goal:** As in 2D, construct a retraction

$$* \simeq \text{Met}(M) \longrightarrow \text{Met}_{K \equiv 1}(M)$$

**Strategy:**

$$\begin{array}{ccc} \text{Met}(M) & \longrightarrow & \{\text{singular RFs}\} \overset{?}{\dashrightarrow} \text{Met}_{K \equiv 1}(M) \\ \text{cont. family} & & \text{cont. family of} \\ \text{of metrics} & & \text{singular RFs} \end{array}$$

**Short proof (30 pages):** if  $M \not\approx S^3, \mathbb{R}P^3$  and assuming the Smale Conj. for  $S^3$

## 2 Observations:

- 1 Distinguished end  
 $\mathcal{C} \approx M \times [T_1, T_2) \subset \mathcal{M}$  on which metric converges to  $K \equiv 1$  metric modulo rescaling.
- 2 Every component of every time-slice only has finitely many bad points.

(**bad point**  $x \in \mathcal{M}$ :  
 trajectory of  $-\partial_t$  through  $x$   
 does not exist up to time 0.)

$(W, \bar{g}_t) :=$  push-forward of  $(\mathcal{C}_t, g_t)$   
 via flow of  $-\partial_t$

$(W, \bar{g} := \lim_{t \nearrow T_2} \lambda_t \bar{g}_t)$   
 $\cong S^3 / \Gamma \setminus \{p_1, \dots, p_N\}$

$\rightsquigarrow$  continuous, canonical map

$$\text{Met}(M) \longrightarrow \text{PartMet}_{K \equiv 1}(M), \quad g \longmapsto (W, \bar{g})$$

Obstruction theory  $\rightsquigarrow$   $\text{PartMet}_{K \equiv 1}(M) \longrightarrow \text{Met}_{K \equiv 1}(M)$ .

