

**Math 16B – F05 – Supplementary Notes 1**  
**The Derivative and Local Linear Approximation**

For a function  $g(x)$  of one variable, the value  $g'(a)$  of the derivative of  $g$  at the point  $x = a$  is the slope of the tangent line to the graph of  $g$  at the point  $(a, g(a))$ . Near that point, the tangent line is a “good” approximation to the graph, in the following sense. The equation of the tangent line is

$$y = g(a) + g'(a)(x - a).$$

The difference

$$(1) \quad R(x) = g(x) - g(a) - g'(a)(x - a)$$

is the error you get at  $x$  when you approximate  $g$  by the function whose graph is the tangent line (see Figure 1.1). The approximation is good near  $a$  in the sense that  $R(x)$  is small for  $x$  near  $a$  compared to  $x - a$ , vanishingly small in the limit:

$$(2) \quad \lim_{x \rightarrow a} \frac{R(x)}{x - a} = 0.$$

To see this, simply divide (1), the equality that defines  $R(x)$ , by  $x - a$ , to get

$$\frac{R(x)}{x - a} = \frac{g(x) - g(a) - g'(a)(x - a)}{x - a} = \frac{g(x) - g(a)}{x - a} - g'(a).$$

As  $x$  approaches  $a$ , the difference quotient on the right side of the equality approaches  $g'(a)$ , so the ratio on the left side tends to 0.

Among all straight lines through the point  $(a, g(a))$ , the tangent line gives the best linear approximation near  $a$  to the graph of  $g$ . For the line with slope  $c \neq g'(a)$ , the error in the approximation at  $x$ , divided by  $x - a$ , has the limit  $g'(a) - c$ , not 0, as  $x$  approaches  $a$ .

The relation (2), while it says you can expect the error in a certain approximation to be small, does not tell you how small. It does not tell you, for example, how close  $x$  should be to  $a$  to guarantee that the approximation is accurate to, say, three decimal places. Later in the course, when we study Taylor polynomials, we will learn of a method for obtaining useful error estimates. (For linear approximations of  $g$  the estimate involves the second derivative of  $g$ .) At this point one can obtain a good feel for the situation by looking at a few examples. Here is one example. We take  $g(x) = \sqrt{x}$  and  $a = 4$ . Then  $g(4) = 2$ ,  $g'(x) = 1/2\sqrt{x}$ , and  $g'(4) = 1/4$ . The approximating linear function is the function  $2 + \frac{1}{4}(x - 4)$ , which we expect to give a good approximation to  $\sqrt{x}$  for  $x$  near 4. The following table compares  $\sqrt{4}$ , given to four decimal places, to the approximate value  $2 + \frac{1}{4}(x - 4)$ , for various values of  $x$  near 4.

| $x$ | $2 + \frac{1}{4}(x - 4)$ | $\sqrt{x}$ | $R(x)$ |
|-----|--------------------------|------------|--------|
| 3   | 1.75                     | 1.7321     | -.0179 |
| 3.5 | 1.875                    | 1.8708     | -.0042 |
| 3.6 | 1.9                      | 1.8974     | -.0026 |
| 3.7 | 1.925                    | 1.9235     | -.0015 |
| 3.8 | 1.95                     | 1.9494     | -.0006 |
| 3.9 | 1.975                    | 1.9748     | -.0002 |
| 4.1 | 2.025                    | 2.0248     | -.0002 |
| 4.2 | 2.05                     | 2.0494     | -.0006 |
| 4.3 | 2.075                    | 2.0736     | -.0019 |
| 4.4 | 2.1                      | 2.0976     | -.0024 |
| 4.5 | 2.125                    | 2.1213     | -.0037 |
| 5   | 2.25                     | 2.2361     | -.0139 |

One sees that for  $|x - 4| \leq .5$  the approximation is good to two decimals, for  $|x - 4| \leq .1$  it is good to three.

**Question.** Why in this example does the approximation always produce an overestimate?

### More Variables

The considerations above, with partial derivatives in place of derivatives, apply to functions of several variables. Consider a function  $f(x, y)$  of two variables. Near a point  $(a, b)$  in the plane consider the two one-variable functions  $f(x, b)$  and  $g(a, y)$ . We then have the approximation

$$(3) \quad f(x, b) \approx f(a, b) + \left( \frac{\partial f}{\partial x}(a, b) \right) (x - a)$$

$$(4) \quad f(a, y) \approx f(a, b) + \left( \frac{\partial f}{\partial y}(a, b) \right) (y - b),$$

the first holding for  $x$  near  $a$ , the second for  $y$  near  $b$ .

The relations (3) and (4) are special cases of a more general linear approximation property, namely

$$(5) \quad f(x, y) \approx f(a, b) + \left( \frac{\partial f}{\partial x}(a, b) \right) (x - a) + \left( \frac{\partial f}{\partial y}(a, b) \right) (y - b),$$

where the error in the approximation, the quantity

$$R(x, y) = f(x, y) - f(a, b) - \left( \frac{\partial f}{\partial x}(a, b) \right) (x - a) - \left( \frac{\partial f}{\partial y}(a, b) \right) (y - b),$$

is small compared to the distance of  $(x, y)$  from  $(a, b)$ :

$$(6) \quad \lim_{(x,y) \rightarrow (a,b)} \frac{R(x, y)}{\sqrt{(x - a)^2 + (y - b)^2}} = 0.$$

This is more subtle than its one-variable forebear in that its validity requires more than the mere existence of the partial derivatives  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  at  $(a, b)$ . The kinds of functions for which it fails, however, do not arise in Math 16B.

## Tangent Lines to Level Curves

Consider a level curve  $f(x, y) = c$  of the function  $f$  and a point  $(a, b)$  on it. The equation for the tangent line to the level curve at  $(a, b)$  can (usually) be written in terms of the partial derivatives of  $f$  at  $(a, b)$ . To simplify the notation we shall write  $f_x$  for  $\frac{\partial f}{\partial x}$  and  $f_y$  for  $\frac{\partial f}{\partial y}$ .

**Theorem.** *If  $(a, b)$  is not a critical point of  $f$ , then the tangent line at  $(a, b)$  to the level curve of  $f$  through  $(a, b)$  has the equation*

$$(7) \quad f_x(a, b)(x - a) + f_y(a, b)(y - b) = 0.$$

To get an idea of why this is true, consider a point  $(x_1, y_1)$  on the level curve close to but distinct from  $(a, b)$ . We write down the approximate equality (5) for  $(x, y) = (x_1, y_1)$ . Since we have  $f(x_1, y_1) = f(a, b)$ , we get

$$f_x(a, b)(x_1 - a) + f_y(a, b)(y_1 - b) \approx 0,$$

which seems to be saying that the point  $(x_1, y_1)$  is nearly on the line (7). To make this more precise we employ the limit relation (6), which in our case reads

$$(8) \quad \lim_{(x_1, y_1) \rightarrow (a, b)} \frac{f_x(a, b)(x_1 - a) + f_y(a, b)(y_1 - b)}{\sqrt{(x_1 - a)^2 + (y_1 - b)^2}} = 0,$$

it being understood that the limit is taken as  $(x_1, y_1)$  approaches  $(a, b)$  along the level curve. To interpret (8) we introduce the point  $(\tilde{x}_1, \tilde{y}_1)$  defined by

$$\tilde{x}_1 = a + \frac{x_1 - a}{\sqrt{(x_1 - a)^2 + (y_1 - b)^2}}, \quad \tilde{y}_1 = b + \frac{y_1 - b}{\sqrt{(x_1 - a)^2 + (y_1 - b)^2}}.$$

The point  $(\tilde{x}_1, \tilde{y}_1)$  lies on the secant line to the level curve determined by the points  $(a, b)$  and  $(x_1, y_1)$ , and it has distance 1 from  $(a, b)$  (see Figure 1.2). In terms of  $(\tilde{x}_1, \tilde{y}_1)$  we can rewrite (8) as

$$(9) \quad \lim_{(x_1, y_1) \rightarrow (a, b)} f_x(a, b)(\tilde{x}_1 - a) + f_y(a, b)(\tilde{y}_1 - b) = 0.$$

As  $(x_1, y_1)$  approaches  $(a, b)$  along the level curve, the secant line determined by  $(a, b)$  and  $(x_1, y_1)$  will rotate and approach the tangent line at  $(a, b)$ . The point  $(\tilde{x}_1, \tilde{y}_1)$  will converge to a point  $(\tilde{x}, \tilde{y})$  on the tangent line, at distance 1 from  $(a, b)$ , hence different from  $(a, b)$ . The limit relation (9) tells us that

$$f_x(a, b)(\tilde{x} - a) + f_y(a, b)(\tilde{y} - b) = 0,$$

in other words,  $(\tilde{x}, \tilde{y})$  lies on the line (7). Since a line in the plane is uniquely determined by any two of its points, we conclude that the tangent line is in fact the line (7).

The case where  $(a, b)$  is a critical point of  $f$ , ignored by the theorem above, is more complicated and will not be discussed here. It will not arise in later developments.