

Lecture 2

NEWTON POLYHEDRA AND
VECTOR-VALUED LAURENT POLYNOMIALS

Askold Khovanskii
*Department of Mathematics,
University of Toronto*

March 19, 2025

NOTATIONS

T is the n -dimensional complex torus $(\mathbb{C}^*)^n$

$$\mathbf{x} = (x_1, \dots, x_n) \in T \Leftrightarrow x_1 \neq 0, \dots, x_n \neq 0.$$

$$\mathbf{x}^\alpha = x_1^{a_1} \cdots x_n^{a_n}$$

is a **character** (monomial) **of the power**


$$\alpha = (a_1, \dots, a_n) \in \mathbb{Z}^n.$$

The *support*, $\text{Supp}(P)$, of a **Laurent polynomial** P is the finite set $A \in \mathbb{Z}^n$, such that $P = \sum_{\alpha \in A} c_\alpha \mathbf{x}^\alpha$, where $c_\alpha \neq 0$.

The **Newton polyhedron** $\Delta(P)$ is the convex hull of $\text{Supp}(P)$.

EXAMPLE

Let P be $y^2 + a_0 + a_1x + a_2x^2 + a_3x^3$, where $a_0 \neq 0$, $a_1 \neq 0$, $a_2 \neq 0$, $a_3 \neq 0$. Then $\Delta(P)$ is



and $\text{Supp}(P) = \{(0, 0), (0, 1), (0, 2), (0, 3), (2, 0)\}$.

FINITE DIMENSIONAL T -INVARIANT SPACES

Let $\mathbb{C}[T]$ be the ring of regular functions on the torus T , i.e., let $\mathbb{C}[T]$ be the ring of Laurent polynomials in $x \in T$.

For a finite set \mathcal{A} in the lattice \mathbb{Z}^n , let $L_{\mathcal{A}}$ be the space spanned by characters x^{α} for $\alpha \in \mathcal{A}$.

Thus a Laurent polynomial P belongs to the space $L_{\mathcal{A}}$ if and only if the inclusion $\text{Supp}(P) \subset \mathcal{A}$ holds.

The torus T acts naturally on the ring $\mathbb{C}(T)$. Each space $L_{\mathcal{A}}$ is a finite dimensional T -invariant subspace of the ring $\mathbb{C}(T)$.

Vice versa, any T -invariant finite dimensional subspace of $\mathbb{C}[T]$ is the space $L_{\mathcal{A}}$ for some finite $\mathcal{A} \subset \mathbb{Z}^n$.

MAIN PROBLEM OF THE THEORY

Problem 1. *For generic $P_1 \in L_{\mathcal{A}_1}, \dots, P_r \in L_{\mathcal{A}_r}$ determine the discrete invariants of $X \subset T$ defined by the system*

$$P_1 = \dots = P_r = 0.$$

It turns out that the invariants from Problem 1 depend not on supports \mathcal{A}_i of the Laurent polynomials P_i but only on their Newton polyhedra $\Delta_i = \Delta(P_i)$. Thus Problem 1 is equivalent to the following

Problem 1'. *For generic P_1, \dots, P_r with Newton polyhedra $\Delta_1, \dots, \Delta_r$ determine the discrete invariants of $X \subset T$ defined by*

$$P_1 = \dots = P_r = 0.$$

CURVE X DEFINED IN $(\mathbb{C}^*)^2$ BY

$$P = 0 \text{ WITH } \Delta(P) = \Delta$$

Theorem. *For generic P , let $\overline{X} = X \cup X_\infty$ be a smooth compact curve, where X_∞ is a finite set. Then:*

- 1) the **genus** $g(\overline{X})$ is equal to the number $B(\Delta)$ of integral points in the interior of $\Delta = \Delta(P)$;
- 2) the number $\#X_\infty$ is equal to the number of integral points in the boundary of Δ ;
- 3) the **Euler characteristic** $\chi(X)$ of X is equal to the volume $V(\Delta)$ of Δ multiplied by $-2!$.

TOY GEOMETRIC APPLICATION

The following formula from planar geometry is well-known:

The Pick formula. *The area of a planar integral polygon Δ is equal to the number of integral points inside Δ plus half of the number of integral points in its boundary $\partial\Delta$ minus one.*


The invariants 1)-3) of an algebraic curve X are related:

$$\chi(\overline{X}) = \chi(X) + \#(X_\infty) = 2 - 2g(\overline{X}).$$

Thus Theorem from the previous slide implies the **Pick formula** for a planar integral polyhedron Δ :

$$V(\Delta) = \# \left((\Delta \setminus \partial\Delta) \cap \mathbb{Z}^2 \right) + \frac{1}{2} \# \left(\partial\Delta \cap \mathbb{Z}^2 \right) - 1.$$

PROJECTIVE PLANE AND POLYNOMIALS



On the left: $\Delta(P)$ for generic polynomial P of degree 5.

On the right: diagram of projective plane.

Terms on the horizontal side of $\Delta(P)$ determine $P(x, 0)$.

Terms on the vertical side of $\Delta(P)$ determine $P(0, y)$.

Terms on the hypotenuse of $\Delta(P)$ determine asymptotic of P about the infinite line.

TORIC VARIETIES

Toric variety M of dimension n is a normal irreducible variety equipped with an action of $(\mathbb{C}^*)^n$ with one orbit $O \sim (\mathbb{C}^*)^n$. Under this action M is broken up into a finite number of orbits isomorphic to tori of different dimensions.

Fan $\Delta^\perp \subset (\mathbb{R}^n)^*$ **dual to** $\Delta \subset \mathbb{R}^n$ is a decomposition of $(\mathbb{R}^n)^*$ into polyhedral cones Γ^\perp dual to faces Γ of Δ . A covector ξ lies in the interior of Γ^\perp if the function $\langle \xi, x \rangle$ attains minima on Δ at Γ .

To Δ^\perp **one associates a projective toric variety** M_Δ . Each face $\Gamma \subset \Delta$ corresponds to an orbit $O_\Gamma \subset M_\Delta$ such that:

- 1) $\dim_{\mathbb{C}} O_\Gamma = \dim_{\mathbb{R}} \Gamma$;
- 2) if $\Gamma_1 \subset \Gamma_2$, then O_{Γ_1} belongs to the closure of O_{Γ_2} .

**Subvariety X IN $(\mathbb{C}^*)^2$ DEFINED BY $P_1 = \cdots = P_r = 0$
with $\Delta(P_1) = \Delta_1, \dots, \Delta(P_r) = \Delta_r$.**

Let $M_{\mathcal{F}} \supset (\mathbb{C}^*)^n$ be a smooth toric variety whose fan \mathcal{F} is a subdivision of the dual fan Δ^\perp to $\Delta = \Delta_1 + \cdots + \Delta_r$.

Theorem A. *The closure of X in $M_{\mathcal{F}}$ is a smooth variety which is transversal to all orbits of the toric variety $M_{\mathcal{F}}$.*

Theorem A helps to determine many invariants of X . Thus, with V. Danilov we discovered an algorithm for computing Hodge numbers for the mixed Hodge structure on a cohomology ring of X .

Eventually, this algorithm helped to obtain closed formulae for these numbers. V. Batyrev made use of these formulas in his works on mirror symmetry.

THE BKK THEOREM

Theorem (Kouchnirenko). *For generic P_1, \dots, P_n with the same Newton polyhedron Δ , the system*

$$P_1(x) = \dots = P_n(x) = 0, \quad x \in (\mathbb{C}^*)^n$$

has simple roots only and their number is $n!V(\Delta)$, where V is the standard volume in \mathbb{R}^n .

Theorem (D. Bernstein). *For generic P_1, \dots, P_n with Newton polyhedra $\Delta_1, \dots, \Delta_n$, the system*

$$P_1(x) = \dots = P_n(x) = 0, \quad x \in (\mathbb{C}^*)^n$$

has simple roots only and their number is $n!V_n(\Delta_1, \dots, \Delta_n)$, where V_n is the Minkowski mixed volume in \mathbb{R}^n .

These theories are often referred to as Bernstein–Kouchnirenko–Khovanskii (BKK) theorem. In the last 50 year, many generalizations of the BKK theorem were discovered.

THE MINKOWSKI MIXED VOLUME

(\exists !) $V_n(\Delta_1, \dots, \Delta_n)$, on n -tuples of convex bodies in $\Delta_i \subset \mathbb{R}^n$, such that:

1. $V_n(\Delta, \dots, \Delta)$ is the volume of Δ ;

2. V_n is symmetric;

3. V_n is multi-linear, for example:

$$V_n(\Delta'_1 + \Delta''_1, \Delta_2, \dots) = V_n(\Delta'_1, \Delta_2, \dots) + V_n(\Delta''_1, \Delta_2, \dots);$$

4. $0 \leq V_n(\Delta_1, \dots, \Delta_n)$;

5. $\Delta'_1 \subseteq \Delta_1, \dots, \Delta'_n \subseteq \Delta_n \Rightarrow V_n(\Delta'_1, \dots, \Delta'_n) \leq V_n(\Delta_1, \dots, \Delta_n)$;

6. Alexandrov–Fenchel inequality

$$V_n^2(\Delta_1, \Delta_2, \dots, \Delta_n) \geq V_n(\Delta_1, \Delta_1, \dots, \Delta_n)V_n(\Delta_2, \Delta_2, \dots, \Delta_n);$$

7. isoperimetric inequality ($n = 2$, Δ_1 is the unite ball, $\Delta = \Delta_2$)

$$\left(\frac{1}{2} \text{ length of } \partial\Delta\right)^2 \geq \pi V(\Delta).$$

VECTOR-VALUED VERSION OF THE THEORY

A **vector-valued Laurent polynomial**, whose values lie in a finite dimensional complex vector space \mathcal{E} , is an element of the space

$$\mathcal{E}_T = \mathcal{E} \otimes \mathbb{C}[T].$$

The torus T acts on \mathcal{E}_T by acting on $\mathbb{C}[T]$. For $\alpha \in \mathbb{Z}^n$, let E_α be a subspace in \mathcal{E} which is nonzero only for finitely many α .

The set $\{E_\alpha\}$, $\alpha \in \mathbb{Z}^n$, defines a subspace

$$\mathbf{L} = \bigoplus_{\alpha} E_{\alpha} \otimes x^{\alpha}$$

of \mathcal{E}_T which is T -invariant and finite dimensional.

Conversely, *any finite dimensional T -invariant $\mathbf{L} \subset \mathcal{E}_T$ can be defined by some set $\{E_\alpha\}$.*

MAIN PROBLEM OF THE THEORY

The *rank of \mathbf{L}* is the dimension of the space $E = \bigoplus_{\alpha} \{E_{\alpha}\}$.

For $\mathbf{f} \in \mathbf{L}$, let $Y(\mathbf{f}) \subset T$ be the variety defined by the vector equation $\mathbf{f}(x) = 0$.

Problem 2. For generic $\mathbf{f} \in \mathbf{L}$, determine the discrete invariants of the variety $Y(\mathbf{f})$ in terms of the discrete invariants of the T -invariant space \mathbf{L} .

If $\{E_{\alpha}\}$ contains only coordinate subspaces of E (with respect to some bases), then Problem 2 is equivalent to Problem 1. Otherwise, it provides its wide generalization.

We'll start with discrete invariants of \mathbf{L} .

CHARACTERISTIC SEQUENCE OF \mathbf{L}

Let \mathbf{L} be a space of the rank r defined by the set $\{E_\alpha\}$.

An i -tuple $(\alpha_1, \dots, \alpha_i)$ of integral points is **admissible for \mathbf{L}** if one can pick

$$e_j \in E_{\alpha_j} \quad 1 \leq j \leq i,$$

such that $\{e_1, \dots, e_i\}$ is linearly independent.

For $1 \leq i \leq r$, the **characteristic polyhedron** Δ_i is the convex hull of all points α representable as the sum $\alpha_1 + \dots + \alpha_i$ of elements of an admissible i -tuple for \mathbf{L} .

The sequence $\Delta_1, \dots, \Delta_r$ of characteristic polyhedra form the **characteristic sequence of \mathbf{L}** . Its role for Problem 2 is similar to the role of the set of Newton polyhedra for Problem 1'.

SUPPORT FUNCTION OF \mathbf{L}

Let \mathbf{L} be the space defined by a set $\{E_\alpha\}$. For $\xi \in (\mathbb{R}^n)^*$ and $c \in \mathbb{R}$, let $E_{\xi,c}$ be the space defined by the following identity:

$$\bigoplus_{\langle \xi, \alpha \rangle \leq c} E_\alpha = E_{\xi,c}.$$

For fixed $\xi \in (\mathbb{R}^n)^*$, the **spaces $E_{\xi,c}$ form a filtration in E** . A number c_0 is **critical of multiplicity $k(c_0)$ for ξ** if

$$\dim_{\mathbb{C}} E_{\xi,c_0} / \bigoplus_{c < c_0} E_{\xi,c} = k(c_0) > 0.$$

For each $\xi \neq 0$, we have $\sum_{c_0 \in \mathbb{R}} k(c_0) = r = \dim_{\mathbb{C}} E$.

A **value of r -valued support function $\mathbf{h}_{\mathbf{L}}$ of \mathbf{L} at ξ** is the multiset of critical numbers c_0 of ξ , where c_0 are repeated $k(c_0)$ times.

REPRESENTATION OF SUPPORT FUNCTION

A sequence h_1, \dots, h_r of piecewise linear functions *represents* an r -valued support function $\mathbf{h}_{\mathbf{L}}$ if for each ξ the multisets

$$\{h_i(\xi)\} \quad \text{and} \quad \mathbf{h}(\xi)$$

coincide. A sequence h_1, \dots, h_r , representing $\mathbf{h}_{\mathbf{L}}$, is not unique, but many invariants of such sequences are equal.

Example. If the rank r of \mathbf{L} is equal to n , then one can define **mixed volume** $V_n(\mathbf{h})$ of the space \mathbf{L} as the mixed volume of a sequence of virtual polyhedra whose support functions h_1, \dots, h_n represent $\mathbf{h}_{\mathbf{L}}$.

One can show that the mixed volume of $\mathbf{h}_{\mathbf{L}}$ is well-defined, i.e., is independent of a choice of the sequence $\{h_1, \dots, h_n\}$ which represents $\mathbf{h}_{\mathbf{L}}$.

DISCRETE INVARIANTS OF A SPACE \mathbf{L}

The characteristic sequence $\Delta_1, \dots, \Delta_r$ and the support function $\mathbf{h}_{\mathbf{L}}$ of \mathbf{L} carry the same discrete information about the space \mathbf{L} .

The support function $\mathbf{h}_{\mathbf{L}}$ of a space \mathbf{L} of rank r has a unique representation as a sequence $h_1(\mathbf{L}), \dots, h_r(\mathbf{L})$ of increasing piecewise linear functions, i.e., $h_1(\mathbf{L}) \leq \dots \leq h_r(\mathbf{L})$.

Theorem. *The increasing sequence $h_1(\mathbf{L}), \dots, h_r(\mathbf{L})$ of piecewise linear functions representing the support function $\mathbf{h}_{\mathbf{L}}$ of a space \mathbf{L} of rank r can be defined as follows:*

$$h_i(\mathbf{L}) = h_{\Delta_i} - h_{\Delta_{i-1}},$$

where h_{Δ_i} for $1 \leq i \leq r$ is the support function of the characteristic polyhedron Δ_i , and $h_{\Delta_0} = 0$.

TORIC VARIETY $M_{\mathcal{F}}$ CONVENIENT FOR \mathbf{L}

A toric variety $M_{\mathcal{F}} \supset T$ is **convenient** for \mathbf{L} if its fan \mathcal{F} is a subdivision of the normal fan Δ^\perp of

$$\Delta = \Delta_1 + \cdots + \Delta_r,$$

where $\Delta_1, \dots, \Delta_r$ is the characteristic sequence for the space \mathbf{L} .

Theorem B. *If a toric variety $M_{\mathcal{F}}$ is smooth and convenient for the space \mathbf{L} , then for generic vector-function $\mathbf{f} \in \mathbf{L}$, the closure in $M_{\mathcal{F}}$ of the variety $Y(\mathbf{f}) \subset T$ is smooth and transversal to all orbits of $M_{\mathcal{F}}$.*

Theorem B relates vector-valued Laurent polynomials with the theory of toric varieties. It has a version which can be stated in terms of T -equivariant vector bundles over toric varieties.

T-EQUIVARIANT TORIC VECTOR BUNDLES

Theorem C. *If $M_{\mathcal{F}}$ is convenient for \mathbf{L} , then there is a unique T -equivariant vector bundle \mathcal{E} over $M_{\mathcal{F}}$, such that:*

- 1) any $\mathbf{f} \in \mathbf{L}$ corresponds to a section of \mathcal{E} over T which can be extended to the global regular section $s(\mathbf{f})$ of \mathcal{E} ;*
- 2) for any $x \in M_{\mathcal{F}}$ and any vector v in the fiber over x , there is $\mathbf{f} \in \mathbf{L}$ such that $v = s(\mathbf{f})(x)$;*
- 3) the multi-valued function associated to the T -invariant vector bundle \mathcal{E} in the Klyachko's classification is equal to $\mathbf{h}_{\mathbf{L}}$.*

Corollary. *The equivariant Chern roots of the vector bundle \mathcal{E} over $M_{\mathcal{F}}$ from the previous theorem are represented by the functions $\{h_1, \dots, h_r\}$.*

GENERALIZATION OF THE BKK THEOREM

Theorem (generalized BKK theorem). *Assume that the space \mathbf{L} has rank $r = n$. Then, for generic vector-function $\mathbf{f} \in \mathbf{L}$, all points in $Y(\mathbf{f})$ are non-degenerate and their number is equal to $n!MV_n(\mathbf{h}_{\mathbf{L}})$.*

In particular, it is equal to

$$n!MV_n(\Delta_1, \Delta_2 - \Delta_1, \dots, \Delta_n - \Delta_{n-1}).$$

Corollary. *Assume that \mathbf{L} has rank $r \leq n$. Then, for generic $\mathbf{f} \in \mathbf{L}$, one can obtain an explicit formula for the class of the variety $Y(\mathbf{f})$ in the ring of conditions of the torus T in terms of the support function $\mathbf{h}_{\mathbf{L}}$.*

HYPERPLANES ARRANGEMENT

Let $\{H_1, \dots, H_N\}$ be a hyperplane arrangement in the projective space \mathbb{P}^n , where $N > n$. Each hyperplane H_i is defined by $u_i = 0$, where u_i is a linear function in the homogeneous coordinates in \mathbb{P}^n .

A Laurent polynomial P in u_1, \dots, u_N whose monomials have degree zero is a rational function on \mathbb{P}^n which is regular on $\mathbb{P}^n \setminus \cup H_i$.

Problem 3. How many solutions has a generic system

$$P_1 = \dots = P_n = 0$$

in $\mathbb{P}^n \setminus \cup H_i$, where P_i are Laurent polynomials in u_1, \dots, u_N , whose Newton polyhedra $\tilde{\Delta}_1, \dots, \tilde{\Delta}_n \subset \mathbb{R}^N$ lie in three hyperplane

$$x_1 + \dots + x_N = 0.$$

CHARACTERISTIC SEQUENCE FOR $\{H_1, \dots, H_N\}$

A subsets $J \subset \{1, \dots, N\}$ is *i-admissible* for some i , such that $1 \leq i < N - n$ if it contains i elements and the following condition holds:

$$\bigcap_{j \notin J} H_j = \emptyset.$$

For an i -admissible set J , let $A_J \in \mathbb{R}^N$ be the point with the coordinate $x_j(A_J) = 1$ if $j \in J$, and $x_j(A_J) = 0$ if $j \notin J$.

The *characteristic sequence* for H_1, \dots, H_N is the sequence of polyhedra $\Delta_1, \dots, \Delta_{N-n-1} \subset \mathbb{R}^N$, where Δ_i is the convex hull of the set of points A_J for all i -admissible sets J . The polyhedron Δ_i lies in the hyperplane

$$x_1 + \dots + x_N = i.$$

SOLUTION TO PROBLEM 3

Theorem. *The number of solutions of a generic system*

$$P_1 = \cdots = P_n = 0$$

in $\mathbb{P}^n \setminus \cup H_i$, where P_i are Laurent polynomials in u_1, \dots, u_N , whose Newton polyhedra $\tilde{\Delta}_1, \dots, \tilde{\Delta}_n \subset \mathbb{R}^N$ lie in the hyperplane $x_1 + \cdots + x_N = 0$, is equal to:

$$m!V_m(\Delta_1, \Delta_2 - \Delta_1, \dots, \Delta_{m-n} - \Delta_{m-n-1}, \tilde{\Delta}_1, \dots, \tilde{\Delta}_n),$$

where $m = N - 1$ and $\Delta_1, \dots, \Delta_{N-n-1}$ is the characteristic sequence for the hyperplanes arrangement.

Mixed volume in the formula makes sense, since all involved polyhedra belong to affine spaces which are parallel to the $(N - 1)$ -dimensional space $x_1 + \cdots + x_N = 0$.

LITERATURE

There are many papers dedicated to the Newton polyhedra theory. The most related to the lecture are the following papers:

[1] Khovanskii, A. *Newton polyhedra, and toroidal varieties*. *Funct. Anal. Appl.* 11 (1977), no. 4, 289–296 (1978).

[2] A. Khovanskii. *Newton Polyhedra and the genus of complete intersections*. *Funct. Anal. Appl.* 12 (1978), no. 1, 38–46 (1978).

[3] Danilov, V.I. & Khovanskii, A. *Newton polyhedra and an algorithm for computing Hodge–Deligne numbers*. *Math. USSR – Izv.* 29(1987), No. 2, 279–298.

The theory of vector valued Laurent polynomials was initiated very recently, see the following paper in preparation:

[4] Kaveh, K. Khovanskii A., and hunter Spink, H. *Vector-valued Laurent polynomial equations, toric vector bundles and matroids.*