

On Fraïssé’s conjecture for linear orders of finite Hausdorff rank

Alberto Marcone

*Dipartimento di Matematica e Informatica, Università di Udine, viale delle
Scienze 206, 33100 Udine, Italy*

Antonio Montalbán

Department of Mathematics, University of Chicago, Chicago, IL 60637, USA

Abstract

We prove that the maximal order type of the wqo of linear orders of finite Hausdorff rank under embeddability is $\varphi_2(0)$, the first fixed point of the ε -function. We then show that Fraïssé’s conjecture restricted to linear orders of finite Hausdorff rank is provable in $\text{ACA}_0^+ + \text{“}\varphi_2(0)\text{ is well-ordered”}$ and, over RCA_0 , implies $\text{ACA}'_0 + \text{“}\varphi_2(0)\text{ is well-ordered”}$.

1 Introduction

Let LO be the class of countable linear orders. If $L, L' \in \text{LO}$ let $L \preceq L'$ mean that L is *embeddable* into L' , i.e. there exists an order preserving injective map from L to L' . $L \sim L'$ abbreviates $L \preceq L'$ and $L' \preceq L$. In this case we say that L and L' are *equimorphic*. It is immediate that \preceq is a quasi-order (i.e. a reflexive and transitive binary relation) and that \sim is an equivalence relation.

Fraïssé’s conjecture (FRA) is the statement that LO is well-quasi-ordered by \preceq , i.e. that there are neither infinite descending chains nor infinite antichains. Roland Fraïssé formulated this conjecture in 1948 ([1]). Richard Laver ([2]) established FRA in 1971 by proving a stronger statement using Nash-Williams’ notion of better-quasi-order ([3]). *Laver’s Theorem* states that LO

¹ This research was done while the second author was visiting the first in Udine during the summer of 2006. The second author was partially supported by NSF Grant DMS-0600824.

is better-quasi-ordered by \preceq (actually Laver proved even more, considering σ -scattered linear orders with labels from a better-quasi-order). All known proofs of Fraïssé’s conjecture (see e.g. [4] for a descriptive set-theoretic one) actually establish Laver’s theorem.

It is easy to state FRA in the language of second order arithmetic, and it is a longstanding open problem in reverse mathematics to establish its exact axiomatic strength. ([5] is the main reference for the reverse mathematics program, and [6] includes a survey on the reverse mathematics of problems related to well- and better-quasi-orderings.) Laver’s proof can be carried out within the strong system $\Pi_2^1\text{-CA}_0$, and Richard Shore ([7]) proved that FRA implies ATR_0 . Since FRA is a Π_2^1 statement, standard model theoretic arguments (see e.g. [8, Corollary 1.10]) yield that FRA does not imply $\Pi_1^1\text{-CA}_0$ (and, a fortiori, $\Pi_2^1\text{-CA}_0$) over ATR_0 . More recently the second author ([9]) showed that FRA is equivalent (over RCA_0 or slightly stronger theories) to other statements about linear orders.

An easy observation is that to establish FRA it suffices to consider *scattered* linear orders, i.e. linear orders $L \in \text{LO}$ such that $\mathbb{Q} \not\preceq L$ (all non-scattered countable linear orders are equimorphic to \mathbb{Q}). Scattered linear orders were first studied by Hausdorff a century ago ([10]), and his results lead to the notion of Hausdorff rank.

Definition 1 *For every ordinal α , \mathbb{Z}^α is the linear order with domain*

$$\{f : \alpha \rightarrow \mathbb{Z} \mid \{\beta < \alpha \mid f(\beta) \neq 0\} \text{ is finite}\}$$

ordered by $f \sqsubset g$ iff $f(\beta) < g(\beta)$ for the largest $\beta < \alpha$ such that $f(\beta) \neq g(\beta)$.

We say that a linear order L has Hausdorff rank less than α , and write $\text{rk}_H(L) < \alpha$, if L embeds into a proper segment of \mathbb{Z}^α , or equivalently if $\mathbf{1} + L + \mathbf{1} \preceq \mathbb{Z}^\alpha$. We say that L has finite Hausdorff rank if it has Hausdorff rank less than ω . (See [11, Chapter 5] for equivalent definitions of \mathbb{Z}^α and rk_H . There are slightly different definitions of Hausdorff rank in the literature, but they are off by at most one.)

Hausdorff proved that $\text{rk}_H(L)$ exists (i.e. $\text{rk}_H(L) < \alpha$ for some ordinal α) if and only if L is scattered. Moreover if L is countable scattered then $\text{rk}_H(L)$ is less than a countable ordinal. The reverse mathematics of the properties of Hausdorff rank was studied by Clote in [12].

If α is a countable ordinal let $\text{LO}_\alpha = \{L \in \text{LO} \mid \text{rk}_H(L) < \alpha\}$. We denote by FRA_α the statement that LO_α is well-quasi-ordered by \preceq . Hence, FRA_α is the restriction of FRA to a subset of LO . Our long term goal is to obtain information on the strength of FRA by looking at the strength of various FRA_α . In this paper we carry out the first step in this project by studying FRA_ω , i.e.

Fraïssé's conjecture for linear orders of finite Hausdorff rank.

A key ingredient for all known proofs of FRA is the notion of indecomposable linear order.

Definition 2 *A linear order L is indecomposable if whenever $L = L_1 + L_2$ then either $L \preceq L_1$ or $L \preceq L_2$. Let ILO (resp. ILO_α) be the class of countable indecomposable linear orders (resp. indecomposable linear orders of Hausdorff rank less than α).*

It follows from FRA that every countable scattered linear order is the finite sum of indecomposable linear orders. The second author showed in [9] that the latter statement is indeed equivalent to FRA over RCA_0 .

We use signed trees ([9,13], see Section 2 below for the definition and the basic properties of these objects) to study indecomposable scattered linear orders. We denote by ST be the set of all signed trees. Using the appropriate notion of homomorphism between signed tree we define equimorphism between signed trees. The key fact about ST is that signed trees represent indecomposable countable linear orders in such a way that homomorphism of signed trees is equivalent to embeddability of the corresponding linear orders. Moreover for each indecomposable countable linear order L there exist a signed tree representing a linear order equimorphic to L . (Indeed, in [9] the second author proved that, over RCA_0 , FRA is equivalent to the statement that ST is well-quasi-ordered by homomorphism.) In this paper we exploit the fact that indecomposable linear orders of finite Hausdorff rank are represented by signed trees of finite height. Since each of these trees is equimorphic to a finite signed tree, to study LO_ω we consider ST_ω , the set of finite signed trees.

It is well-known that a quasi-order (Q, \leq) is a well-quasi-order (wqo for short) if and only if all its linear extensions are well-orders (see [14] for an analysis of the logical strength of the equivalence between this and other characterizations of wqo). Here a linear extension of (Q, \leq) is a quasi-order \sqsubseteq which is linear (i.e. either $q \sqsubseteq r$ or $r \sqsubseteq q$ holds for every $q, r \in Q$), and satisfies $q \leq r \Rightarrow q \sqsubseteq r$ for all $q, r \in Q$. The linear extension is a well-order if it is well-founded and in this case its order type is the unique ordinal α isomorphic to it.

Definition 3 *If (Q, \leq) is a well-quasi-order the maximal order type of (Q, \leq) is*

$$o(Q, \leq) = \sup\{\alpha \mid \alpha \text{ is the order type of a linear extension of } (Q, \leq)\}.$$

Whenever the quasi-order \leq is clear from the context we write $o(Q)$.

De Jongh and Parikh ([15]) showed that the supremum in the definition of maximal order type is actually a maximum, i.e. every well-quasi-order (Q, \leq)

has a linear extension of order type $o(Q)$. Schmidt continued the study of maximal order types in her Habilitationsschrift ([16]). There, she computed the maximal order type of the wqo investigated by Higman ([17]), and gave upper bounds for the maximal order types of the wqo's investigated by Kruskal ([18]) and Nash-Williams ([19]). Harvey Friedman (see [20]) used the maximal order type of the class of finite trees with embeddability preserving g.l.b.s to prove that Kruskal's theorem cannot be proved in ATR_0 . Further extensions of Friedman's method were then used to show that Robertson and Seymour's celebrated result about graph minors is not provable in $\Pi_1^1\text{-CA}_0$ ([21]).

The starting point of our results is the computation of $o(\text{LO}_\omega, \preceq)$. Recall that the Veblen functions give an ordinal notation system for ordinals below Γ_0 . They are defined by letting $\varphi_0(\alpha) = \omega^\alpha$ and, for $\beta > 0$, $\varphi_\beta(\alpha)$ = the α -th common fixed point of all φ_γ with $\gamma < \beta$. In particular φ_1 enumerates the ε -numbers, i.e. the fixed points of the function $\beta \mapsto \omega^\beta$, and we will write ε_α in place of $\varphi_1(\alpha)$.

Using finite signed trees in Section 3 we prove:

Theorem 4 $o(\text{LO}_\omega, \preceq) = \varphi_2(0)$, i.e. the least fixed point of the ε function.

Up to and including Section 3 the paper does not deal with subsystems of second order arithmetic and can be read also by those interested only in wqo theory.

In Section 4 we introduce the subsystems of second order arithmetic we will need and in particular ACA_0^+ and ACA'_0 . ACA_0^+ consists of RCA_0 plus the statement “for every X , $X^{(\omega)}$ (the arithmetic jump of X) exists” (see [22] for recent results about ACA_0^+). ACA_0^+ is strictly weaker than ATR_0 and the ordinal $\varphi_2(0)$ mentioned in Theorem 4 is precisely the proof-theoretic ordinal of ACA_0^+ ([23]). This implies that ACA_0^+ does not prove that $\varphi_2(0)$ is a well-ordering. In Section 4 we make the latter statement precise by introducing, in RCA_0 , an ordinal notation system for ordinal below $\varphi_2(0)$. ACA'_0 consists of RCA_0 plus the statement “for every X and k , $X^{(k)}$ exists”. ACA'_0 is strictly weaker than ACA_0^+ and strictly stronger than ACA_0 .

Formalizing the proof of Theorem 4 in Section 5 we obtain:

Theorem 5 $\text{ACA}_0^+ + \text{“}\varphi_2(0) \text{ is well-ordered”}$ proves FRA_ω .

In Section 6, building on ideas from [7,22], we obtain a lower bound for the complexity of FRA_ω .

Theorem 6 RCA_0 proves that FRA_ω implies $\text{ACA}'_0 + \text{“}\varphi_2(0) \text{ is well-ordered”}$.

There is still a gap between the upper and lower bounds since $\text{ACA}_0^+ + \text{“}\varphi_2(0) \text{ is well-ordered”}$ is strictly stronger than $\text{ACA}'_0 + \text{“}\varphi_2(0) \text{ is well-ordered”}$, as the ω -model consisting of all arithmetic sets is a model of the latter theory but not of the former.

2 Signed trees

Let us start by establishing our notation for finite sequences and trees.

Definition 7 We denote by $X^{<\omega}$ the set of finite sequences of elements of a set X . We use \emptyset to denote the empty sequence. Let $x \in X$ and $\sigma, \tau \in X^{<\omega}$: $\langle x \rangle$ is the sequence of length 1 with sole element x ; $|\sigma|$ is the length of σ and for $i < |\sigma|$, $\sigma(i)$ is the i -th element of σ ; $\sigma \subseteq \tau$ means that σ is an initial segment of τ and $\sigma \subset \tau$ that σ is a proper initial segment of τ ; $\sigma \hat{\ } \tau$ is the sequence of length $|\sigma| + |\tau|$ obtained by concatenating σ and τ ; $\sigma * x$ and $x * \sigma$ abbreviate $\sigma \hat{\ } \langle x \rangle$ and $\langle x \rangle \hat{\ } \sigma$ respectively.

Definition 8 A nonempty set $T \subseteq X^{<\omega}$ is a tree if $\sigma \in T$ and $\tau \subset \sigma$ imply $\tau \in T$. We often refer to members of a tree as nodes and to \emptyset (which belongs to every tree) as the root. A subtree of T is a set of the form $\{\tau \in X^{<\omega} \mid \sigma \hat{\ } \tau \in T\}$ for some $\sigma \in T$; when $\sigma = \langle x \rangle$ we have an immediate subtree of T and we use T_x to denote it.

A child of the node σ of the tree T is a node of T of the form $\sigma * x$. A leaf of T is a node of T with no children. Let $L(T) = \{\sigma \mid \sigma \text{ is a leaf of } T\}$.

A tree T is well-founded if there is no infinite sequence $\sigma_0 \subset \sigma_1 \subset \dots$ of elements of T . When T is well-founded we can define $\text{ht}(\sigma, T) = \sup\{\text{ht}(\sigma * n, T) + 1 \mid \sigma * n \in T\}$ for every $\sigma \in T$. The height of T is $\text{ht}(T) = \text{ht}(\emptyset, T)$.

We can now introduce signed trees, which are our main tool in the study of indecomposable linear orders.

Definition 9 A signed tree is a pair (T, s_T) where $T \subseteq \omega^{<\omega}$ is a well-founded tree and $s_T : T \rightarrow \{+, -\}$.

Let \mathbf{ST} be the set of all signed trees and $\mathbf{ST}_\omega = \{(T, s_T) \in \mathbf{ST} \mid T \text{ is finite}\}$.

If $(T, s_T) \in \mathbf{ST}$ and $\langle n \rangle \in T$ let $s_{T_n} : T_n \rightarrow \{+, -\}$ be defined by $s_{T_n}(\tau) = s_T(n * \tau)$. Obviously $(T_n, s_{T_n}) \in \mathbf{ST}$.

Definition 10 If $(T, s_T), (T', s_{T'}) \in \mathbf{ST}$ a map $f : T \rightarrow T'$ is a homomorphism if

- $\sigma \subset \tau$ implies $f(\sigma) \subset f(\tau)$ for every $\sigma, \tau \in T$;
- $s_{T'}(f(\sigma)) = s_T(\sigma)$ for every $\sigma \in T$.

Note that f is not required to be one-to-one and that $f(\sigma) \subset f(\tau)$ might occur even when $\sigma \not\subset \tau$. If there exists a homomorphism from (T, s_T) to $(T', s_{T'})$ we write $(T, s_T) \preceq (T', s_{T'})$. (T, s_T) and $(T', s_{T'})$ are equimorphic (and we write $(T, s_T) \sim (T', s_{T'})$) if $(T, s_T) \preceq (T', s_{T'})$ and $(T', s_{T'}) \preceq (T, s_T)$.

Lemma 11 *Let $(T, s_T), (T', s_{T'}) \in \mathbf{ST}$ be such that $s_T(\emptyset) = s_{T'}(\emptyset)$. If for all n with $\langle n \rangle \in T$ there exists m such that $\langle m \rangle \in T'$ and $(T_n, s_{T_n}) \preceq (T'_m, s_{T'_m})$ then $(T, s_T) \preceq (T', s_{T'})$.*

PROOF. Immediate from the definition of homomorphism. \square

Lemma 12 *For every $(T, s_T) \in \mathbf{ST}$ with $\text{ht}(T) < \omega$ there exists $(T', s_{T'}) \in \mathbf{ST}_\omega$ such that $(T, s_T) \sim (T', s_{T'})$.*

PROOF. Easy induction on $\text{ht}(T)$, using Lemma 11. \square

The following definition, which was introduced in [9] and [13], connects signed trees with indecomposable linear orders.

Definition 13 *To each $(T, s_T) \in \mathbf{ST}$ we associate a countable linear order $\text{lin}(T, s_T)$ as follows (the definition is by recursion on the height of the tree):*

- if $T = \{\emptyset\}$ and $s_T(\emptyset) = +$ let $\text{lin}(T, s_T) = \omega$;
- if $T = \{\emptyset\}$ and $s_T(\emptyset) = -$ let $\text{lin}(T, s_T) = \omega^*$;
- if $T \neq \{\emptyset\}$ and $s_T(\emptyset) = +$ let

$$\text{lin}(T, s_T) = \sum_{k \in \omega} \left(\sum_{n \in \{0, \dots, k\}: \langle n \rangle \in T} \text{lin}(T_n, s_{T_n}) \right);$$

- if $T \neq \{\emptyset\}$ and $s_T(\emptyset) = -$ let

$$\text{lin}(T, s_T) = \sum_{k \in \omega^*} \left(\sum_{n \in \{0, \dots, k\}: \langle n \rangle \in T} \text{lin}(T_n, s_{T_n}) \right).$$

Lemma 14 ([9]) *For each $(T, s_T), (T', s_{T'}) \in \mathbf{ST}$ we have*

- $\text{lin}(T, s_T)$ is a countable scattered indecomposable linear order;
- $(T, s_T) \preceq (T', s_{T'})$ if and only if $\text{lin}(T, s_T) \preceq \text{lin}(T', s_{T'})$, and in particular $(T, s_T) \sim (T', s_{T'})$ if and only if $\text{lin}(T, s_T) \sim \text{lin}(T', s_{T'})$;
- $\text{rk}_H(\text{lin}(T, s_T))$ and $\text{ht}(T)$ differ by at most one.

Moreover each countable scattered indecomposable linear order different from $\mathbf{1}$ is equimorphic to $\text{lin}(T, s_T)$ for some $(T, s_T) \in \text{ST}$.

Corollary 15 $o(\text{ILO}, \preceq) = o(\text{ST}, \preceq)$ and $o(\text{ILO}_\alpha, \preceq) = o(\text{ST}_\alpha, \preceq)$ whenever α is a limit ordinal. In particular $o(\text{ILO}_\omega, \preceq) = o(\text{ST}_\omega, \preceq)$.

3 The ordinal of $(\text{LO}_\omega, \preceq)$

In this section we prove Theorem 4, i.e. $o(\text{LO}_\omega, \preceq) = \varphi_2(0)$. Our idea is to compute $o(\text{ILO}_\omega)$ using Corollary 15: therefore we concentrate on $o(\text{ST}_\omega)$. We will prove in Propositions 20 and 27 the two inequalities needed to establish $o(\text{ST}_\omega) = \varphi_2(0)$. This yields $o(\text{LO}_\omega) \geq o(\text{ILO}_\omega) = \varphi_2(0)$. We will later use the fact that every countable linear order is the finite sum of indecomposable linear orders to show that $o(\text{LO}_\omega) \leq \varphi_2(0)$.

Here is some notation we will be using. We use \oplus to denote Hessenberg's natural sum of ordinals: recall that given ordinals written in Cantor normal form $\alpha = \omega^{\alpha_0} + \dots + \omega^{\alpha_{n-1}}$ and $\beta = \omega^{\beta_0} + \dots + \omega^{\beta_{m-1}}$,

$$\alpha \oplus \beta = \omega^{\gamma_0} + \omega^{\gamma_1} + \dots + \omega^{\gamma_{n+m-1}},$$

where $\gamma_0, \dots, \gamma_{n+m-1}$ are such that $\gamma_0 \geq \gamma_1 \geq \dots \geq \gamma_{n+m-1}$, and there exists a partition $\{\{a_0, \dots, a_{n-1}\}, \{b_0, \dots, b_{m-1}\}\}$ of $\{0, \dots, n+m-1\}$ such that $\gamma_{a_i} = \alpha_i$ and $\gamma_{b_i} = \beta_i$. Hessenberg's natural sum is commutative and $\alpha + \beta \leq \alpha \oplus \beta$.

We use $\omega_i(\alpha)$ to denote an exponential tower of i ω 's with α on top: $\omega_0(\alpha) = \alpha$ and $\omega_{i+1}(\alpha) = \omega^{\omega_i(\alpha)}$.

We begin by proving or recalling some general facts about the function o . We start by making explicit a widely used technique (implicit in [15] and [16]) for computing an upper bound for the maximal order type of a wqo.

Definition 16 Let (Q, \leq) and (Q', \leq') be quasi-orders. The function $F : Q' \rightarrow Q$ embeds (Q', \leq') in (Q, \leq) , and we write $F : (Q', \leq') \hookrightarrow (Q, \leq)$, if

$$\forall q, r \in Q' (F(q) \leq F(r) \Rightarrow q \leq' r).$$

We write $(Q', \leq') \hookrightarrow (Q, \leq)$ (or $Q' \hookrightarrow Q$ when the binary relations are understood) if there exists a such function F .

Lemma 17 Let (Q, \leq) be a well-quasi-order and (Q', \leq') be a quasi-order. If $Q' \hookrightarrow Q$ then (Q', \leq') is a well-quasi-order and $o(Q') \leq o(Q)$.

PROOF. Let F witness $Q' \hookrightarrow Q$ and denote the range of F by \tilde{Q} . Obviously $o(\tilde{Q}) \leq o(Q)$.

Let \sqsubseteq' be any linear extension of (Q', \leq') . Define a binary relation \sqsubseteq on \tilde{Q} by setting, for $q, r \in Q'$,

$$F(q) \sqsubseteq F(r) \iff q \sqsubseteq' r.$$

This definition is well-posed because $F(q) = F(q')$ implies $q \sqsubseteq' q' \sqsubseteq' q$.

It is straightforward to check that \sqsubseteq is a linear extension of (\tilde{Q}, \leq) and hence (since Q is a well-quasi-order) is a well-order. Hence (Q', \sqsubseteq') is a well-order. This suffices to show that (Q', \leq') is a well-quasi-order.

Moreover the order types of (Q', \sqsubseteq') and (\tilde{Q}, \sqsubseteq) are the same. Since the latter is not greater than $o(\tilde{Q})$ and \sqsubseteq' is arbitrary we have $o(Q') \leq o(\tilde{Q}) \leq o(Q)$. \square

Definition 18 *If (Q, \leq) is a quasi-order and $r \in Q$ let $Q_{\not\leq r} = \{q \in Q \mid q \not\leq r\}$.*

Lemma 19 *Let (Q, \leq) be a well-quasi-order and suppose that $\hat{Q} \subseteq Q$ is cofinal (i.e. $\forall q \in Q \exists r \in \hat{Q} q \leq r$). Then*

$$o(Q) = \sup\{o(Q_{\not\leq r}) + 1 \mid r \in \hat{Q}\}.$$

PROOF. This is essentially Lemma 2.6 of [15]. \square

3.1 The lower bound

Proposition 20 $o(\text{ST}_\omega) \geq \varphi_2(0)$.

PROOF. By Lemma 17 it suffices to define a function $F : \varphi_2(0) \rightarrow \text{ST}_\omega$ witnessing $\varphi_2(0) \hookrightarrow \text{ST}_\omega$.

We define $F(\alpha)$ by recursion on α . For notational convenience let $\varepsilon_{-1} = \omega^{-1} = 0$ and let $F(-1)$ be the signed tree consisting only of the root labeled $+$. With this convention for every $\alpha < \varphi_2(0)$ there exists $\beta < \alpha$ such that $\varepsilon_\beta \leq \alpha < \varepsilon_{\beta+1}$. If $\alpha = \varepsilon_\beta + \delta$, we can write δ in Cantor normal form as $\omega^{\gamma_0} + \dots + \omega^{\gamma_{k-1}}$ with the convention that if $\delta = 0$ we have $k = 1$ and $\gamma_0 = -1$, while if $\delta > 0$ then $\gamma_i \neq -1$ for every $i < k$. We thus have

$$\alpha = \varepsilon_\beta + \omega^{\gamma_0} + \dots + \omega^{\gamma_{k-1}}$$

with $k > 0$, $\beta < \alpha$, and $\alpha > \gamma_0 \geq \dots \geq \gamma_{k-1}$. We can assume that $F(\beta)$, $F(\gamma_0)$, \dots , $F(\gamma_{k-1})$ are already defined.




Fig. 1. A sample $F(\alpha)$ with $k = 2$.

Let $n = \text{ht}(F(\beta)) + 3$, $m_0 = n + \text{ht}(F(\gamma_0)) + 2$, and $m_i = m_{i-1} + \text{ht}(F(\gamma_i)) + 1$ for $0 < i < k$. Then $F(\alpha)$ is the following signed tree. (A sample $F(\alpha)$ with $k = 2$ is depicted in Figure 1.) The root of $F(\alpha)$ is labeled $+$, and has k children, all labeled $+$. The i -th child of the root has two children, both labeled $+$: one of them is the starting point of a linear branch of m_i nodes all labeled $+$ (i.e. it has a single child, which has a single child, etc. for m_i times). The other child of the i -th child of the root has two children, one labeled $-$ and the other labeled $+$. The node labeled $-$ has a single child, which is the root of a copy of $F(\beta)$. The node labeled $+$ is the starting point of a linear branch of n nodes all labeled $+$ and the last node of this branch is the root of a copy of $F(\gamma_i)$.

Notice that for every α the root of $F(\alpha)$ is labeled $+$. It is easy to show inductively that if a node labeled $-$ appears in $F(\alpha)$ then it has a single child which is the root of a copy of some $F(\xi)$ with $\varepsilon_\xi \leq \alpha$. Notice also that if $\alpha \neq -1$ then $F(\alpha)$ has at least one node labeled $-$.

Now we prove by induction on α that if $\alpha, \alpha' < \varphi_2(0)$ are such that $F(\alpha) \preceq F(\alpha')$ then $\alpha \leq \alpha'$. Fix α, α' and a homomorphism f witnessing $F(\alpha) \preceq F(\alpha')$. We can write

$$\begin{aligned}\alpha &= \varepsilon_\beta + \omega^{\gamma_0} + \cdots + \omega^{\gamma_{k-1}} \quad \text{and} \\ \alpha' &= \varepsilon_{\beta'} + \omega^{\gamma'_0} + \cdots + \omega^{\gamma'_{k'-1}}\end{aligned}$$

as above, with $\delta = \omega^{\gamma_0} + \cdots + \omega^{\gamma_{k-1}}$ and $\delta' = \omega^{\gamma'_0} + \cdots + \omega^{\gamma'_{k'-1}}$. Let also $n, m_0, \dots, m_{k-1}, n', m'_0, \dots, m'_{k'-1}$ be defined as above for each of the two ordinals.

The nodes in $F(\alpha')$ labeled $-$ are the one immediately above the root of $F(\beta')$, and nodes in the interior of $F(\beta')$ or of some $F(\gamma'_i)$ with $i < k'$. Since the node in $F(\alpha)$ immediately above the root of one of the copies of $F(\beta)$ is mapped by f to one of these nodes, the restriction of f to any copy of $F(\beta)$ shows that either (1) $F(\beta) \preceq F(\beta')$, or (2) $F(\beta) \prec F(\beta')$, or (3) $F(\beta) \prec F(\gamma'_i)$. In case (1), by induction hypothesis, we have $\beta \leq \beta'$. Case (2) yields immediately $\beta < \beta'$. In case (3) we have $F(\beta) \leq F(\xi)$ for some ξ with $\varepsilon_\xi \leq \gamma'_i$: since $\gamma'_i \leq \gamma'_0 < \alpha' < \varepsilon_{\beta'+1}$ we have $\xi \leq \beta'$. Moreover, by induction hypothesis, $\beta \leq \xi$ and hence $\varepsilon_\beta \leq \varepsilon_\xi \leq \varepsilon_{\beta'}$.

Thus in every case we have $\beta \leq \beta'$: if $\beta < \beta'$ we are done because $\alpha < \varepsilon_{\beta+1} \leq \varepsilon_{\beta'} \leq \alpha'$. We now assume $\beta = \beta'$ and hence also $n = n'$. We need to show that $\delta \leq \delta'$. If $\delta = 0$ this is obvious. Hence we may assume $\gamma_0 \neq -1$.

Since the root of the copy of $F(\gamma_0)$ in $F(\alpha)$ has height $n + 2 = \text{ht}(F(\beta)) + 5$, this node is not mapped by f to a node in a copy of $F(\beta') = F(\beta)$ in $F(\alpha')$. Since $F(\gamma_0)$ has nodes labeled $-$, f does not map $F(\gamma_0)$ into one of the linear branches of $F(\alpha')$ where all nodes are labeled $+$. Thus there exists $j_0 < k'$ such that the restriction of f to $F(\gamma_0)$ witnesses $F(\gamma_0) \preceq F(\gamma'_{j_0})$. By induction hypothesis, $\gamma_0 \leq \gamma'_{j_0} \leq \gamma'_0$. If at least one of the inequalities is strict then $\gamma_0 < \gamma'_0$ and hence $\delta < \delta'$. Thus we assume $\gamma_0 = \gamma'_0$ and therefore $m_0 = m'_0$.

Repeating the same argument we obtain $j_1 < k'$ such that $F(\gamma_1) \preceq F(\gamma'_{j_1})$ and hence $\gamma_1 \leq \gamma'_{j_1}$. The linear branch with all nodes labeled $+$ which is in the same immediate subtree of $F(\alpha)$ as $F(\gamma_1)$ has a node of height $m_1 + 1 = m_0 + \text{ht}(F(\gamma_0)) + 2$: this node cannot be mapped into the first immediate subtree of $F(\alpha')$, and this implies $j_1 > 0$. If either $\gamma_1 < \gamma'_{j_1}$ or $j_1 > 1$ then $\delta < \delta'$ follows immediately, and otherwise we iterate the argument. If we have to iterate the argument k times, then $k \leq k'$ and $\gamma_i = \gamma'_i$ for every $i < k$, which implies $\delta \leq \delta'$. \square

3.2 The upper bound

There are various ways of proving upper bounds for maximal order type for some well-quasi-orders. We choose one that can be carried out in RCA_0 , but we will not worry about subsystems of second order arithmetic until the next section. The general format for this type of proof was used by Simpson in [24] and by Rathjen and Weiermann in [25]. The main tool is the notion of reification of a quasi-order by an ordinal.

Definition 21 *Let (Q, \leq) be a quasi-ordering. We say that a finite sequence $\langle q_0, \dots, q_k \rangle \in Q^{<\omega}$ is bad if $\forall i < j \ q_i \not\leq q_j$. We define $\text{Bad}(Q) \subseteq Q^{<\omega}$ to be the tree of finite bad sequences from Q .*

Notice that if $q * \sigma \in \text{Bad}(Q)$ the $\sigma(i) \in Q_{\not\leq q}$ for all $i < |\sigma|$, so that $\sigma \in \text{Bad}(Q_{\not\leq q})$.

Note that Q is wqo if and only if $\text{Bad}(Q)$ is a well-founded tree. The results of de Jongh and Parikh ([15]) imply that if Q is a wqo, then $o(Q) = \text{ht}(\text{Bad}(Q))$.

Definition 22 *Let (Q, \leq) be a quasi-ordering. A reification of (Q, \leq) by α is a map $G : \text{Bad}(Q) \rightarrow \alpha + 1$ such that $\sigma \subset \tau \Rightarrow G(\sigma) > G(\tau)$.*

Lemma 23 *If there exists a reification of (Q, \leq) by α , then (Q, \leq) is a wqo and $o(Q) \leq \alpha$.*

We now define some operations on quasi-orderings which preserve wqo. Some of these operations are well-known. See [15] and [16] for more results and proofs.

Disjoint union If (Q_1, \leq_1) and (Q_2, \leq_2) are quasi-orders with $Q_1 \cap Q_2 = \emptyset$ we denote by $Q_1 \sqcup Q_2$ the quasi-order on $Q_1 \cup Q_2$ with no comparabilities between elements of Q_1 and Q_2 .

Finite parts If (Q, \leq) is a quasi-order, let $\mathcal{P}_f(Q)$ be the set of finite subset of Q with quasi order defined by

$$X \leq' Y \iff \forall x \in X \exists y \in Y \ x \leq y.$$

Finite sequences Let $Q^{<\omega}$ be the set of finite sequences of elements of Q with quasi-order \leq' defined by

$$\begin{aligned} \langle x_0, \dots, x_{n-1} \rangle \leq' \langle y_0, \dots, y_{m-1} \rangle &\iff \\ &\exists f : n \rightarrow m \text{ strictly increasing } \forall i < n \ x_i \leq y_{f(i)}. \end{aligned}$$

Finite trees with leaves labeled by Q Let $\mathcal{T}(Q)$ be the set of pairs (T, l_T) where T is a nonempty finite trees and $l_T : L(T) \rightarrow Q$ (recall that $L(T)$ is the set of the leaves of T).

The notion of homomorphism between elements of $\mathcal{T}(Q)$ is an adaptation of the notion of homomorphism between signed trees (definition 10): if $(T, l_T), (T', l_{T'}) \in \mathcal{T}(Q)$ a map $f : T \rightarrow T'$ is a *homomorphism* if

- $\sigma \subset \tau$ implies $f(\sigma) \subset f(\tau)$ for every $\sigma, \tau \in T$;
- if $\sigma \in L(T)$ then $f(\sigma) \in L(T')$ and $l_T(\sigma) \leq l_{T'}(f(\sigma))$.

If there exists a homomorphism from (T, l_T) to $(T', l_{T'})$ we write $(T, l_T) \preceq (T', l_{T'})$.

The maximal order types of the wqo's obtained applying some of these operations has already been computed.

Lemma 24 ([15]) *Let (Q, \leq) , (Q_1, \leq_1) and (Q_2, \leq_2) be well-quasi-orders.*

- (1) *If $Q_1 \cap Q_2 = \emptyset$ then $Q_1 \sqcup Q_2$ is a well-quasi-order and $o(Q_1 \sqcup Q_2) = o(Q_1) \oplus o(Q_2)$.*
- (2) *$(Q^{<\omega}, \leq')$ is a well-quasi-order and $o(Q^{<\omega}) \leq \omega_2(o(Q) + 1) = \omega^{\omega^{o(Q)+1}}$.*

If (Q, \leq) is a well-quasi-order it follows from well-known facts in wqo theory (e.g. Kruskal's Theorem) that $(\mathcal{T}(Q), \preceq)$ is also wqo. Our goal is to bound $o(\mathcal{T}(Q))$ in terms of $o(Q)$ (and this will also give another proof that $(\mathcal{T}(Q), \preceq)$ is a wqo). Also, it is not hard to see that $\mathcal{P}_f(Q) \hookrightarrow Q^{<\omega}$, and therefore $\mathcal{P}_f(Q)$ is a wqo.

To prove that $o(\mathbf{ST}_\omega) \leq \varphi_2(0)$, we simultaneously prove upper bounds for the maximal order type of various well-quasi-ordering.

Definition 25 *Let \mathbb{W} be the smallest set of all quasi-orderings such that:*

- (1) $\mathbf{1} = \{0\} \in \mathbb{W}$;
- (2) $Q, P \in \mathbb{W} \Rightarrow P \sqcup Q \in \mathbb{W}$;
- (3) $Q \in \mathbb{W} \Rightarrow \mathcal{P}_f(Q) \in \mathbb{W}$;
- (4) $Q \in \mathbb{W} \Rightarrow \mathcal{T}(Q) \in \mathbb{W}$.

Let $F : \mathbb{W} \rightarrow \varphi_2(0)$ be defined by

- (1) $F(\mathbf{1}) = 1$;
- (2) $F(P \sqcup Q) = F(P) \oplus F(Q)$;
- (3) $F(\mathcal{P}_f(Q)) = \omega^{F(Q)}$;
- (4) $F(\mathcal{T}(Q)) = \varepsilon_{F(Q)}$.

Lemma 26 *For every $Q \in \mathbb{W}$, $o(Q) \leq F(Q)$.*

PROOF. By Lemma 23, for each $Q \in \mathbb{W}$, it suffices to define a reification $G_Q : \text{Bad}(Q) \rightarrow F(Q) + 1$ of Q . The idea used in the definition of G_Q is the following. On the empty sequence, we define $G_Q(\emptyset) = F(Q)$. Now, if

$\sigma \in \text{Bad}(Q)$ is not empty, we have to define $G_Q(\sigma) < F(Q)$. Suppose that $\sigma = q * \tau$ where $q \in Q$ and $\tau \in \text{Bad}(Q_{\neq q})$. The plan is to embed $Q_{\neq q}$ into some member of \mathbb{W} , and use recursion. So, to each $q \in Q \in \mathbb{W}$ we assign a partial ordering $H_{Q,q} \in \mathbb{W}$ and an embedding

$$h_{Q,q} : Q_{\neq q} \hookrightarrow H_{Q,q}.$$

The assignment of partial ordering has to satisfy $F(H_{Q,q}) < F(Q)$ so that, using recursion on $|\sigma|$, we can define

$$G_Q(q * \tau) = G_{H_{Q,q}}(h_{Q,q}(\tau)),$$

where $h_{Q,q}(\langle q_0, \dots, q_{k-1} \rangle) = \langle h_{Q,q}(q_0), \dots, h_{Q,q}(q_{k-1}) \rangle \in \text{Bad}(H_{Q,q})$. An easy induction on $|\sigma|$ shows that if $\gamma \subset \sigma$, then $G_Q(\gamma) > G_Q(\sigma)$.

All is left is to define $H_{Q,q}$ and $h_{Q,q}$ for each $q \in Q \in \mathbb{W}$. The definition is by recursion on how many operations are needed to build $Q \in \mathbb{W}$. At the same time we will prove by induction that $F(H_{Q,q}) < F(Q)$.

- (1) Case $Q = \mathbf{1} = \{0\}$: In this case we skip the definition of $H_{Q,q}$ and $h_{Q,q}$ and directly define G_Q . $G_{\mathbf{1}}(\emptyset) = 1$, $G_{\mathbf{1}}(\langle 0 \rangle) = 0$.
- (2) Case $Q = P_0 \sqcup P_1$: First consider $q \in P_0$. Let

$$H_{Q,q} = H_{P_0,q} \sqcup P_1$$

and $h_{Q,q}(r) = h_{P_0,q}(r) \in H_{P_0,q}$ if $r \in P_0_{\neq q}$ and $h_{Q,q}(r) = r$ if $r \in P_1$. Note that

$$F(H_{Q,q}) = F(H_{P_0,q}) \oplus F(P_1) < F(P_0) \oplus F(P_1) = F(Q).$$

If $q \in P_1$, we define $H_{Q,q}$ and $h_{Q,q}$ analogously.

- (3) Case $Q = \mathcal{P}_f(P)$: Consider $q = \{q_0, \dots, q_{k-1}\} \subseteq Q$. Let

$$H_{Q,q} = \bigsqcup_{i < k} \mathcal{P}_f(H_{P,q_i}).$$

Given $r = \{r_0, \dots, r_{l-1}\} \in Q_{\neq q}$, there exists $i < k$ such that $\forall j < l$ $q_i \not\subseteq r_j$. Consider the least such i . Then let $h_{Q,q}(r) = \{h_{P,q_i}(r_0), \dots, h_{P,q_i}(r_{l-1})\} \in \mathcal{P}_f(H_{P,q_i})$. Note that

$$F(H_{Q,q}) = \bigoplus_{i < k} \omega^{F(H_{P,q_i})} < \omega^{F(P)} = F(Q).$$

- (4) Case $Q = \mathcal{T}(P)$: We need the following auxiliary operation. Let (J, \leq) be a quasi-order. For every $k \in \omega$ let $\mathcal{T}_k(J) = \{(T, l_T) \in \mathcal{T}(J) \mid \text{ht}(T) \leq k\}$. Let $\{q_n \mid n \in \omega\} \subseteq P$ be such that $\forall q \in P \exists n$ $q \leq q_n$. Let $(T_n, l_n) \in \mathcal{T}_n(P)$ consist of a single branch of n nodes with the last node splitting

into $n + 1$ leaves labeled q_0, \dots, q_n . Let

$$\mathcal{S}^n = \{ (T, l_T) \in \mathcal{T}(P) \mid (T_n, l_n) \not\preceq (T, l_T) \}.$$

For each n we define $H_n \in \mathbb{W}$ with $F(H_n) < F(Q)$ and $h_n : \mathcal{S}^n \hookrightarrow H_n$.

Before defining H_n and h_n let us show how to use them to complete the proof. Notice that the set $\{ (T_n, l_n) \mid n \in \omega \}$ is cofinal in $\mathcal{T}(P)$. In fact given $(T, l_T) \in \mathcal{T}(P)$ pick n so large that $\text{ht}(T) \leq n$ and the finite range of l_T is a subset of $\bigcup_{i \leq n} \{ q \in P \mid q \leq q_i \}$: then $(T, l_T) \preceq (T_n, l_n)$. Therefore, for every $q \in Q$, there exists a least n such that $Q_{\not\preceq q} \subseteq \mathcal{S}^n$, so it suffices to let $H_{Q,q} = H_n$ and $h_{Q,q} = h_n \upharpoonright Q_{\not\preceq q}$, the restriction of h_n to $Q_{\not\preceq q}$.

To construct H_n let us start from

$$\hat{H}_n = \mathcal{T}_n(P \sqcup \bigsqcup_{i \leq n} \mathcal{T}(P_{\not\preceq q_i})).$$

Consider $(T, l_T) \in \mathcal{S}^n$. Notice that if $\sigma \in T$ is such that $|\sigma| = n$ and $\sigma \notin L(T)$ then there exists $i \leq n$ such that $\forall \tau \in L(T) (\tau \supset \sigma \Rightarrow q_i \not\preceq l_T(\tau))$, i.e. $l_T(\tau) \in P_{\not\preceq q_i}$. Now let $\hat{T} = \{ \tau \in T \mid |\tau| \leq n \}$ so that $\text{ht}(\hat{T}) \leq n$. Define also $l_{\hat{T}} : L(\hat{T}) \rightarrow P \sqcup \bigsqcup_{i < n} \mathcal{T}(P_{\not\preceq q_i})$ by

$$l_{\hat{T}}(\sigma) = \begin{cases} l_T(\sigma), & \text{if } |\sigma| < n; \\ \text{the subtree of } (T, l_T) \text{ rooted at } \sigma, & \text{if } |\sigma| = n. \end{cases}$$

In the first case $\sigma \in L(T)$ and $l_{\hat{T}}(\sigma) \in P$, while in the second case $l_{\hat{T}}(\sigma) \in \mathcal{T}(P_{\not\preceq q_i})$ for some $i \leq n$ by the observation above.

Set $\hat{h}_n(T, l_T) = (\hat{T}, l_{\hat{T}})$. It is not hard to check that $\hat{h}_n : \mathcal{S}^n \hookrightarrow \hat{H}_n$.

Since $\hat{H}_n \notin \mathbb{W}$, we have to modify our construction a bit. Let $J = P \sqcup \bigsqcup_{i \leq n} \mathcal{T}(H_{P, q_i}) \in \mathbb{W}$ and $j : P \sqcup \bigsqcup_{i \leq n} \mathcal{T}(P_{\not\preceq q_i}) \hookrightarrow J$ be the map defined in the obvious way using the h_{P, q_i} 's. We can then extend j to $j : \hat{H}_n \hookrightarrow \mathcal{T}_n(J)$.

Define $f : \mathcal{T}_n(J) \rightarrow J \sqcup \mathcal{P}_f(\mathcal{T}_{n-1}(J))$ as follows:

$$f(T, l_T) = \begin{cases} l_T(\emptyset) & \text{if } T = \{\emptyset\}; \\ \{ (T_n, l_{T_n}) \mid \langle n \rangle \in T \} & \text{otherwise.} \end{cases}$$

Checking that f witnesses $\mathcal{T}_n(J) \hookrightarrow J \sqcup \mathcal{P}_f(\mathcal{T}_{n-1}(J))$ is straightforward. Iterating n times f we obtain

$$f^n : \mathcal{T}_n(J) \hookrightarrow J \sqcup \mathcal{P}_f(J \sqcup \mathcal{P}_f(\dots (J \sqcup \mathcal{P}_f(J)) \dots))$$

Let

$$H_n = J \sqcup \mathcal{P}_f(J \sqcup \mathcal{P}_f(\dots (J \sqcup \mathcal{P}_f(J)) \dots))$$

and $h_n = f^n \circ j \circ \hat{h}_n : \mathcal{S}^n \hookrightarrow H_n$.

Since $F(Q) = \varepsilon_{F(P)}$ is closed under \oplus and $F(H_{P,q_i}) < F(P)$, we have $F(J) = F(P) \oplus \bigoplus_{i \leq n} \varepsilon_{F(H_{P,q_i})} < \varepsilon_{F(P)}$. Since $F(Q)$ is also closed under exponentiation with base ω , we obtain $F(H_{Q,q}) < F(Q)$. \square

Proposition 27 $o(\mathbf{ST}_\omega) \leq \varphi_2(0)$.

PROOF. We employ the same technique of the proof of Lemma 26: we keep the same notation and define, for $q \in Q = \mathbf{ST}_\omega$, $H_{Q,q} \in \mathbb{W}$ with $F(H_{Q,q}) < \varphi_2(0)$ and $h_{Q,q} : Q \not\leq q \hookrightarrow H_{Q,q}$. As in the preceding proof this yields a reification of \mathbf{ST}_ω by $\varphi_2(0)$, and by Lemma 23 the proof is complete.

For every n let $(T_n, s_n), (T_n, \hat{s}_n) \in \mathbf{ST}_\omega$ be defined as follows: T_n consists of a single branch of $n + 1$ nodes; s_n labels the nodes of T_n with even length $+$, and the nodes with odd length $-$; \hat{s}_n acts dually, labeling nodes of T_n with even length $-$, and nodes with odd length $+$. Let $\mathbf{ST}_\omega^n = \{(T, s_T) \in \mathbf{ST}_\omega \mid (T_n, s_n) \not\leq (T, s_T)\}$ and similarly $\widehat{\mathbf{ST}}_\omega^n = \{(T, s_T) \in \mathbf{ST}_\omega \mid (T_n, \hat{s}_n) \not\leq (T, s_T)\}$. It is obvious that $o(\mathbf{ST}_\omega^n) = o(\widehat{\mathbf{ST}}_\omega^n)$.

The set $\{(T_n, s_n) \mid n \in \omega\}$ is cofinal in \mathbf{ST}_ω : if $\text{ht}(T) = k$ then $(T, s_T) \preceq (T_{2k+1}, s_{2k+1})$. Therefore, for every $q = (T, s_T) \in \mathbf{ST}_\omega$ there exists n such that

$$Q \not\leq q \subseteq \mathbf{ST}_\omega^n.$$

For every n we define $H_n \in \mathbb{W}$ with $F(H_n) < \varphi_2(0)$ and $h_n : \mathcal{S}^n \hookrightarrow H_n$. Once this is done the proof is completed exactly as in Case (4) of the previous proof.

When $n = 0$ notice that an element of \mathbf{ST}_ω^0 is a tree with all nodes labeled $-$. Therefore \mathbf{ST}_ω^0 is order isomorphic to $\mathcal{T}(\mathbf{1}) \in \mathbb{W}$. Thus we can set $H_0 = \mathcal{T}(\mathbf{1})$ (so that $F(H_0) = \varepsilon_1 < \varphi_2(0)$) and let h_0 be the order isomorphism mentioned above.

Fix $(T, s_T) \in \mathbf{ST}_\omega^{n+1}$. Since $(T_n, s_n) \not\leq (T, s_T)$, T does not contain a sequence $\sigma_0 \subset \sigma_1 \subset \dots \subset \sigma_n$ with $s_T(\sigma_i) = +$ for even i , and $s_T(\sigma_i) = -$ for odd i . Therefore if σ_0 is such that $s_T(\sigma_0) = +$ the subtree of (T, s_T) rooted at σ_0 belongs to $\widehat{\mathbf{ST}}_\omega^n$ (because it contains no sequence $\sigma_1 \subset \sigma_2 \subset \dots \subset \sigma_n$ with $s_T(\sigma_i) = +$ for even i , and $s_T(\sigma_i) = -$ for odd i). Let $U_T = \{\sigma \in T \mid \forall \tau \subset \sigma s_T(\tau) = -\}$. U_T is a tree and if $\sigma \in L(U_T)$ then $\sigma \in L(T)$ or $s_T(\sigma) = +$. Let $l_T(\sigma)$ be the subtrees of (T, s_T) rooted at σ . By the observation above $l_T(\sigma) \in \widehat{\mathbf{ST}}_\omega^n$ when $s_T(\sigma) = +$. When $s_T(\sigma) = -$ then $\sigma \in L(T)$ and $l_T(\sigma)$ is the tree consisting of the root labeled $-$: if $n > 0$ we have $l_T(\sigma) \in \widehat{\mathbf{ST}}_\omega^n$ also in this case, while when $n = 0$ we identify this tree with the lone element of $\mathbf{1}$. Therefore $l_T : L(U_T) \rightarrow \widehat{\mathbf{ST}}_\omega^n$ when $n > 0$ and $l_T : L(U_T) \rightarrow \mathbf{1} \sqcup \widehat{\mathbf{ST}}_\omega^n$ when $n = 0$.

If we set $f_n(T, s_T) = (U_T, l_T)$ we have a function $f_n : \mathbf{ST}_\omega^{n+1} \rightarrow \mathcal{T}(\widehat{\mathbf{ST}}_\omega^n)$ or $f_n : \mathbf{ST}_\omega^{n+1} \rightarrow \mathcal{T}(\mathbf{1} \sqcup \widehat{\mathbf{ST}}_\omega^n)$, depending on whether $n > 0$ or not. It is easy to check that f_n witnesses $\mathbf{ST}_\omega^{n+1} \hookrightarrow \mathcal{T}(\widehat{\mathbf{ST}}_\omega^n)$ (or $\mathbf{ST}_\omega^{n+1} \hookrightarrow \mathcal{T}(\mathbf{1} \sqcup \widehat{\mathbf{ST}}_\omega^n)$). Then

$$f_0 \circ \cdots \circ f_n : \mathbf{ST}_\omega^{n+1} \hookrightarrow Q_n.$$

where $Q_n = \mathcal{T}(\mathcal{T}(\mathcal{T}(\dots(\mathbf{1} \sqcup \mathcal{T}(\mathbf{1}))\dots)))$ with $n + 1$ occurrences of \mathcal{T} . Since $Q_n \in \mathbb{W}$ and $F(Q_n) = \varepsilon_{\varepsilon \dots \varepsilon_{1+1}} < \varphi_2(0)$ the proof is complete. \square

3.3 Maximal order type of \mathbf{LO}_ω

We are now ready to prove Theorem 4 that says that $o(\mathbf{LO}_\omega, \preceq) = \varphi_2(0)$. Since $o(\mathbf{ST}_\omega) = o(\mathbf{ILO}_\omega) \leq o(\mathbf{LO}_\omega)$, by Proposition 20 we have that $o(\mathbf{LO}_\omega, \preceq) \geq \varphi_2(0)$. The other inequality requires the following observation.

Let $L_n = \mathbb{Z}^n \cdot \omega$. Note that for every $L \in \mathbf{LO}_\omega$, there exists n such that $L \preceq L_n$. Therefore $o(\mathbf{LO}_\omega) = \sup_n o((\mathbf{LO}_\omega)_{\not\preceq L_n})$. Every $L \in (\mathbf{LO}_\omega)_{\not\preceq L_n}$ can be decomposed as a finite sum of indecomposable linear orders, say $J_0 + \cdots + J_k$. Since L_n is indecomposable, we have that for every $i \leq k$, $J_i \in (\mathbf{ILO}_\omega)_{\not\preceq L_n}$. This gives us an embedding $(\mathbf{LO}_\omega)_{\not\preceq L_n} \hookrightarrow ((\mathbf{ILO}_\omega)_{\not\preceq L_n})^{<\omega}$. Since $o((\mathbf{ILO}_\omega)_{\not\preceq L_n}) < \varphi_2(0)$, we have that $o(((\mathbf{ILO}_\omega)_{\not\preceq L_n})^{<\omega}) < \varphi_2(0)$. It follows that $o((\mathbf{LO}_\omega)_{\not\preceq L_n}) < \varphi_2(0)$ and hence $o(\mathbf{LO}_\omega, \preceq) \leq \varphi_2(0)$.

4 Subsystems of second order arithmetic

We refer the reader to [5] for background information on subsystems of second order arithmetic. From now on, when we write the name of a subsystems of second order arithmetic in parenthesis at the beginning of a definition or of a theorem, it means that the definition or the theorem is being carried out in that system.

Recall that \mathbf{RCA}_0 is the basis system: it consists of axioms stating that the natural numbers form an ordered semi-ring, plus the axiom schemes of Σ_1^0 -induction and Δ_1^0 -comprehension.

Definition 28 (\mathbf{RCA}_0) *We say that $X = Y'$ if $\forall e(e \in X \iff \{e\}^Y(e) \downarrow)$, where $\{e\}^Y$ is the e -th Turing functional with oracle Y .*

We say that $X = Y^{(k)}$ if $X = \langle X_0, \dots, X_k \rangle$, $X_0 = Y$ and for every $n < k$, $X_{n+1} = (X_n)'$. We say that $X = Y^{(\omega)}$ if $X = \langle X_0, \dots \rangle$, $X_0 = Y$ and for every n , $X_{n+1} = (X_n)'$.

Recall that ACA_0 (the system obtained by adding arithmetic comprehension to RCA_0) is equivalent to RCA_0 plus the statement $\forall Y \exists X X = Y'$.

Definition 29 ACA'_0 is RCA_0 plus the statement $\forall Y \forall k \exists X X = Y^{(k)}$. ACA_0^+ is RCA_0 plus the statement $\forall Y \exists X X = Y^{(\omega)}$.

As in the proof of [9, Lemma 3.4] we define Hausdorff rank using powers of \mathbb{Z} . Since we are interested only in finite Hausdorff rank we do not need any transfinite recursion.

Remark 30 (RCA_0) *A linear order L has finite Hausdorff rank if $L \preceq \mathbb{Z}^n$ for some n .*

The proof-theoretic ordinals of ACA_0 , ACA'_0 and ACA_0^+ are ε_0 , ε_ω and $\varphi_2(0)$ respectively. Since ACA_0 is conservative over Peano Arithmetic the first result follows from Gentzen's analysis of that theory. The result for ACA'_0 is due originally to Jäger (unpublished notes) and a proof appears in [26]; recently Michael Rathjen gave a new proof sketch: details will appear in the forthcoming Ph.d. thesis of Bahareh Afshari at the University of Leeds. The computation of the proof-theoretic ordinal of ACA_0^+ is due to Rathjen ([23]).

In particular, the consistency of ACA_0^+ is equivalent to the statement saying that $\varphi_2(0)$ is well-ordered, and hence this statement cannot be proved in ACA_0^+ . To make this statement more precise, we now explain how “ $\varphi_2(0)$ is well-ordered” is expressed within RCA_0 .

Let us introduce a system of notations for the ordinals below $\varphi_2(0)$. The idea is to formalize the way of writing an ordinal $\alpha < \varphi_2(0)$ used in the proof of Proposition 20: with the convention $\varepsilon_{-1} = \omega^{-1} = 0$ we have

$$\alpha = \varepsilon_\beta + \omega^{\gamma_0} + \cdots + \omega^{\gamma_{k-1}}$$

with $k > 0$, $\beta < \alpha < \varepsilon_{\beta+1}$, and $\alpha > \gamma_0 \geq \cdots \geq \gamma_{k-1}$. We can then write β , and $\gamma_0, \dots, \gamma_{k-1}$ the same way, and this process will eventually stop when α is written in *normal form*.

The formal definition is the following.

Definition 31 (RCA_0) *Let \mathcal{L} be the language consisting of the constant -1 , the unary operations $\omega^{(\cdot)}$ and $\varepsilon_{(\cdot)}$, and the binary operation $(\cdot) + (\cdot)$.*

We simultaneously define a set of terms Φ of \mathcal{L} and an ordering on Φ as follows:

- $-1 \in \Phi$ and $-1 \leq t$ for every $t \in \Phi$;
- $\varepsilon_t + \omega^{t_0} + \cdots + \omega^{t_{k-1}} \in \Phi$ whenever $k > 0$, $t, t_0, \dots, t_{k-1} \in \Phi$, and $\varepsilon_t + \omega^{-1} \geq t_0 \geq \cdots \geq t_{k-1}$. If $t_i = -1$ then we require $k = 1$ and $i = 0$;

•

$$\varepsilon_{t'} + \omega^{t'_0} + \cdots + \omega^{t'_{k'-1}} \leq \varepsilon_t + \omega^{t_0} + \cdots + \omega^{t_{k-1}}$$

if and only if either $t' < t$ or $t' = t$ and (either there exists i such that $t'_i \neq t_i$, and for the first such i , $t'_i < t_i$, or $t'_i = t_i$ for every $i < k'$ and $k' \leq k$).

Notice that the ordering (Φ, \leq) is primitive recursive. When we write a term in Φ , we are thinking of an ordinal below $\varphi_2(0)$. Conversely, as in Proposition 20 we can prove that for every ordinal below $\varphi_2(0)$ there is a term representing it. It is straightforward to show that the ordering relation between elements of Φ is isomorphic to $\varphi_2(0)$.

When we refer to the statement “ $\varphi_2(0)$ is well-ordered” in the language of second order arithmetic, we are actually saying “ (Φ, \leq) is well-ordered”. In terms of reverse mathematics, RCA_0 suffices to show that (Φ, \leq) is a linear ordering. It is much harder to prove that is a well-ordering (as noticed above, ACA_0^+ does not suffice).

Within RCA_0 it is straightforward to define the set ST_ω and the notion of homomorphism between finite signed trees. We can also define ST , LO_ω and ILO_ω , but notice that these are classes and not sets. Within RCA_0 we also define the function lin and prove the following Lemma ([9, Proposition 2.13]), showing that Lemma 14.b is provable in ACA_0 .

Lemma 32 (ACA_0) *Let $T, T' \in \text{ST}$. Then*

$$T \preceq T' \iff \text{lin}(T) \preceq \text{lin}(T').$$

When we say within RCA_0 that a quasi-ordering (Q, \leq) is wqo we mean that there exists no infinite bad sequences (i.e. for every $f : \omega \rightarrow Q$ there exist $i < j$ with $f(i) \leq f(j)$): in [14] it is proved that in RCA_0 this implies, but is not equivalent to, the statement that (Q, \leq) has no infinite descending chains and antichains. When we deal with a quasi-ordering which is a class some simple coding formalizes the same definition within RCA_0 : this leads to the statement of FRA and FRA_ω in RCA_0 .

5 Upper bound for FRA_ω

In this section we prove Theorem 5, i.e. that ACA_0^+ together with the statement “ $\varphi_2(0)$ is well-ordered” suffices to prove FRA_ω .

Proposition 33 (RCA_0) *The following are equivalent:*

- (i) ST_ω is well-quasi-ordered by \preceq ;
- (ii) $\varphi_2(0)$ is well-ordered.

PROOF. The proof consists of formalizing in RCA_0 the proofs of Section 3.

For the implication (i) \Rightarrow (ii), we need to look at the proof of Proposition 20. There we defined a function $F : \varphi_2(0) \rightarrow \text{ST}_\omega$ witnessing $\varphi_2(0) \hookrightarrow \text{ST}_\omega$. The definition of F and the proof that it satisfies the required properties are by Δ_0^0 -induction on the length of the normal form of the ordinal notations below $\varphi_2(0)$ and hence they can be easily be carried out in RCA_0 . Then, if we had an infinite descending sequence in $\varphi_2(0)$, we could map it through F and get an infinite bad sequence through ST_ω .

For the implication (ii) \Rightarrow (i), we first need to look at the proof of Lemma 26. We can easily name all the partial orderings that are in \mathbb{W} . So, in RCA_0 we can work with \mathbb{W} as a set, by using a set of names for its elements. Then, we can think of $F : \mathbb{W} \rightarrow \varphi_2(0)$ as a second order object which is defined by recursion. The definition of $H_{Q,q}$ and $h_{Q,q}$ is computably uniform in $q \in Q \in \mathbb{W}$ and is done by recursion on number of operations needed to define $Q \in \mathbb{W}$ (i.e. on the complexity of the name for Q). $G_Q(\sigma)$ is also computably uniform in $Q \in \mathbb{W}$ and is defined by recursion on $|\sigma|$. By induction on number of operations needed to define $Q \in \mathbb{W}$ we can prove that for every $q \in Q \in \mathbb{W}$ we have $F(H_{Q,q}) < F(Q)$. Then, in RCA_0 , we can prove that $\sigma \subset \tau \Rightarrow G_Q(\sigma) > G_Q(\tau)$. Therefore, for all $Q \in \mathbb{W}$, if $F(Q)$ is well-ordered $\text{Bad}(Q)$ is well-founded and hence Q is well-quasi-ordered.

The argument applies also to ST_ω (proof of Proposition 27) and shows that if $\varphi_2(0)$ is well-ordered then ST_ω is well-quasi-ordered. \square

Lemma 34 (ACA_0^+) *Given a sequence $\{L_i \mid i \in \omega\}$ of linear orderings of finite rank, there is a sequence $\{\langle T_{i,1}, \dots, T_{i,k_i} \rangle \mid i \in \omega\}$ of finite sequences of finite signed trees such that for every i*

$$L_i \sim \text{lin}(T_{i,1}) + \dots + \text{lin}(T_{i,k_i}).$$

PROOF. In [9, Lemma 3.4] it is proved that if ST_α is well-quasi-ordered, then every computable linear order L of rank α is equimorphic to a finite sum of linear orders of the form $\text{lin}(T)$ and of rank $\leq \alpha$ via an equimorphism computable in $0^{(2(\alpha+1)^2)}$.

When $\alpha < \omega$, RCA_0 proves that ST_α is well-quasi-ordered just because it is finite. Noting that the proof of Lemma 3.4 in [9] is uniform, we obtain that given a computable sequence $\{L_i \mid i \in \omega\}$ of linear orders of finite rank there

is a sequence $\{ \langle T_{i,1}, \dots, T_{i,k_i} \rangle \mid i \in \omega \}$ computable in $0^{(\omega)}$ which satisfies the statement of the Lemma.

If the sequence we are given is not computable, it suffices to relativize the proof, and ACA_0^+ suffices. \square

Proposition 35 (ACA_0^+) *If ST_ω is well-quasi-ordered by \preceq then FRA_ω holds.*

PROOF. Consider an infinite sequence $\{ L_i \mid i \in \omega \}$ of linear orderings of finite rank: we want to show it is not a bad sequence. By Lemma 34 there exists a sequence $\{ \langle T_{i,1}, \dots, T_{i,k_i} \rangle \mid i \in \omega \}$ of members of $\text{ST}_\omega^{<\omega}$ such that for every i

$$L_i \sim \text{lin}(T_{i,1}) + \dots + \text{lin}(T_{i,k_i}).$$

By Higman's theorem, which is provable in ACA_0 ([24,27,6]), we have that $\text{ST}_\omega^{<\omega}$ is well-quasi-ordered by \preceq . Therefore, there exists $i < j$ such that $\langle T_{i,1}, \dots, T_{i,k_i} \rangle \preceq \langle T_{j,1}, \dots, T_{j,k_j} \rangle$. By Lemma 32 we have

$$L_i \sim \text{lin}(T_{i,1}) + \dots + \text{lin}(T_{i,k_i}) \preceq \text{lin}(T_{j,1}) + \dots + \text{lin}(T_{j,k_j}) \sim L_j.$$

Thus $\{ L_i \mid i \in \omega \}$ is not a bad sequence. \square

Propositions 33 and 35 obviously imply Theorem 5.

6 Lower bound for FRA_ω

In this section we prove Theorem 6, i.e. that $\text{RCA}_0 + \text{FRA}$ implies ACA'_0 plus the statement “ $\varphi_2(0)$ is well-ordered”.

We start by noticing that some of the intermediate steps in Shore's proof [7] that FRA implies ATR_0 show that $\text{RCA}_0 \vdash \text{FRA}_\omega \Rightarrow \text{ACA}_0$. Indeed the proofs of [7, Theorem 3.1] (establishing that $\text{RCA}_0 \vdash \text{FRA} \Rightarrow \Sigma_2^0$ -induction) and of [7, Theorem 1.1] (showing that $\text{RCA}_0 + \Sigma_2^0$ -induction $\vdash \text{FRA} \Rightarrow \text{ACA}_0$) make use only of linear orders (actually, well-orders) of finite Hausdorff rank. We thus have:

Proposition 36 (RCA_0) *FRA_ω implies ACA_0 .*

We will now build on Shore's ideas to prove the following Proposition.

Proposition 37 (ACA_0) *FRA_ω implies ACA'_0 .*

PROOF. Fix k . We want to show that $0^{(k)}$ exists. By relativizing the proof as usual, we will get that for every Y , $Y^{(k)}$ exists.

First, for each $n \leq k$, we construct a sequence of linear orders $\{A_{j,n} \mid j \in \omega\}$, such that, if $0^{(n)}$ exists and $h : A_{j,n} \rightarrow A_{l,n}$ is an embedding for some $j < l$, then h computes $0^{(n+1)}$. Unfortunately, this induction step is not enough to prove ACA'_0 as Σ^1_1 -induction would be required. So we will do the following. For each j we define

$$A_j = \sum_{n \leq k} (\omega^* + \omega + A_{j,n}).$$

Then, we will show that if we have an embedding $A_j \preceq A_l$ for some $j < l$, we can recover embeddings $h_n : A_{j,n} \rightarrow A_{l,n}$ for each $n \leq k$. We use these embeddings together to define a sequence of sets X_0, X_1, \dots, X_k . Then, using arithmetic induction on $n \leq k$, we show that for every n , $X_{n+1} \geq_T (X_n)'$, and thus $\langle X_0, \dots, X_k \rangle$ witnesses the existence of $0^{(k)}$.

The idea of the construction is as follows. We want to define $\{A_{j,n} \mid j \in \omega\}$ such that any embedding $h_n : A_{j,n} \rightarrow A_{l,n}$ for $j < l$ produces a function that dominates the set of true stages for the enumeration of $0^{(n+1)}$ from $0^{(n)}$. (Recall that t is a true stage for $f : \omega \rightarrow \omega$ if $\forall s > t \ f(s) > f(t)$). As noticed in [7], RCA_0 suffices to prove that for any one-to-one f there exists infinitely many true stages, while $\text{RCA}_0 + \Sigma^0_2$ -induction proves that for any one-to-one f and any i there exist the i -th true stage for f .) Let t_i be the i -th true stage for the enumeration of $0^{(n+1)}$ from $0^{(n)}$. Given j , we will let $A_{n,j,t_{j+i}}$ be isomorphic to $\omega^{n+1} + 1$ and, for s not of the form t_{j+i} , $A_{j,n,s}$ be isomorphic to $\omega^n \cdot p + 1$ for some $p \in \omega$. Then, we define

$$A_{j,n} = \sum_s A_{j,n,s}.$$

Working in ACA_0 , we have to be careful in defining the $A_{j,n,s}$'s. We will use sequences of linear orders $\{L_{n,e}^\Sigma \mid n \in \omega\}$ and $\{L_{n,e}^\Pi \mid n \in \omega\}$ (the main idea for this construction is taken from a similar construction in [22, Section 5]): the reader should keep in mind that we would like to have, for every $n, e \in \omega$,

$$L_{n,e}^\Sigma \cong \begin{cases} \omega^{n+1} & \text{if } e \in 0^{(n+1)} \\ \omega^n & \text{if } e \notin 0^{(n+1)} \end{cases} \quad (6.1)$$

and

$$L_{n,e}^\Pi \cong \begin{cases} \omega^n \cdot p \text{ for some } p \in \omega, p > 0 & \text{if } e \in 0^{(n+1)} \\ \omega^{n+1} & \text{if } e \notin 0^{(n+1)}. \end{cases} \quad (6.2)$$

For each n denote by Ω^{n+1} the natural presentation of the linear order of order type ω^{n+1} consisting of the ordered n -tuples of natural numbers ordered

antilexicographically. $L_{n,e}^\Sigma$ and $L_{n,e}^\Pi$ will be subsets of Ω^{n+1} under this order. Since ACA_0 proves that Ω^{n+1} is a well-order for each n , we have that $L_{n,e}^\Sigma$ and $L_{n,e}^\Pi$ are well-orders.

We define $L_{n,e}^\Sigma$ and $L_{n,e}^\Pi$ by recursion on n . Recall the existence of a computable function f such that for all $n \in \omega$

- (a) for all $e \in \omega$, $e \in 0^{(n+1)}$ if and only if there exists x such that $f(n, e, x) \notin 0^{(n)}$;
- (b) for all $e, x, y \in \omega$, if $x < y$ and $f(n, e, x) \notin 0^{(n)}$ then $f(n, e, y) \notin 0^{(n)}$.

The function f can be defined in RCA_0 , and for each n , if $0^{(n)}$ exists then RCA_0 proves (a) and (b) above (here, since $0^{(n+1)}$ might not exist, $e \in 0^{(n+1)}$ is to be understood as an abbreviation for $\{e\}^{0^{(n)}}(e) \downarrow$).

Let $L_{0,e}^\Sigma$ (resp. $L_{0,e}^\Pi$) be the set $\{0\} \cup \{s \mid \{e\}_s(e) \downarrow\}$ (resp. $\{0\} \cup \{s \mid \{e\}_s(e) \uparrow\}$), which is indeed a subset of Ω^1 . If we have already defined $L_{n,e}^\Sigma$ and $L_{n,e}^\Pi$ for every e , let

$$\begin{aligned} L_{n+1,e}^\Sigma &= \{(\vec{y}, x) \in \Omega^{n+2} \mid \vec{y} \in L_{n,f(n,e,x)}^\Pi\} \quad \text{and} \\ L_{n+1,e}^\Pi &= \{(\vec{y}, x) \in \Omega^{n+2} \mid \vec{y} \in L_{n,f(n,e,x)}^\Sigma\}, \end{aligned}$$

so that the order types of $L_{n+1,e}^\Sigma$ and $L_{n+1,e}^\Pi$ are respectively

$$\sum_{x \in \omega} L_{n,f(n,e,x)}^\Pi \quad \text{and} \quad \sum_{x \in \omega} L_{n,f(n,e,x)}^\Sigma.$$

Claim 38 *For every $n, e \in \omega$, $\text{rk}_H(L_{n,e}^\Sigma) \leq n + 1$ and $\text{rk}_H(L_{n,e}^\Pi) \leq n + 1$.*

PROOF. This is obvious, since $L_{n,e}^\Sigma \preceq \Omega^{n+1} \preceq \mathbb{Z}^{n+1}$ and the same for $L_{n,e}^\Pi$. \square

Claim 39 *Let $m \in \omega$ be such that $0^{(m)}$ exists. Then for every $n \leq m$ and $e \in \omega$ (6.1) and (6.2) hold.*

PROOF. Fix m and assume $0^{(m)}$ exists. We use induction on $n \leq m$ to prove an apparently stronger statement: for every e , $L_{n,e}^\Sigma$ (resp. $L_{n,e}^\Pi$) satisfies (6.1) (resp. (6.2)) with an isomorphism computable in $0^{(n)}$. This statement is arithmetical, so we have enough induction in ACA_0 to carry out the proof.

The base and the induction steps are both straightforward using that f satisfies (a) and (b). \square

When we do not know that $0^{(n)}$ exists, we know very little about the order type of $L_{n,e}^\Sigma$ and $L_{n,e}^\Pi$. However, as noticed above, we do know they are well-ordered.

Now we are ready to formally define the $A_{j,n,s}$'s, and then the $A_{j,n}$'s and the A_j 's. Notice that “ s is a true stage for f and there are at least $j - 1$ true stages for f smaller than s ” is a Π_1^0 formula. Hence there exists a computable function g such that for each n, s, j , $g(n, j, s) \notin 0^{(n+1)}$ if and only if s is the $(j + i)$ -th true stage for the enumeration of $0^{(n+1)}$ from $0^{(n)}$, for some i . Define $A_{j,n,s} = L_{n,g(n,j,s)}^\Pi + 1$. We may assume that the domains of the $A_{j,n,s}$'s are pairwise disjoint and denote by $m_{j,n,s}$ the first element of $A_{j,n,s}$.

As announced let $A_{j,n} = \sum_s A_{j,n,s}$ and $A_j = \sum_{n \leq k} (\omega^* + \omega + A_{j,n})$. By Claim 38 we have $\text{rk}_H(A_{j,n}) \leq n + 2$ and hence $\text{rk}_H(A_j) \leq k + 3$ for every j . Let $\pi_{j,n} : A_{j,n} \rightarrow \omega$ be the function which assigns to every $z \in A_{j,n}$ the unique s such that $z \in A_{j,n,s}$.

By FRA_ω there exists $j < l$ and an embedding $h : A_j \rightarrow A_l$. For $n \leq k$ let h_n be the restriction of h to $A_{j,n}$.

Claim 40 *For each $n \leq k$, h_n maps $A_{j,n}$ into $A_{l,n}$.*

PROOF. For every j we have $\omega^* \cdot (k + 1) \preceq A_j$, while the fact that each $L_{n,e}^\Pi$ is well-ordered implies that the $A_{j,n}$'s are well-ordered and hence $\omega^* \cdot (k + 2) \not\preceq A_j$. In other words, both A_j and A_l contain exactly $k + 1$ copies of ω^* .

Thus, for each $n \leq k$, an initial segment of the copy of ω^* in A_j immediately preceding $\omega + A_{j,n}$ is mapped by h coinitially to the copy of ω^* in A_l which precedes immediately $\omega + A_{l,n}$. Therefore the copy of ω immediately preceding $A_{j,n}$ is mapped to $\omega^* + \omega + A_{l,n}$, and this implies that h_n maps $A_{j,n}$ into $A_{l,n}$. \square

Claim 41 *Suppose $0^{(n)}$ exists. If $j < l$ and $h_n : A_{j,n} \rightarrow A_{l,n}$ is an embedding, then $0^{(n+1)} \leq_T h_n \oplus 0^{(n)}$.*

PROOF. Recursively in h_n we define $k : \omega \rightarrow \omega$ by setting $k(0) = t_j$ and $k(x + 1) = \pi_{l,n}(h_n(m_{j,n,k(x)}))$.

We prove by induction that $k(x) \geq t_{j+x}$ for every $x \in \omega$. The case $x = 0$ is trivial. Assuming $k(x) \geq t_{j+x}$ notice that $\omega^{n+1} \cdot (x + 1)$ embeds into the initial segment of $A_{j,n}$ bounded by $m_{j,n,k(x)}$, and hence also into the initial segment of $A_{l,n}$ bounded by $h_n(m_{j,n,k(x)})$. This implies that $t_{l+x} \leq \pi_{l,n}(h_n(m_{j,n,k(x)})) = k(x + 1)$. Since $j < l$ we have $t_{j+x+1} \leq t_{l+x}$ and the induction step is complete.

Hence $x \in 0^{(n+1)}$ if and only if it appears among the first $k(x)$ elements of the enumeration of $0^{(n+1)}$ from $0^{(n)}$. Thus $0^{(n+1)} \leq_T h_n \oplus 0^{(n)}$. \square

Let $X_0 = \emptyset$, and, for $n < k$, $X_{n+1} = h_n \oplus X_n$. Using the previous claim, we can prove by arithmetic induction on n , that for every $n \leq k$ there exist an index e_n such that $\{e_n\}^{X_n} = 0^{(n)}$. Therefore $\langle \{e_0\}^{X_0}, \dots, \{e_k\}^{X_k} \rangle$ witnesses the existence of $0^{(k)}$. \square

Remark 42 *Proposition 37 cannot be strengthened by replacing ACA'_0 with ACA_0^+ . In fact FRA_ω holds in the ω -model consisting of all arithmetic sets, where ACA_0^+ fails. The reason is that if we are given a sequence $\{L_n : n \in \omega\}$ of arithmetic linear orderings of finite Hausdorff rank, then there exists $i < j$ such that $L_i \preceq L_j$. At first, we do not know that this embedding is in the model of arithmetic sets. But it follows from the proofs of [9, Lemmas 3.4 and 2.5] that, if there is an embedding between two linear orderings of finite Hausdorff rank, then the embedding is arithmetic in the linear orderings and hence belongs to the model.*

Proposition 43 (ACA_0) FRA_ω implies that ST_ω is well-quasi-ordered by \preceq .

PROOF. Consider an infinite sequence $\{(T_i, s_{T_i}) \mid i \in \omega\}$ of elements of ST_ω ; we want to show that it is not a bad sequence. The sequence $\{\text{lin}(T_i, s_{T_i}) \mid i \in \omega\}$ is a sequence in LO_ω . By FRA_ω , there exists $i < j$ such that $\text{lin}(T_i, s_{T_i}) \preceq \text{lin}(T_j, s_{T_j})$. By Lemma 32, we have $(T_i, s_{T_i}) \preceq (T_j, s_{T_j})$. \square

Propositions 36, 37, 43, and 33 imply Theorem 6.

References

- [1] R. Fraïssé, Sur la comparaison des types d'ordres, C. R. Acad. Sci. Paris 226 (1948) 1330–1331.
- [2] R. Laver, On Fraïssé's order type conjecture, Ann. of Math. (2) 93 (1971) 89–111.
- [3] C. S. J. A. Nash-Williams, On better-quasi-ordering transfinite sequences, Proc. Cambridge Philos. Soc. 64 (1968) 273–290.
- [4] S. G. Simpson, Bqo-theory and Fraïssé's conjecture, Chapter 9 of [28] (1985).
- [5] S. G. Simpson, Subsystems of second order arithmetic, Springer-Verlag, Berlin, 1999.
- [6] A. Marcone, Wqo and bqo theory in subsystems of second order arithmetic, in: S. G. Simpson (Ed.), Reverse mathematics 2001, Lecture Notes in Logic, Association for Symbolic Logic, La Jolla, CA, 2005, pp. 303–330.

- [7] R. A. Shore, On the strength of Fraïssé’s conjecture, in: J. N. Crossley, J. B. Remmel, R. A. Shore, M. E. Sweedler (Eds.), *Logical methods*, Birkhäuser, Boston, MA, 1993, pp. 782–813.
- [8] A. Marcone, On the logical strength of Nash-Williams’ theorem on transfinite sequences, in: W. Hodges, M. Hyland, C. Steinhorn, J. Truss (Eds.), *Logic: from foundations to applications*, Oxford Sci. Publ., Oxford Univ. Press, New York, 1996, pp. 327–351.
- [9] A. Montalbán, Equivalence between Fraïssé’s conjecture and Jullien’s theorem, *Ann. Pure Appl. Logic* 139 (1-3) (2006) 1–42.
- [10] F. Hausdorff, Grundzüge einer Theorie der geordneten Mengen, *Math. Ann.* 65 (4) (1908) 435–505.
- [11] J. G. Rosenstein, *Linear Orderings*, Academic Press, New York, 1982.
- [12] P. Clote, The metamathematics of scattered linear orderings, *Arch. Math. Logic* 29 (1) (1989) 9–20.
- [13] A. Montalbán, On the equimorphism types of linear orderings, *Bull. Symb. Log.* 13 (1) (2007) 71–99.
- [14] P. Cholak, A. Marcone, R. Solomon, Reverse mathematics and the equivalence of definitions for well and better quasi-orders, *J. Symbolic Logic* 69 (3) (2004) 683–712.
- [15] D. H. J. d. Jongh, R. Parikh, Well-partial orderings and hierarchies, *Nederl. Akad. Wetensch. Proc. Ser. A* **80**=*Indag. Math.* 39 (3) (1977) 195–207.
- [16] D. Schmidt, *Well-partial orderings and their maximal order types*, Habilitationsschrift, Heidelberg University (1979).
- [17] G. Higman, Ordering by divisibility in abstract algebras, *Proc. London Math. Soc.* (3) 2 (1952) 326–336.
- [18] J. B. Kruskal, Well-quasi-ordering, the Tree Theorem, and Vazsonyi’s conjecture, *Trans. Amer. Math. Soc.* 95 (1960) 210–225.
- [19] C. S. J. A. Nash-Williams, On well-quasi-ordering transfinite sequences, *Proc. Cambridge Philos. Soc.* 61 (1965) 33–39.
- [20] S. G. Simpson, Nonprovability of certain combinatorial properties of finite trees, in: L. A. Harrington, M. D. Morley, A. Ščedrov, S. G. Simpson (Eds.), *Harvey Friedman’s research on the foundations of mathematics*, North-Holland, Amsterdam, 1985, pp. 87–117.
- [21] H. Friedman, N. Robertson, P. Seymour, The metamathematics of the graph minor theorem, in: S. G. Simpson (Ed.), *Logic and combinatorics*, American Mathematical Society, Providence, R.I., 1987, pp. 229–261.
- [22] R. A. Shore, Invariants, Boolean algebras and ACA_0^+ , *Trans. Amer. Math. Soc.* 358 (3) (2006) 989–1014.

- [23] M. Rathjen, The role of parameters in bar rule and bar induction, *J. Symbolic Logic* 56 (2) (1991) 715–730.
- [24] S. G. Simpson, Ordinal numbers and the Hilbert basis theorem, *J. Symbolic Logic* 53 (3) (1988) 961–974.
- [25] M. Rathjen, A. Weiermann, Proof-theoretic investigations on Kruskal’s theorem, *Ann. Pure Appl. Logic* 60 (1) (1993) 49–88.
- [26] K. McAloon, Paris-Harrington incompleteness and progressions of theories, in: A. Nerode, R. A. Shore (Eds.), *Proceedings of the AMS-ASL summer institute held in Ithaca, N.Y., June 28–July 16, 1982*, Vol. 42 of *Proceedings of Symposia in Pure Mathematics*, American Mathematical Society, Providence, RI, 1985, pp. 447–460.
- [27] P. Clote, The metamathematics of Fraïssé’s order type conjecture, in: K. Ambos-Spies, G. H. Müller, G. E. Sacks (Eds.), *Recursion theory week*, Springer-Verlag, Berlin, 1990, pp. 41–56.
- [28] R. Mansfield, G. Weitkamp, *Recursive aspects of descriptive set theory*, Oxford University Press, New York, 1985, with a chapter by Stephen Simpson.