

**DO NOT TURN OVER UNTIL
INSTRUCTED TO DO SO.**

Formulae

$$\begin{aligned}\int \tan(x) dx &= \ln |\sec(x)| + C & \int \sec(x) dx &= \ln |\sec(x) + \tan(x)| + C \\ \int \frac{1}{1+x^2} dx &= \arctan(x) + C & \int \frac{1}{\sqrt{1-x^2}} dx &= \arcsin(x) + C \\ \frac{d \tan(x)}{dx} &= \sec^2(x) & \frac{d \sec(x)}{dx} &= \tan(x) \sec(x) \\ 1 &= \sin^2(x) + \cos^2(x) & 1 + \tan^2(x) &= \sec^2(x) \\ \cos^2(x) &= \frac{1 + \cos(2x)}{2} & \sin^2(x) &= \frac{1 - \cos(2x)}{2} \\ |E_M| &\leq \frac{K(b-a)^3}{24n^2} & |E_S| &\leq \frac{K(b-a)^5}{180n^4}\end{aligned}$$

CALCULATORS ARE NOT PERMITTED

**YOU MAY USE YOUR OWN BLANK
PAPER FOR ROUGH WORK**

**SO AS NOT TO DISTURB OTHER
STUDENTS, EVERYONE MUST STAY
UNTIL THE EXAM IS FINISHED**

**REMEMBER THIS EXAM IS GRADED BY
A HUMAN BEING. WRITE YOUR
SOLUTIONS NEATLY AND
COHERENTLY, OR THEY RISK NOT
RECEIVING FULL CREDIT**

This exam consists of 5 questions. Answer the questions in the spaces provided.

Name and section: _____

GSI's name: _____

1. Compute the following integrals:

(a) (10 points)

$$\int x \arctan(x) dx$$

Solution:

(b) (10 points)

$$\int x^3 \sqrt{1-x^2} dx$$

Solution:

2. Evaluate the following improper integrals (if divergent, write divergent and explain your reasoning):

(a) (10 points)

$$\int_0^{\frac{\pi}{4}} \frac{\sec(x)}{x^{\frac{3}{2}}} dx$$

(Hint: use the Comparison test)

Solution:

(b) (10 points)

$$\int_0^{\infty} \frac{x^3}{\sqrt{7+x^4}} dx$$

3. (a) (10 points) Express the following rational function

$$\frac{2x^3 + x^2 + 4x + 1}{(x^2 + 1)^2}$$

as a sum of partial fractions.

Solution:

- (b) (10 points) Hence evaluate the integral

$$\int \frac{2x^3 + x^2 + 4x + 1}{(x^2 + 1)^2} dx$$

Solution:

4. (20 points) Find the arc length of the curve given by the function

$$y = x^2 - 2x + 6 - \frac{1}{8} \ln(x - 1),$$

between $x = 2$ and $x = 4$.

Solution:

5. (a) (10 points) Assume that $f(0) = 4$. Use the Midpoint Rule with $n = 5$ to approximate the value of $f(10)$ where $f'(x)$ takes the following values:

x	0	1	2	3	4	5	6	7	8	9	10
$f'(x)$	2	4	3	3	7	6	4	1	5	6	3

Solution:

- (b) (10 points) Assuming that $|f^{(3)}(x)| \leq 1$, for all $0 \leq x \leq 10$, how large an n would you need to choose to guarantee that the above estimate is within 0.001 of the true value $f(10)$? You do not need to give an exact value, just a rough bound.

Solution: