

Math-113, Homework 11, non-textbook problems

The following set of problems is a continuation of exercise 26.30. Please complete textbook exercise 26.30 first. Below we assume that all rings are commutative and we denote by $\text{Nil}(X)$, the nilradical of a ring X .

A. Let R be a ring. For every ideal $I \subseteq R$ define the **radical** of I , denoted as \sqrt{I} , to be the set

$$\sqrt{I} = \{r : r^k \in I, \text{ for some } k \in \mathbb{N}\}.$$

- Show that \sqrt{I} is an ideal.
- Show that $I \subseteq \sqrt{I}$.
- Show that $\text{Nil}(R) = \sqrt{(0)}$. Here (0) stands for the zero ideal of R .
- Let I_1, I_2 be ideals, is it true that $\sqrt{I_1 \cap I_2} = \sqrt{I_1} \cap \sqrt{I_2}$? Prove or provide a counterexample.

B. An ideal I is said to be **radical** if $I = \sqrt{I}$. Let $R = \mathbb{C}[x]$ be the ring of polynomials in one variable with complex coefficients.

- Find at least one ideal $I \subseteq R$, which is not radical.
- For the example constructed above, is it true that R/I has zero divisors? Give an example of a zero divisor, or prove that there are none.
- Consider the quotient ring \sqrt{I}/I for the example you have constructed. It is, in particular, a vector space over \mathbb{C} . Is it finite dimensional? If yes, find a basis and describe multiplication of basis elements.

C. (Optional, not for the grade) Show that $\sqrt{I}/I \cong \text{Nil}(R/I)$.