

# Math 53 Homework 10

Due Wednesday 11/13/13 in section

(The problems in parentheses are for extra practice and optional. Only turn in the underlined problems.)

## Monday 11/4: Line integrals

- **Read:** section 16.2.
- **Work:** 16.2: 1, 3, (7), (11), (15), 17, 20, (29), 32\*, (39), (41), 42.

Problem 1 below.

\* For 16.2 # 32: for part (b), try to find a geometric argument instead! What is the direction of  $\vec{F}$ ? Observe:  $\vec{F} = x(x\hat{i} + y\hat{j})$ .

## Wednesday 11/6: Gradient fields, fundamental theorem for line integrals

- **Read:** section 16.3.
- **Work:** 16.3: 3, (5), (7), 10, (11), (13), 15, (17), (19), 23, 25, (29).<sup>1</sup>

Problems 2 and 3 below.

## Friday 11/8: Green's theorem

- **Read:** section 16.4.
- **Work:** 16.4: (1), 2, 3, (4), (7), 9, (11), 13<sup>2</sup>, (17), 19, (21), 25, (26).

---

### Problem 1.

Consider the vector field  $\vec{F} = (x^2y + \frac{1}{3}y^3)\hat{i}$ , and let  $C$  be the portion of the graph  $y = f(x)$  running from  $(x_1, f(x_1))$  to  $(x_2, f(x_2))$  (assume that  $x_1 < x_2$ , and  $f$  takes positive values). Show that the line integral  $\int_C \vec{F} \cdot d\vec{r}$  is equal to the polar moment of inertia of the region  $R$  lying below  $C$  and above the  $x$ -axis (with density  $\rho = 1$ ).

**Problem 2.** Consider the vector field  $\vec{F}(x, y) = \frac{-y\hat{i} + x\hat{j}}{x^2 + y^2}$ .

a) Show that  $\vec{F}$  is the gradient of the polar angle function  $\theta(x, y) = \tan^{-1}(y/x)$  defined over the right half-plane  $x > 0$ . (Note: this formula for  $\theta$  does not make sense for  $x = 0$ !)

b) Suppose that  $C$  is a smooth curve in the right half-plane  $x > 0$  joining two points  $A : (x_1, y_1)$  and  $B : (x_2, y_2)$ . Express  $\int_C \vec{F} \cdot d\vec{r}$  in terms of the polar coordinates  $(r_1, \theta_1)$  and  $(r_2, \theta_2)$  of  $A$  and  $B$ .

---

<sup>1</sup>6th ed: 16.3: 3, (5), (7), 10, (11), (13), 15, (17), (19), 21, 23, (27).

<sup>2</sup>6th ed: 16.4 # 13 is different. Please do the problem from 7th ed:  $\vec{F}(x, y) = \langle y - \cos y, x \sin y \rangle$ ,  $C$  is the circle  $(x - 3)^2 + (y + 4)^2 = 4$  oriented clockwise.

c) Compute directly from the definition the line integrals  $\int_{C_1} \vec{F} \cdot d\vec{r}$  and  $\int_{C_2} \vec{F} \cdot d\vec{r}$ , where  $C_1$  is the upper half of the unit circle running from  $(1, 0)$  to  $(-1, 0)$ , and  $C_2$  is the lower half of the unit circle, also going from  $(1, 0)$  to  $(-1, 0)$ .

d) Using the results of parts (a)-(c), is  $\vec{F}$  conservative (path-independent) over its entire domain of definition? Is it conservative over the right half-plane  $x > 0$ ? Justify your answers.

e) Show that the components  $P$  and  $Q$  of  $\vec{F}$  satisfy the equation  $\partial P/\partial y = \partial Q/\partial x$  at any point of the plane where  $\vec{F}$  is defined (not just in the right half-plane  $x > 0$ ).

f) (After Friday's lecture) Show that  $\int_C \vec{F} \cdot d\vec{r} = 0$  for every simple closed curve that does not pass through or enclose the origin. Does this remain true if  $C$  encloses the origin?

*Note:* in fact it is true that  $\vec{F} = \nabla\theta$  everywhere. However, the polar angle  $\theta$  cannot be defined as a single-valued differentiable function everywhere (if you try, you will find that it is only well-defined up to adding multiples of  $2\pi$ ). This is why in parts (a) and (b) we only consider the right half-plane; any other region over which  $\theta$  can be defined unambiguously in a continuous manner would be equally suitable.

### Problem 3.

a) For which values of  $n$  do the components  $P$  and  $Q$  of  $\vec{F} = r^n(x\hat{i} + y\hat{j})$  satisfy  $\partial P/\partial y = \partial Q/\partial x$ ? (Here  $r = \sqrt{x^2 + y^2}$ ; start by finding formulas for  $r_x$  and  $r_y$ ).

b) Whenever possible, find a function  $g$  such that  $\vec{F} = \nabla g$ . (Hint: look for a function of the form  $g = g(r)$ , with  $r = \sqrt{x^2 + y^2}$ . Watch out for a certain negative value of  $n$  for which the general formula doesn't work.)