

Math 53 Homework 10

Due Wednesday 11/1/17 in section

(The problems in parentheses are for extra practice and optional. Only turn in the underlined problems.)

Monday 10/23: Vector fields

- **Read:** section 16.1.
- **Work:** 16.1: (5), 11, 13, (15), 18, (21), 23*, (26), 31.

* For 16.1 # 23: also describe geometrically the gradient vector field.

Wednesday 10/25: Line integrals

- **Read:** section 16.2.
- **Work:** 16.2: 1¹, 3, (7), (11), (15), 17, 20², (29), 32*, (39), (41), 42.

Problem 1 below.

* For 16.2 # 32: for part (b), try to find a geometric argument instead! What is the direction of \vec{F} ? Observe: $\vec{F} = x(x\hat{i} + y\hat{j})$.

Friday 10/27: Gradient fields, fundamental theorem for line integrals

- **Read:** section 16.3.
- **Work:** 16.3: (3), (5), 7, (10), (11), (13), 15, (17), 19³, (23), 25, (29).

Problems 2 and 3 below.

Problem 1.

Consider the vector field $\vec{F} = (x^2y + \frac{1}{3}y^3)\hat{i}$, and let C be the portion of the graph $y = f(x)$ running from $(x_1, f(x_1))$ to $(x_2, f(x_2))$ (assume that $x_1 < x_2$, and f takes positive values). Show that the line integral $\int_C \vec{F} \cdot d\vec{r}$ is equal to the polar moment of inertia of the region R lying below C and above the x -axis (with density $\rho = 1$).

Problem 2. The goal of this problem is to show the importance of the condition that the domain under consideration be a simply connected region (i.e., without holes) in the criterion for a vector field to be conservative.

Consider the vector field $\vec{F}(x, y) = \frac{-y\hat{i} + x\hat{j}}{x^2 + y^2}$.

a) Show that \vec{F} is the gradient of the polar angle function $\theta(x, y) = \tan^{-1}(y/x)$ defined over the right half-plane $x > 0$. (Note: this formula for θ does not make sense for $x = 0$!)

¹**7th ed:** do the 8th ed problem: $\int_C y ds$, $C : x = t^2, y = 2t, 0 \leq t \leq 3$.

²**7th ed:** do 8th ed problem: $\vec{F}(x, y, z) = (x + y^2)\hat{i} + xz\hat{j} + (y + z)\hat{k}$, $\vec{r}(t) = t^2\hat{i} + t^3\hat{j} - 2t\hat{k}$, $0 \leq t \leq 2$.

³**7th ed:** do the 8th ed problem: $\int_C 2xe^{-y} dx + (2y - x^2e^{-y}) dy$, C any path from $(1, 0)$ to $(2, 1)$.

b) Suppose that C is a smooth curve in the right half-plane $x > 0$ joining two points $A : (x_1, y_1)$ and $B : (x_2, y_2)$. Express $\int_C \vec{F} \cdot d\vec{r}$ in terms of the polar coordinates (r_1, θ_1) and (r_2, θ_2) of A and B .

c) Compute directly from the definition the line integrals $\int_{C_1} \vec{F} \cdot d\vec{r}$ and $\int_{C_2} \vec{F} \cdot d\vec{r}$, where C_1 is the upper half of the unit circle running from $(1, 0)$ to $(-1, 0)$, and C_2 is the lower half of the unit circle, also going from $(1, 0)$ to $(-1, 0)$.

d) Using the results of parts (a)-(c), is \vec{F} conservative (path-independent) over its entire domain of definition? Is it conservative over the right half-plane $x > 0$? Justify your answers.

Note: in fact it is true that $\vec{F} = \nabla\theta$ everywhere. However, the polar angle θ cannot be defined as a single-valued differentiable function everywhere (if you try, you will find that it is only well-defined up to adding multiples of 2π). This is why in parts (a) and (b) we only consider the right half-plane; any other region over which θ can be defined unambiguously in a continuous manner would be equally suitable.

Problem 3.

a) For which values of n do the components P and Q of $\vec{F} = r^n(x\hat{i} + y\hat{j})$ satisfy $\partial P/\partial y = \partial Q/\partial x$? (Here $r = \sqrt{x^2 + y^2}$; start by finding formulas for r_x and r_y).

b) Whenever possible, find a function g such that $\vec{F} = \nabla g$. (Hint: look for a function of the form $g = g(r)$, with $r = \sqrt{x^2 + y^2}$. Watch out for a certain negative value of n for which the formula is different.)