

Math 53 – Practice Midterm 1 A – 70 minutes

Problem 1. (15 points)

- Find the area of the space triangle with vertices $P_0 : (2, 1, 0)$, $P_1 : (1, 0, 1)$, $P_2 : (2, -1, 1)$.
- Find the equation of the plane containing the three points P_0, P_1, P_2 .
- Find the intersection of this plane with the line parallel to the vector $\vec{V} = \langle 1, 1, 1 \rangle$ and passing through the point $S : (-1, 0, 0)$.

Problem 2. (15 points)

- Let $\vec{r} = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ be the position vector of a path. Give a simple intrinsic formula for $\frac{d}{dt}(\vec{r} \cdot \vec{r})$ in vector notation (not using coordinates).
- Show that if \vec{r} has constant length, then \vec{r} and \vec{v} are perpendicular.
- let \vec{a} be the acceleration: still assuming that \vec{r} has constant length, and using vector differentiation, express the quantity $\vec{r} \cdot \vec{a}$ in terms of the velocity vector only.

Problem 3. (10 points)


Find the area which lies between the x -axis and the curve $x = \sin t$, $y = \sin t \cos t$, $0 \leq t \leq \pi/2$.

Problem 4. (10 points)

On the topographical map below, the level curves for the height function $h(x, y)$ are marked (in feet); adjacent level curves represent a difference of 100 feet in height. A scale is given.

a) Estimate to the nearest .1 the value at the point P of the directional derivative $D_{\hat{u}}h$, where \hat{u} is the unit vector in the direction of $\hat{i} + \hat{j}$.

b) Mark on the map a point Q at which $h = 2200$, $\frac{\partial h}{\partial x} = 0$ and $\frac{\partial h}{\partial y} < 0$. Estimate to the nearest .1 the value of $\frac{\partial h}{\partial y}$ at Q .



Problem 5. (10 points)

Let $f(x, y) = xy - x^4$.

a) Find the gradient of f at $P : (1, 1)$.

b) Give an approximate formula telling how small changes Δx and Δy produce a small change Δw in the value of $w = f(x, y)$ at the point $(x, y) = (1, 1)$.

Problem 6. (5 points)

Find the equation of the tangent plane to the surface $x^3y + z^2 = 3$ at the point $(-1, 1, 2)$.

Problem 7. (5 points)

Let $w = f(u, v)$, where $u = xy$ and $v = x/y$. Express $\frac{\partial w}{\partial x}$ and $\frac{\partial w}{\partial y}$ in terms of x, y, f_u and f_v .


Problem 8. (20 points)

A rectangular box is placed in the first octant as shown, with one corner at the origin and the three adjacent faces in the coordinate planes. The opposite point $P : (x, y, z)$ is constrained to lie on the paraboloid $x^2 + y^2 + z = 1$. Which P gives the box of greatest volume?

a) Show that the problem leads one to maximize $f(x, y) = xy - x^3y - xy^3$, and write down the equations for the critical points of f .

b) Find a critical point of f which lies in the first quadrant ($x > 0, y > 0$), and determine its nature by using the second derivative test.

c) Find the maximum of f in the first quadrant (justify your answer).



Problem 9. (10 points)

In Problem 8 above, instead of substituting for z , one could also use Lagrange multipliers to maximize the volume $V = xyz$ with the same constraint $x^2 + y^2 + z = 1$.

a) Write down the Lagrange multiplier equations for this problem.

b) Solve the equations (still assuming $x > 0, y > 0$).