

PROJECTIVE SPACES AS ORTHOGONAL MODULAR VARIETIES

HAOWU WANG AND BRANDON WILLIAMS

ABSTRACT. We construct 16 reflection groups Γ acting on symmetric domains \mathcal{D} of Cartan type IV, for which the graded algebras of modular forms are freely generated by forms of the same weight, and in particular the Satake–Baily–Borel compactification of \mathcal{D}/Γ is isomorphic to a projective space. Four of these are previously known results of Freitag–Salvati Manni, Matsumoto, Perna and Runge. In addition we find several new modular groups of orthogonal type whose algebras of modular forms are freely generated.

1. INTRODUCTION

In this paper we realize the projective spaces $\mathbb{P}^3(\mathbb{C})$ and $\mathbb{P}^4(\mathbb{C})$ in several ways as orthogonal modular varieties using the theory of Borchers products. Let \mathcal{D}_n be a type IV Hermitian symmetric domain of dimension n with $n \geq 3$, i.e. $\mathcal{D}_n \cong \mathbb{O}_{n,2}^+ / (\mathrm{SO}_n \times \mathrm{O}_2)$ where $\mathbb{O}_{n,2}^+$ is the orthogonal group that preserves \mathcal{D}_n , and let Γ be an arithmetic subgroup of $\mathbb{O}_{n,2}^+$. Then Γ acts properly discontinuously on \mathcal{D}_n and the quotient \mathcal{D}_n/Γ is a quasi-projective variety of dimension n . Orthogonal modular forms, i.e. automorphic forms on \mathcal{D}_n for Γ , are a powerful tool in the study of these varieties. An orthogonal modular form of weight k is a holomorphic function on the affine cone over \mathcal{D}_n which has homogenous degree $-k$ and is invariant under Γ . Orthogonal modular forms of all weights for Γ form a graded algebra $M_*(\Gamma)$. By [2] the graded algebra $M_*(\Gamma)$ is finitely generated over \mathbb{C} and the Satake-Baily-Borel compactification $(\mathcal{D}_n/\Gamma)^*$ of the modular variety \mathcal{D}/Γ is a projective variety isomorphic to $\mathrm{Proj}(M_*(\Gamma))$. In particular, if $M_*(\Gamma)$ is freely generated by $n + 1$ forms of weights k_1, k_2, \dots, k_{n+1} , then $(\mathcal{D}_n/\Gamma)^* = \mathrm{Proj}(M_*(\Gamma))$ is a weighted projective space with weights k_1, k_2, \dots, k_{n+1} .

Free algebras of modular forms are rare. Many of the known examples are related to irreducible root systems as in [21]. It is known that the group Γ must be generated by reflections if $M_*(\Gamma)$ is free (see [16]). (This immediately rules out any modular groups Γ except subgroups of $\mathrm{U}(n, 1)$ and of $\mathrm{O}(n, 2)$. In this paper Γ is always an arithmetic subgroup of $\mathrm{O}(n, 2)$.) The first named author found a necessary and sufficient condition for $M_*(\Gamma)$ to be free in [19] and used it to construct 16 new free algebras of modular forms in [20]. This condition is essentially the existence of a distinguished modular form which vanishes precisely on all mirrors of reflections in Γ with multiplicity one and equals the Jacobian of the $n + 1$ generators. In this paper, we use that criterion to find reflection groups Γ such that $M_*(\Gamma)$ is freely generated by some forms of the same weight. For any such Γ the Satake-Baily-Borel compactification of \mathcal{D}_n/Γ is a projective space.

Modular varieties isomorphic to projective spaces are very exceptional. The authors are aware of only four examples of dimension $n \geq 3$ in the literature. The first was found by Runge in 1993 [17]. Runge’s theorem states that the algebra of Siegel modular forms of genus 2 on the level 4 subgroup $\Gamma_2[2, 4]$ is the polynomial algebra in 4 theta constants of second order, which implies that the corresponding modular variety is isomorphic to $\mathbb{P}^3(\mathbb{C})$. The second example was found

Date: August 19, 2020.

2010 Mathematics Subject Classification. 11F55, 51F15, 32N15.

Key words and phrases. Symmetric domains of type IV, modular forms on orthogonal groups, projective spaces, reflection groups, Borchers products.

by Matsumoto [13] in the same year. Matsumoto proved that the algebra of symmetric Hermitian modular forms of degree 2 over the Gaussian numbers for the principal congruence group of level $1+i$ is freely generated by five forms of weight 2, which implies that the corresponding modular variety is isomorphic to $\mathbb{P}^4(\mathbb{C})$. A different proof of this was given by Hermann [10]. The third example was constructed by Freitag and Salvati Manni in 2006. They proved in [7] that the algebra of symmetric Hermitian modular forms of degree 2 over the Eisenstein integers for the congruence group of level $\sqrt{-3}$ is freely generated by five forms of weight 1 and therefore the modular variety is also isomorphic to $\mathbb{P}^4(\mathbb{C})$. Finally, Perna [15] proved that the algebra of Siegel modular forms for a certain congruence subgroup containing $\Gamma_2[2, 4]$ is freely generated by the squares of 4 theta constants of second order.

We will give a simple and largely uniform proof of the above results in the context of orthogonal modular forms. Runge's and Perna's theorems can be interpreted in terms of orthogonal modular forms for the lattice $2U(4) \oplus A_1$. (Here and below, U is an even unimodular lattice of signature $(1, 1)$, and A_1, A_2 denote the lattice generated by the root system of the same name.) Matsumoto's theorem is treated using the lattice model $2U(2) \oplus 2A_1$, and for this lattice we also find two new free algebra of modular forms for smaller groups. The first of these is freely generated by five forms of weight 1 and the second is freely generated by four forms of weight 2 and one form of weight 1, such that the modular varieties associated to both of these groups are isomorphic to $\mathbb{P}^4(\mathbb{C})$. The theorem of Freitag–Salvati Manni corresponds to the lattice $2U(3) \oplus A_2$, and we also find two new free algebras of modular forms for larger groups related to this lattice. The first of these is freely generated by five forms of weight 2 and the second is freely generated by four forms of weight 2 and one form of weight 1. The modular varieties associated to the two groups are then isomorphic to $\mathbb{P}^4(\mathbb{C})$. In addition, we determine eight new reflection groups whose associated modular varieties are isomorphic to $\mathbb{P}^3(\mathbb{C})$. Three of them are related to $2U(2) \oplus A_1$ and the weights of generators of the three algebras are $\{2, 2, 2, 2\}$, $\{1, 1, 1, 1\}$ and $\{1, 2, 2, 2\}$ respectively. Also we determine an algebra related to $2U(3) \oplus A_1$ which is generated by four forms of weight 1, and two algebras related to $U(2) \oplus U(4) \oplus A_1$ which are generated by four forms of weight 1 and four forms of weight $1/2$ respectively. Finally we obtain two algebras related to $U \oplus U(2) \oplus A_1(2)$ generated by modular forms of weights $\{2, 2, 2, 2\}$ and $\{1, 2, 2, 2\}$ respectively.

Altogether, we will realize $\mathbb{P}^3(\mathbb{C})$ and $\mathbb{P}^4(\mathbb{C})$ in sixteen ways as orthogonal modular varieties associated to seven lattices in Theorems 3.6, 3.7, 4.3, 4.6, 4.7, 5.4, 5.5, 5.6, 6.2, 6.4, 6.5, 7.2, 8.2, 8.4, 9.2, 9.3. The generators of these algebras of modular forms are all constructed as Borcherds products [3], with one exception where we require an additive theta lift. Along the way we find a simple proof of the famous theorem of Igusa [11] that the algebra of Siegel modular forms of genus 2 on the level 8 subgroup $\Gamma_2[4, 8]$ is generated by the ten theta constants (Theorem 3.8), as well as its analogue for Hermitian modular forms over the Gaussian integers (Theorem 4.5). We also find several other interesting free algebras of modular forms; for example, we will find a tower of eight congruence subgroups of $\mathrm{Sp}_4(\mathbb{Z})$, all of whose algebras of modular forms are described explicitly and six of which are free.

The layout of this paper is as follows. In §2 we recall the necessary and sufficient condition for free algebras mentioned above (and a straightforward generalization to half-integral weight modular forms). In §3–9 we study the algebras of modular forms attached to the seven lattices respectively. In the appendices we give some data related to the input forms for certain Borcherds products; the Fourier expansions of the Borcherds products, and the products themselves as SAGE objects, are available in the ancillary material.

At several points in this paper, we work directly with the Fourier expansions of Borcherds products and therefore need a significant number of coefficients of their input forms. These were computed in SAGE [18] using the algorithm outlined in [22], an implementation of which is available

on GitHub. We also used SAGE for certain graph computations, including much of the data in the appendices.

2. PRELIMINARIES

Let M be an even lattice of signature $(n, 2)$ with $n \geq 3$ and dual lattice M^\vee . The type IV symmetric domain $\mathcal{D}_n = \mathcal{D}(M)$ is one of the two connected components of the space

$$\{[\mathcal{Z}] \in \mathbb{P}(M \otimes \mathbb{C}) : (\mathcal{Z}, \mathcal{Z}) = 0, (\mathcal{Z}, \bar{\mathcal{Z}}) < 0\}.$$

We define the affine cone over $\mathcal{D}(M)$ as

$$\mathcal{A}(M) = \{\mathcal{Z} \in M \otimes \mathbb{C} : [\mathcal{Z}] \in \mathcal{D}(M)\}.$$

Let us denote by $O^+(M)$ the integral orthogonal group of M preserving the connected component $\mathcal{D}(M)$. The discriminant kernel $\tilde{O}^+(M)$ is the subgroup of $O^+(M)$ which acts trivially on the discriminant form of M . Fix a finite index subgroup Γ of $O^+(M)$.

Definition 2.1. Let k be a non-negative integer. A *modular form* of weight k and character $\chi : \Gamma \rightarrow \mathbb{C}^*$ for Γ is a holomorphic function $F : \mathcal{A}(M) \rightarrow \mathbb{C}$ satisfying

$$\begin{aligned} F(t\mathcal{Z}) &= t^{-k}F(\mathcal{Z}), \quad \forall t \in \mathbb{C}^*, \\ F(g\mathcal{Z}) &= \chi(g)F(\mathcal{Z}), \quad \forall g \in \Gamma. \end{aligned}$$

The graded algebra of modular forms of integral weight is denoted by

$$M_*(\Gamma) = \bigoplus_{k=0}^{\infty} M_k(\Gamma).$$

One can realize the symmetric domain $\mathcal{D}(M)$ as a tube domain at a zero-dimensional cusp. The above modular form can be viewed as an automorphic form on a tube domain for Γ with respect to an automorphy factor. Using the tube domain model one can define modular forms of half-integral weight with a multiplier system. We refer to [4, §3.3] for more details.

For any $r \in M^\vee$ of positive norm, the hyperplane

$$\mathcal{D}_r(M) = r^\perp \cap \mathcal{D}(M) = \{[\mathcal{Z}] \in \mathcal{D}(M) : (\mathcal{Z}, r) = 0\}$$

is called the *rational quadratic divisor* associated to r . The reflection fixing $\mathcal{D}_r(M)$ is

$$\sigma_r(x) = x - \frac{2(r, x)}{(r, r)}r, \quad x \in M.$$

The hyperplane $\mathcal{D}_r(M)$ is called the *mirror* of σ_r . For a non-zero vector $r \in M^\vee$ we denote its order in M^\vee/M by $\text{ord}(r)$. A primitive vector $r \in M^\vee$ of positive norm is called *reflective* if $\sigma_r \in O^+(M)$, or equivalently, if there exists a positive integer d such that $(r, r) = \frac{2}{d}$ and $\text{ord}(r) = d$ or $\frac{d}{2}$. In this case we call $\mathcal{D}_r(M)$ a *reflective divisor*. A modular form F for $\Gamma < O^+(M)$ is called *reflective* if its divisor is a sum of reflective divisors. F is called *2-reflective* if its divisor is a sum of divisors $\mathcal{D}_r(M)$ for $r \in M$ with $(r, r) = 2$. We remark that if $r \in M$ is primitive then the reflection σ_r belongs to $\tilde{O}^+(M)$ if and only if $(r, r) = 2$.

For convenience, we recall some results of [19] which will be used in this paper. The modular Jacobian, or Rankin–Cohen–Ibukiyama differential operator, was first introduced in [1] for Siegel modular forms.

Theorem 2.2 (Theorem 2.5 in [19]). *Let M be an even lattice of signature $(n, 2)$ with $n \geq 3$, and let $\Gamma < O^+(M)$ be a finite index subgroup. Let $f_i \in M_{k_i}(\Gamma)$ for $1 \leq i \leq n + 1$. We view f_i*

as modular forms on the tube domain at a given zero-dimensional cusp and let z_i , $1 \leq i \leq n$, be coordinates on the tube domain. We define

$$J := J(f_1, \dots, f_{n+1}) = \begin{vmatrix} k_1 f_1 & k_2 f_2 & \cdots & k_{n+1} f_{n+1} \\ \frac{\partial f_1}{\partial z_1} & \frac{\partial f_2}{\partial z_1} & \cdots & \frac{\partial f_{n+1}}{\partial z_1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_1}{\partial z_n} & \frac{\partial f_2}{\partial z_n} & \cdots & \frac{\partial f_{n+1}}{\partial z_n} \end{vmatrix}.$$

- (1) J is a cusp form of weight $n + \sum_{i=1}^{n+1} k_i$ for Γ with the determinant character \det .
- (2) J is not identically zero if and only if the $n+1$ modular forms f_i are algebraically independent over \mathbb{C} .
- (3) Let $r \in M$. If the reflection σ_r belongs to Γ , then J vanishes on the hyperplane $\mathcal{D}_r(M)$.

If $M_*(\Gamma)$ is a free algebra then the Jacobian of its generators satisfies some remarkable properties:

Theorem 2.3 (Theorem 3.5 in [19]). *Assume that $M_*(\Gamma)$ is a free algebra with generators f_1, \dots, f_{n+1} of weights k_1, \dots, k_{n+1} .*

- (1) *The Jacobian $J = J(f_1, \dots, f_{n+1})$ is not identically zero and it is a cusp form of weight $n + \sum_{i=1}^{n+1} k_i$ for Γ with the character \det .*
- (2) *The divisor of J is the sum of all mirrors of reflections in Γ , each with multiplicity 1. In particular, J is a reflective cusp form.*
- (3) *Let $\{\Gamma\pi_1, \dots, \Gamma\pi_s\}$ denote the Γ -equivalence classes of mirrors of reflections in Γ . Then for each $1 \leq i \leq s$ there exists a modular form J_i for Γ with trivial character and divisor $\text{div}(J_i) = 2\Gamma\pi_i$, and $J^2 = \prod_{i=1}^s J_i$. The forms J_i are irreducible in $M_*(\Gamma)$.*
- (4) *There exist polynomials P, P_i , $1 \leq i \leq s$, in $n+1$ variables over \mathbb{C} such that $J^2 = P(f_1, \dots, f_{n+1})$ and $J_i = P_i(f_1, \dots, f_{n+1})$. Thus $P = \prod_{i=1}^s P_i$ and these P_i are irreducible.*

The following sufficient condition for a graded algebra of modular forms to be free will play a vital role in this paper.

Theorem 2.4 (Theorem 5.1 in [19]). *Let $\Gamma < O^+(M)$ be a finite index subgroup. Suppose there exist modular forms f_1, \dots, f_{n+1} with trivial character whose Jacobian*

$$J = J(f_1, \dots, f_{n+1})$$

vanishes exactly on the mirrors of reflections in Γ with multiplicity one. Then the graded algebra $M_(\Gamma)$ is freely generated by f_1, \dots, f_{n+1} and Γ is generated by the reflections whose mirrors lie in the divisor of J .*

This theorem also holds for modular forms of half-integral weight and the proof is nearly the same. We replace $M_*(\Gamma)$ with the graded algebra of half-integral weight modular forms with respect to a fixed multiplier system v of weight $1/2$:

$$M_*(\Gamma, v) = \bigoplus_{k=0}^{\infty} M_{\frac{k}{2}}(\Gamma, v^k).$$

Theorem 2.5. *Suppose there exist modular forms $f_1, \dots, f_{n+1} \in M_*(\Gamma, v)$ whose Jacobian $J = J(f_1, \dots, f_{n+1})$ vanishes exactly on the mirrors $\mathcal{D}_r(M)$ of reflections σ_r in Γ that satisfy $v(\sigma_r) = 1$ with multiplicity one. Then $M_*(\Gamma, v)$ is freely generated by f_i .*

Proof. Suppose that $M_*(\Gamma, v)$ is not generated by f_i . Then $\mathbb{C}[f_1, \dots, f_{n+1}] \neq M_*(\Gamma, v)$, and we choose a modular form $f_{n+2} \in M_{k_{n+2}}(\Gamma, v^{2k_{n+2}})$ of minimal weight such that $f_{n+2} \notin \mathbb{C}[f_1, \dots, f_{n+1}]$. For $1 \leq t \leq n+2$ we define

$$J_t = J(f_1, \dots, \hat{f}_t, \dots, f_{n+2})$$

as the Jacobian of the $n + 1$ modular forms f_i omitting f_t (so in particular $J = J_{n+2}$). Similarly to Theorem 2.2, one can show that the Jacobian J_t is a modular form of weight $k = n + \sum_{i \neq t} k_i$ and multiplier system $v^{2k} \det$ on Γ , and that J_t vanishes on all mirrors of reflections in Γ satisfying $v(\sigma_r) = 1$. Therefore the quotient $g_t := J_t/J$ is a holomorphic modular form in $M_*(\Gamma, v)$.

We compute

$$0 = \det \begin{pmatrix} k_1 f_1 & k_2 f_2 & \cdots & k_{n+2} f_{n+2} \\ k_1 f_1 & k_2 f_2 & \cdots & k_{n+2} f_{n+2} \\ \nabla_z f_1 & \nabla_z f_2 & \cdots & \nabla_z f_{n+2} \end{pmatrix} = \sum_{t=1}^{n+2} (-1)^t k_t f_t J_t = \left(\sum_{t=1}^{n+2} (-1)^t k_t f_t g_t \right) \cdot J,$$

and therefore

$$(-1)^{n+1} k_{n+2} f_{n+2} = \sum_{t=1}^{n+1} (-1)^t k_t f_t g_t$$

because $g_{n+2} = 1$. In particular, each g_t has weight strictly less than that of f_{n+2} . By construction of f_{n+2} , this implies $g_t \in \mathbb{C}[f_1, \dots, f_{n+1}]$, and therefore $f_{n+2} \in \mathbb{C}[f_1, \dots, f_{n+1}]$, a contradiction. Therefore $M_*(\Gamma, v)$ is generated by the f_i . Since $M_*(\Gamma, v)$ has Krull dimension $n+1$, these generators are algebraically independent. \square

Unlike the integral-weight case, we cannot conclude in general that Γ is generated by the reflections whose mirrors are contained in the divisor of the Jacobian. However, this does hold if $v(\sigma_r) = 1$ for all reflections $\sigma_r \in \Gamma$.

We can investigate the modular variety \mathcal{D}_n/Γ using algebras of half-integral weight modular forms because the Proj is unchanged under Veronese embeddings:

$$\text{Proj}(M_*(\Gamma, v)^{(d)}) = \text{Proj}(M_*(\Gamma, v)) = \text{Proj}(M_*(\Gamma)^{(d)}) = \text{Proj}(M_*(\Gamma)) = (\mathcal{D}_n/\Gamma)^*,$$

for any $d \in \mathbb{N}$, where

$$M_*(\Gamma, v)^{(d)} = \bigoplus_{k \in \mathbb{N}} M_{\frac{dk}{2}}(\Gamma, v^{dk}), \quad M_*(\Gamma)^{(d)} = \bigoplus_{k \in \mathbb{N}} M_{dk}(\Gamma).$$

3. THE $2U(4) \oplus A_1$ LATTICE

In this section we recover Runge's theorem and Perna's theorem mentioned in the introduction. Runge's theorem asserts that the algebra of Siegel modular forms of genus 2 on the level 4 subgroup $\Gamma_2[2, 4]$ is freely generated by 4 theta constants of second order. We will find that the Jacobian of four theta constants of second order is exactly Igusa's cusp form Φ_{5, A_1} up to a multiple, where Φ_{5, A_1} is the product of ten even theta constants. The Igusa cusp form Φ_{5, A_1} is a reflective modular form of weight 5 for $O^+(2U \oplus A_1)$ whose divisor is the sum of all \mathcal{D}_r for primitive vectors $r \in 2U \oplus A_1^\vee$ with $(r, r) = \frac{1}{2}$ and $\text{ord}(r) = 2$, each with multiplicity one. In view of the isomorphisms

$$(3.1) \quad O^+(M) = O^+(M^\vee) = O^+(M^\vee(m)),$$

we have $O^+(2U \oplus A_1) = O^+(2U(4) \oplus A_1)$ and the function Φ_{5, A_1} can also be viewed as a 2-reflective modular form of weight 5 for $2U(4) \oplus A_1$ whose divisor is the sum of \mathcal{D}_s with multiplicity one for primitive vectors $s \in 2U(\frac{1}{4}) \oplus A_1^\vee$ with $(s, s) = \frac{1}{2}$ and $\text{ord}(s) = 2$. Unlike $\tilde{O}^+(2U \oplus A_1)$, these account for all 2-reflections in $\tilde{O}^+(2U(4) \oplus A_1)$. Moreover, the discriminant kernel $\tilde{O}^+(2U(4) \oplus A_1)$ is a subgroup of $\tilde{O}^+(2U \oplus A_1)$, and there are many more Borcherds products on $\tilde{O}^+(2U(4) \oplus A_1)$. For these reasons it is natural to work with the lattice $2U(4) \oplus A_1$.

We will first recall the Borcherds products of singular weight on $2U(4) \oplus A_1$.

Lemma 3.1.

- (1) *There are 70 holomorphic Borcherds products of singular weight 1/2 on $2U(4) \oplus A_1$. Their product is the Igusa cusp form of weight 35.*

- (2) Ten of the singular weight products have inputs with principal part $q^{-1/4}(e_v + e_w)$ where v and w have order 2 in M^\vee/M . These are the ten theta constants and their product is Φ_{5,A_1} . We call them the Type I products and denote them θ_i , $i = 1, 2, \dots, 10$.
- (3) The remaining 60 singular weight products have inputs with principal part $q^{-1/4}(e_u + e_{-u})$, where u has order 4. We call them the Type II products.

Proof. These facts can be read off of the Fourier coefficients of a weight 5/2 Eisenstein series. This is carried out in detail in section 4 of [14]. With respect to the table of section 4 of [14], the principal parts of the input into the Type I products consist of one of ten vectors v from the 7th orbit and one of six vectors w from the 8th orbit. The vector-valued modular forms with principal part $q^{-1/4}e_v$ alone have non-integral Fourier coefficients so they are, strictly speaking, not valid inputs into the Borcherds lift. However, the $q^{-1/4}e_w$ term does not contribute to their divisor. \square

Let $O_1(2U(4) \oplus A_1)$ be the subgroup of $O^+(2U(4) \oplus A_1)$ generated by all reflections associated to the divisor of Φ_{5,A_1} . It is obvious that $O_1(2U(4) \oplus A_1)$ is contained in the discriminant kernel $\tilde{O}^+(2U(4) \oplus A_1)$. The 70 products satisfy the following basic properties.

Lemma 3.2.

- (1) The 10 Type I products θ_i are linearly independent over \mathbb{C} .
- (2) The 55 forms $\theta_i\theta_j$ are linearly independent over \mathbb{C} .
- (3) The 60 Type II products are modular forms of weight 1/2 for $O_1(2U(4) \oplus A_1)$, all of which have the same multiplier system which we denote v_Θ . Moreover, $v_\Theta = 1$ for all reflections in $O_1(2U(4) \oplus A_1)$.

Proof. (1) The Type I products are modular forms on $\tilde{O}^+(2U(4) \oplus A_1)$. Let σ_i be a reflection associated to the divisor of some θ_i . Since $\sigma_i \in \tilde{O}^+(2U(4) \oplus A_1)$, we obtain $\sigma_i(\theta_i) = -\theta_i$ and $\sigma_i(\theta_j) = \theta_j$ for $j \neq i$. This implies the linear independence of the 10 products.

(2) Using the argument in (1), the linear independence of the 55 functions $\theta_i\theta_j$ reduces to the linear independence of the 10 squares θ_i^2 , which can be shown by computing Fourier coefficients.

(3) Consider the quotient of any two products of type II. This is a weight zero meromorphic modular form on $O_1(2U(4) \oplus A_1)$. Since it does not vanish on any divisors contained in $\text{div } \Phi_{5,A_1}$, it has trivial character on $O_1(2U(4) \oplus A_1)$, so the type II products have the same multiplier system. From the principal parts of their input forms we see that no type II product vanishes on the divisor \mathcal{D}_r for any reflection $\sigma_r \in O_1(2U(4) \oplus A_1)$, which forces $v_\Theta(\sigma_r) = 1$. We remark that $v_\Theta^2 = 1$ on $O_1(2U(4) \oplus A_1)$. \square

Throughout this section we will consider sets of four type II products $\Theta_1, \Theta_2, \Theta_3, \Theta_4$ that transform under larger modular groups than $O_1(2U(4) \oplus A_1)$. For this the following definition is useful.

Definition 3.3. A **-set* (of type $2U(4) \oplus A_1$) is a set of four Type II products, each of whose input forms is invariant under all reflections associated to the divisors of the three other products.

In other words, if the type II product Θ_i has input with principal part $q^{-1/4}(e_{u_i} + e_{-u_i})$ then $\{\Theta_1, \Theta_2, \Theta_3, \Theta_4\}$ forms a *-set if and only if $\sigma_{u_i}(u_j) \in \pm u_j + (2U(4) \oplus A_1)$ for $1 \leq i, j \leq 4$.

*-sets have several interesting properties.

Lemma 3.4.

- (1) There are exactly 105 *-sets.
- (2) Every product of type II belongs to exactly seven *-sets.
- (3) There are 1320 pairs of type II products that do not belong to a *-set; there are 360 pairs that belong to exactly one *-set; and the remaining 90 pairs belong to exactly three *-sets.
- (4) Any four type II products that form a *-set are linearly independent over \mathbb{C} .

Proof. Parts (1)-(3) were checked by computer. We constructed a graph with the Type II products as vertices and an edge between two products if and only if their input forms are invariant under the reflections associated to each other's divisors. (See Appendix A for an image.) The $*$ -sets are the maximal cliques in this graph.

(4) Let f_j , $1 \leq j \leq 4$, be a $*$ -set and let Γ_* be the group generated by $O_1(2U(4) \oplus A_1)$ and the reflections associated to the divisors of the four products, such that these products are modular forms on Γ_* . Let σ_i be a reflection associated to the divisor of f_i . Then $\sigma_i(f_i) = -f_i$ and $\sigma_i(f_j) = f_j$ for $j \neq i$, which implies the linear independence. \square

The extra structure given by the 90 pairs of type II products in part (3) will be useful later in this section.

Lemma 3.5. *There is a $*$ -set $\{\Theta_1, \Theta_2, \Theta_3, \Theta_4\}$ for which the Jacobian $J(\Theta_1, \Theta_2, \Theta_3, \Theta_4)$ equals Igusa's cusp form Φ_{5, A_1} up to a nonzero constant multiple.*

Proof. In the notation of Appendix A we took the products labelled $\Theta_1, \Theta_2, \Theta_{26}, \Theta_{55}$. By computing their Fourier expansions it is straightforward to show that their Jacobian is not identically zero. Since these products are modular forms of weight $1/2$ and multiplier system v_Θ on $O_1(2U(4) \oplus A_1)$, and $v_\Theta = 1$ for all reflections in $O_1(2U(4) \oplus A_1)$, we conclude from Theorem 2.2 (4) that their Jacobian J vanishes on the mirrors of all reflections in $O_1(2U(4) \oplus A_1)$. It follows that $J/\Phi_{5, A_1}$ is a holomorphic modular form of weight 0 and therefore constant. \square

By applying Theorem 2.5, we obtain Runge's theorem in the context of orthogonal groups.

Theorem 3.6. *The type II products span a four-dimensional space over \mathbb{C} . If $\Theta_1, \Theta_2, \Theta_3, \Theta_4$ are any linearly independent Type II products, then they are algebraically independent and generate the algebra of modular forms:*

$$\begin{aligned} M_*(O_1(2U(4) \oplus A_1), v_\Theta) &= \mathbb{C}[\Theta_1, \Theta_2, \Theta_3, \Theta_4], \\ (\mathcal{D}_3/O_1(2U(4) \oplus A_1))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Perna's theorem can be phrased in terms of a larger modular group associated to a $*$ -set.

Theorem 3.7. *Let $\{\Theta_1, \Theta_2, \Theta_3, \Theta_4\}$ be a $*$ -set and let $O_2(2U(4) \oplus A_1)$ be the subgroup generated by $O_1(2U(4) \oplus A_1)$ and the reflections with mirrors in the divisor of $\prod_{j=1}^4 \Theta_j$. Then*

$$\begin{aligned} M_*(O_2(2U(4) \oplus A_1)) &= \mathbb{C}[\Theta_1^2, \Theta_2^2, \Theta_3^2, \Theta_4^2], \\ (\mathcal{D}_3/O_2(2U(4) \oplus A_1))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Proof. By definition of $*$ -sets, the forms Θ_j^2 are modular forms of weight 1 with trivial character on $O_2(2U(4) \oplus A_1)$. By Lemma 3.5 their Jacobian equals $\Phi_{5, A_1} \prod_{j=1}^4 \Theta_j$ up to a non-zero multiple. By Theorem 2.4, the graded algebra of modular forms on $O_2(2U(4) \oplus A_1)$ is freely generated by the four squares. \square

As another application, we determine an algebra of modular forms generated by ten theta constants and reprove a well-known theorem of Igusa.

The Jacobian Φ_{5, A_1} factors as $\prod_{i=1}^{10} \theta_i$ in $M_*(O_1(2U(4) \oplus A_1), v_\Theta)$. By Theorem 2.3, this decomposition determines the character group of $O_1(2U(4) \oplus A_1)$. For $1 \leq i \leq 10$, let χ_i be the character of $O_1(2U(4) \oplus A_1)$ defined as the quotient of the multiplier system of θ_i by v_Θ . These ten basic characters generate the 1024 characters of $O_1(2U(4) \oplus A_1)$. Let $O'_1(2U(4) \oplus A_1)$ be the commutator

subgroup of $O_1(2U(4) \oplus A_1)$. Then we have

$$M_{\frac{k}{2}}(O'_1(2U(4) \oplus A_1), v_{\Theta}^k) = \bigoplus_{\chi} M_{\frac{k}{2}}(O_1(2U(4) \oplus A_1), v_{\Theta}^k \chi),$$

$$M_{\frac{k}{2}}(O_1(2U(4) \oplus A_1), v_{\Theta}^k \chi_i) = M_{\frac{k-1}{2}}(O_1(2U(4) \oplus A_1), v_{\Theta}^{k-1}) \theta_i, \quad 1 \leq i \leq 10,$$

the first sum taken over the character group of $O_1(2U(4) \oplus A_1)$.

Since each θ_i^2 lies in $M_1(O_1(2U(4) \oplus A_1))$ and the ten squares θ_i^2 are linearly independent, we conclude from Theorem 3.6 that the ten θ_i^2 form a basis of $M_1(O_1(2U(4) \oplus A_1))$. We define O_0 as the subgroup on which the basic characters coincide:

$$O_0(2U(4) \oplus A_1) = \{\gamma \in O_1(2U(4) \oplus A_1) : \chi_1(\gamma) = \cdots = \chi_{10}(\gamma)\}.$$

The group $O_0(2U(4) \oplus A_1)$ properly contains $O'_1(2U(4) \oplus A_1)$ because $\sigma_1 \cdots \sigma_{10} \in O_0(2U(4) \oplus A_1)$ where σ_i is any reflection associated to the divisor of θ_i for $1 \leq i \leq 10$. We note that

$$\theta_i \in M_{\frac{1}{2}}(O_1(2U(4) \oplus A_1), v_{\Theta} \chi_i), \quad 1 \leq i \leq 10.$$

Let χ be the common restriction of the χ_i to $O_0(2U(4) \oplus A_1)$. We define a multiplier system v_{ϑ} of weight $1/2$ on $O_0(2U(4) \oplus A_1)$ by

$$v_{\vartheta} = v_{\Theta} \chi.$$

Then

$$(3.2) \quad \theta_i \in M_{\frac{1}{2}}(O_0(2U(4) \oplus A_1), v_{\vartheta}), \quad 1 \leq i \leq 10$$

and

$$(3.3) \quad \Theta_j \in M_{\frac{1}{2}}(O_0(2U(4) \oplus A_1), v_{\vartheta} \chi), \quad 1 \leq j \leq 4.$$

Since the quotient group $O_0(2U(4) \oplus A_1)/O'_1(2U(4) \oplus A_1)$ is abelian, we obtain an eigenspace decomposition

$$(3.4) \quad M_{\frac{k}{2}}(O'_1(2U(4) \oplus A_1), v_{\vartheta}^k) = \bigoplus_{\epsilon} M_{\frac{k}{2}}(O_0(2U(4) \oplus A_1), v_{\vartheta}^k \epsilon),$$

the direct sum taken over the characters of $O_0(2U(4) \oplus A_1)/O'_1(2U(4) \oplus A_1)$. Note that $v_{\vartheta} = v_{\Theta}$ on $O'_1(2U(4) \oplus A_1)$. Therefore, every modular form in $M_{\frac{k}{2}}(O'_1(2U(4) \oplus A_1), v_{\vartheta}^k)$ can be decomposed as a sum

$$P_0 + \sum_{i=1}^4 P_i \Theta_i,$$

where P_0 is a modular form of weight $k/2$, and P_1, P_2, P_3, P_4 are modular forms of weight $(k-1)/2$, and

$$P_0, \dots, P_4 \in \mathbb{C}[\theta_i, 1 \leq i \leq 10].$$

Combining (3.2), (3.3) and (3.4), we obtain

$$M_*(O_0(2U(4) \oplus A_1), v_{\vartheta}) = \mathbb{C}[\theta_i, 1 \leq i \leq 10]$$

and we see that χ is the unique non-trivial character of $O_2(2U(4) \oplus A_1)/O'_1(2U(4) \oplus A_1)$. Altogether, we obtain the generators found by Igusa in [11]:

Theorem 3.8.

$$M_*(O_0(2U(4) \oplus A_1), v_{\vartheta}) = \mathbb{C}[\theta_i, 1 \leq i \leq 10],$$

$$M_*(O'_1(2U(4) \oplus A_1), v_{\Theta}) = \mathbb{C}[\Theta_1, \Theta_2, \Theta_3, \Theta_4, \theta_i, 1 \leq i \leq 10].$$

Remark 3.9. The ten theta constants θ_i are not all modular on $O_2(2U(4) \oplus A_1)$. In fact, Φ_{5,A_1} factors as the product of five forms of weights $1/2, 1/2, 1, 1$ and 2 on $O_2(2U(4) \oplus A_1)$, which are products of some theta constants.

We can now construct some new free algebras of Siegel modular forms associated to $*$ -sets. Let $\Theta_1, \dots, \Theta_4$ be a $*$ -set and let $O_2(2U(4) \oplus A_1)$ be the group generated by its reflections and by $O_1(2U(4) \oplus A_1)$.

Theorem 3.10. For $1 \leq i \leq 4$ let v_i be the multiplier system of Θ_i on $O_2(2U(4) \oplus A_1)$. Then

$$M_*(O_2(2U(4) \oplus A_1), v_i) = \mathbb{C}[\Theta_i, \Theta_j^2, 1 \leq j \leq 4, j \neq i], \quad 1 \leq i \leq 4.$$

This includes Theorem 3.7 which describes the subring of integer-weight forms.

Proof. Without loss of generality we take $i = 1$. It is clear that $v_1 = 1$ for all reflections associated to the divisor of $\prod_{j=2}^4 \Theta_j$ and $v_1 = -1$ for reflections associated to the divisor of Θ_1 . By Lemma 3.5 the Jacobian of Θ_1, Θ_j^2 for $2 \leq j \leq 4$ equals $\Phi_{5,A_1} \prod_{j=2}^4 \Theta_j$ up to a non-zero multiple. The claim follows from Theorem 2.5. \square

Let $O_{1,1}(2U(4) \oplus A_1)$, $O_{1,12}(2U(4) \oplus A_1)$ and $O_{1,123}(2U(4) \oplus A_1)$ denote the subgroups generated by reflections associated to the divisor of $\Phi_{5,A_1} \Theta_1$, $\Phi_{5,A_1} \Theta_1 \Theta_2$ and $\Phi_{5,A_1} \Theta_1 \Theta_2 \Theta_3$, respectively. By a similar argument, we can prove the following results using Theorem 2.5:

$$\begin{aligned} M_*(O_{1,1}(2U(4) \oplus A_1), v_4) &= \mathbb{C}[\Theta_1^2, \Theta_2, \Theta_3, \Theta_4], \\ M_*(O_{1,12}(2U(4) \oplus A_1), v_4) &= \mathbb{C}[\Theta_1^2, \Theta_2^2, \Theta_3, \Theta_4], \\ M_*(O_{1,123}(2U(4) \oplus A_1), v_4) &= \mathbb{C}[\Theta_1^2, \Theta_2^2, \Theta_3^2, \Theta_4]. \end{aligned}$$

We remark that $O_{1,1}(2U(4) \oplus A_1)$ does not contain any reflections associated to the divisor of $\Theta_2 \Theta_3 \Theta_4$. Indeed, the multiplier systems v_2, v_3 and v_4 coincide on $O_{1,1}(2U(4) \oplus A_1)$ because the quotient of any two of them defines a character whose values on generators are always 1. If $O_{1,1}(2U(4) \oplus A_1)$ contained a reflection σ associated to the divisor of Θ_2 then we would find $v_2(\sigma) = -1$ but $v_3(\sigma) = v_4(\sigma) = 1$, a contradiction. Similarly, $O_{1,12}(2U(4) \oplus A_1)$ does not contain any reflections associated to the divisor of $\Theta_3 \Theta_4$.

On the other hand, $O_{1,123}(2U(4) \oplus A_1)$ does contain reflections associated to the divisor of Θ_4 . In fact,

$$(3.5) \quad O_{1,123}(2U(4) \oplus A_1) = O_2(2U(4) \oplus A_1).$$

(Of course, this agrees with Theorem 3.10.) If this was not true, then the only alternative would be $M_*(O_{1,123}(2U(4) \oplus A_1), v_4^2) = \mathbb{C}[\Theta_i^2, i = 1, 2, 3, 4]$, contradicting the decomposition of the Jacobian in Theorem 2.3.

Finally we obtain some larger groups generated by reflections and free algebras of modular forms associated to triples of $*$ -sets. Suppose Θ_1, Θ_2 is one of the 90 pairs of type II products that can be extended to three distinct $*$ -sets, denoted

$$\{\Theta_1, \Theta_2, \Theta_3, \Theta_4\}, \quad \{\Theta_1, \Theta_2, \Theta_5, \Theta_6\}, \quad \{\Theta_1, \Theta_2, \Theta_7, \Theta_8\}.$$

The principal part into the input function to Θ_i will be denoted $q^{-1/4}(e_{u_i} + e_{-u_i})$.

In this situation the reflection σ_{u_5} fixes the set $\{\pm u_1, \pm u_2, \pm u_3, \pm u_4\}$ and has u_1 and u_2 as eigenvectors. It cannot have u_3 and u_4 as eigenvectors as otherwise the proof of Lemma 3.4 yields five linearly independent type II products. Therefore it maps u_3 to $\pm u_4$ and u_4 to $\pm u_3$. Taking Borcherds lifts and using the fact that all type II products have Fourier coefficients in $\mathbb{Z}[i]$, this implies $\sigma_{u_5}(\Theta_3) = i^k \Theta_4$ for some $k \in \{0, 1, 2, 3\}$. The same statements hold for the reflections σ_{u_i} , $i = 6, 7, 8$.

Let $O_{2,56}(2U(4) \oplus A_1)$ be the group generated by $O_2(2U(4) \oplus A_1)$ and the reflections associated to the divisor of Θ_5 . Since σ_{u_3} swaps u_5 and u_6 (up to multiples), $O_{2,56}(2U(4) \oplus A_1)$ also contains the reflections associated to the divisor of Θ_6 . It is clear that $\Theta_3^2\Theta_4^2 \in M_2(O_{2,56}(2U(4) \oplus A_1))$. From the previous paragraph it follows that $\sigma_{u_5}(\Theta_3^2) = c\Theta_4^2$ where $c \in \{\pm 1\}$. Then $\sigma_{u_5}(\Theta_4^2) = c\Theta_3^2$ and therefore $\Theta_3^2 + c\Theta_4^2 \in M_1(O_{2,56}(2U(4) \oplus A_1))$.

On the other hand, $\Theta_5\Theta_6 \in M_1(O_2(2U(4) \oplus A_1))$ is a \mathbb{C} -linear combination of Θ_i^2 for $i = 1, 2, 3, 4$. By considering the action of σ_{u_5} on these functions, we find that $\Theta_5\Theta_6$ is equal to $\Theta_3^2 - c\Theta_4^2$ up to a non-zero constant multiple. Using Lemma 3.5, we see that the Jacobian of $\Theta_1^2, \Theta_2^2, \Theta_3^2 + c\Theta_4^2$ and $\Theta_3^2\Theta_4^2$ is equal to $\Phi_{5,A_1} \prod_{i=1}^6 \Theta_i$ up to a non-zero multiple. We have therefore proved the following:

$$M_*(O_{2,56}(2U(4) \oplus A_1)) = \mathbb{C}[\Theta_1^2, \Theta_2^2, \Theta_3^2 + c\Theta_4^2, \Theta_3^2\Theta_4^2].$$

Similarly, since $\Theta_7\Theta_8 \in M_1(O_{2,56}(2U(4) \oplus A_1))$, it is equal to $\Theta_3^2 + c\Theta_4^2$ up to a non-zero constant by considering the action of σ_{u_7} . Letting $O_{2,78}(2U(4) \oplus A_1)$ be the group generated by $O_2(2U(4) \oplus A_1)$ and the reflections associated to the divisor of Θ_7 , we obtain by the same argument

$$M_*(O_{2,78}(2U(4) \oplus A_1)) = \mathbb{C}[\Theta_1^2, \Theta_2^2, \Theta_3^2 - c\Theta_4^2, \Theta_3^2\Theta_4^2].$$

Finally, let $O_{2,5678}(2U(4) \oplus A_1)$ be the group generated by $O_2(2U(4) \oplus A_1)$ and reflections associated to the divisor of $\Theta_5\Theta_7$. Then this group also contains the reflections associated to the divisor of $\Theta_6\Theta_8$. Using Lemma 3.5 we derive that the Jacobian of $\Theta_1^2, \Theta_2^2, \Theta_3^2\Theta_4^2, \Theta_5^2\Theta_6^2$ equals $\Phi_{5,A_1} \prod_{i=1}^8 \Theta_i$ up to a non-zero constant multiple, and have proved the following:

$$M_*(O_{2,5678}(2U(4) \oplus A_1)) = \mathbb{C}[\Theta_1^2, \Theta_2^2, \Theta_3^2\Theta_4^2, \Theta_5^2\Theta_6^2].$$

Altogether we have computed the algebras of modular forms for eight subgroups of $O^+(2U \oplus A_1)$:

$$\begin{aligned} O'_1(2U(4) \oplus A_1) &\subsetneq O_0(2U(4) \oplus A_1) \subsetneq O_1(2U(4) \oplus A_1) \subsetneq O_{1,1}(2U(4) \oplus A_1) \subsetneq \\ &\subsetneq O_{1,12}(2U(4) \oplus A_1) \subsetneq O_2(2U(4) \oplus A_1) \subsetneq O_{2,56}(2U(4) \oplus A_1) \subsetneq O_{2,5678}(2U(4) \oplus A_1). \end{aligned}$$

Remark 3.11. There is no multiplier system v of $O_2(U(4) \oplus A_1)$ which equals 1 on all reflections. If such a v existed, then v_4/v would define a character of $O_2(U(4) \oplus A_1)$ which equals 1 on all reflections associated to the divisor of $\Theta_1\Theta_2\Theta_3$ but is -1 on all reflections associated to the divisor of Θ_4 , which contradicts (3.5).

Remark 3.12. The Baily-Borel compactification of the modular variety $\mathcal{D}_3/O_1(2U(4) \oplus A_1)$ is a projective space, but the algebra $M_*(O_1(2U(4) \oplus A_1))$ of modular forms of integral weight is not free. It is well-known that if $M_*(\Gamma)$ is free then $(\mathcal{D}/\Gamma)^*$ is a weighted projective space (see [2]), but the above example shows that the converse does not hold. Eberhard Freitag suggested to the authors that the following statement may hold.

Conjecture 3.13. If the modular variety $(\mathcal{D}/\Gamma)^*$ is a weighted projective space, then there exists a weight k_0 and a multiplier system v_0 of weight k_0 such that the graded algebra

$$M_*(\Gamma, (k_0, v_0)) := \bigoplus_{k \in \mathbb{N}} M_{kk_0}(\Gamma, v_0^k)$$

is freely generated.

Remark 3.14. There are twenty linearly independent modular forms of weight $1/2$ for the Weil representation attached to $2U(4) \oplus A_1$, and their images under the additive theta lift ([3], Theorem 14.3) span the 10-dimensional space of modular forms of weight one for $O_1(2U(4) \oplus A_1)$.

4. THE $2U(2) \oplus 2A_1$ LATTICE

In this section we prove Matsumoto's theorem in the context of modular forms on $2U(2) \oplus 2A_1$. We first work out some Borcherds products on $2U(2) \oplus 2A_1$.

Lemma 4.1.

- (1) *There are 10 holomorphic Borcherds products of singular weight 1. Their inputs have principal parts of the form*

$$q^{-1/4}(e_{v_1} + e_{v_2}) + 2e_0, \quad (v_1, v_1) = (v_2, v_2) = 1/2, \quad \text{ord}(v_1) = \text{ord}(v_2) = 2,$$

where v_1 and v_2 are images of each other under swapping the two A_1 components. In particular they are 2-reflective modular forms. We label the ten forms F_i , $1 \leq i \leq 10$ such that with respect to the Gram matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

their principal parts are as follows:

$$\begin{aligned} F_1 &: q^{-1/4}(e_{(0,1/2,1/2,0,0,0)} + e_{(0,1/2,0,1/2,0,0)}); & F_2 &: q^{-1/4}(e_{(1/2,0,1/2,0,0,0)} + e_{(1/2,0,0,1/2,0,0)}); \\ F_3 &: q^{-1/4}(e_{(1/2,1/2,1/2,0,1/2,1/2)} + e_{(1/2,1/2,0,1/2,1/2,1/2)}); & F_4 &: q^{-1/4}(e_{(1/2,0,1/2,0,1/2,0)} + e_{(1/2,0,0,1/2,1/2,0)}); \\ F_5 &: q^{-1/4}(e_{(0,1/2,1/2,0,0,1/2)} + e_{(0,1/2,0,1/2,0,1/2)}); & F_6 &: q^{-1/4}(e_{(1/2,1/2,1/2,0,0,0)} + e_{(1/2,1/2,0,1/2,0,0)}); \\ F_7 &: q^{-1/4}(e_{(0,0,1/2,0,1/2,1/2)} + e_{(0,0,0,1/2,1/2,1/2)}); & F_8 &: q^{-1/4}(e_{(0,0,1/2,0,1/2,0)} + e_{(0,0,0,1/2,1/2,0)}); \\ F_9 &: q^{-1/4}(e_{(0,0,1/2,0,0,1/2)} + e_{(0,0,0,1/2,0,1/2)}); & F_{10} &: q^{-1/4}(e_{(0,0,1/2,0,0,0)} + e_{(0,0,0,1/2,0,0)}). \end{aligned}$$

The ten F_i are linearly independent over \mathbb{C} . We define $\Phi_{10,2A_1} := \prod_{j=1}^{10} F_j$.

- (2) *There are 15 reflective Borcherds products of weight 2 with principal parts of the form $q^{-1/2}e_u$, where $(u, u) = 1$, $\text{ord}(u) = 2$ and u is invariant under the swapping of two A_1 components.*
- (3) *There is a holomorphic Borcherds product of weight 4, which we denote $\Phi_{4,2A_1}$, whose input has the principal part $q^{-1/2}e_{(0,0,1/2,1/2,0,0)}$. The reflection σ_1 associated to the vector $(0, 0, 1/2, 1/2, 0, 0)$ swaps the two A_1 components.*

Let $O_1(2U(2) \oplus 2A_1)$ be the subgroup of $O^+(2U(2) \oplus 2A_1)$ generated by all reflections associated to the divisors of $\Phi_{10,2A_1}$ and $\Phi_{4,2A_1}$. The inputs of all F_i and of $\Phi_{4,2A_1}$ are invariant under the reflection σ_1 , so F_i and $\Phi_{4,2A_1}$ are modular forms for $O_1(2U(2) \oplus 2A_1)$. Each F_i has a quadratic character on $O_1(2U(2) \oplus 2A_1)$ so their squares F_i^2 have trivial character.

Lemma 4.2. *The Jacobian of F_i^2 , $1 \leq i \leq 5$ equals $\Phi_{10,2A_1}\Phi_{4,2A_1}$ up to a non-zero constant multiple. In particular, the forms F_i for $1 \leq i \leq 5$ are algebraically independent over \mathbb{C} .*

Proof. By computing the first few terms of the Fourier expansion we find that $J = J(F_i^2, 1 \leq i \leq 5)$ is not identically zero. By Theorem 2.2, J is a modular form of weight 14 and vanishes on mirrors of all reflections in $O_1(2U(2) \oplus 2A_1)$. In particular $J/(\Phi_{10,2A_1}\Phi_{4,2A_1})$ is a holomorphic modular form of weight 0 and therefore constant. \square

By Theorem 2.4, we obtain the following result which is equivalent to Matsumoto's theorem.

Theorem 4.3.

$$\begin{aligned} M_*(O_1(2U(2) \oplus 2A_1)) &= \mathbb{C}[F_i^2, 1 \leq i \leq 5], \\ (\mathcal{D}_4/O_1(2U(2) \oplus 2A_1))^* &\cong \mathbb{P}^4(\mathbb{C}). \end{aligned}$$

Corollary 4.4. *The squares F_i^2 span a five-dimensional space over \mathbb{C} and they satisfy the four-term relations*

$$\begin{aligned} F_1^2 - F_5^2 + F_6^2 - F_{10}^2 &= 0, \\ F_2^2 - F_5^2 - F_8^2 - F_9^2 &= 0, \\ F_3^2 - F_5^2 + F_6^2 - F_9^2 &= 0, \\ F_4^2 - F_6^2 - F_8^2 + F_{10}^2 &= 0, \\ F_7^2 - F_8^2 - F_9^2 + F_{10}^2 &= 0. \end{aligned}$$

The vector space spanned by the 15 weight two products in Lemma 4.1 and the ten squares F_i^2 has dimension 5.

Proof. The exact form of the relations among the F_i^2 can be read off of their Fourier expansions. The 15 weight two products have trivial character on $O_1(2U(2) \oplus 2A_1)$ and therefore lie in the span of F_1^2, \dots, F_5^2 . \square

We will now obtain an analogue of Igusa's theorem for Hermitian modular forms of degree 2 over the Gaussian numbers. The Jacobian $J = J(F_i^2, 1 \leq i \leq 5)$ is the product of the ten F_i and $\Phi_{4,2A_1}$. To apply Theorem 2.3 we have to show that $\Phi_{4,2A_1}$ is irreducible as a modular form on $O_1(2U(2) \oplus 2A_1)$, i.e. it is not a product of two non-constant modular forms with characters (or multiplier systems) on $O_1(2U(2) \oplus 2A_1)$. Since $\Phi_{4,2A_1}^2 \in M_8(O_1(2U(2) \oplus 2A_1))$, it can be expressed as a polynomial P in terms of the five F_i^2 . We computed this polynomial and found that it is irreducible. From Theorem 2.3, it follows that $\Phi_{4,2A_1}$ is irreducible on $O_1(2U(2) \oplus 2A_1)$.

By Theorem 2.3, there are exactly 2048 characters of $O_1(2U(2) \oplus 2A_1)$ and they are generated by the basic characters χ_i of F_i and the character of $\Phi_{4,2A_1}$. Let $O'_1(2U(2) \oplus 2A_1)$ be the commutator subgroup of $O_1(2U(2) \oplus 2A_1)$. Then

$$M_*(O'_1(2U(2) \oplus 2A_1)) = \mathbb{C}[\Phi_{4,2A_1}, F_i, 1 \leq i \leq 10].$$

Let O_0 be the subgroup

$$O_0(2U(2) \oplus 2A_1) = \{\gamma \in O_1(2U(2) \oplus 2A_1) : \chi_i(\gamma) = 1, 1 \leq i \leq 10\}.$$

We have the following result immediately.

Theorem 4.5.

$$M_*(O_0(2U(2) \oplus 2A_1)) = \mathbb{C}[F_i, 1 \leq i \leq 10].$$

Similarly, we define another subgroup of $O_1(2U(2) \oplus 2A_1)$ via

$$O_{1'}(2U(2) \oplus 2A_1) = \{\gamma \in O_1(2U(2) \oplus 2A_1) : \chi_i(\gamma) = 1, 1 \leq i \leq 5\}.$$

The group $O_{1'}(2U(2) \oplus 2A_1)$ contains all reflections associated to the divisor of $\Phi_{4,2A_1} \prod_{j=6}^{10} F_j$. The forms F_i , $1 \leq i \leq 5$, are modular with trivial character on $O_{1'}(2U(2) \oplus 2A_1)$ and their Jacobian equals $\Phi_{4,2A_1} \prod_{j=6}^{10} F_j$ up to a non-zero constant multiple by Lemma 4.2. Therefore we obtain the following result.

Theorem 4.6.

$$\begin{aligned} M_*(O_{1'}(2U(2) \oplus 2A_1)) &= \mathbb{C}[F_i, 1 \leq i \leq 5], \\ (\mathcal{D}_4 / O_{1'}(2U(2) \oplus 2A_1))^* &\cong \mathbb{P}^4(\mathbb{C}). \end{aligned}$$

The above theorem also implies that $O_{1'}(2U(2) \oplus 2A_1)$ is generated by all reflections associated to the divisor of $\Phi_{4,2A_1} \prod_{j=6}^{10} F_j$. (Note that the same argument applies with F_1, \dots, F_5 replaced by

any of the 162 linearly independent sets of five squares of products F_i^2 . These sets can be read off of the four-term relations of Corollary 4.4.)

We define a larger subgroup of $O_1(2U(2) \oplus 2A_1)$ via

$$O_{1''}(2U(2) \oplus 2A_1) = \{\gamma \in O_1(2U(2) \oplus 2A_1) : \chi_1(\gamma) = 1\},$$

i.e. the subgroup generated by all reflections associated to the divisor of $\Phi_{4,2A_1} \prod_{j=2}^{10} F_j$. A similar argument to the above theorem yields the following result.

Theorem 4.7.

$$\begin{aligned} M_*(O_{1''}(2U(2) \oplus 2A_1)) &= \mathbb{C}[F_1, F_2^2, F_3^2, F_4^2, F_5^2], \\ (\mathcal{D}_4/O_{1''}(2U(2) \oplus 2A_1))^* &\cong \mathbb{P}(1, 2, 2, 2, 2) \cong \mathbb{P}^4(\mathbb{C}). \end{aligned}$$

The ring of even-weight modular forms for $O_{1''}(2U(2) \oplus 2A_1)$ is freely generated in weight two (and indeed is exactly the ring from Theorem 4.3):

$$M_{2*}(O_{1''}(2U(2) \oplus 2A_1))^{\text{even}} = \mathbb{C}[F_1^2, F_2^2, F_3^2, F_4^2, F_5^2].$$

Remark 4.8. There is a six-dimensional space of modular forms of weight 1 for the Weil representation attached to $2U(2) \oplus 2A_1$, and these map to a five-dimensional space of modular forms of weight 2 with trivial character on the discriminant kernel $\tilde{O}^+(2U(2) \oplus 2A_1)$ under the additive theta lift. Applying Theorem 2.4 to $\tilde{O}^+(2U(2) \oplus 2A_1)$, we find that $M_*(\tilde{O}^+(2U(2) \oplus 2A_1))$ is freely generated by the five additive lifts. Moreover, $\tilde{O}^+(2U(2) \oplus 2A_1)$ is generated by all 2-reflections and then

$$\tilde{O}^+(2U(2) \oplus 2A_1) = O_1(2U(2) \oplus 2A_1).$$

By the Eichler criterion (cf. Proposition 3.3 of [9]), the divisor of $\Phi_{4,2A_1}$ is irreducible on $\tilde{O}^+(2U(2) \oplus 2A_1)$. In this way we obtain a different proof that $\Phi_{4,2A_1}$ is irreducible on $O_1(2U(2) \oplus 2A_1)$. We also see that the square of each of the singular weight products is an additive lift.

Remark 4.9. By [8, Lemma 6.1], we have

$$O^+(2U(2) \oplus 2A_1) \cong O^+(2U \oplus 2A_1).$$

Thus $\Phi_{10,2A_1}$ can be regarded as a modular form on $2U \oplus 2A_1$. In the interpretation as Hermitian modular forms this is the Borchers product ϕ_{10} in [5, Corollary 4], which was realized earlier by Freitag [6] as the product of ten theta constants. Indeed, our F_i are exactly the ten theta constants when interpreted as Hermitian modular forms.

5. THE $2U(3) \oplus A_2$ LATTICE

In this section we reprove the theorem of Freitag and Salvati Manni in the context of modular forms on $2U(3) \oplus A_2$. We first work out some Borchers products on $2U(3) \oplus A_2$:

- (1) There are 45 holomorphic Borchers products of singular weight 1. Their inputs have principal parts of the form

$$q^{-1/3}(e_v + e_{-v}) + 2e_0, \quad (v, v) = 2/3, \text{ ord}(v) = 3.$$

The product of these 45 forms is a reflective modular form Φ_{45,A_2} of weight 45 which can be viewed as a 2-reflective modular form for $O^+(2U \oplus A_2)$.

- (2) There is a holomorphic Borchers product of weight 9 whose input has principal part

$$(q^{-1} + 18)e_0.$$

We label this form Φ_{9,A_2} . It is a 2-reflective modular form on $O^+(2U(3) \oplus A_2)$ and can be regarded as a reflective modular form for $O^+(2U \oplus A_2)$.

Let $O_1(2U(3) \oplus A_2)$ be the subgroup of $O^+(2U(3) \oplus A_2)$ generated by all 2-reflections, i.e. reflections associated to the divisor of Φ_{9,A_2} . It is clear that $O_1(2U(3) \oplus A_2)$ is a subgroup of $\tilde{O}^+(2U(3) \oplus A_2)$. By considering their divisors, we see that the 45 weight 1 products have trivial character on $O_1(2U(3) \oplus A_2)$.

It will again be convenient to use the notion of $*$ -sets.

Definition 5.1. A $*$ -set (of type $2U(3) \oplus A_2$) is a set of five products of weight 1 on $2U(3) \oplus A_2$ whose inputs are invariant under all reflections associated to the divisors of any of the five products.

$*$ -sets of type $2U(3) \oplus A_2$ satisfy properties analogous to Lemma 3.4:

Lemma 5.2.

- (1) *There are exactly 27 $*$ -sets.*
- (2) *Every product of weight 1 belongs to exactly three $*$ -sets.*
- (3) *There are 720 pairs of weight 1 products that do not belong to a $*$ -set. The remaining 270 pairs belong to a unique $*$ -set.*
- (4) *The five elements of any $*$ -set are linearly independent over \mathbb{C} .*

Lemma 5.3. *There is a $*$ -set $\{G_1, G_2, G_3, G_4, G_5\}$ whose Jacobian equals Φ_{9,A_2} up to a non-zero constant multiple.*

Proof. In the notation of Appendix B, we used the forms $G_1, G_{14}, G_{15}, G_{23}, G_{27}$ which form a $*$ -set. We computed their Jacobian to precision 10 and found that it is not identically zero (and indeed agrees with the Fourier expansion of Φ_{9,A_2}). By the same argument as Lemma 3.5, the quotient $J/\Phi_{9,A_2}$ is holomorphic of weight zero and therefore constant. \square

By applying Theorem 2.4 we obtain the theorem of Freitag and Salvati Manni in the context of orthogonal groups.

Theorem 5.4. *The 45 singular-weight products on $2U(3) \oplus A_2$ span a five-dimensional space over \mathbb{C} . Any five linearly independent products G_1, \dots, G_5 are algebraically independent and generate the algebra of modular forms:*

$$M_*(O_1(2U(3) \oplus A_2)) = \mathbb{C}[G_1, G_2, G_3, G_4, G_5],$$

$$(\mathcal{D}_4/O_1(2U(3) \oplus A_2))^* \cong \mathbb{P}^4(\mathbb{C}).$$

We also have the following analogue of Perna's theorem.

Theorem 5.5. *Let $\{G_1, \dots, G_5\}$ be a $*$ -set and let $O_2(2U(3) \oplus A_2)$ be the subgroup generated by $O_1(2U(3) \oplus A_2)$ and the reflections associated to the divisor of $\prod_{j=1}^5 G_j$. Then*

$$M_*(O_2(2U(3) \oplus A_2)) = \mathbb{C}[G_1^2, G_2^2, G_3^2, G_4^2, G_5^2],$$

$$(\mathcal{D}_4/O_2(2U(3) \oplus A_2))^* \cong \mathbb{P}^4(\mathbb{C}).$$

In Remark 5.7 below, we will see that the squared Jacobian Φ_{9,A_2}^2 is irreducible in $M_*(O_1(2U(3) \oplus A_2))$. From Theorem 2.3 it follows that \det is the only non-trivial character of $O_1(2U(3) \oplus A_2)$. Let $O'_1(2U(3) \oplus A_2)$ be the commutator subgroup of $O_1(2U(3) \oplus A_2)$. Then

$$M_*(O'_1(2U(3) \oplus A_2)) = \mathbb{C}[\Phi_{9,A_2}, G_i, 1 \leq i \leq 5].$$

As in §3 we can construct a tower of free algebras of modular forms. We fix a $*$ -set $\{G_1, \dots, G_5\}$. Let $O_{1,1}(2U(3) \oplus A_2)$, $O_{1,12}(2U(3) \oplus A_2)$, $O_{1,123}(2U(3) \oplus A_2)$ and $O_{1,1234}(2U(3) \oplus A_2)$ be the subgroups generated by reflections associated to the divisors of $\Phi_{9,A_2} G_1$, $\Phi_{9,A_2} \prod_{j=1}^2 G_j$, $\Phi_{9,A_2} \prod_{j=1}^3 G_j$

and $\Phi_{9,A_2} \prod_{j=1}^4 G_j$ respectively. It is easy to derive the following structure results:

$$\begin{aligned} M_*(\mathcal{O}_{1,1}(2U(3) \oplus A_2)) &= \mathbb{C}[G_1^2, G_2, G_3, G_4, G_5], \\ M_*(\mathcal{O}_{1,12}(2U(3) \oplus A_2)) &= \mathbb{C}[G_1^2, G_2^2, G_3, G_4, G_5], \\ M_*(\mathcal{O}_{1,123}(2U(3) \oplus A_2)) &= \mathbb{C}[G_1^2, G_2^2, G_3^2, G_4, G_5], \\ M_*(\mathcal{O}_{1,1234}(2U(3) \oplus A_2)) &= \mathbb{C}[G_1^2, G_2^2, G_3^2, G_4^2, G_5]. \end{aligned}$$

From the last of these we obtain another realization of $\mathbb{P}^4(\mathbb{C})$ as an orthogonal modular variety:

Theorem 5.6.

$$(\mathcal{D}_4/\mathcal{O}_{1,1234}(2U(3) \oplus A_2))^* \cong \mathbb{P}(1, 2, 2, 2, 2) \cong \mathbb{P}^4(\mathbb{C}).$$

Remark 5.7. The space of invariants of the Weil representation attached to $2U(3) \oplus A_2$ is ten-dimensional, and these are mapped to a five-dimensional space of modular forms of weight one with trivial character on the full discriminant kernel $\tilde{\mathcal{O}}^+(2U(3) \oplus A_2)$ under the additive theta lift. In particular the 45 singular weight products are all additive lifts. Using the argument in Remark 4.8 we conclude that $M_*(\tilde{\mathcal{O}}^+(2U(3) \oplus A_2))$ is freely generated by the five additive lifts and that

$$\tilde{\mathcal{O}}^+(2U(3) \oplus A_2) = \mathcal{O}_1(2U(3) \oplus A_2),$$

and the Eichler criterion implies that Φ_{9,A_2} is irreducible in $M_*(\mathcal{O}_1(2U(3) \oplus A_2))$.

6. THE $2U(2) \oplus A_1$ LATTICE

In this section we will find three interesting free algebras of modular forms associated to the lattice $2U(2) \oplus A_1$. We first describe some Borcherds products on $2U(2) \oplus A_1$ of small weight.

- (1) There are nine holomorphic Borcherds products of weight 1 and one holomorphic product of weight 2 on $2U(2) \oplus A_1$. All have principal parts of the form

$$q^{-1/4}e_v, \quad (v, v) = 1/2, \quad \text{ord}(v) = 2.$$

The product of these ten forms, which we denote $\Phi_{11,A_1(2)}$, is a 2-reflective modular form of weight 11 on $2U(2) \oplus A_1$. This can also be viewed as a modular form on $2U \oplus A_1(2)$ by [8, Lemma 6.1] because

$$\mathcal{O}^+(2U(2) \oplus A_1) \cong \mathcal{O}^+(2U \oplus A_1(2)).$$

- (2) There are six holomorphic Borcherds products f_1, \dots, f_6 of weight 2 with principal parts

$$q^{-1/2}e_v, \quad (v, v) = 1, \quad \text{ord}(v) = 2.$$

We fix the Gram matrix $\begin{pmatrix} 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 \end{pmatrix}$ and label the six products f_1, \dots, f_6 above such that their principal parts are as follows:

$$\begin{aligned} f_1 : v &= (0, 1/2, 0, 1/2, 0); & f_2 : v &= (0, 1/2, 0, 1/2, 1/2); \\ f_3 : v &= (1/2, 0, 0, 1/2, 1/2); & f_4 : v &= (1/2, 0, 0, 0, 1/2); \\ f_5 : v &= (1/2, 1/2, 0, 0, 1/2); & f_6 : v &= (1/2, 1/2, 0, 1/2, 0). \end{aligned}$$

Let $\mathcal{O}_1(2U(2) \oplus A_1)$ be the subgroup of $\mathcal{O}^+(2U(2) \oplus A_1)$ generated by all reflections associated to the divisor of $\Phi_{11,A_1(2)}$.

Lemma 6.1.

- (1) *The six products f_i are modular forms with trivial character on $\mathcal{O}_1(2U(2) \oplus A_1)$.*

(2) The Jacobian $J(f_1, f_2, f_3, f_4)$ equals $\Phi_{11, A_1(2)}$ up to a non-zero multiple.

Proof. (1) This can be seen from the divisor of the f_i .

(2) We checked by computer that $J = J(f_1, \dots, f_4)$ is nonzero (and indeed equals $768\Phi_{11, A_1(2)}$ up to precision $O(q, s)^{10}$). Applying Theorem 2.2 (4) as in the earlier sections shows that $J/\Phi_{11, A_1(2)}$ is holomorphic of weight zero and therefore a constant. \square

Therefore we can apply Theorem 2.4 to this situation. This yields another realization of $\mathbb{P}^3(\mathbb{C})$ as a modular variety.

Theorem 6.2. *The six products f_1, \dots, f_6 span a four-dimensional space and satisfy the relations*

$$f_1 + f_3 + f_5 = f_2 + f_4 + f_6 = 0.$$

Any four that are linearly independent are algebraically independent and generate the ring of modular forms for $O_1(2U(2) \oplus A_1)$:

$$\begin{aligned} M_*(O_1(2U(2) \oplus A_1)) &= \mathbb{C}[f_1, f_2, f_3, f_4], \\ (\mathcal{D}_3/O_1(2U(2) \oplus A_1))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Proof. Theorem 2.4 implies everything except the exact form of the relations among the f_i , which can be determined from their Fourier expansions. \square

Remark 6.3. The squares of the nine Borcherds products b_i of weight one are modular forms without character for $O_1(2U(2) \oplus A_1)$ and therefore lie in the span of f_1, \dots, f_4 . Indeed they also span this space and they satisfy five three-term linear relations of the form $b_i^2 + b_j^2 = b_k^2$ as one can check from their Fourier expansions.

Choose any linearly independent squares of weight one Borcherds products b_1^2, b_2^2, b_3^2 and b_4^2 as in the remark. Let $O_{1'}(2U(2) \oplus A_1)$ be the subgroup of $O_1(2U(2) \oplus A_1)$ generated by all reflections associated to the divisor of $\Phi_{11, A_1(2)}/(\prod_{j=1}^4 b_j)$. Similarly to the case of $2U(2) \oplus 2A_1$, the four forms b_j are modular with trivial character on $O_{1'}(2U(2) \oplus A_1)$ and their Jacobian equals $\Phi_{11, A_1(2)}/(\prod_{j=1}^4 b_j)$ up to a nonzero multiple. From this we obtain another realization of $\mathbb{P}^3(\mathbb{C})$ as a modular variety:

Theorem 6.4.

$$\begin{aligned} M_*(O_{1'}(2U(2) \oplus A_1)) &= \mathbb{C}[b_1, b_2, b_3, b_4], \\ (\mathcal{D}_3/O_{1'}(2U(2) \oplus A_1))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Furthermore, we define $O_{1''}(2U(2) \oplus A_1)$ as the subgroup of $O_1(2U(2) \oplus A_1)$ generated by all reflections associated to the divisor of $\Phi_{11, A_1(2)}/b_1$. Then

Theorem 6.5.

$$\begin{aligned} M_*(O_{1''}(2U(2) \oplus A_1)) &= \mathbb{C}[b_1, b_2^2, b_3^2, b_4^2], \\ (\mathcal{D}_3/O_{1''}(2U(2) \oplus A_1))^* &\cong \mathbb{P}(1, 2, 2, 2) \cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Remark 6.6. The space of modular forms of weight $3/2$ for the Weil representation attached to $2U(2) \oplus A_1$ is five-dimensional, and these forms map to a four-dimensional space of modular forms of weight 2 under the additive theta lift. In particular, every modular form in $M_2(O_1(2U(2) \oplus A_1))$ is an additive lift and therefore has trivial character on the full discriminant kernel. Similarly to Remark 4.8, we find that $M_*(\tilde{O}^+(2U(2) \oplus A_1))$ is freely generated by the four additive lifts and that

$$\tilde{O}^+(2U(2) \oplus A_1) = O_1(2U(2) \oplus A_1).$$

Similarly to the previous sections, the decomposition of the Jacobian $\Phi_{11,A_1(2)}$ determines the structure of the algebra of modular forms for the commutator group $O'_1(2U(2) \oplus A_1)$ and some related groups. We omit the details.

7. THE $2U(3) \oplus A_1$ LATTICE

In this section we determine an interesting free algebra of modular forms on $2U(3) \oplus A_1$. We first work out Borcherds products on $2U(3) \oplus A_1$:

- (1) There are 29 holomorphic Borcherds products of weight 1 on $2U(3) \oplus A_1$. Sixteen of them have principal parts of the form

$$q^{-1/4}(e_v + e_{-v}), \quad (v, v) = 1/2, \text{ ord}(v) = 6,$$

and they are not reflective modular forms. Twelve of them have principal parts

$$q^{-1/3}(e_v + e_{-v}), \quad (v, v) = 2/3, \text{ ord}(v) = 3,$$

and they are reflective modular forms. The last one is a 2-reflective modular form denoted by Δ_1 with principal part

$$q^{-1/4}e_{(0,0,1/2,0,0)}$$

with respect to the Gram matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 3 & 0 & 0 & 0 \\ 3 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

- (2) There is a holomorphic Borcherds product of weight 7 with principal part $q^{-1}e_0$. We label this 2-reflective form $\Phi_{7,A_1(3)}$.

Let $O_1(2U(3) \oplus A_1)$ be the subgroup of $O^+(2U(3) \oplus A_1)$ generated by all reflections associated to the divisor of $\Phi_{7,A_1(3)}$. The 28 products of weight 1 other than Δ_1 are modular forms of trivial character on $O_1(2U(3) \oplus A_1)$.

Lemma 7.1. *There are four products g_1, \dots, g_4 of weight 1 whose Jacobian $J(g_1, g_2, g_3, g_4)$ equals $\Phi_{7,A_1(3)}$ up to a non-zero multiple.*

Proof. We used the products g_i whose inputs have principal parts $q^{-1/3}(e_{v_i} + e_{-v_i})$, where

$$\begin{aligned} v_1 &= (1/3, 2/3, 0, 2/3, 0), & v_2 &= (1/3, 2/3, 0, 1/3, 2/3), \\ v_3 &= (1/3, 1/3, 0, 2/3, 2/3), & v_4 &= (1/3, 1/3, 0, 1/3, 0), \end{aligned}$$

(although it will turn out that any four linearly independent forms not including Δ_1 will do) and using Fourier series computed that their Jacobian J is nonzero (and equals $(12\zeta_3 + 6)\Phi_{7,A_1(3)}$ up to precision $O(q, s)^{10}$). As in the previous sections $J/\Phi_{7,A_1(3)}$ is holomorphic of weight zero and therefore constant. \square

By applying Theorem 2.4 we obtain the following theorem.

Theorem 7.2. *Every linearly independent set g_1, \dots, g_4 of weight one products that does not include Δ_1 is algebraically independent and generates the ring of modular forms for $O_1(2U(3) \oplus A_1)$:*

$$\begin{aligned} M_*(O_1(2U(3) \oplus A_1)) &= \mathbb{C}[g_1, g_2, g_3, g_4], \\ (\mathcal{D}_3/O_1(2U(3) \oplus A_1))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

In particular, the 28 products of weight 1 other than Δ_1 span a 4-dimensional space. (Δ_1 has a nontrivial character on $O_1(2U(3) \oplus A_1)$ and in particular does not lie in their span.)

Remark 7.3. The space of modular forms of weight $1/2$ for the Weil representation attached to $2U(3) \oplus A_1$ is 8-dimensional, and these map to a four-dimensional space of modular forms of weight one under the additive theta lift. It follows that all of the 28 products of weight 1 other than Δ_1 are additive lifts. Similarly to $2U(2) \oplus 2A_1$, we conclude that $M_*(\tilde{O}^+(2U(3) \oplus A_1))$ is freely generated by the four additive lifts and that

$$\tilde{O}^+(2U(3) \oplus A_1) = \mathcal{O}_1(2U(3) \oplus A_1).$$

By the Eichler criterion, the forms Δ_1 and $\Phi_{7,A_1(3)}/\Delta_1$ are irreducible on $\mathcal{O}_1(2U(3) \oplus A_1)$.

8. THE $U(4) \oplus U(2) \oplus A_1$ LATTICE

In this section we determine two interesting free algebras of modular forms on $U(4) \oplus U(2) \oplus A_1$. We first work out Borcherds products on $U(4) \oplus U(2) \oplus A_1$:

- (1) There are 8 holomorphic Borcherds products of weight $1/2$ and 3 holomorphic products of weight 1 on $U(4) \oplus U(2) \oplus A_1$, all with principal parts of the form

$$q^{-1/4}e_v, \quad (v, v) = 1/2, \quad \text{ord}(v) = 2.$$

Their product is a 2-reflective modular form of weight 7 which we label $\Phi_{7,A_1(4)}$.

- (2) There are 16 other holomorphic Borcherds products of weight 1. They are all reflective. Twelve of them have input forms with principal parts

$$q^{-1/4}(e_v + e_{-v}), \quad (v, v) = 1/2, \quad \text{ord}(v) = 4.$$

The remaining four have input forms with principal parts

$$q^{-1/2}e_v, \quad (v, v) = 1, \quad \text{ord}(v) = 2.$$

Let $\mathcal{O}_1(U(4) \oplus U(2) \oplus A_1)$ be the subgroup of $\mathcal{O}^+(U(4) \oplus U(2) \oplus A_1)$ generated by all reflections associated to the divisor of $\Phi_{7,A_1(4)}$. Then the 16 products of weight 1 in (2) are modular forms of trivial character on $\mathcal{O}_1(U(4) \oplus U(2) \oplus A_1)$.

Let h_1, \dots, h_4 denote the four products in (2) whose input forms have principal part $q^{-1/2}e_v$ with v of order two.

Lemma 8.1. *The Jacobian $J = J(h_1, h_2, h_3, h_4)$ is equals $\Phi_{7,A_1(4)}$ up to a non-zero constant multiple. The four forms h_1, h_2, h_3 and h_4 are algebraically independent over \mathbb{C} .*

Proof. As in the previous sections the quotient $J/\Phi_{7,A_1(4)}$ is holomorphic of weight zero and therefore constant. We checked by computer that this constant is not zero. \square

Theorem 2.4 yields the following structure theorem:

Theorem 8.2.

$$\begin{aligned} M_*(\mathcal{O}_1(U(4) \oplus U(2) \oplus A_1)) &= \mathbb{C}[h_1, h_2, h_3, h_4], \\ (\mathcal{D}_3/\mathcal{O}_1(U(4) \oplus U(2) \oplus A_1))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Corollary 8.3. *The 16 products of weight one in (2) span a 4-dimensional space. The 8 squares of weight $1/2$ products also span this space.*

We label the 8 weight $1/2$ products d_i , $1 \leq i \leq 8$. Let v_i be the multiplier system of d_i on $\mathcal{O}_1(U(4) \oplus U(2) \oplus A_1)$. Assume that d_i^2 for $1 \leq i \leq 4$ are linearly independent over \mathbb{C} . Let $\mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1)$ be the subgroup of $\mathcal{O}_1(U(4) \oplus U(2) \oplus A_1)$ generated by all reflections associated to the divisor of $\Phi_{7,A_1(4)}/(\prod_{j=1}^4 d_j)$. The four multiplier systems v_i for $1 \leq i \leq 4$ coincide on $\mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1)$ because the quotient of any two of them is a character of $\mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1)$ which equals one on the generator reflections of that group. Their common

restriction defines a multiplier system of $\mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1)$ and we denote it by v . This implies that $\mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1)$ does not contain reflections associated to the divisor of $\prod_{j=1}^4 d_j$, and that $v = 1$ for all reflections in $\mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1)$. We obtain the following result:

Theorem 8.4.

$$\begin{aligned} M_*(\mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1), v) &= \mathbb{C}[d_1, d_2, d_3, d_4], \\ (\mathcal{D}_3 / \mathcal{O}_{1'}(U(4) \oplus U(2) \oplus A_1))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Remark 8.5. There are four linearly independent additive theta lifts of weight 1 for the Weil representation attached to $U(4) \oplus U(2) \oplus A_1$. This implies that every weight 1 modular form of trivial character on $\mathcal{O}_1(U(4) \oplus U(2) \oplus A_1)$ is an additive lift and therefore is modular under the full discriminant kernel. Similarly to Remark 4.8, we conclude that $M_*(\tilde{\mathcal{O}}^+(U(4) \oplus U(2) \oplus A_1))$ is freely generated by the four additive lifts and that

$$\tilde{\mathcal{O}}^+(U(4) \oplus U(2) \oplus A_1) = \mathcal{O}_1(U(4) \oplus U(2) \oplus A_1).$$

9. THE $U \oplus U(2) \oplus A_1(2)$ LATTICE

In this section we find two interesting free algebras of modular forms for the $U \oplus U(2) \oplus A_1(2)$ lattice. We first describe some Borcherds products:

- (1) There are 3 holomorphic Borcherds products t_1, t_2, t_3 of weight 1. They have principal parts of the form

$$q^{-1/8}(e_v + e_{-v}), \quad (v, v) = 1/4, \text{ ord}(v) = 4.$$

- (2) There is a holomorphic Borcherds product $\Psi_{8, A_1(2)}$ of weight 8 whose principal part with respect to the Gram matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 4 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

is

$$\begin{aligned} &-q^{-1/8}(e_{(0,0,1/4,0,0)} + e_{(0,0,3/4,0,0)} + e_{(0,0,1/4,1/2,0)} \\ &\quad + e_{(0,0,3/4,1/2,0)} + e_{(0,1/2,1/4,0,0)} + e_{(0,1/2,3/4,0,0)}) \\ &+ q^{-1/2}e_{(0,0,1/2,0,0)} + q^{-1}e_{(0,0,0,0,0)}. \end{aligned}$$

In addition, we let m_2 be the additive theta lift of the (unique) modular form of weight 3/2 for the Weil representation attached to $U \oplus U(2) \oplus A_1(2)$ with constant term $1\epsilon_0$; in particular, m_2 has weight two.

Let $\mathcal{O}_1(U \oplus U(2) \oplus A_1(2))$ and $\mathcal{O}_2(U \oplus U(2) \oplus A_1(2))$ be the subgroups of $\mathcal{O}^+(U \oplus U(2) \oplus A_1(2))$ generated by the reflections associated to the divisors of $\Psi_{8, A_1(2)}$ and $\Psi_{8, A_1(2)}t_1t_2t_3$ respectively. By considering the actions of these reflections on the input forms, we see that t_1, t_2, t_3 are modular forms with a character of order two on $\mathcal{O}_2(U \oplus U(2) \oplus A_1(2))$, and m_2 is modular on $\mathcal{O}_2(U \oplus U(2) \oplus A_1(2))$ without character. Moreover, t_1, t_2, t_3 have trivial character on $\mathcal{O}_1(U \oplus U(2) \oplus A_1(2))$.

Lemma 9.1. *The Jacobian $J = J(t_1, t_2, t_3, m_2)$ equals $\Psi_{8, A_1(2)}$ up to a nonzero multiple.*

Proof. Using Fourier series we computed that J is not identically zero (and equals $(-1/3)\Psi_{8, A_1(2)}$ up to precision $O(q, s)^{10}$). As in the previous sections $J/\Psi_{8, A_1(2)}$ is holomorphic of weight zero and therefore constant. \square

Therefore, we can apply Theorem 2.4:

Theorem 9.2.

$$\begin{aligned} M_*(\mathrm{O}_1(U \oplus U(2) \oplus A_1(2))) &= \mathbb{C}[t_1, t_2, t_3, m_2], \\ M_*(\mathrm{O}_2(U \oplus U(2) \oplus A_1(2))) &= \mathbb{C}[t_1^2, t_2^2, t_3^2, m_2], \\ (\mathcal{D}_3/\mathrm{O}_2(U \oplus U(2) \oplus A_1(2)))^* &\cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Furthermore, let $\mathrm{O}_{2'}(U \oplus U(2) \oplus A_1(2))$ be the subgroup of $\mathrm{O}^+(U \oplus U(2) \oplus A_1(2))$ generated by the reflections associated to the divisor of $\Psi_{8,A_1(2)}t_2t_3$. We then have another realization of $\mathbb{P}^3(\mathbb{C})$.

Theorem 9.3.

$$\begin{aligned} M_*(\mathrm{O}_{2'}(U \oplus U(2) \oplus A_1(2))) &= \mathbb{C}[t_1, t_2^2, t_3^2, m_2], \\ (\mathcal{D}_3/\mathrm{O}_{2'}(U \oplus U(2) \oplus A_1(2)))^* &\cong \mathbb{P}(1, 2, 2, 2) \cong \mathbb{P}^3(\mathbb{C}). \end{aligned}$$

Remark 9.4. The following eight groups constructed in the paper are all subgroups of $\mathrm{O}^+(2U \oplus A_1)$:

$$\begin{aligned} \mathrm{O}_1(2U(4) \oplus A_1), \quad \mathrm{O}_2(2U(4) \oplus A_1), \quad \mathrm{O}_1(2U(2) \oplus A_1), \quad \mathrm{O}_{1'}(2U(2) \oplus A_1), \\ \mathrm{O}_{1''}(2U(2) \oplus A_1), \quad \mathrm{O}_1(2U(3) \oplus A_1), \quad \mathrm{O}_1(U(4) \oplus U(2) \oplus A_1), \quad \mathrm{O}_{1'}(U(4) \oplus U(2) \oplus A_1). \end{aligned}$$

For any of them, the Satake–Baily–Borel compactification of modular variety is isomorphic to \mathbb{P}^3 . The decomposition into irreducibles of the square of the Jacobian of any generators corresponds to the Γ -equivalence classes of mirrors of reflections in Γ (see Theorem 2.3). In particular, this decomposition is an invariant of the groups up to conjugacy. We give the decomposition for the 16 reflection groups in Table 1, which shows that the eight subgroups above are pairwise non-conjugate except for the possible cases

$$\mathrm{O}_1(U(4) \oplus U(2) \oplus A_1) \quad \text{and} \quad \mathrm{O}_{1'}(2U(2) \oplus A_1)$$

and

$$\mathrm{O}_{1'}(U(4) \oplus U(2) \oplus A_1) \quad \text{and} \quad \mathrm{O}_1(2U(4) \oplus A_1).$$

Remark 9.5. The groups of type $\mathrm{O}_1(M)$ in our paper are finite index subgroups of the full integral orthogonal group $\mathrm{O}^+(M)$. This follows easily from the Margulis normal subgroup theorem [12]. It can also be proved using the basic argument in the proof of [20, Theorem 3.1].

Remark 9.6. Many examples in this paper support Conjecture 5.2 in [19] which states that if $M_*(\Gamma)$ is a free algebra for a finite index subgroup Γ of $\mathrm{O}^+(M)$ then $M_*(\Gamma_1)$ is also free for any other reflection subgroup Γ_1 satisfying $\Gamma < \Gamma_1 < \mathrm{O}^+(M)$.

Acknowledgements The authors would like to thank Eberhard Freitag and Riccardo Salvati Manni for proposing this topic and for helpful discussions. H. Wang is grateful to Max Planck Institute for Mathematics in Bonn for its hospitality and financial support. B. Williams is supported by a fellowship of the LOEWE research group Uniformized Structures in Algebra and Geometry.

TABLE 1. 16 reflection groups Γ for which $(\mathcal{D}/\Gamma)^*$ is a projective space. The symbol $k(J)$ stands for the weight of the Jacobian J . For a group Γ , the entry $\mathbf{a}_1 \times x_1 + \cdots + \mathbf{a}_t \times x_t$ in the rightmost column means that the decomposition of J^2 into irreducibles in $M_*(\Gamma)$ consists of x_i forms of weight \mathbf{a}_i , $1 \leq i \leq t$.

group	$k(J)$	weights of generators	decomposition of J^2
$O_1(2U(4) \oplus A_1)$	5	$\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$	$\mathbf{1} \times 10$
$O_2(2U(4) \oplus A_1)$	7	1, 1, 1, 1	$\mathbf{1} \times 6 + \mathbf{2} \times 2 + \mathbf{4}$
$O_1(2U(2) \oplus A_1)$	11	2, 2, 2, 2	$\mathbf{2} \times 9 + \mathbf{4}$
$O_{1'}(2U(2) \oplus A_1)$	7	1, 1, 1, 1	$\mathbf{1} \times 8 + \mathbf{2} \times 3$
$O_{1''}(2U(2) \oplus A_1)$	10	1, 2, 2, 2	$\mathbf{2} \times 8 + \mathbf{4}$
$O_1(2U(3) \oplus A_1)$	7	1, 1, 1, 1	$\mathbf{2} + \mathbf{12}$
$O_1(U(4) \oplus U(2) \oplus A_1)$	7	1, 1, 1, 1	$\mathbf{1} \times 8 + \mathbf{2} \times 3$
$O_{1'}(U(4) \oplus U(2) \oplus A_1)$	5	$\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$	$\mathbf{1} \times 10$
$O_2(U \oplus U(2) \oplus A_1(2))$	11	2, 2, 2, 2	$\mathbf{2} \times 7 + \mathbf{8}$
$O_{2'}(U \oplus U(2) \oplus A_1(2))$	10	1, 2, 2, 2	$\mathbf{2} \times 6 + \mathbf{8}$
$O_1(2U(2) \oplus 2A_1)$	14	2, 2, 2, 2, 2	$\mathbf{2} \times 10 + \mathbf{8}$
$O_{1'}(2U(2) \oplus 2A_1)$	9	1, 1, 1, 1, 1	$\mathbf{2} \times 5 + \mathbf{8}$
$O_{1''}(2U(2) \oplus 2A_1)$	13	1, 2, 2, 2, 2	$\mathbf{2} \times 9 + \mathbf{8}$
$O_1(2U(3) \oplus A_2)$	9	1, 1, 1, 1, 1	$\mathbf{18}$
$O_{1,1234}(2U(3) \oplus A_2)$	13	1, 2, 2, 2, 2	$\mathbf{2} \times 4 + \mathbf{18}$
$O_2(2U(3) \oplus A_2)$	14	2, 2, 2, 2, 2	$\mathbf{2} \times 5 + \mathbf{18}$

APPENDIX A: TYPE II PRODUCTS AND *-SETS OF TYPE $2U(4) \oplus A_1$

The sixty type II singular-weight products attached to the lattice $M = 2U(4) \oplus A_1$ have input forms with principal part $q^{-1/4}(e_v + e_{-v})$ for certain cosets $v \in M^\vee/M$ of order 4. In the table below, we list one coset representative v for each product Θ_i , with respect to the Gram matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 4 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 \\ 4 & 0 & 0 & 0 & 0 \end{pmatrix} :$$

Θ_1	$(1/4, 1/2, 0, 1/2, 1/4)$	Θ_2	$(1/4, 0, 0, 0, 1/4)$
Θ_3	$(0, 0, 1/2, 1/4, 1/4)$	Θ_4	$(1/2, 1/2, 1/2, 1/4, 1/4)$
Θ_5	$(1/4, 0, 0, 1/4, 1/4)$	Θ_6	$(1/4, 1/2, 0, 1/4, 3/4)$
Θ_7	$(1/2, 0, 1/2, 1/4, 0)$	Θ_8	$(0, 0, 1/2, 1/4, 1/2)$
Θ_9	$(0, 1/4, 1/2, 0, 1/4)$	Θ_{10}	$(1/2, 1/4, 1/2, 1/2, 3/4)$
Θ_{11}	$(1/4, 1/2, 1/2, 0, 0)$	Θ_{12}	$(1/4, 0, 1/2, 0, 0)$
Θ_{13}	$(1/4, 1/2, 1/2, 1/4, 1/2)$	Θ_{14}	$(1/4, 1/2, 1/2, 3/4, 1/2)$
Θ_{15}	$(1/4, 3/4, 1/2, 0, 0)$	Θ_{16}	$(1/4, 1/4, 1/2, 0, 0)$
Θ_{17}	$(1/4, 3/4, 0, 1/2, 3/4)$	Θ_{18}	$(1/4, 3/4, 0, 0, 1/4)$
Θ_{19}	$(0, 1/4, 0, 1/4, 1/4)$	Θ_{20}	$(1/2, 1/4, 0, 3/4, 1/4)$
Θ_{21}	$(1/4, 3/4, 0, 1/4, 1/2)$	Θ_{22}	$(1/4, 3/4, 0, 3/4, 0)$
Θ_{23}	$(0, 1/2, 1/2, 1/2, 1/4)$	Θ_{24}	$(0, 0, 1/2, 1/2, 1/4)$
Θ_{25}	$(1/2, 1/4, 0, 1/4, 0)$	Θ_{26}	$(1/2, 1/4, 0, 1/4, 1/2)$
Θ_{27}	$(1/2, 1/4, 1/2, 0, 0)$	Θ_{28}	$(1/2, 1/4, 1/2, 0, 1/2)$
Θ_{29}	$(1/4, 1/4, 1/2, 1/4, 3/4)$	Θ_{30}	$(1/4, 1/4, 1/2, 3/4, 1/4)$
Θ_{31}	$(1/4, 0, 0, 1/2, 1/4)$	Θ_{32}	$(1/4, 1/2, 0, 0, 1/4)$
Θ_{33}	$(0, 0, 1/2, 1/4, 3/4)$	Θ_{34}	$(1/2, 1/2, 1/2, 1/4, 3/4)$
Θ_{35}	$(1/4, 1/2, 0, 3/4, 3/4)$	Θ_{36}	$(1/4, 0, 0, 3/4, 1/4)$
Θ_{37}	$(1/2, 0, 1/2, 1/4, 1/2)$	Θ_{38}	$(0, 0, 1/2, 1/4, 0)$
Θ_{39}	$(1/2, 1/4, 1/2, 1/2, 1/4)$	Θ_{40}	$(0, 1/4, 1/2, 0, 3/4)$
Θ_{41}	$(1/4, 0, 1/2, 1/2, 0)$	Θ_{42}	$(1/4, 1/2, 1/2, 1/2, 0)$
Θ_{43}	$(1/4, 0, 1/2, 1/4, 0)$	Θ_{44}	$(1/4, 0, 1/2, 3/4, 0)$
Θ_{45}	$(1/4, 1/4, 1/2, 1/2, 1/2)$	Θ_{46}	$(1/4, 3/4, 1/2, 1/2, 1/2)$
Θ_{47}	$(1/4, 1/4, 0, 1/2, 3/4)$	Θ_{48}	$(1/4, 1/4, 0, 0, 1/4)$
Θ_{49}	$(1/2, 1/4, 0, 3/4, 3/4)$	Θ_{50}	$(0, 1/4, 0, 1/4, 3/4)$
Θ_{51}	$(1/4, 1/4, 0, 3/4, 1/2)$	Θ_{52}	$(1/4, 1/4, 0, 1/4, 0)$
Θ_{53}	$(0, 1/2, 1/2, 0, 1/4)$	Θ_{54}	$(0, 0, 1/2, 0, 1/4)$
Θ_{55}	$(0, 1/4, 0, 1/4, 0)$	Θ_{56}	$(0, 1/4, 0, 1/4, 1/2)$
Θ_{57}	$(0, 1/4, 1/2, 0, 1/2)$	Θ_{58}	$(0, 1/4, 1/2, 0, 0)$
Θ_{59}	$(1/4, 3/4, 1/2, 1/4, 1/4)$	Θ_{60}	$(1/4, 3/4, 1/2, 3/4, 3/4)$

The *-sets are the maximum cliques in the graph formed by connecting products Θ_i and Θ_j by an edge if the reflections associated to the divisor of Θ_j also preserve the divisor of Θ_i . With respect to the ordering above, the 105 *-sets are $\{\Theta_{i_1}, \Theta_{i_2}, \Theta_{i_3}, \Theta_{i_4}\}$ where (i_1, i_2, i_3, i_4) is one of the following:

- $(1, 2, 26, 55), (3, 17, 18, 34), (6, 19, 36, 49), (10, 14, 40, 44), (13, 19, 20, 44), (17, 27, 47, 57), (23, 24, 53, 54),$
 $(1, 2, 29, 60), (3, 21, 22, 34), (7, 8, 37, 38), (10, 30, 40, 60), (13, 30, 43, 60), (17, 48, 49, 50), (23, 24, 57, 58),$
 $(1, 2, 31, 32), (3, 23, 33, 53), (7, 11, 38, 42), (10, 39, 53, 54), (14, 18, 43, 47), (18, 19, 20, 47), (23, 54, 55, 56),$
 $(1, 5, 32, 36), (4, 8, 34, 38), (7, 13, 14, 38), (10, 39, 57, 58), (14, 29, 44, 59), (18, 28, 48, 58), (24, 25, 26, 53),$




FIGURE 1. There is an edge between the type II products Θ_i and Θ_j if Θ_i is modular under the reflections associated to the divisor of Θ_j . This is a 15-regular graph with 23,040 automorphisms. The edge between Θ_i and Θ_j is colored red, green or blue depending on $i + j \pmod 3$.

- $(1, 7, 8, 32), (4, 24, 34, 54), (7, 24, 37, 54), (11, 12, 28, 57), (14, 43, 49, 50), (19, 20, 49, 50), (25, 26, 55, 56),$
 $(1, 18, 31, 48), (4, 33, 47, 48), (8, 12, 37, 41), (11, 12, 41, 42), (15, 16, 28, 57), (19, 23, 50, 54), (25, 29, 56, 60),$
 $(1, 27, 31, 57), (4, 33, 51, 52), (8, 23, 38, 53), (11, 12, 45, 46), (15, 16, 41, 42), (19, 25, 26, 50), (25, 31, 32, 56),$
 $(2, 6, 31, 35), (5, 6, 22, 51), (8, 37, 43, 44), (11, 21, 41, 51), (15, 16, 45, 46), (20, 24, 49, 53), (26, 30, 55, 59),$
 $(2, 17, 32, 47), (5, 6, 35, 36), (9, 10, 22, 51), (11, 26, 41, 56), (15, 19, 45, 49), (20, 49, 55, 56), (27, 28, 53, 54),$
 $(2, 28, 32, 58), (5, 6, 39, 40), (9, 10, 35, 36), (11, 42, 43, 44), (15, 29, 30, 46), (21, 22, 47, 48), (27, 28, 57, 58),$
 $(2, 31, 37, 38), (5, 15, 35, 45), (9, 10, 39, 40), (12, 13, 14, 41), (15, 33, 34, 46), (21, 22, 51, 52), (27, 41, 42, 58),$
 $(3, 4, 16, 45), (5, 20, 35, 50), (9, 13, 39, 43), (12, 22, 42, 52), (16, 20, 46, 50), (21, 25, 51, 55), (27, 45, 46, 58),$
 $(3, 4, 29, 30), (5, 36, 37, 38), (9, 23, 24, 40), (12, 25, 42, 55), (16, 45, 59, 60), (21, 35, 36, 52), (29, 30, 59, 60),$
 $(3, 4, 33, 34), (6, 7, 8, 35), (9, 27, 28, 40), (13, 14, 43, 44), (17, 18, 47, 48), (21, 39, 40, 52), (30, 31, 32, 59),$
 $(3, 7, 33, 37), (6, 16, 36, 46), (9, 29, 39, 59), (13, 17, 44, 48), (17, 18, 51, 52), (22, 26, 52, 56), (33, 34, 59, 60).$

The ordering is chosen such that the 90 exceptional pairs of type II forms that extend in three ways to $*$ -sets are precisely those of the form $(\Theta_{i+j}, \Theta_{i+k})$ where $i < 30$ is odd and $j, k \in \{0, 1, 30, 31\}$ are distinct.




FIGURE 2. Contracting the 90 exceptional pairs in Figure 1 yields the strongly regular graph $\text{srg}(15, 6, 1, 3)$ on 15 vertices. The edges are colored red, green, blue by the same rule as Figure 1.

APPENDIX B: SINGULAR-WEIGHT PRODUCTS AND *-SETS OF TYPE $2U(3) \oplus A_2$

The 45 singular-weight products attached to the lattice $M = 2U(3) \oplus A_2$ have input forms with principal part $q^{-1/3}(e_v + e_{-v})$ for cosets $v \in M^\vee/M$ of order 3. In the table below, for each product G_i we list one coset representative v with respect to the Gram matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 \\ 0 & 3 & 0 & 0 & 0 & 0 \\ 3 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} :$$

G_1	$(0, 0, 2/3, 2/3, 0, 1/3)$	G_2	$(0, 0, 2/3, 2/3, 1/3, 1/3)$
G_3	$(0, 1/3, 2/3, 2/3, 0, 0)$	G_4	$(1/3, 2/3, 1/3, 1/3, 1/3, 1/3)$
G_5	$(0, 1/3, 1/3, 1/3, 0, 1/3)$	G_6	$(0, 0, 2/3, 2/3, 2/3, 1/3)$
G_7	$(0, 1/3, 1/3, 1/3, 0, 0)$	G_8	$(1/3, 0, 1/3, 1/3, 2/3, 0)$
G_9	$(1/3, 1/3, 1/3, 1/3, 0, 0)$	G_{10}	$(1/3, 1/3, 0, 0, 0, 1/3)$
G_{11}	$(0, 0, 2/3, 2/3, 0, 2/3)$	G_{12}	$(1/3, 2/3, 1/3, 1/3, 2/3, 2/3)$
G_{13}	$(1/3, 2/3, 0, 0, 1/3, 2/3)$	G_{14}	$(1/3, 0, 2/3, 2/3, 1/3, 0)$
G_{15}	$(1/3, 1/3, 2/3, 2/3, 0, 0)$	G_{16}	$(0, 0, 2/3, 2/3, 2/3, 2/3)$
G_{17}	$(1/3, 1/3, 0, 0, 1/3, 0)$	G_{18}	$(1/3, 2/3, 1/3, 1/3, 0, 0)$
G_{19}	$(0, 1/3, 1/3, 1/3, 0, 2/3)$	G_{20}	$(0, 1/3, 2/3, 2/3, 0, 2/3)$
G_{21}	$(0, 0, 2/3, 2/3, 1/3, 0)$	G_{22}	$(0, 1/3, 0, 0, 1/3, 1/3)$
G_{23}	$(0, 1/3, 0, 0, 1/3, 2/3)$	G_{24}	$(0, 0, 2/3, 2/3, 0, 0)$
G_{25}	$(1/3, 2/3, 0, 0, 0, 1/3)$	G_{26}	$(0, 1/3, 2/3, 2/3, 0, 1/3)$
G_{27}	$(1/3, 2/3, 2/3, 2/3, 2/3, 2/3)$	G_{28}	$(1/3, 0, 0, 0, 2/3, 1/3)$
G_{29}	$(1/3, 1/3, 1/3, 1/3, 1/3, 2/3)$	G_{30}	$(1/3, 0, 0, 0, 0, 1/3)$
G_{31}	$(0, 0, 2/3, 2/3, 1/3, 2/3)$	G_{32}	$(1/3, 0, 0, 0, 1/3, 1/3)$
G_{33}	$(1/3, 2/3, 2/3, 2/3, 0, 0)$	G_{34}	$(1/3, 0, 1/3, 1/3, 1/3, 0)$
G_{35}	$(0, 1/3, 0, 0, 1/3, 0)$	G_{36}	$(1/3, 2/3, 0, 0, 2/3, 0)$
G_{37}	$(1/3, 1/3, 0, 0, 2/3, 2/3)$	G_{38}	$(1/3, 0, 1/3, 1/3, 0, 0)$
G_{39}	$(1/3, 1/3, 2/3, 2/3, 1/3, 2/3)$	G_{40}	$(1/3, 0, 2/3, 2/3, 0, 0)$
G_{41}	$(0, 0, 2/3, 2/3, 2/3, 0)$	G_{42}	$(1/3, 2/3, 2/3, 2/3, 1/3, 1/3)$
G_{43}	$(1/3, 1/3, 2/3, 2/3, 2/3, 1/3)$	G_{44}	$(1/3, 0, 2/3, 2/3, 2/3, 0)$
G_{45}	$(1/3, 1/3, 1/3, 1/3, 2/3, 1/3)$		

With respect to this ordering the 27 *-sets are the sets $\{G_i, i \in I\}$ where I is one of the index sets:

- $(1, 14, 15, 23, 27), (1, 22, 33, 39, 44), (1, 35, 40, 42, 43), (2, 3, 14, 25, 45), (2, 9, 13, 20, 44),$
 $(2, 26, 29, 36, 40), (3, 8, 10, 31, 42), (3, 12, 21, 30, 39), (4, 6, 7, 10, 44), (4, 11, 35, 38, 45),$
 $(4, 15, 21, 26, 28), (5, 6, 14, 18, 37), (5, 9, 28, 41, 42), (5, 16, 36, 38, 39), (6, 12, 17, 19, 40),$
 $(7, 16, 25, 34, 43), (7, 27, 29, 30, 41), (8, 11, 18, 22, 29), (8, 13, 15, 16, 19), (9, 11, 12, 23, 34),$
 $(10, 23, 24, 32, 36), (13, 24, 30, 35, 37), (17, 20, 27, 31, 38), (17, 22, 24, 25, 28), (18, 20, 21, 32, 43),$
 $(19, 32, 33, 41, 45), (26, 31, 33, 34, 37).$

These can be computed as the maximal cliques in the graph depicted in Figure 3.




FIGURE 3. There is an edge between the singular-weight products G_i and G_j if G_i is modular under the reflections associated to the divisor of G_j . This is a strongly regular graph with parameters $(45, 12, 3, 3)$ and has 51,840 automorphisms.

REFERENCES

- [1] Hiroki Aoki and Tomoyoshi Ibukiyama. Simple graded rings of Siegel modular forms, differential operators and Borchers products. *Internat. J. Math.*, 16(3):249–279, 2005.
- [2] Walter Baily and Armand Borel. Compactification of arithmetic quotients of bounded symmetric domains. *Ann. of Math. (2)*, 84:442–528, 1966.
- [3] Richard Borcherds. Automorphic forms with singularities on Grassmannians. *Invent. Math.*, 132(3):491–562, 1998.
- [4] Jan Hendrik Bruinier. *Borchers products on $O(2, 1)$ and Chern classes of Heegner divisors*, volume 1780 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002.
- [5] Tobias Dern and Aloys Krieg. Graded rings of Hermitian modular forms of degree 2. *Manuscripta Math.*, 110(2):251–272, 2003.
- [6] Eberhard Freitag. Modulformen zweiten Grades zum rationalen und Gaußschen Zahlkörper. *S.-B. Heidelberger Akad. Wiss. Math.-Natur. Kl.*, 1967:3–49, 1967.
- [7] Eberhard Freitag and Riccardo Salvati Manni. Hermitian modular forms and the Burkhardt quartic. *Manuscripta Math.*, 119(1):57–59, 2006.
- [8] Valery Gritsenko and Viacheslav Nikulin. Lorentzian Kac-Moody algebras with Weyl groups of 2-reflections. *Proc. Lond. Math. Soc. (3)*, 116(3):485–533, 2018.
- [9] Valery Gritsenko, Klaus Hulek, and Gregory Sankaran. Abelianisation of orthogonal groups and the fundamental group of modular varieties. *J. Algebra*, 322(2):463–478, 2009.

- [10] Carl Friedrich Hermann. Some modular varieties related to \mathbf{P}^4 . In *Abelian varieties (Egloffstein, 1993)*, pages 105–129. de Gruyter, Berlin, 1995.
- [11] Jun-ichi Igusa. On Siegel modular forms genus two. II. *Amer. J. Math.*, 86:392–412, 1964.
- [12] Grigory Margulis. *Discrete subgroups of semisimple Lie groups*, volume 17 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1991.
- [13] Keiji Matsumoto. Theta functions on the bounded symmetric domain of type $I_{2,2}$ and the period map of a 4-parameter family of $K3$ surfaces. *Math. Ann.*, 295(3):383–409, 1993.
- [14] Sebastian Opitz and Markus Schwagenscheidt. Holomorphic Borcherds products of singular weight for simple lattices of arbitrary level. *Proc. Amer. Math. Soc.*, 147, 2019.
- [15] Sara Perna. On isomorphisms between Siegel modular threefolds. *Abh. Math. Semin. Univ. Hambg.*, 86(1):55–68, 2016.
- [16] Vladimir Popov and Ernest Vinberg. Invariant theory. In *Algebraic geometry, 4 (Russian)*, Itogi Nauki i Tekhniki, pages 137–314, 315. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989.
- [17] Bernhard Runge. On Siegel modular forms. I. *J. Reine Angew. Math.*, 436:57–85, 1993.
- [18] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.1)*, 2020. <https://www.sagemath.org>.
- [19] Haowu Wang. The classification of free algebras of orthogonal modular forms. 2020. URL [arXiv:2006.02291](https://arxiv.org/abs/2006.02291).
- [20] Haowu Wang. On some free algebras of orthogonal modular forms II. 2020. URL [arXiv:2006.02680](https://arxiv.org/abs/2006.02680).
- [21] Haowu Wang and Brandon Williams. On some free algebras of orthogonal modular forms. *Adv. Math.*, 373, 2020. URL [arXiv:2003.05374](https://arxiv.org/abs/2003.05374). To appear.
- [22] Brandon Williams. Poincaré square series for the Weil representation. *Ramanujan J.*, 47(3):605–650, 2018.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY
Email address: haowu.wangmath@gmail.com

FACHBEREICH MATHEMATIK, TECHNISCHE UNIVERSITÄT DARMSTADT, 64289 DARMSTADT, GERMANY
Email address: bwilliams@mathematik.tu-darmstadt.de