9.2: Direction Fields and Euler's Method Wednesday, April 8

Warmup

- 1. Show that every function of the form $x(t) = t^2 + Bt + C$ is a solution to the differential equation x'' = 2.
- 2. Show that every function of the form $y = Ke^{-ct}$ is a solution to the differential equation y' = -cy.
- 3. Show that every function of the form $y = A \sin x + B \cos x$ solves y'' = -y.

Say that the positon x of a particle over time is described by $\frac{dx}{dt} = x(x-1)(x-2)$.

- 1. For what values of x is the particle moving in the positive direction?
- 2. For what starting values of x will the particle not move at all?
- 3. Suppose the particle starts at x = 0.1. What will happen to it as $t \to \infty$?
- 4. What if the particle starts at x = 1.5?
- 5. What if the particle starts at x = 2.5?


Direction Fields

Draw a vector field for the differential equation in the previous question. Use this to sketch some possible paths for the particle in question.

Match each of the following differential equations with the proper vector fields.

1. $y' = y^2$	2. $y' = x - y$	3. $y' = 2 - y$	4. $y' = x$
		, , , , , , , , , , , , , , , , , , ,	
	$ \begin{array}{c} 0 \\ - \\ - \\ - \\ - \\ - \\ - \\ - \\ - \\ - \\$		04 - 0 - 04 - 09 - 12 - 18 - 2 - 21
******		1111111111	

~																	~										
2			2	2	1	C	2	R	2	2	2	2	3	2	2	1	2	2	R	2	2	2	2	2	2	2	
-	-	. ~	-	-	-	-	~	-	-	~	-	-	-	~	-	-	-	~	-	-	-	-	-	-	-	~	-
	-	• •	-	-	+	-	-	-	-	-	-	+	2		-	+	-	-	-	-	+	-	+	-	+	-	-
1							1			1	1	-	1	1	1	1	1	1	1	1	1	1	1	1	1	1	-
1	1	1	1	1	1	1	1	6	2	6	2				6	1	1	2	6	2	1	2	1	2	1	1	ĺ,
7	1	1	1	1	1	2	1	1	1	1	2	2	1	1	2	7	1	1	1	2	1	1	1	2	1	1	-
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
, 7	7	7	.7	7	43		1	1				1		1	7	4		7			1	37	1	7	1	7	7
							1																1	1	1	1	1
							1							1									4		4		7
								,											6								
					1			1		1									1								
1	t	1	1	1	1	1	1							1			1	1	1	1	1	1	1	t	1	1	1
1	1	1	1	1	1	1	1	1	1								1	1	1	1	1	1	1	1	1	1	1
1	1	1	1	1	1	1	1	1	1	1	1	1	\$	1	1	1	1	1	1	1	1	1	1	1	1	1	1



Particle Simulation/Euler's Method

Consider the solution to the differential equation y' = y with y(0) = 1.

- 1. Estimate y(1) using Euler's Method with a step size of 1.
- 2. Estimate y(1) using Euler's Method with a step size of 1/2.
- 3. Show that $y = e^x$ is a solution to y' = y.
- 4. What will happen to your prediction for y(1) as the step size shrinks to zero?

Air Resistance

Let the velocity of a falling object be described by the equation $\frac{dv}{dt} = -4 - v$, where up is considered to be the positive direction.

- 1. Sketch the direction field. What will happen to the object's velocity as $t \to \infty$?
- 2. If the object starts at x = 20, v = 0, estimate its position and velocity at t = 4 using Euler's Method.

Nullclines

Sketch the direction fields describing the differential equations y' = y(5 - x - y) and $y' = 1 - y - x^2$.