

# Worksheet 1: January 22

## 1 Propositional Logic

1. Determine whether each of the following sentences is a proposition or not. If it is a proposition, determine its truth value.
  - (a) Take BART from Downtown Berkeley to Embarcadero.
  - (b)  $1 + 1 = 2$ .
  - (c) San Francisco is the largest city in California.
  - (d)  $x^2 + y^2 = z^2$ .
  - (e) This sentence is false.

**Solution:**

2. Describe the difference between  $p \vee q$  and  $p \oplus q$ . How do these relate to the English word “or”?

**Solution:**

3. Draw a truth table for the following:

- (a)  $p \wedge \neg p$
- (b)  $p \rightarrow (p \vee q)$
- (c)  $(p \rightarrow q) \rightarrow (p \rightarrow r)$

**Solution:**

4. Consider the following three propositions:

- (a)  $p =$  I go shopping for food
- (b)  $q =$  I order at Cheeseboard
- (c)  $r =$  I make dinner

Rephrase the following English sentences using logical operators:

- (a) If I go shopping for food, then I make dinner.
- (b) If I don't go shopping for food, then I order at Cheeseboard.
- (c) I either make dinner or order at Cheeseboard.

Write each English sentence above using different wording, so that each retains the same meaning when expressed using logical operators.

**Solution:**

5. Show that the converse and the inverse of a conditional statement are equivalent.

**Solution:**

## 2 Propositional Equivalences

6. Use De Morgan's laws to find the negation of each of the following statements:

- (a) Kwame will take a job in industry or go to graduate school.
- (b) Yoshiko knows Java and calculus.
- (c) Celia has been to a Giants game and an A's game.

**Solution:**

7. Use a truth table to verify the distributive law  $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$ .

**Solution:**

8. Determine whether each of these compound propositions is satisfiable.

- (a)  $(p \leftrightarrow q) \wedge (\neg p \leftrightarrow q)$
- (b)  $(p \vee q \vee \neg r) \wedge (p \vee \neg q \vee \neg s) \wedge (p \vee \neg s \vee \neg r) \wedge (\neg p \vee \neg q \vee \neg s) \wedge (p \vee q \vee \neg s)$

**Solution:**

9. How many different truth tables of compound propositions are there that involve the propositional variables  $p$  and  $q$ ?

**Solution:**

10. The  $n$ -Queens problem (see section 1.3.6 in the textbook). The  $n$ -queens problem asks for an arrangement of  $n$  queens on an  $n \times n$  chessboard so that no queen can attack another queen (no two queens can be placed in the same row, column, or diagonal). Introduce variables  $p(i, j)$  for each location  $(i, j)$  in the chessboard, such that  $p(i, j)$  is true if there is a queen at  $(i, j)$  and false otherwise.

In row  $i$ , there exists a queen can be expressed as  $\bigvee_{j=1}^n p(i, j)$ . Then as there is a queen in each row, that can be expressed as

$$\bigwedge_{i=1}^n \bigvee_{j=1}^n p(i, j).$$

That row  $i = 1$  has no more than one queen can be expressed as  $\bigwedge_{j=1}^{n-1} \bigwedge_{k=j+1}^n \neg p(i, j) \vee \neg p(i, k)$ . Then as there is no more than one queen in each row, that can be expressed as

$$\bigwedge_{i=1}^n \bigwedge_{j=1}^{n-1} \bigwedge_{k=j+1}^n \neg p(i, j) \vee \neg p(i, k).$$

Write similar assertions that each column and diagonal must contain at least and no more than one queen. Compute the number of possible solutions to the  $n$ -queens problem when  $n = 2, 3, 4$ .

**Solution:**