

Worksheet 2: January 24

1 Predicates and Quantifiers

1. What are the truth values of these statements?

(a) $\exists x P(x) \rightarrow \forall x P(x)$

(b) $\forall x P(x) \rightarrow \exists x P(x)$

(c) $\exists x \neg P(x) \rightarrow \neg \forall x P(x)$

(d) $\neg \forall x P(x) \rightarrow \neg \exists x P(x)$

(e) $\neg \forall x P(x) \rightarrow \exists x \neg P(x)$

(f) $\neg \exists x P(x) \rightarrow \forall x \neg P(x)$

Solution:

2. Find a counterexample to the following statements:

(a) All prime numbers are even.

(b) The square of a positive integer x is larger than x .

Solution:

2 Nested Quantifiers

3. Determine the truth value of each of these statements if the domain for all variables consists of all integers.

(a) $\forall n \exists m (n^2 < m)$

(b) $\exists n \forall m (n < m^2)$

(c) $\forall n \exists m (n + m = 0)$

(d) $\exists n \forall m (nm = m)$

(e) $\exists n \exists m (n^2 + m^2 = 5)$

(f) $\exists n \exists m (n^2 + m^2 = 6)$

Solution:

3 Rules of Inference

4. Determine whether these are valid arguments.

(a) If x is a positive real number, then x^2 is a positive real number. Therefore, if a^2 is positive, where a is a real number, then a is a positive real number.

(b) If $x^2 \neq 0$, where x is a real number, then $x \neq 0$. Let a be a real number with $a^2 \neq 0$; then $a \neq 0$.

Solution:

5. Justify the rule of universal transitivity, which states that if $\forall x (P(x) \rightarrow Q(x))$ and $\forall x (Q(x) \rightarrow R(x))$ are true, then $\forall x (P(x) \rightarrow R(x))$ is true, where the domains of all quantifiers are the same.

Solution: