

This is the final preprint version of a paper which appeared at
Indagationes Mathematicae, 18 (2007) 349–403. The published version is accessible
in PDF at http://math.berkeley.edu/~gbergman/papers/embed_reprt.pdf
or [http://dx.doi.org/10.1016/S0019-3577\(07\)00018-3](http://dx.doi.org/10.1016/S0019-3577(07)00018-3) .

Some results on embeddings of algebras, after de Bruijn and McKenzie*

George M. Bergman

April 7, 2011

Abstract

In 1957, N. G. de Bruijn showed that the symmetric group $\text{Sym}(\Omega)$ on an infinite set Ω contains a free subgroup on $2^{\text{card}(\Omega)}$ generators, and proved a more general statement, a sample consequence of which is that for any group A of cardinality $\leq \text{card}(\Omega)$, the group $\text{Sym}(\Omega)$ contains a coproduct of $2^{\text{card}(\Omega)}$ copies of A , not only in the variety of all groups, but in any variety of groups to which A belongs. His key lemma is here generalized to an arbitrary variety of algebras \mathbf{V} , and formulated as a statement about functors $\mathbf{Set} \rightarrow \mathbf{V}$. From this one easily obtains analogs of the results stated above with “group” and $\text{Sym}(\Omega)$ replaced by “monoid” and the monoid $\text{Self}(\Omega)$ of endomaps of Ω , by “associative K -algebra” and the K -algebra $\text{End}_K(V)$ of endomorphisms of a K -vector-space V with basis Ω , and by “lattice” and the lattice $\text{Equiv}(\Omega)$ of equivalence relations on Ω . It is also shown, extending another result from de Bruijn’s 1957 paper, that each of $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$ and $\text{End}_K(V)$ contains a coproduct of $2^{\text{card}(\Omega)}$ copies of itself.

That paper also gave an example of a group of cardinality $2^{\text{card}(\Omega)}$ that was *not* embeddable in $\text{Sym}(\Omega)$, and R. McKenzie subsequently established a large class of such examples. Those results are shown here to be instances of a general property of the lattice of solution sets in $\text{Sym}(\Omega)$ of sets of equations with constants in $\text{Sym}(\Omega)$. Again, similar results – this time of varying strengths – are obtained for $\text{Self}(\Omega)$, $\text{End}_K(V)$, and $\text{Equiv}(\Omega)$, and also for the monoid $\text{Rel}(\Omega)$ of binary relations on Ω .

Many open questions and areas for further investigation are noted.

1 Conventions, and outline.

Throughout this note, Ω will be an infinite set. Each ordinal (in particular, each natural number) is understood to be the set of all smaller ordinals; the set of all natural numbers is denoted ω . Functions, including elements of permutation groups, will be written to the left of their arguments and composed accordingly. The word “algebra” will be used in the sense of general algebra (universal algebra), except in the combination “ K -algebra”, which will always mean an associative unital algebra in the sense of ring theory, over a field K assumed fixed throughout this note. In those contexts, V will denote a vector space with basis Ω over that field K .

In §§2-3 we develop results to the effect that algebras arising as values of certain sorts of functors can be embedded in certain infinite direct product algebras, and obtain, as immediate corollaries, results on embeddability of groups, monoids, K -algebras, and lattices in the group $\text{Sym}(\Omega)$, the monoid $\text{Self}(\Omega)$, the K -algebra $\text{End}_K(V)$, and the lattice $\text{Equiv}(\Omega)$ respectively (all defined as in the abstract). The remaining sections obtain results specific to embeddings in one or another of those four structures, and in the monoid $\text{Rel}(\Omega)$. In §4 (and two appendices, §§10-11) it is shown that one can embed into each of the first three of these algebras a coproduct of $2^{\text{card}(\Omega)}$ copies of that same algebra, while §§5-8 obtain restrictions on algebras A embeddable in these five algebras, in terms of order-properties of chains of solution sets of systems of equations in A . §9 suggests some ways in which the results of this note might be extended.

*2000 Mathematics Subject Classifications. Primary: 08B25. Secondary: 06Bxx, 08B20, 16S50, 18A99, 20B07, 20M20, 54Hxx. This preprint is readable online at <http://math.berkeley.edu/~gbergman/papers/>, and arXiv:math/0606407. The former version is likely to be updated more frequently than the latter.

For some further unusual properties of $\text{Sym}(\Omega)$ and of some of the other structures here considered, cf. [4], [5, §6], [6], and works referred to in those papers.

I am indebted to Anatole Khelif for inadvertently bringing these questions to my attention, to Zachary Mesyan for a careful reading of the first draft, to Andreas Blass and Vladimir Tolstykh for helpful information on the literature, and to the referee for several useful suggestions and corrections.

2 Free algebras.

Recall that by Cayley's Theorem, every group of cardinality $\leq \text{card}(\Omega)$ can be embedded in the symmetric group $\text{Sym}(\Omega)$ on Ω ; in particular, $\text{Sym}(\Omega)$ contains free groups of all ranks $\leq \text{card}(\Omega)$. An obvious question is whether it contains larger free groups, for example, a free group of rank $\text{card}(\text{Sym}(\Omega)) = 2^{\text{card}(\Omega)}$. In [7], de Bruijn answered this question affirmatively by a method which also gave embeddings of many interesting nonfree groups in $\text{Sym}(\Omega)$. We will begin by illustrating his key trick, concerning subalgebras of direct products, in the case of free algebras, making the trivial generalization from the variety of groups to arbitrary varieties of algebras. In the next section, his more general statement will be motivated, reformulated in functorial terms, and generalized still further.

In these two sections, \mathbf{V} will be any variety of finitary algebras. ("Finitary" means that every operation has finite arity, but does not exclude varieties with infinitely many operations; for example, modules over an infinite ring.)

Proposition 2.1 (cf. [7]). *Let H be the free algebra on \aleph_0 generators in a variety \mathbf{V} . Then the direct product algebra $H^{\text{card}(\Omega)}$ has a subalgebra free on $2^{\text{card}(\Omega)}$ generators.*

Proof. Let $\mathbf{P}(\Omega)$ denote the power set of Ω , and $\mathbf{P}_{\text{fin}}(\Omega) \subseteq \mathbf{P}(\Omega)$ the set of finite subsets of Ω . Then $\text{card}(\mathbf{P}(\Omega)) = 2^{\text{card}(\Omega)}$ and $\text{card}(\mathbf{P}_{\text{fin}}(\Omega)) = \text{card}(\Omega)$, so it will suffice to find a $\mathbf{P}(\Omega)$ -tuple of elements of $H^{\mathbf{P}_{\text{fin}}(\Omega)}$ that satisfies no relations other than the identities of \mathbf{V} .

For each $s \in \mathbf{P}_{\text{fin}}(\Omega)$, let H_s denote the factor indexed by s in our product $H^{\mathbf{P}_{\text{fin}}(\Omega)}$, and let us pick $2^{\text{card}(s)}$ of the \aleph_0 free generators of H_s , denoting these $x_{s,t}$, with t ranging over the subsets of s . For every $r \in \mathbf{P}(\Omega)$, let X_r be the element of $H^{\mathbf{P}_{\text{fin}}(\Omega)}$ which, for each $s \in \mathbf{P}_{\text{fin}}(\Omega)$, has H_s -component $x_{s,r \cap s}$.

We claim that the $\mathbf{P}(\Omega)$ -tuple of elements $(X_r)_{r \in \mathbf{P}(\Omega)}$ satisfies no relations other than identities of \mathbf{V} . Indeed, since \mathbf{V} is finitary, any relation satisfied by these elements involves only finitely many of them; let $R(X_{r_1}, \dots, X_{r_n})$ be such a relation, where r_1, \dots, r_n are distinct elements of $\mathbf{P}(\Omega)$. Choose a finite subset $s \subseteq \Omega$ such that $r_1 \cap s, \dots, r_n \cap s$ are distinct. Then the s -components of X_{r_1}, \dots, X_{r_n} , namely $x_{s,r_1 \cap s}, \dots, x_{s,r_n \cap s}$, are independent indeterminates in H_s , so projecting the relation $R(X_{r_1}, \dots, X_{r_n})$ that we assumed to hold in $H^{\mathbf{P}_{\text{fin}}(\Omega)}$ onto the component H_s of that product, we see that it is an identity of \mathbf{V} . \square

Corollary 2.2. (i) *The symmetric group $\text{Sym}(\Omega)$ on Ω has subgroups free on $2^{\text{card}(\Omega)}$ generators in every variety \mathbf{V} of groups.*

(ii) *The monoid $\text{Self}(\Omega)$ of endomaps of Ω has submonoids free on $2^{\text{card}(\Omega)}$ generators in every variety \mathbf{V} of monoids.*

(iii) *The endomorphism ring $\text{End}_K(V)$ of the K -vector space V with basis Ω has K -subalgebras free on $2^{\text{card}(\Omega)}$ generators in every variety \mathbf{V} of associative K -algebras.*

(iv) *The lattice $\text{Equiv}(\Omega)$ of equivalence relations on Ω has sublattices free on $2^{\text{card}(\Omega)}$ generators in every variety \mathbf{V} of lattices.*

Proof. Cayley's Theorem shows that every group of cardinality $\leq \text{card}(\Omega)$ is embeddable in $\text{Sym}(\Omega)$, and similar arguments give embeddings of all monoids with $\leq \text{card}(\Omega)$ elements in $\text{Self}(\Omega)$ and of all associative K -algebras of vector-space dimension $\leq \text{card}(\Omega)$ in $\text{End}_K(V)$. The corresponding statement for embeddability of lattices in $\text{Equiv}(\Omega)$ is Whitman's Theorem [21]. (Whitman does not explicitly say there that if the given lattice L is infinite, then the set on which he represents it has cardinality $\leq \text{card}(L)$, but this can be verified from his construction; or one can deduce the possibility of an embedding with this cardinality condition from the embeddability result without it.) In particular, each of these structures contains a copy of the free algebra H on \aleph_0 generators in any subvariety \mathbf{V} of the given variety, since that free algebra is countable, or in the K -algebra case, countable dimensional.

Moreover, each of the four algebras named contains a $\text{card}(\Omega)$ -fold direct product of copies of itself. To see this, let us write the given set Ω as a disjoint union of $\text{card}(\Omega)$ subsets of cardinality $\text{card}(\Omega)$, $\Omega = \bigcup_{i \in \text{card}(\Omega)} \Omega_i$. Then within $\text{Sym}(\Omega)$, the subgroup of consisting those permutations that respect each Ω_i is such a direct product, and the analogous statement holds in the monoid $\text{Self}(\Omega)$. In the K -algebra case we similarly use the algebra of endomorphisms that carry the span of each subset Ω_i of our basis Ω of V into itself, and in the lattice case, the sublattice of equivalence relations that relate members of each Ω_i only with other members of Ω_i .

Since we saw in the first paragraph that each of our objects contains a free algebra H on countably many generators in the (arbitrary) subvariety \mathbf{V} , it follows that it will contain a copy of the product algebra $H^{\text{card}(\Omega)}$, and Proposition 2.1 now gives the desired conclusions. \square

Note that the embedding of $\text{Equiv}(\Omega)^{\text{card}(\Omega)}$ in $\text{Equiv}(\Omega)$ used in the second paragraph of the above proof takes the least element of $\text{Equiv}(\Omega)^{\text{card}(\Omega)}$ to the least element of $\text{Equiv}(\Omega)$, but does not take the greatest element to the greatest element. (It takes that element to the relation whose equivalence classes are the sets Ω_i .) With a little more work, however, one can get the an embedding that respects both greatest and least elements.

Namely, fix any $p \in \Omega$, and let $\Omega' \subseteq \Omega^{\text{card}(\Omega)}$ be the set of elements $(x_i)_{i \in \text{card}(\Omega)}$ such that $x_i = p$ for all but finitely many i . We see that $\text{card}(\Omega') = \text{card}(\Omega)$, so it will suffice to embed $\text{Equiv}(\Omega)^{\text{card}(\Omega)}$ in $\text{Equiv}(\Omega')$. We do this by taking each $\text{card}(\Omega)$ -tuple $(\alpha_i)_{i \in \text{card}(\Omega)}$ ($\alpha_i \in \text{Equiv}(\Omega)$) to the relation $\alpha \in \text{Equiv}(\Omega')$ such that $((x_i), (y_i)) \in \alpha$ if and only if $(x_i, y_i) \in \alpha_i$ for all i . It is straightforward to show that this is a lattice embedding which indeed respects least and greatest elements.

So Corollary 2.2(iv) also holds for lattices with greatest and/or least element, and mappings respecting these elements. For brevity, I will not mention lattices with this additional structure in subsequent sections, except when points come up where I notice that what we can prove about lattices with such structure differs from what we can prove for lattices without it. (Incidentally, both the above embedding and the one in the proof of Corollary 2.2(iv) also respect *infinitary* meets and joins; but this is not relevant to our embedding results, since those require that the algebra operations used be finitary.)

3 Coproducts and functors.

The free algebra on a set Ω in a variety \mathbf{V} is the *coproduct* in \mathbf{V} of an Ω -tuple of copies of the free algebra on one generator. To start the ball of generalization rolling, let us note how to extend the proof of Proposition 2.1 to the case where free algebras are replaced by coproducts of copies of an arbitrary algebra.

In our proof of Proposition 2.1, we chose in each copy H_s of H a $\mathbf{P}(s)$ -tuple $(x_{s,t})_{t \subseteq s}$ of distinct members of our \aleph_0 -tuple of free generators. This time, let H be the coproduct in \mathbf{V} of \aleph_0 copies of a fixed algebra A , and let us take for each $s \in \mathbf{P}_{\text{fin}}(\Omega)$ a $\mathbf{P}(s)$ -tuple $(p_{s,t})_{t \subseteq s}$ of distinct members of the \aleph_0 -tuple of coprojection maps $A \rightarrow H_s$ defining the coproduct structure. We can then define, for each $r \in \mathbf{P}(\Omega)$, a map $P_r : A \rightarrow H^{\mathbf{P}_{\text{fin}}(\Omega)}$ by letting the composite of P_r with each projection $H^{\mathbf{P}_{\text{fin}}(\Omega)} \rightarrow H_s$ be $p_{s,r \cap s}$. With these adjustments, the proof of Proposition 2.1 goes over, and we likewise get the corollary that if A is any group, monoid, associative K -algebra, or lattice, of cardinality, respectively K -dimension, $\leq \text{card}(\Omega)$, and \mathbf{V} any variety of groups, monoids, K -algebras or lattices containing A , then the group $\text{Sym}(\Omega)$, the monoid $\text{Self}(\Omega)$, the K -algebra $\text{End}_K(V)$ or the lattice $\text{Equiv}(\Omega)$ contains a coproduct in \mathbf{V} of $2^{\text{card}(\Omega)}$ copies of A .

To suggest the next level of generalization, let me give a more or less random concrete example. Let A be the group presented by two generators, x and y , and the relations saying that the generator x has exponent 2, and commutes with the element obtained by conjugating it by the square of the generator y :

- (1) $x^2 = 1$,
- (2) $x(y^2 x y^{-2}) = (y^2 x y^{-2})x$.

Now if I is any index-set, let $F(I)$ be the group presented by generators x_i, y_i ($i \in I$) subject to the relations

- (3) $x_i^2 = 1 \quad (i \in I)$,

$$(4) \quad x_i((y_j y_k) x_i (y_j y_k)^{-1}) = ((y_j y_k) x_i (y_j y_k)^{-1}) x_i \quad (i, j, k \in I, \text{ not necessarily distinct}).$$

Looking at the $i = j = k$ case of these relations, we see that for each i , there is a homomorphism $A \rightarrow F(I)$ acting by $x \mapsto x_i$, $y \mapsto y_i$. This is in fact an embedding, for we also see from (1)-(4) that there exists a homomorphism $F(I) \rightarrow A$ mapping all x_i to x and all y_i to y , which gives a left inverse to each of the preceding homomorphisms. (Note, incidentally, that the choices made in (3) and (4), to turn x^2 to x_i^2 , but y^2 to $y_j y_k$, were somewhat arbitrary: other choices would have led to these same conclusions, so the relations (1) and (2) did not uniquely determine (3) and (4).)

Though the group $F(I)$ is generated by an I -tuple of embedded homomorphic images of A , our presentation does not make it a coproduct of those subgroups, since the relations (4) relate elements from *different* copies of A ; nor does it make it their coproduct in some subvariety of groups, since (3) and (4) do not describe identities satisfied by all elements of $F(I)$. We see, however, as for coproducts, that any map of index-sets $I \rightarrow J$ induces a group homomorphism $F(I) \rightarrow F(J)$, making F a functor from sets to groups.

We shall find below that the idea of Proposition 2.1 can be used to show that $F(\aleph_0)^{\text{card}(\Omega)}$ contains a copy of $F(2^{\text{card}(\Omega)})$, and that the corresponding statement holds with the variety of groups replaced by any variety \mathbf{V} of finitary algebras, and (3) and (4) by any such system of “relations parametrized by families of indices”.

We could give a careful formulation of this concept of a “parametrized system of generators and relations”. Fortunately, we do not have to, for we shall see that the concept is equivalent to one that can be defined in a simpler way. We noted above that the construction F was a functor $\mathbf{Set} \rightarrow \mathbf{V}$; and it clearly satisfies

$$(5) \quad \text{For every set } I, \text{ the algebra } F(I) \text{ is generated by the union of the images of } A = F(1) \text{ under the homomorphisms induced by all maps } 1 \rightarrow I.$$

I claim, conversely, that any functor $\mathbf{Set} \rightarrow \mathbf{V}$ satisfying (5) corresponds to a system of algebras determined by “generators and relations with parameters” in the sense suggested by the above discussion. Indeed, given F , let us take for generator-symbols (corresponding to the x and y in our group-theoretic example) any generating set X for $F(1)$. For every set I , every $i \in I$, and every $x \in X$, let us write x_i for the image of x under the map $F(1) \rightarrow F(I)$ induced by the map $1 \rightarrow I$ taking 0 to i . Then (5) shows that $F(I)$ is generated by

$$(6) \quad \{x_i \mid x \in X, i \in I\}.$$

To get relations, let us, for each natural number n , choose a set of relations presenting $F(n)$ in terms of the generators x_i ($x \in X, i \in n$), and let us turn each of these into a “system of relations with parameters” by replacing the subscripts $0, \dots, n-1 \in n$ on the generators appearing in each relation with symbols i_0, \dots, i_{n-1} ranging over a general index-set I .

We see from the functoriality of F that for any I , the generators (6) of $F(I)$ satisfy all instances of the system of relations so obtained. To see that no more relations are needed, note that any relation satisfied in $F(I)$ by the elements (6) can involve only finitely many of these elements, say those coming from the image of $F(n)$ under some one-to-one map $n \rightarrow I$, for some $n \in \omega$. If $I \neq \emptyset$, we can take $n > 0$, so that we may choose a left inverse $I \rightarrow n$ to this map, and applying F to it, we see that the corresponding relation indeed holds in $F(n)$, and so is a consequence of the system of relations we have chosen. If $I = \emptyset = 0$, then n will also equal 0, and such a map likewise exists, yielding the same conclusion. Thus, the indicated system of generators and relations indeed determines $F(I)$ for all I . (Equations satisfied by the empty set of generators correspond to relations on the set of zeroary operations of \mathbf{V} , which hold in all $F(I)$ including $F(0)$. However, for the arguments below, we only need the values of $F(I)$ for nonempty index-sets I , so nothing is lost if the reader prefers to consider F a functor from the category of nonempty sets to \mathbf{V} , and so avoid dealing with the case $I = \emptyset$.)

De Bruijn [7] proves his embeddability results for what he calls “symmetrically generated groups”. On examination, these turn out to be precisely the values $F(I)$ of group-valued functors F satisfying (5). However, rather than stopping here, we may ask whether, in addition to allowing relations like (4) that depend on more than one parameter, we could allow this in our *generators* as well. For example, suppose we associate to each set I the group $F(I)$ with generators

$$(7) \quad x_{ij} \quad (i, j \in I),$$

subject to relations

$$(8) \quad x_{ij}x_{jk} = x_{jk}x_{ij} \quad (i, j, k \in I).$$

It is again clear that maps among index-sets induce homomorphisms among these groups, giving a functor $F : \mathbf{Set} \rightarrow \mathbf{Group}$, and that the corresponding statement is true for systems of algebras of any variety \mathbf{V} presented by generators and relations similarly parametrized by multiple subscripts. The resulting functors will not in general satisfy (5), but assuming the string of subscripts on each generator is finite, they will satisfy

$$(9) \quad \text{For every set } I, \text{ the algebra } F(I) \text{ is the union of the images of the homomorphisms } F(a) : F(n) \rightarrow F(I), \text{ where } n \text{ ranges over } \omega, \text{ and } a \text{ over all set-maps } n \rightarrow I.$$

Conversely, it is straightforward to show, as before, that the values of any functor satisfying (9) arise from this sort of presentation-with-parameters.

We can now give our generalization of Proposition 2.1. As indicated in the second paragraph of §2, \mathbf{V} denotes an arbitrary fixed variety of finitary algebras.

Theorem 3.1 (cf. [7, Theorem 3.1]). *Let F be a functor $\mathbf{Set} \rightarrow \mathbf{V}$ satisfying (9). Then $F(\aleph_0)^{\text{card}(\Omega)}$ has a subalgebra isomorphic to $F(2^{\text{card}(\Omega)})$.*

Proof. As in the proof of Proposition 2.1, it suffices to construct an embedding $h : F(\mathbf{P}(\Omega)) \rightarrow F(\aleph_0)^{\mathbf{P}_{\text{fin}}(\Omega)}$; and again, we may specify such an h by giving its composites with the projections of $F(\aleph_0)^{\mathbf{P}_{\text{fin}}(\Omega)}$ onto the factors $F(\aleph_0)$ corresponding to each $s \in \mathbf{P}_{\text{fin}}(\Omega)$. For each such s , let $c_s : \mathbf{P}(\Omega) \rightarrow \mathbf{P}(s)$ be defined by $r \mapsto r \cap s$, and let us choose an embedding $e_s : \mathbf{P}(s) \rightarrow \aleph_0$, and take the composite of h with the s th projection to be $F(e_s c_s) : F(\mathbf{P}(\Omega)) \rightarrow F(\aleph_0)$.

To show that h is an embedding, consider any two elements $u \neq v \in F(\mathbf{P}(\Omega))$. We claim there exists a component of $F(\aleph_0)^{\mathbf{P}_{\text{fin}}(\Omega)}$ at which $h(u)$ and $h(v)$ have distinct coordinates.

Indeed, from (9) we can see that the images in $F(\mathbf{P}(\Omega))$ of homomorphisms $F(a) : F(n) \rightarrow F(\mathbf{P}(\Omega))$ induced by maps $a : n \rightarrow \mathbf{P}(\Omega)$ ($n \in \omega$) form a directed system of subalgebras with union $F(\mathbf{P}(\Omega))$. Hence u and v will together lie in such an image; so let $u = F(a)(u_0)$, $v = F(a)(v_0)$ for some $a : n \rightarrow \mathbf{P}(\Omega)$ and $u_0, v_0 \in F(n)$, necessarily distinct; here we may assume $n > 0$.

Now choose $s \in \mathbf{P}_{\text{fin}}(\Omega)$ such that $a(0) \cap s, \dots, a(n-1) \cap s$ are distinct. Then the composite map $e_s c_s a : n \rightarrow \mathbf{P}(\Omega) \rightarrow \mathbf{P}(s) \rightarrow \aleph_0$ is one-to-one, hence it has a left inverse. Hence so does $F(e_s c_s a) : F(n) \rightarrow F(\aleph_0)$; hence that is also one-to-one. In particular, the images of u_0 and v_0 under the latter map, which are the s -coordinates of $h(u)$ and $h(v)$, are distinct, as required. \square

As before, we immediately get the particular embeddability results:

Theorem 3.2 (cf. [7, Theorem 3.1]). *Suppose F is a functor from \mathbf{Set} to (i) the category of groups, respectively (ii) the category of monoids, (iii) the category of associative algebras over a field K , or (iv) the category of lattices; and suppose that F satisfies (9), and has the property that the cardinality of $F(\aleph_0)$ in case (i), (ii) or (iv), or its K -dimension in case (iii), is $\leq \text{card}(\Omega)$. (For instance, starting with an algebra A of cardinality or K -dimension $\leq \text{card}(\Omega)$, one might define F to be the functor associating to every set I the I -fold coproduct of copies of A in some fixed variety or quasivariety containing A .)*

Then $F(2^{\text{card}(\Omega)})$ is embeddable in (i) $\text{Sym}(\Omega)$, (ii) $\text{Self}(\Omega)$, (iii) $\text{End}_K(V)$, or (iv) $\text{Equiv}(\Omega)$, respectively. \square

The reader may have noticed when we first proved Proposition 2.1 that we did not really need the factors in our product to be free of rank \aleph_0 ; free objects of finite nonzero ranks would do, as long as there were at least $\text{card}(\Omega)$ such factors of rank greater than or equal to each natural number N ; and, similarly, that in the proof of Theorem 3.1, we could have used a direct product (with enough repetitions) of objects $F(n)$ for n finite, instead of a power of $F(\aleph_0)$. However, it is not hard to verify in each of these cases that the product of such a family would contain an embedded copy of $F(\aleph_0)^{\text{card}(\Omega)}$, reducing these situations to that of Theorem 3.1. Let us record here the observation from which this follows.

Lemma 3.3 (cf. [7]). *Let F be a functor $\mathbf{Set} \rightarrow \mathbf{V}$ satisfying (9). Then $\prod_{0 < n < \omega} F(n)$ has a subalgebra isomorphic to $F(\aleph_0)$.*

Proof. For each $n > 0$, let $f_n : \aleph_0 = \omega \rightarrow n$ be the map taking each natural number r to $\min(r, n-1)$, and define $f : F(\aleph_0) \rightarrow \prod_{0 < n < \omega} F(n)$ to have $F(f_n)$ as its n th coordinate, for each n . An argument of the sort used in the proof of Theorem 3.1 shows that any two distinct elements of $F(\aleph_0)$ have distinct projections in some $F(n)$, so f is an embedding. \square

It would be interesting to look for results similar to those of this section for functors on categories other than **Set**. I leave these investigations to others, but give below one such result I have noticed, and a couple of examples of how it can be applied.

Theorem 3.4. *Let **T.ord** be the category whose objects are totally ordered sets, and whose morphisms are isotone maps (maps satisfying $x \leq y \implies a(x) \leq a(y)$), and let every ordinal, and likewise the set \mathbb{R} of real numbers, be regarded as objects of **T.ord** via their standard orderings. Suppose $F : \mathbf{T.ord} \rightarrow \mathbf{V}$ is a functor satisfying the analog of (9) with “totally ordered set” for “set”, and “isotone maps” for “set-maps”. Then $F(\omega)^{\aleph_0}$ has a subalgebra isomorphic to $F(\mathbb{R})$.*

Proof. The set $\mathbf{P}_{\text{fin}}(\mathbb{Q})$ of finite sets of rational numbers is countable, so it suffices to embed $F(\mathbb{R})$ in $F(\omega)^{\mathbf{P}_{\text{fin}}(\mathbb{Q})}$. Given $s \in \mathbf{P}_{\text{fin}}(\mathbb{Q})$ whose distinct elements are $q_1 < \dots < q_n$, let $a_s : \mathbb{R} \rightarrow \omega$ be the isotone map which sends each $r \in \mathbb{R}$ to the greatest $i \in \{1, \dots, n\}$ such that $q_i \leq r$ if such an i exists, and sends all $r < q_1$ to 0. Let $h : F(\mathbb{R}) \rightarrow F(\omega)^{\mathbf{P}_{\text{fin}}(\mathbb{Q})}$ be the map whose composite with the projection indexed by each $s \in \mathbf{P}_{\text{fin}}(\mathbb{Q})$ is $F(a_s)$.

It is not hard to see that we will be able to complete the proof as we did that of Theorem 3.1 if for every finite set of real numbers $r_0 < \dots < r_n$, we can find an $s \in \mathbf{P}_{\text{fin}}(\mathbb{Q})$ and an isotone map $b : \omega \rightarrow \mathbb{R}$ such that the map $ba_s : \mathbb{R} \rightarrow \omega \rightarrow \mathbb{R}$ fixes r_0, \dots, r_n . To do this, choose $q_1, \dots, q_n \in \mathbb{Q}$ so that $r_{i-1} < q_i \leq r_i$ ($i = 1, \dots, n$), let $s = \{q_1, \dots, q_n\}$, and let b take i to r_i for $i = 0, \dots, n$, and be extended in an arbitrary isotone manner to larger i . Thus, a_s takes r_i ($i = 0, \dots, n$) to i , which b takes back to r_i , as required. \square

For a functor F as in the above theorem, the algebras $F(I)$ will have presentations by systems of generators and relations indexed by finite sequences of subscripts from I , where the indices occurring in each generator or relation may be constrained by inequalities of the form $i \leq j$. A simple example is the functor associating to each totally ordered set I the group presented by generators x_i and y_i ($i \in I$) subject to the relations

$$(10) \quad x_i y_j = y_j x_i \text{ for } i \leq j \text{ in } I.$$

Another is the functor taking each I to the (commutative) monoid presented by generators x_i ($i \in I$) and relations

$$(11) \quad x_i x_j = x_i = x_j x_i \text{ for } i \leq j \text{ in } I.$$

In each of these cases, the object $F(\omega)$ is countable, hence embeddable in $\text{Sym}(\aleph_0)$, respectively $\text{Self}(\aleph_0)$; hence, combining Theorem 3.4 with the method of proof of Theorem 3.2, we see that $F(\mathbb{R})$ is also embeddable in $\text{Sym}(\aleph_0)$, respectively $\text{Self}(\aleph_0)$. Likewise, the group algebra $KF(\mathbb{R})$ for F determined by (10), and the monoid algebra $KF(\mathbb{R})$ for F determined by (11), are the values at \mathbb{R} of K -algebra-valued functors satisfying the analog of (9), and so are embeddable in $\text{End}_K(V)$ for V countable-dimensional. I do not see any way to obtain these results from Theorems 3.1 and 3.2 themselves.

Let us note in connection with the group-theoretic construction (10) (and for some later uses) that if X is any set, and R any symmetric reflexive binary relation on X , and we form the group G presented by the generating set X and the relations

$$(12) \quad x x' = x' x \text{ } ((x, x') \in R),$$

then distinct subsets of X generate distinct subgroups of G (clear by looking at the abelianization of G), and elements $x, x' \in X$ commute in G if and only if $(x, x') \in R$. To see the latter statement, consider any $x \in X$, and let G_0 be the group presented as above, but using the set $X - \{x\}$ and the restriction of R to that set. Then G can be described as an HNN extension [15] of G_0 , obtained by adjoining an additional generator x , whose conjugation action is specified on the subgroup generated by $\{y \in X - \{x\} \mid (x, y) \in R\}$ as the identity map. By the structure of HNN extensions, conjugation by x fixes precisely the elements of that subgroup, giving the asserted characterization of the commuting pairs of elements of X .

4 Each of $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$ and $\text{End}_K(V)$ contains a coproduct of copies of itself.

In the case of Theorem 3.2 where $F(I)$ is the I -fold coproduct of copies of an algebra A , the number of copies of A in the conclusion, $2^{\text{card}(\Omega)}$, is, in general, as large as it can be, embeddability of larger coproducts being precluded by the size (i.e., cardinality or K -dimension) of the object we are trying to embed in. But the assumption that A itself has cardinality or K -dimension $\leq \text{card}(\Omega)$ is not forced in that way; we assumed it so that we could be sure that the coproduct of countably many copies of A would be embeddable in the object in question.

Can we prove results of the same sort for any larger algebras A ?

We shall sketch in the next few paragraphs a proof that the symmetric group $\text{Sym}(\Omega)$ contains a coproduct of two copies of *itself*. Hence, by iteration, it contains coproducts of all finite numbers of copies of itself, hence, by Lemma 3.3, a coproduct of countably many copies of itself, hence, by Theorem 3.1, a coproduct of $2^{\text{card}(\Omega)}$ copies of itself. This result, like those that we generalized in preceding sections, was proved by de Bruijn in [7]. We will then see how to adapt our argument to the case of the monoid $\text{Self}(\Omega)$, and, with more work, the associative algebra $\text{End}_K(V)$.

(In an earlier version of this note, I asked whether the corresponding result held for the lattice $\text{Equiv}(\Omega)$. An affirmative answer has been given by F. Wehrung [20].)

As indicated above, the hard step, for each of these objects, is to show that it contains the coproduct of two copies of itself. Note that to do this for the group $\text{Sym}(\Omega)$ is equivalent to finding two faithful actions of $\text{Sym}(\Omega)$ on Ω (or on some set of the same cardinality) such that there is no nontrivial “interaction” between the permutations giving these actions. In its most naive form, the idea behind the construction we shall describe is to take the natural representation of $\text{Sym}(\Omega)$ on Ω , and the same representation conjugated by a “random” permutation t of Ω , and hope that elements of the two representations will not interact.

As stated, this is much too naive: no matter how we choose t to eliminate interaction among certain permutations in our two representations, it will inevitably lead to interaction among others. However, suppose we replace the set Ω by the disjoint union of $\text{card}(\Omega)$ copies of itself, on each of which we start with the natural representation of $\text{Sym}(\Omega)$, and on each of which we perturb this representation by a different “ t ”. Then we can hope that any given interaction among finitely many elements of our original and perturbed images of $\text{Sym}(\Omega)$ will be avoided in at least *one* of these copies. If this is so, then the representation of $\text{Sym}(\Omega) \amalg \text{Sym}(\Omega)$ on our union of copies of Ω will be faithful.

In particular, we might index our set of copies of Ω by the group $\text{Sym}_{\text{fin}}(\Omega)$ of all permutations of Ω that move only finitely many elements, and on the copy indexed by each t in that group, let that t be our perturbing permutation.

There is still one difficulty: When we construct a t to prevent interaction in some long expression w in elements from our two groups, the behavior of t that we need at one step may be different from the behavior we want at a later step. To get around this, each copy of Ω in the above sketch will be replaced by a disjoint union of countably many copies of itself, $\Omega \times \omega$, and t will range over $\text{Sym}_{\text{fin}}(\Omega \times \omega)$. Given a group relation $w = v$ that we want to cause to fail, we will find that we can select our t and an element $(p_1, 0) \in \Omega \times \{0\}$ so that as we apply w or v to $(p_1, 0)$, that element is moved by successive occurrences of t into $\Omega \times \{1\}$, $\Omega \times \{2\}$, etc., and on each of those copies, we shall be able to independently control what t does.

As mentioned, the above technique can also be adapted to the monoid $\text{Self}(\Omega)$, and to the K -algebra $\text{End}_K(V)$. In the next lemma, the group, monoid, and K -algebra cases are all stated, and the proof is given for the first two. I have relegated the longer proof for $\text{End}_K(V)$ to an appendix, §10, so as not to interrupt the flow of the paper. (Another appendix, §11, gives an alternative construction in the $\text{Sym}(\Omega)$ case, which I found before encountering de Bruijn’s papers, but was not able to adapt to the monoid or K -algebra cases. It may, however, be of independent group-theoretic interest.)

Recall that we are writing functions to the left of their arguments (in contrast to the usage in many papers in the theory of infinite symmetric groups).

- Lemma 4.1** (cf. [7]). (i) $\text{Sym}(\Omega)$ contains a coproduct of two copies of itself as a group.
(ii) $\text{Self}(\Omega)$ contains a coproduct of two copies of itself as a monoid.
(iii) $\text{End}_K(V)$ contains a coproduct of two copies of itself as an associative K -algebra.

Proof of (i) and (ii). We shall verify (ii), then deduce (i) from it.

Recall that the normal form for an element of the coproduct $M \amalg N$ of two monoids is

$$(13) \quad \dots \alpha(g_i) \beta(g_{i-1}) \alpha(g_{i-2}) \beta(g_{i-3}) \dots,$$

where $\alpha : M \rightarrow M \amalg N$, $\beta : N \rightarrow M \amalg N$ are the coprojection maps, the elements g_k with k of one parity (in (13), the parity of i) are elements of $M - \{1\}$, and those with k of the other parity are elements of $N - \{1\}$. In (13), I do not explicitly show the first and last factors, because each may involve either α or β . The identity element is given by the empty product (13).

Thus, elements of $\text{Self}(\Omega) \amalg \text{Self}(\Omega)$ can be written uniquely as products (13) in which all g_i come from $\text{Self}(\Omega) - \{1\}$.

To prove (ii), it suffices to construct a faithful action of $\text{Self}(\Omega) \amalg \text{Self}(\Omega)$ on a set of the same cardinality as Ω . As suggested in the above discussion, that set will be the disjoint union of a family of copies of $\Omega \times \omega$ indexed by the group $\text{Sym}_{\text{fin}}(\Omega \times \omega)$. On every copy of $\Omega \times \omega$, we let elements $\alpha(g)$ ($g \in \text{Self}(\Omega)$) act in the ‘‘natural’’ manner, $g((p, k)) = (g(p), k)$ ($p \in \Omega$, $k \in \omega$), while on the copy of $\Omega \times \omega$ indexed by $t \in \text{Sym}_{\text{fin}}(\Omega \times \omega)$, we let $\beta(g)$ act by $t g t^{-1}$, i.e., the conjugate by t of that same natural action.

To prove that the resulting action of $\text{Self}(\Omega) \amalg \text{Self}(\Omega)$ on our union of copies of $\Omega \times \omega$ is faithful, assume we are given two distinct elements of that monoid, say (13) and

$$(14) \quad \dots \alpha(h_j) \beta(h_{j-1}) \alpha(h_{j-2}) \beta(h_{j-3}) \dots$$

We shall show below how to obtain a $t \in \text{Sym}_{\text{fin}}(\Omega \times \omega)$ such that the induced actions of (13) and (14) on $\Omega \times \omega$, namely

$$(15) \quad \dots g_i (t g_{i-1} t^{-1}) g_{i-2} (t g_{i-3} t^{-1}) \dots \quad \text{and} \\ \dots h_j (t h_{j-1} t^{-1}) h_{j-2} (t h_{j-3} t^{-1}) \dots,$$

act differently on a certain element of $\Omega \times \omega$.

The t we shall construct will be of order 2, so the above two expressions take the forms

$$(16) \quad \dots g_i t g_{i-1} t g_{i-2} t g_{i-3} t \dots \quad \text{and} \\ \dots h_j t h_{j-1} t h_{j-2} t h_{j-3} t \dots$$

We may assume, by interchanging (13) and (14) if necessary, that the former expression involves at least as many factors from $\text{Sym}(\Omega)$ as the latter, and, moreover, that *if* they have the same number of such factors, and have α s and β s in the same places, then for the least value k such that $g_k \neq h_k$, some element of Ω on which g_k and h_k disagree is moved by the former.

If the rightmost term of our original expression (13) is an α term, rather than a β term, let us multiply both lines of (16) on the right by t , and likewise if the left-hand term of (13) is an α term, let us multiply both lines on the left by t . Since t is going to be invertible, the non-equality of the new expressions, which we will prove, is equivalent to the non-equality of the old ones. The first of the new expressions can now be written more precisely; the two products have become

$$(17) \quad t g_n t \dots t g_i t g_{i-1} t g_{i-2} t g_{i-3} t \dots t g_1 t, \quad \text{and} \\ \dots t h_i t h_{i-1} t h_{i-2} t h_{i-3} t \dots$$

Clearly, by the assumptions we have made, the first line of (17) has at least as many occurrences of t as the second.

To construct our promised t , let us now choose, for each $k \in \{1, \dots, n\}$, an element $p_k \in \Omega$ that is moved by g_k ; moreover, we take it to be an element at which g_k and h_k disagree whenever this is possible; i.e., we require this for every value of k such that there exists an element moved by g_k at which g_k and h_k differ. We then define t to fix all elements of $\Omega \times \omega$ except the following $2(n+1)$ elements, which we let it transpose in pairs, as shown:

$$(18) \quad (p_1, 0) \leftrightarrow (p_1, 1), \quad (g_k(p_k), k) \leftrightarrow (p_{k+1}, k+1) \quad (1 \leq k < n), \quad (g_n(p_n), n) \leftrightarrow (g_n(p_n), n+1).$$

Note that (18) is consistent: For $1 \leq k \leq n$, g_k moves p_k , hence the two elements of $\Omega \times \{k\}$ on which (18) prescribes (in different ways) the behavior of t , namely (p_k, k) and $(g_k(p_k), k)$, are distinct.

We now see that when the element shown on the first line of (17) is applied to $(p_1, 0)$, the successive factors of that element (reading from the right), namely t, g_1, t, g_2, t, \dots , move it as follows

$$(19) \quad (p_1, 0) \mapsto (p_1, 1) \mapsto (g_1(p_1), 1) \mapsto (p_2, 2) \mapsto (g_2(p_2), 2) \mapsto (p_3, 3) \mapsto \dots \mapsto (g_{n-1}(p_{n-1}), n-1) \mapsto (p_n, n) \mapsto (g_n(p_n), n) \mapsto (g_n(p_n), n+1).$$

In particular, $(p_1, 0)$ is carried from $\Omega \times \{0\}$ into $\Omega \times \{n+1\}$.

When we instead apply the *second* line of (17) to $(p_1, 0)$ there are several possible cases. If there are fewer factors h_j than g_i , there will be fewer factors t in that second line than in the first line, so there is no way the permutation represented by the second line can move an element from $\Omega \times \{0\}$ into $\Omega \times \{n+1\}$. If there are the same number, n , of h s as of g s, but if the α s and β s don't appear on the same factors, then since the first line of (17) was adjusted to have a t at each end, the second line will not; so again there will be fewer factors t , and $(p_1, 0)$ cannot be moved all the way into $\Omega \times \{n+1\}$.

Finally, if there are the same number of factors and the α s and β s appear in the same positions, then by assumption, for the least k such that $g_k \neq h_k$, the element g_k moves some element of Ω at which these elements disagree (see sentence after (16)), and by our choice of p_k , the latter will be such an element (first sentence of paragraph containing (18)). When we apply the second line of (17) to $(p_1, 0)$, the input to the factor h_k will be (p_k, k) (since the terms have agreed up to this point), so the output will be $(h_k(p_k), k) \neq (g_k(p_k), k)$. Thus, our element will fail to be in the unique position (cf. (18)) from which it can “catch the boat” to be shifted by t from $\Omega \times \{k\}$ to $\Omega \times \{k+1\}$; and since t moves elements by only one level at a time, our element will not be able to catch up later on. So the second line of (17) does not move $(p_1, 0)$ into $\Omega \times \{n+1\}$, hence the two lines represent distinct elements of $\text{Self}(\Omega \times \omega)$, completing the proof of (ii).

To deduce (i) from (ii), note that the normal forms of coproducts of groups and of monoids are formally the same, hence the inclusion of $\text{Sym}(\Omega)$ in $\text{Self}(\Omega)$ induces an embedding of $\text{Sym}(\Omega) \amalg \text{Sym}(\Omega)$ into $\text{Self}(\Omega) \amalg \text{Self}(\Omega)$. Since monoid homomorphisms carry invertible elements to invertible elements, the image of this copy of $\text{Sym}(\Omega) \amalg \text{Sym}(\Omega)$ under the embedding of statement (ii) lies in the group $\text{Sym}(\Omega)$ of invertible elements of $\text{Self}(\Omega)$, so we have indeed embedded $\text{Sym}(\Omega) \amalg \text{Sym}(\Omega)$ in $\text{Sym}(\Omega)$, as required.

As mentioned earlier, the proof of (iii) will be given in an appendix, §10. □

By the reasoning sketched at the beginning of this section, we deduce

Theorem 4.2. (i) $\text{Sym}(\Omega)$ contains a coproduct of $2^{\text{card}(\Omega)}$ copies of itself as a group.

(ii) $\text{Self}(\Omega)$ contains a coproduct of $2^{\text{card}(\Omega)}$ copies of itself as a monoid.

(iii) $\text{End}_K(V)$ contains a coproduct of $2^{\text{card}(\Omega)}$ copies of itself as an associative K -algebra. □

Hence, for instance, if A is any group, not necessarily of cardinality $\leq \text{card}(\Omega)$, that is embeddable in $\text{Sym}(\Omega)$, then the coproduct of $2^{\text{card}(\Omega)}$ copies of A in the category of groups is also embeddable in $\text{Sym}(\Omega)$.

I should mention that at the beginning of this section, when I said that the cardinal $2^{\text{card}(\Omega)}$ appearing in our results was, in general, the best we could hope for, the phrase “in general” was a hedge. There is an exception, concerning the algebras $\text{End}_K(V)$ when K is a field of cardinality $> 2^{\text{card}(\Omega)}$. We will see at the end of §10 that in that case, we can get a stronger conclusion than Theorem 4.2(iii).

Note that Theorem 4.2, unlike the results of previous sections, says nothing about coproducts in subvarieties of our varieties. So we ask,

Question 4.3. *Suppose A is a group, monoid or associative K -algebra which belongs to a subvariety \mathbf{V} of the variety of all such algebras, and which is embeddable in $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$ or $\text{End}_K(V)$ respectively.*

Must the same be true of the coproduct in \mathbf{V} of two copies of A ? (If this is indeed true for all such A , the corresponding statement will hold for coproducts in \mathbf{V} of $2^{\text{card}(\Omega)}$ copies of such A , by Lemma 3.3 and Theorem 3.1.)

In the case where \mathbf{V} is the variety of *abelian groups*, or any subvariety thereof, one has an affirmative answer, for de Bruijn [8, Theorem 4.3] shows that *every* abelian group of cardinality $\leq 2^{\text{card}(\Omega)}$ is embeddable in $\text{Sym}(\Omega)$. However, the analog of this stronger statement fails for all varieties of groups not contained in the variety of abelian groups, by a result of McKenzie that will be recalled in the next section.

A question similar to the preceding, but concerning additional constants rather than additional identities, is

Question 4.4. Suppose B is a subgroup of $\text{Sym}(\Omega)$, a submonoid of $\text{Self}(\Omega)$, or a sub- K -algebra of $\text{End}_K(V)$.

Must $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$ or $\text{End}_K(V)$ respectively have a subalgebra containing B , and isomorphic over B to the coproduct of two copies of $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$ or $\text{End}_K(V)$ with amalgamation of B (i.e., isomorphic over B to the pushout, in the variety of all groups, semigroups, or K -algebras, of the diagram formed by B and two copies of the indicated algebra; equivalently, to the coproduct of two copies of that algebra in the variety of groups, monoids or K -algebras with distinguished constants corresponding to the elements of B)?

If this is not true in general, does it become true when B has some “good” form; e.g., in the case of $\text{Self}(\Omega)$, when B is a group of invertible elements, or in the case of $\text{End}_K(V)$ when B is a division algebra?

Turning back to the argument we used to get Theorem 4.2 from Lemma 4.1, we should note that a certain fact was implicitly called on which is true of the varieties of all groups, all monoids, and all associative K -algebras, and in many other familiar varieties of algebras, but not in all – namely that, given inclusions of algebras $A' \subseteq A$ and $B' \subseteq B$, the induced homomorphism of coproducts in our variety,

$$(20) \quad A' \amalg B' \rightarrow A \amalg B,$$

is also injective. It is this that allows us to say that if an algebra A contains a coproduct of two copies of itself, it contains a coproduct of any finite number of such copies.

An example of a variety \mathbf{V} where the injectivity of maps (20) fails is the variety of groups generated by the infinite dihedral group. To see this, note that \mathbf{V} satisfies the identity

$$(21) \quad (x^2, y^2) = 1,$$

but no identity $x^n = 1$ ($n > 0$). Let A and B be infinite cyclic groups $\langle x \rangle$ and $\langle y \rangle$; these are each free on one generator in \mathbf{V} . Let A', B' be the subgroups $\langle x^2 \rangle \subseteq A$ and $\langle y^2 \rangle \subseteq B$, which are isomorphic to A and B . The coproduct $A \amalg B$ in \mathbf{V} is the free algebra on $\{x, y\}$ in that variety, hence is noncommutative, and so the same is true of $A' \amalg B'$. But the image of $A' \amalg B'$ in $A \amalg B$ is generated by x^2 and y^2 , which commute by (21), so the map $A' \amalg B' \rightarrow A \amalg B$ is not an embedding.

Though this shows that the principle we used in the proof of Theorem 4.2 is not valid in all varieties, it does not show that the consequence of that principle that we used, concerning objects containing coproducts of copies of themselves, can fail. So we ask

Question 4.5. Does there exist an algebra A in a variety \mathbf{V} such that A contains the coproduct in \mathbf{V} of two copies of itself, but not the coproduct of three such copies?

The preceding results about fitting into $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$ and $\text{End}_K(V)$ multiple copies of themselves suggest questions about fitting these objects into each other, in various ways. We record two easy results in this direction:

Lemma 4.6. (i) $\text{End}_K(V)$ contains an embedded copy of $K \text{Self}(\Omega)$, the monoid algebra over K on the monoid $\text{Self}(\Omega)$.

(ii) $\text{Self}(\Omega)$ contains an embedded copy of $\text{Equiv}(\Omega)_\wedge$, i.e., $\text{Equiv}(\Omega)$ made a monoid under the meet operation \wedge .

Proof. (i) Let $K\langle\Omega\rangle$ denote the free associative K -algebra on Ω . Then the action of $\text{Self}(\Omega)$ on Ω induces an action of $\text{Self}(\Omega)$ on $K\langle\Omega\rangle$ by K -algebra endomorphisms, which extends to an action of the monoid algebra $K \text{Self}(\Omega)$ by vector-space endomorphisms of $K\langle\Omega\rangle$. Since $K\langle\Omega\rangle$ has the same vector-space dimension, $\text{card}(\Omega)$, as V , our assertion will follow if we can show that the endomorphisms of $K\langle\Omega\rangle$ induced by any finite family g_1, \dots, g_n of distinct elements of $\text{Self}(\Omega)$ are K -linearly independent.

Given such g_1, \dots, g_n , let us choose $p_1, \dots, p_m \in \Omega$ such that no two of g_1, \dots, g_n behave the same on all of these elements. Regarding p_1, \dots, p_m as members of the free generating set Ω of $K\langle\Omega\rangle$, we can form the product $p_1 \dots p_m$ therein, and observe that the actions of g_1, \dots, g_n take this monomial to distinct monomials, hence are indeed K -linearly independent.

(ii) If we regard equivalence relations on Ω as subsets of $\Omega \times \Omega$, then the meet operation on $\text{Equiv}(\Omega)$ is the restriction of the intersection operation on $\mathbf{P}(\Omega \times \Omega)$, hence it will suffice to embed $\mathbf{P}(\Omega \times \Omega)_\cap$ in

$\text{Self}(\Omega)$. Since $\text{card}(\Omega \times \Omega) = \text{card}(\Omega) = \text{card}(\Omega \times 2)$, we can do this, in turn, if we can embed $\mathbf{P}(\Omega)_\cap$ in $\text{Self}(\Omega \times 2)$. To do this, let us send each $S \subseteq \Omega$ to the endomap of $\Omega \times 2$ that fixes all elements $(p, 0)$, and also all elements $(p, 1)$ with $p \in S$, but sends $(p, 1)$ to $(p, 0)$ if $p \notin S$. The verification that this is a monoid homomorphism, and indeed an embedding, is straightforward. \square

We will see in §6 that the analog of statement (ii) above with “meet” replaced by “join” is false.

It is also interesting to note that the analog of (i) fails if Ω replaced by a finite set with $n \geq 2$ elements. Indeed, for $n \geq 3$, even the group algebra $K \text{Sym}(\Omega)$ cannot be embedded in $\text{End}_K(V)$, for it has dimension $n!$, while $\text{End}_K(V)$ has dimension only n^2 . In particular, the $n!$ permutation matrices do not generate a copy of the group algebra – they are not linearly independent. To get the nonembeddability statement for $K \text{Self}(\Omega)$ when $n = 2$, let $R = K \text{Self}(2)$, and note by comparing dimensions that a K -algebra embedding of R in $M_2(K)$ would have to be an isomorphism. Let $z \in \text{Self}(2)$ be the map taking both elements of 2 to 0, and note that it satisfies the left-zero identity $(\forall a) za = z$. Hence $zR \subseteq Kz$, hence $zR(1 - z) = 0$; but $M_2(K)$ has no idempotent z with this property other than 0 and 1.

5 Restrictions on groups embeddable in $\text{Sym}(\Omega)$ and monoids embeddable in $\text{Self}(\Omega)$.

With such vast classes of groups, monoids, associative K -algebras and lattices embeddable in $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$, $\text{End}_K(V)$, and $\text{Equiv}(\Omega)$, it is natural to ask whether there are groups, etc., of cardinality, respectively K -dimension, $\leq 2^{\text{card}(\Omega)}$, that are not so embeddable.

For the case of groups, de Bruijn [7] showed, in effect, that for any set I of cardinality $> \text{card}(\Omega)$, the group presented by generators x_i ($i \in I$) and relations

$$(22) \quad x_i^2 = 1 \quad (i \in I),$$

$$(23) \quad (x_i x_j)^3 = 1 \quad (i, j \in I, \text{ distinct}),$$

$$(24) \quad (x_i x_j x_k x_l)^5 = 1 \quad (i, j, k, l \in I, \text{ distinct}),$$

cannot be embedded in $\text{Sym}(\Omega)$. Note that the fact that the indices in (23) and (24) are required to be distinct keeps this system of groups from having the form to which the results of §3 apply. (If those indices were not required to be distinct, then setting $k = i$, $l = j$ in (24) would give $(x_i x_j)^{10} = 1$, which, combined with (23), would give $x_i x_j = 1$, making the group collapse to Z_2 .)

On the other hand, de Bruijn claimed in [7] that his result corresponding to Theorem 3.1 showed that the restricted direct product (called in [7] the direct product) of $2^{\text{card}(\Omega)}$ copies of any group A of cardinality $\leq \text{card}(\Omega)$ could be embedded in $\text{Sym}(\Omega)$ – not noticing that because the commutativity relations which the restricted direct product construction imposes on elements of different copies of A fail to hold among elements of a single copy (unless A is commutative), that result is not applicable. In [8] he corrected this error, noting that the argument is only valid when A is abelian, and posed his earlier assertion as an open question.

That question was answered in the negative by McKenzie [16], who showed that if G is a group such that for some index set I with $\text{card}(I) > \text{card}(\Omega)$ there are elements $x_i, y_i \in G$ ($i \in I$) satisfying

$$(25) \quad x_i y_j = y_j x_i \text{ whenever } i \neq j, \text{ but } x_i y_i \neq y_i x_i,$$

then G cannot be embedded in $\text{Sym}(\Omega)$.

We shall see that McKenzie’s criterion is an instance of more general facts. By a *centralizer subgroup* in a group G , let us understand a subgroup of the form

$$(26) \quad C_G(X) = \{g \in G \mid (\forall x \in X) gx = xg\}$$

for some subset $X \subseteq G$. A subgroup $H < G$ is clearly a centralizer subgroup if and only if $H = C_G(C_G(H))$. Recall also that a *jump* in a totally ordered set means a pair of elements $x < y$ such that $\{z \mid x < z < y\}$ is empty. A totally ordered set without jumps can have subsets with jumps; for instance, the set of reals or of rationals has none, but their subset \mathbb{Z} has countably many. We shall see below that the lattice of centralizer

subgroups of $\text{Sym}(\Omega)$, and hence of any group embeddable therein, contains no chains with $> \text{card}(\Omega)$ jumps, while the lattice of centralizer subgroups of a group with a family of elements satisfying (25) does have such chains.

In fact, we shall prove the former result not only for centralizer subgroups, but for subsets of $\text{Sym}(\Omega)$ defined by arbitrary systems of equations (in several variables) with constants in $\text{Sym}(\Omega)$, which will also yield a quick proof of de Bruijn's example. Our result will follow from the fact that such solution subsets are closed in the function topology on $\text{Sym}(\Omega)$, together with the following lemma in general topology.

Note that unless explicitly stated, we do not assume topologies to be Hausdorff. The function topology, to which we will apply the lemma in this section, is Hausdorff, but in the next section we will apply the same lemma to both Hausdorff and non-Hausdorff topologies.

Lemma 5.1. *Let T be a topological space having an infinite basis (or more generally, subbasis) B of open sets. Then the lattice of open subsets of T contains no chain with $> \text{card}(B)$ jumps. Hence its opposite, the lattice of closed subsets of T , also has no such chains. In particular, that lattice contains no well-ordered or reverse-well-ordered chains of cardinality $> \text{card}(B)$.*

Proof. The case where B is a subbasis reduces to that in which it is a basis, since in the former case, a basis is given by the set of intersections of finite subsets of B , and for B infinite there are only $\text{card}(B)$ of these. So we assume B a basis.

Suppose C is a chain of open subsets of T . For each jump $U \subset V$ in C , let us choose a point $p_{(U,V)} \in V - U$. Since V is a neighborhood of $p_{(U,V)}$, our basis B contains some subneighborhood $N_{(U,V)} \subseteq V$ of $p_{(U,V)}$; let $N_{(U,V)}$ be so chosen for each jump $U \subset V$. Then if $U \subset V$ and $U' \subset V'$ are distinct jumps, say with $U \subset V \subseteq U' \subset V'$, we must have $N_{(U,V)} \neq N_{(U',V')}$, since $N_{(U,V)} \subseteq V$, while $N_{(U',V')}$ contains $p_{(U',V')} \notin U'$. Hence distinct jumps in C give distinct elements $N_{(U,V)} \in B$, so the number of jumps in C does not exceed $\text{card}(B)$. \square

Recall next that if Ω and Ω' are sets, and we give Ω' the discrete topology, then the *function topology* on the set of all maps $\Omega \rightarrow \Omega'$ has for a subbasis of open sets the sets $U_{y,x} = \{f : \Omega \rightarrow \Omega' \mid f(x) = y\}$ ($x \in \Omega$, $y \in \Omega'$), since a basis for the open sets of Ω' is given by the singletons $\{y\}$. In particular, if Ω is infinite, the function topology on the set $\text{Self}(\Omega)$ has a subbasis of cardinality $\text{card}(\Omega \times \Omega) = \text{card}(\Omega)$. Hence for any set J , the direct product of a J -tuple of copies of this space has a subbasis of cardinality $\text{card}(\Omega) \text{card}(J)$. So we get

Corollary 5.2. *If $\text{card}(J) \leq \text{card}(\Omega)$, then the lattice of subsets of $\text{Self}(\Omega)^J$ closed in the product topology on that set induced by the function topologies on the factors $\text{Self}(\Omega)$ has no chains with $> \text{card}(\Omega)$ jumps. \square*

Let us now connect this topology with our algebraic structure. It is straightforward to verify that the operation of composition on $\text{Self}(\Omega)$ is continuous in the function topology; moreover, on its subset $\text{Sym}(\Omega)$, the operation of functional inverse is also continuous, since it simply interchanges $U_{y,x} \cap \text{Sym}(\Omega)$ and $U_{x,y} \cap \text{Sym}(\Omega)$ for all x and y . (We remark, however, that $\text{Sym}(\Omega)$ is not closed in $\text{Self}(\Omega)$.) Hence given a pair of monoid words (respectively group words) v , w , in a variable t and constants from $\text{Self}(\Omega)$ (respectively, from $\text{Sym}(\Omega)$), the solution set $\{a \mid v(a) = w(a)\}$ will be closed in the function topology on $\text{Self}(\Omega)$ (respectively, $\text{Sym}(\Omega)$). More generally, we may look at the solution set of any family of such pairs of words in any number of variables. Let us set up notation for such sets in an arbitrary algebra.

Definition 5.3. *For A an algebra in a variety \mathbf{V} , and J a set, we shall understand a principal solution set in A^J to mean a set of the form*

$$(27) \quad S_{v=w} = \{a = (a_j)_{j \in J} \in A^J \mid v(a) = w(a)\} \subseteq A^J,$$

where v and w are words in a J -tuple of variables $(t_j)_{j \in J}$, constants from A , and the operations of \mathbf{V} .

A solution set in A^J will mean the intersection of an arbitrary family of principal solution sets. We shall denote by $L_{A,J}^-$ the complete lattice of all solution sets in A^J .

Here we understand the intersection of the *empty* family of principal solution sets to be the whole set A^J . Thus, $L_{A,J}^-$ is, as asserted, a complete lattice, the join of any $X \subseteq L_{A,J}^-$ being the intersection of those principal solution sets that contain all members of X . (The superscript “ $-$ ” in $L_{A,J}^-$ indicates that

our solution sets are defined by equations, as in (27). In the next section we shall also make use of solution sets defined by inequalities.)

In the next result, though J is allowed to have cardinality up to $\text{card}(\Omega)$, the most common cardinality in our applications will be 1.

Theorem 5.4. *Let J be any set of cardinality $\leq \text{card}(\Omega)$. Then $L_{\text{Sym}(\Omega), J}^{\bar{\bar{}}}$ contains no chains with $> \text{card}(\Omega)$ jumps. Hence the same is true of $L_{G, J}^{\bar{\bar{}}}$ for any group G embeddable in $\text{Sym}(\Omega)$.*

Likewise, $L_{\text{Self}(\Omega), J}^{\bar{\bar{}}}$ contains no chains with $> \text{card}(\Omega)$ jumps; hence the same is true of $L_{M, J}^{\bar{\bar{}}}$ for any monoid M embeddable in $\text{Self}(\Omega)$.

In particular, for G a group embeddable in $\text{Sym}(\Omega)$ or M a monoid embeddable in $\text{Self}(\Omega)$, the lattice $L_{G, J}^{\bar{\bar{}}}$, respectively $L_{M, J}^{\bar{\bar{}}}$ contains no well-ordered or reverse-well-ordered chain of cardinality $> \text{card}(\Omega)$.

Proof. The assertions about chains in $L_{\text{Sym}(\Omega), J}^{\bar{\bar{}}}$ and $L_{\text{Self}(\Omega), J}^{\bar{\bar{}}}$ are clear from Corollary 5.2 and the continuity of our operations; it remains to deduce the corresponding statements for objects embeddable in $\text{Sym}(\Omega)$ and $\text{Self}(\Omega)$.

If G is a group embeddable in $\text{Sym}(\Omega)$, let us assume for notational convenience that it is a subgroup, and map $L_{G, J}^{\bar{\bar{}}}$ to $L_{\text{Sym}(\Omega), J}^{\bar{\bar{}}}$ by sending each solution set S in G^J to the solution set in $\text{Sym}(\Omega)^J$ of the set of all equations (in a J -tuple of variables, with constants in G) that are satisfied on S . This map is easily seen to be an embedding of partially ordered sets, hence the result on chains in $L_{\text{Sym}(\Omega), J}^{\bar{\bar{}}}$ implies the same conclusion for chains in $L_{G, J}^{\bar{\bar{}}}$. The same argument works for monoids. \square

Some observations on the above proof: Given groups $G < H$, one cannot embed $L_{G, J}^{\bar{\bar{}}}$ in $L_{H, J}^{\bar{\bar{}}}$ by simply sending the solution set of every system of equations in G to the solution set of the same system in H . This does not give a well-defined function, since equations over G having the same solution set in G^J may have different solution sets in H^J . (Consider, for instance, centralizer subgroups of various sets in an abelian group, and of the same sets in a nonabelian overgroup.) The construction of the above proof does give an order-embedding of $L_{G, J}^{\bar{\bar{}}}$ into $L_{H, J}^{\bar{\bar{}}}$, but in general this respects neither meets nor joins; the former because the set H^J is larger than G^J ; the latter because the set of equations with constants in H is larger than the set of equations with constants in G . Another order-embedding of $L_{G, J}^{\bar{\bar{}}}$ in $L_{H, J}^{\bar{\bar{}}}$ is gotten by sending every $S \in L_{G, J}^{\bar{\bar{}}}$ to the set of elements of H^J satisfying all equations *with constants in H* satisfied on S ; it also respects neither meets nor joins, in general.

For our first application of Theorem 5.4, note that the centralizer subgroups (26) in a group G form a complete lattice, which as a partially ordered set (and indeed, as a complete lower semilattice) is embedded in $L_{G, 1}^{\bar{\bar{}}}$. Hence we have

Corollary 5.5. *No group having a chain of centralizer subgroups with $> \text{card}(\Omega)$ jumps is embeddable in $\text{Sym}(\Omega)$. In particular (McKenzie, [16]), if a group G contains, for some set I with $\text{card}(I) > \text{card}(\Omega)$, elements x_i, y_i ($i \in I$) satisfying (25), then G is not embeddable in $\text{Sym}(\Omega)$.*

Proof. The first statement is clear from the first paragraph of Theorem 5.4. In the situation of the second statement, we may, by reindexing, assume I to be a cardinal $\kappa > \text{card}(\Omega)$. For each $\alpha \in \kappa$, let $X_\alpha = \{x_\beta \mid \beta > \alpha\}$. The X_α form a descending chain of subsets, hence their centralizers $C_G(X_\alpha)$ form an ascending chain of centralizer subgroups. Note that each $C_G(X_\alpha)$ contains those elements y_γ with $\gamma \leq \alpha$ and no other y_γ , hence the $C_G(X_\alpha)$ are distinct. Thus we have a well-ordered chain of centralizer subgroups of cardinality $\kappa > \text{card}(\Omega)$; hence G is not embeddable in $\text{Sym}(\Omega)$. \square

The cardinality conditions in the above result are sharp: If we take any set I of cardinality $\leq \text{card}(\Omega)$, and any I -tuple G_i of nonabelian groups each of cardinality $\leq \text{card}(\Omega)$, then their restricted direct product has cardinality $\leq \text{card}(\Omega)$, hence is embeddable in $\text{Sym}(\Omega)$, though it contains elements x_i, y_i satisfying (25).

Turning to de Bruijn's relations (22)-(24), note that for any index set I of cardinality $\leq \text{card}(\Omega)$, if we take an element $p_0 \in \Omega$ and an I -tuple of elements $p_i \in \Omega$ distinct from p_0 and from each other, and for each $i \in I$ let $x_i \in \text{Sym}(\Omega)$ be the transposition that interchanges p_0 and p_i and fixes all other elements, then any product of n distinct elements x_i is an $(n+1)$ -cycle in $\text{Sym}(\Omega)$, from which we see that the

I -tuple $(x_i)_{i \in I}$ satisfies (22)-(24). This shows that the cardinality conditions in the next result are sharp. That result, in fact, does without (22), at the small price of adding two sentences at the start of the proof.

Corollary 5.6 (cf. [7, Theorem 5.1]). *No group G containing a family $(x_i)_{i \in I}$ of distinct elements satisfying (23) and (24), where $\text{card}(I) > \text{card}(\Omega)$, is embeddable in $\text{Sym}(\Omega)$.*

Proof. If there are any pairs $i \neq i' \in I$ such that x_i and $x_{i'}$ are inverse to one another, then dropping one member of each such pair does not decrease $\text{card}(I)$. Hence we may assume there are no such pairs.

Let us also assume, by reindexing, that I has the form $\kappa \times 2$, where κ is a cardinal $> \text{card}(\Omega)$, so that our given elements have the form $x_{\beta,i}$ ($\beta \in \kappa$, $i = 0, 1$). For each $\alpha \in \kappa$, let us define the solution set $S_\alpha = \{(y, z) \in G^2 \mid (\forall \beta > \alpha) (x_{\beta,0} x_{\beta,1} y z)^5 = 1\}$. By (24), this set contains the pair $(x_{\gamma,0}, x_{\gamma,1})$ whenever $\gamma \leq \alpha$.

However, it contains no pair $(x_{\beta,0}, x_{\beta,1})$ with $\beta > \alpha$. Indeed, if it did, we would have

$$(28) \quad 1 = (x_{\beta,0} x_{\beta,1} x_{\beta,0} x_{\beta,1})^5 = (x_{\beta,0} x_{\beta,1})^{10}.$$

But by (23), $(x_{\beta,0} x_{\beta,1})^3 = 1$. Combining these equations we get $x_{\beta,0} x_{\beta,1} = 1$, contradicting our assumption that for i and i' distinct, x_i and $x_{i'}$ are not inverses.

Hence, the sets S_α are distinct, and so form a well-ordered chain of cardinality κ in $L_{G,2}^-$, from which nonembeddability of G in $\text{Sym}(\Omega)$ follows by Theorem 5.4. \square

What about applications of Theorem 5.4 to monoids? Well, the monoid homomorphisms from a group G to a monoid M are the group homomorphisms from G to the group of invertible elements of M ; hence the above two corollaries can also be viewed as giving monoids that are not embeddable in $\text{Self}(\Omega)$.

Here is a more genuinely monoid-theoretic application. Consider again an element $p_0 \in \Omega$ and a family of distinct elements $p_i \in \Omega - \{p_0\}$ indexed by a set I of the same cardinality as Ω . Let $y \in \text{Self}(\Omega)$ be the map sending all elements to p_0 , while for each $i \in I$, let x_i be the map sending everything except p_i to p_0 , and fixing p_i . Then we see that

$$(29) \quad \text{for } i, j \in I, \quad x_i x_j = y \text{ if and only if } i \neq j.$$

This gives $\text{card}(\Omega)$ such elements x_i ; but an application of Theorem 5.4, following the same pattern as the two preceding results, shows that we cannot get a family of $> \text{card}(\Omega)$ such elements; hence

Corollary 5.7. *No monoid containing an element y , and a family of elements x_i , distinct from y , indexed by a set I of cardinality $> \text{card}(\Omega)$, and satisfying (29), is embeddable in $\text{Self}(\Omega)$.* \square

In our examples of nonembeddability using the group conditions (23)-(24) and (25), and the monoid conditions (29), we could have asserted much more than the existence of a chain with $\text{card}(I)$ jumps. For example, given an I -tuple of elements satisfying (25), distinct subsets of $\{x_i \mid i \in I\}$ have centralizers containing distinct subsets of the y_i (indexed by the complementary subsets of I), so we in fact get a copy of the whole partially ordered set $\mathbf{P}(I)$ in $L_{G,1}^-$; and the corresponding observations hold for the other two examples.

However, there are examples that give large chains of solution sets without (as far as I can see) giving so much more as well. If we take the group or monoid presented by (10), respectively (11), with I a cardinal κ , then it will have a chain of centralizers, respectively fixed sets, order-isomorphic to κ (as well as one of the opposite order type), but there is no apparent reason why it should have, say, any large antichain of solution sets.

It is interesting that while the constructions of (10) and (11) with $I = \mathbb{R}$ give, as we saw earlier, groups and monoids embeddable in $\text{Sym}(\omega)$, respectively $\text{Self}(\omega)$, the above paragraph shows that the contrary is true for the same constructions with $I = \text{card}(\mathbb{R})$ as a well-ordered set. We likewise get nonembeddability when $I = \mathbb{R} \times 2$ under lexicographic order (since it also has uncountably many jumps), and for $\mathbb{R} \times \mathbb{R}$ under lexicographic order (since this contains the preceding ordered set).

Here is another application of our observations on (10). For every real number c , let G_c be the group presented by generators x_r, y_r ($r \in \mathbb{R}$) and relations

$$(30) \quad x_r y_s = y_s x_r \text{ for all } r, s \in \mathbb{R} \text{ such that } s \geq r + c.$$

Clearly, each G_c is isomorphic to G_0 , by an isomorphism that fixes the x_r and takes each y_s to y_{s-c} . But we have noted that (by Theorem 3.4) G_0 is embeddable in $\text{Sym}(\aleph_0)$; hence so is every G_c . Now let G_{0+} denote the group with the same generators, but having for relations the union of the sets of relations defining G_c for all $c > 0$; in other words,

$$(31) \quad x_r y_s = y_s x_r \text{ for all } r, s \in \mathbb{R} \text{ with } s > r,$$

and let us define therein, for every $c \in \mathbb{R}$, the centralizer subgroups

$$(32) \quad \begin{aligned} S_c &= \{g \in G_{0+} \mid (\forall s \geq c) g y_s = y_s g\}, \\ S_{c+} &= \{g \in G_{0+} \mid (\forall s > c) g y_s = y_s g\}. \end{aligned}$$

I claim that these form a chain, with jumps $S_c \subset S_{c+}$, and with inclusions $S_{c+} \subset S_d$ whenever $c < d$. Indeed, $S_c \neq S_{c+}$ because $x_c \in S_{c+} - S_c$, and similarly $S_{c+} \neq S_d$ by considering x_e for any e with $c < e < d$. This chain is isomorphic to $\mathbb{R} \times 2$, so G_{0+} is not embeddable in $\text{Sym}(\aleph_0)$, though it is a direct limit, via surjective homomorphisms, of the groups G_c ($c > 0$), which are so embeddable.

Though a positive answer seems implausible, let us ask

Question 5.8. *Is the criterion of the first paragraph of Theorem 5.4 also sufficient for a group of cardinality $\leq 2^{\text{card}(\Omega)}$ to be embeddable in $\text{Sym}(\Omega)$? If so, is it sufficient that it hold for all finite J ? For $J = 1$?*

In an earlier version of this note, I asked the same questions for $\text{Self}(\Omega)$; but the possibility of an affirmative answer is now precluded by a result of Wehrung [19], showing that $\text{Self}(\Omega)^{\text{op}}$ cannot be embedded in $\text{Self}(\Omega)$; indeed, since the lattices $L_{\text{Self}(\Omega), J}^-$ and $L_{\text{Self}(\Omega)^{\text{op}}, J}^-$ are isomorphic, that result shows that no condition on the lattices $L_{M, J}^-$ can be equivalent to embeddability of M in $\text{Self}(\Omega)$. Conceivably, however, one could construct lattices of solution sets not using all the sets $S_{v=w}$, but some subfamilies of these that are not invariant under reversing the orders of factors in the words u and v , such that conditions on these solution sets would characterize embeddability in $\text{Self}(\Omega)$.

It is easy to *formally* strengthen Theorem 5.4 in several ways. First, since the sets $S_{v=w}$ of (27) are closed in the function topology, so are finite unions of such sets, which we might write

$$(33) \quad S_{(v_1=w_1) \vee \dots \vee (v_n=w_n)} = S_{v_1=w_1} \cup \dots \cup S_{v_n=w_n}.$$

So if we let $L_{\text{Sym}(\Omega), J}^{\bar{=}, \vee}$ denote the lattice of arbitrary intersections of families of finite unions (33), these will also satisfy the conditions on chains given by Theorem 5.4.

Secondly, the conclusion of Theorem 5.4 only states one particular consequence of embeddability of our lattice of solution sets in the lattice of closed sets of a topology generated by $\leq \text{card}(\Omega)$ elements. We will examine the latter condition further in an appendix, §12. Meanwhile, we ask

Question 5.9. *If the criteria of Theorem 5.4 are not sufficient for a group X of cardinality $\leq 2^{\text{card}(\Omega)}$ to be embeddable in $\text{Sym}(\Omega)$, do they become so if we replace the lattices $L_{X, J}^-$ of that theorem by the larger lattices $L_{X, J}^{\bar{=}, \vee}$, and/or strengthen the condition on jumps in chains to the condition that our lattice of solution sets be embeddable as a partially ordered set in the lattice of closed subsets of a topological space with a basis of cardinality $\leq \text{card}(\Omega)$ (cf. §12)?*

Do there, at least, exist groups whose embeddability in $\text{Sym}(\Omega)$ is precluded by one of these strengthened conditions, but not by the conditions of Theorem 5.4? Here we may ask the same question for embeddability of monoids in $\text{Self}(\Omega)$.

We mentioned (following Question 4.3) de Bruijn's result that every abelian group of cardinality $\leq 2^{\text{card}(\Omega)}$ is embeddable in $\text{Sym}(\Omega)$. However, not every commutative monoid of cardinality $\leq 2^{\text{card}(\Omega)}$ embeds in $\text{Self}(\Omega)$: the presentations (11) give commutative monoids, but we saw that for I a cardinal $> \text{card}(\Omega)$, the resulting monoid is not so embeddable. So we ask

Question 5.10. *Which varieties \mathbf{V} of monoids have the property that every monoid in \mathbf{V} of cardinality $\leq 2^{\text{card}(\Omega)}$ is embeddable in $\text{Self}(\Omega)$?*

By de Bruijn's result on abelian groups, this is true of every variety of commutative monoids satisfying an identity $x^n = 1$, since such monoids are essentially abelian groups of exponent n , hence embeddable in $\text{Sym}(\Omega)$ as groups. I don't know any other examples.

Returning to groups, suppose we write $\text{Sym}_{<}(\Omega) \subset \text{Sym}(\Omega)$ for the normal subgroup of permutations that move fewer than $\text{card}(\Omega)$ elements. De Bruijn [7, Theorem 4.4] showed that $\text{Sym}(\Omega)$ could be embedded in $\text{Sym}(\Omega)/\text{Sym}_{<}(\Omega)$, while McKenzie [16, Corollary 3] showed that $\text{Sym}(\Omega)/\text{Sym}_{<}(\Omega)$ contains a restricted direct product of $> \text{card}(\Omega)$ copies of itself, and hence, by Corollary 5.5, cannot be embedded in $\text{Sym}(\Omega)$.

Question 5.11. *What restrictions does embeddability in $\text{Sym}(\Omega)/\text{Sym}_{<}(\Omega)$ imply for a group of cardinality $\leq 2^{\text{card}(\Omega)}$?*

Under the assumption of the General Continuum Hypothesis, Jónsson [14] shows, inter alia, that for every uncountable cardinal κ there exists a group of cardinality κ which contains isomorphic copies of *all* groups of cardinality κ . We have seen that $\text{Sym}(\Omega)$ is *not* such a group for $\kappa = 2^{\text{card}(\Omega)}$. Felgner and Haug [12] show that under certain set-theoretic hypothesis, neither is $\text{Sym}(\Omega)/\text{Sym}_{<}(\Omega)$.

6 Restrictions on lattices embeddable in $\text{Equiv}(\Omega)$.

If we want to adapt the technique of the preceding section to get restrictions on lattices embeddable in the lattice $\text{Equiv}(\Omega)$ of equivalence relations on Ω , we must decide what topology on that lattice to use in place of the function topology. One approach is to regard binary relations on Ω as elements of $\mathbf{P}(\Omega \times \Omega) = 2^{\Omega \times \Omega}$, i.e., as functions $\Omega \times \Omega \rightarrow 2 = \{0, 1\}$, and use the function topology on that set induced by the discrete topology on 2. A subbasis of open subsets of $\mathbf{P}(\Omega \times \Omega)$ under this topology is given by the sets

$$(34) \quad \begin{aligned} U_{p,q} &= \{R \in \mathbf{P}(\Omega \times \Omega) \mid (p, q) \in R\} \quad \text{and} \\ {}^c U_{p,q} &= \{R \in \mathbf{P}(\Omega \times \Omega) \mid (p, q) \notin R\}. \end{aligned}$$

We see that each of these sets is clopen (closed and open) in the topology so defined, and that this subbasis has cardinality $\text{card}(\Omega)$.

By abuse of notation, in speaking of subsets of $\text{Equiv}(\Omega)$ let us write $U_{p,q}$ for $U_{p,q} \cap \text{Equiv}(\Omega)$ and ${}^c U_{p,q}$ for ${}^c U_{p,q} \cap \text{Equiv}(\Omega)$ (just as, in introducing the function topology on $\text{Self}(\Omega)$, we earlier wrote $U_{p,q}$ for what we would now describe as $U_{p,q} \cap \text{Self}(\Omega)$).

In this topology, one finds that the meet operation, i.e., intersection as subsets of $\Omega \times \Omega$, is continuous, but that the join operation is not. To see the first fact, note that under the map

$$(35) \quad \wedge : \text{Equiv}(\Omega) \times \text{Equiv}(\Omega) \rightarrow \text{Equiv}(\Omega),$$

the inverse image of every $U_{p,q}$ is the open rectangle $U_{p,q} \times U_{p,q}$, while the inverse image of ${}^c U_{p,q}$ is the union $({}^c U_{p,q} \times \text{Equiv}(\Omega)) \cup (\text{Equiv}(\Omega) \times {}^c U_{p,q})$, and both these sets are open. Under the operation

$$(36) \quad \vee : \text{Equiv}(\Omega) \times \text{Equiv}(\Omega) \rightarrow \text{Equiv}(\Omega),$$

the inverse image of $U_{p,q}$ is still open: it is an infinite union of finite intersections of sets of the forms $U_{r,s} \times \text{Equiv}(\Omega)$ and $\text{Equiv}(\Omega) \times U_{r,s}$, one such intersection for each finite chain of relations which, if they hold in a pair of equivalence relations on Ω , witness the conclusion that (p, q) belongs to the join of those equivalence relations. But the inverse image of ${}^c U_{p,q}$ becomes, by the same reasoning, an infinite *intersection* of clopen sets, which will not be open. (Essentially because no *finite* set of relations and negations of relations can witness the absence of (p, q) from the join of two equivalence relations.)

There are two ways to respond to this difficulty. One is to formulate a criterion in terms of words in the meet operation alone. This corresponds to regarding $\text{Equiv}(\Omega)$ as a meet-semilattice, $\text{Equiv}(\Omega)_{\wedge}$. In the notation of Definition 5.3, we then look at the lattices $L_{\text{Equiv}(\Omega)_{\wedge}, J}^{\equiv}$, and get

Theorem 6.1. *Let J be any set of cardinality $\leq \text{card}(\Omega)$. Then $L_{\text{Equiv}(\Omega)_{\wedge}, J}^{\equiv}$ contains no chains with $> \text{card}(\Omega)$ jumps. Hence for any lower semilattice A embeddable in $\text{Equiv}(\Omega)_{\wedge}$, the lattice $L_{A, J}^{\equiv}$ contains no chains with $> \text{card}(\Omega)$ jumps. In particular, for any lattice A embeddable in $\text{Equiv}(\Omega)$, the lattice $L_{A_{\wedge}, J}^{\equiv}$ has no such chains.*

Proof. This can be gotten by the same method as Theorem 5.4, or deduced therefrom using Lemma 4.6(ii). \square

The following application of this result shows that the meet-join asymmetry we have come up against is real; and the second statement gives the promised example showing that the analog of Lemma 4.6(ii) with $\text{Equiv}(\Omega)_\vee$ in place of $\text{Equiv}(\Omega)_\wedge$ is false.

Proposition 6.2. *The largest cardinality of a set I such that $\text{Equiv}(\Omega)$ contains an element z and elements $x_i \neq z$ ($i \in I$) satisfying*

$$(37) \quad x_i \wedge x_j = z \text{ for all distinct } i, j \in I$$

is $\text{card}(\Omega)$.

However, $\text{Equiv}(\Omega)$ contains an element w and $2^{\text{card}(\Omega)}$ elements $y_i \neq w$ such that

$$(38) \quad y_i \vee y_j = w \text{ for all distinct } i, j \in I.$$

Proof. The upper bound in the first assertion follows from the preceding theorem, by the same reasoning used to get our corollaries to Theorem 5.4. To see without calling on Whitman's Theorem that there does, however, exist such a family of cardinality $\text{card}(\Omega)$, let z be the discrete equivalence relation on Ω , choose an element $p_0 \in \Omega$ and a family of $\text{card}(\Omega)$ distinct elements $p_i \in \Omega - \{p_0\}$, and for each i , let x_i be the equivalence relation that relates p_0 with p_i , but relates no other distinct elements of Ω .

To get the second assertion, let w be the indiscrete equivalence relation, and let the y_i be all the equivalence relations on Ω having exactly two equivalence classes. \square

The other way to deal with the fact that the join operation on $\text{Equiv}(\Omega)$ is not continuous in the topology with subbasis (34) is to weaken the topology. By the discussion following (34), both operations are continuous in the topology having the sets $U_{p,q}$ as a subbasis of open sets. This topology is not T_1 : for every $x \in \text{Equiv}(\Omega)$, we see that the closure of $\{x\}$ is the set of all equivalence relations $\leq x$.

As a consequence, the diagonal subset of $\text{Equiv}(\Omega) \times \text{Equiv}(\Omega)$ is not closed; hence, though arbitrary lattice words v and w with constants in $\text{Equiv}(\Omega)$ still induce continuous operations on $\text{Equiv}(\Omega)$, it does not follow that the sets $S_{v=w} = \{x \in \text{Equiv}(\Omega) \mid v(x) = w(x)\}$ are closed.

However, let us make

Definition 6.3. *For any lattice A , any set J , any lattice word v in a J -tuple of variables with constants in A , and any element $c \in A$, let*

$$(39) \quad S_{v \leq c} = \{a = (a_j)_{j \in J} \in A^J \mid v(a) \leq c\} \subseteq A^J,$$

and let us call such sets principal lower solution sets in A^J . A lower solution set will mean the intersection of an arbitrary family of principal lower solution sets. We shall denote by $L_{A,J}^{\leq \text{const}}$ the lattice of all lower solution sets in A^J .

As in Definition 5.3, we allow the empty intersection, so that $L_{A,J}^{\leq \text{const}}$ is indeed a complete lattice.

Any principal lower solution set $S_{v \leq c}$ in $\text{Equiv}(\Omega)^J$ is the inverse image under the continuous map $v : \text{Equiv}(\Omega)^J \rightarrow \text{Equiv}(\Omega)$ of the closed set $\{d \mid d \leq c\} \subseteq \text{Equiv}(\Omega)$, hence is closed, making Lemma 5.1 applicable. Thus we get

Theorem 6.4. *Let J be any set of cardinality $\leq \text{card}(\Omega)$. Then $L_{\text{Equiv}(\Omega),J}^{\leq \text{const}}$ contains no chains with $> \text{card}(\Omega)$ jumps. Hence for any lattice A embeddable in $\text{Equiv}(\Omega)$, $L_{A,J}^{\leq \text{const}}$ contains no such chains. \square*

To get applications of this result that are not consequences of Theorem 6.1, one has to use sets $S_{v \leq c}$ determined by words v that involve both meets and joins; in fact, that involve meets of joins. For if v does not involve meets of joins, we can write $v = \bigvee_{r=1}^m \bigwedge_{s=1}^{n_r} a_{r,s}$, where the $a_{r,s}$ are variables and/or constants, and m, n_1, \dots, n_m are positive integers (some of which may be 1). Then we see that the relation $v \leq c$ is equivalent to the conjunction of the relations $\bigwedge_{s=1}^{n_r} a_{r,s} \leq c$ ($r = 1, \dots, m$), and each of these can be rewritten $c \wedge \bigwedge_{s=1}^{n_m} a_{r,s} = \bigwedge_{s=1}^{n_m} a_{r,s}$. Hence $S_{v \leq c} \in L_{\text{Equiv}(\Omega)_\wedge, J}^{\leq \text{const}}$, and we are reduced to Theorem 6.1.

Perhaps workers in lattice theory will be able to see interesting applications of Theorem 6.4 that do not reduce to Theorem 6.1.

We remark that while Theorems 6.1 and 6.4 show that $L_{\text{Equiv}(\Omega)_{\wedge}, J}^{\overline{=}}$ and $L_{\text{Equiv}(\Omega), J}^{\leq \text{const}}$ contain no well-ordered chains of cardinality $> \text{card}(\Omega)$, the second paragraph of Proposition 6.2 shows that $L_{\text{Equiv}(\Omega), J}^{\overline{=}}$ and $L_{\text{Equiv}(\Omega)_{\vee}, J}^{\overline{=}}$ do contain such chains.

7 Monoids embeddable in $\text{Rel}(\Omega)$.

In this section we return to monoids, but focus, not on $\text{Self}(\Omega)$, but on its “wild sibling” $\text{Rel}(\Omega)$, the monoid of all binary *relations* on Ω , under relational composition:

$$(40) \quad xy = \{(q, p) \in \Omega \times \Omega \mid (\exists r \in \Omega) (q, r) \in x, (r, p) \in y\}.$$

Note that when specialized to functions, the above definition of composition, together with our convention that functions act on the left and are composed accordingly, requires that we identify each function f with the set of ordered pairs $(f(p), p)$, rather than the more usual $(p, f(p))$.

Clearly, $\text{Rel}(\Omega) \cong \text{Rel}(\Omega)^{\text{op}}$, and this monoid contains $\text{Self}(\Omega)$; hence it also contains an isomorphic copy of $\text{Self}(\Omega)^{\text{op}}$. In fact, it is generated by $\text{Self}(\Omega)$ and the natural copy of $\text{Self}(\Omega)^{\text{op}}$; for if $R \in \text{Rel}(\Omega)$ is nonempty, then since, as a subset of $\Omega \times \Omega$, it has cardinality $\leq \text{card}(\Omega)$, we can find maps $f, g \in \text{Self}(\Omega)$ such that $R = \{(f(p), g(p)) \mid p \in \Omega\}$. Letting $\bar{g} = \{(p, g(p)) \mid p \in \Omega\} \in \text{Self}(\Omega)^{\text{op}}$, we easily verify that $f\bar{g} = R$. The empty relation, on the other hand, equals $\bar{f}g$ for any f and g having disjoint ranges.

The next result shows that monoids embeddable in $\text{Rel}(\Omega)$ are considerably less restricted than those embeddable $\text{Self}(\Omega)$ or $\text{Self}(\Omega)^{\text{op}}$.

Proposition 7.1. *Let w denote the indiscrete equivalence relation on Ω , and (as in the proof of Proposition 6.2) let y_i ($i \in I$) be the $2^{\text{card}(\Omega)}$ distinct equivalence relations on Ω having exactly two equivalence classes. Then as members of $\text{Rel}(\Omega)$, these satisfy*

$$(41) \quad y_i y_i = y_i \quad (i \in I),$$

$$(42) \quad y_i y_j y_i = w \quad (i \neq j \text{ in } I).$$

Hence $L_{\text{Rel}(\Omega), 1}^{\overline{=}}$ contains a well-ordered chain of cardinality $2^{\text{card}(\Omega)}$.

Proof. (41) is immediate. The verification of (42) is routine, but the reader may find the following way of visualizing it helpful. Given y_i and y_j , picture a Venn diagram for Ω , divided by a vertical line representing the partition into the two equivalence classes of y_i and a horizontal line representing the partition into the equivalence classes of y_j . Given $(p, q) \in w = \Omega \times \Omega$, which we wish to show belongs to $y_i y_j y_i$, we may assume by adjusting our diagram that p lies in the upper left-hand box. Since y_i and y_j each have two equivalence classes, at least one of the lower boxes and at least one of the right-hand boxes are nonempty, and since $y_i \neq y_j$, the lower right-hand box is not the only nonempty box other than the upper left-hand one. It is now easy to see that wherever q may lie in our diagram, we can get from p to it by crossing the vertical line at most once and the horizontal line at most once, hence that (p, q) lies in either $y_i y_j$ or $y_j y_i$. In either case, it lies in $y_i y_j y_i$, as claimed.

However, (41) shows that $y_i y_i y_i \neq w$, and the contrast between this inequality and (42) allows us to get, as in the proofs of Corollaries 5.5 and 5.6, a well-ordered chain of cardinality $2^{\text{card}(\Omega)}$ in $L_{\text{Rel}(\Omega), 1}^{\overline{=}}$. \square

The composition operation (40) of $\text{Rel}(\Omega)$ resembles the join operation of $\text{Equiv}(\Omega)$ in being continuous in the topology on $\text{Rel}(\Omega)$ with subbasis of open sets consisting of the sets $U_{p,q}$ (defined in (34)), but not in the Hausdorff topology with subbasis of open sets $U_{p,q}$ and ${}^c U_{p,q}$. Indeed, though composition of relations is simpler to describe than the join of equivalence relations, it is still true that no specification of whether some *finite* number of pairs belong to each of two relations can tell us that (p, q) does not belong to their composite. For $\text{Rel}(\Omega)$, there is no analog of our ploy of restricting attention to the meet operation of $\text{Equiv}(\Omega)$; so let us go directly to the weaker topology. As in the case of $\text{Equiv}(\Omega)$, we can conclude that

Lemma 7.2. For J any set of cardinality $\leq \text{card}(\Omega)$, $L_{\text{Rel}(\Omega), J}^{\leq \text{const}}$ contains no chains with $> \text{card}(\Omega)$ jumps. \square

Unfortunately, I don't see how to use this result in studying the structures of monoids embeddable in the monoid $\text{Rel}(\Omega)$, for the definition of $L_{\text{Rel}(\Omega), J}^{\leq \text{const}}$ makes use of the ordering of $\text{Rel}(\Omega)$ by inclusion, and, unlike the order relation on a lattice, this has no description in terms of its algebra operations. Of course, if we throw in order as additional structure, then Lemma 7.2 yields restrictions on embeddability. Let us record this and another consequence of Lemma 7.2, without looking for examples, then return to the question of embeddability as pure monoids.

By a *partially ordered monoid* (M, \leq) , let us understand a monoid M given with a partial ordering \leq , such that for all $x, y, z \in M$,

$$(43) \quad \begin{aligned} x \leq y &\implies xz \leq yz, \\ x \leq y &\implies zx \leq zy. \end{aligned}$$

An *embedding* of partially ordered monoids $f : (M, \leq_M) \rightarrow (N, \leq_N)$ will mean a monoid embedding $f : M \rightarrow N$ such that the partial ordering on M induced by \leq_N under f is precisely \leq_M . Clearly, Lemma 7.2 gives

Corollary 7.3. If a partially ordered monoid (M, \leq) is embeddable in $(\text{Rel}(\Omega), \subseteq)$, then for any set J of cardinality $\leq \text{card}(\Omega)$, the complete lattice $L_{(M, \leq), J}^{\leq \text{const}}$ contains no chains with $> \text{card}(\Omega)$ jumps. \square

To formulate another consequence of Lemma 7.2, let us define, for M a monoid with a distinguished element z , J a set, and v any monoid word in a J -tuple of variables and arbitrary constants from M , the set $S_{v=z} = \{a = (a_j)_{j \in J} \in M^J \mid v(a) = z\} \subseteq M^J$, and let $L_{(M, z), J}^{\overline{z}}$ denote the lattice of all intersections of families of sets of this sort. If (M, z) and (M', z') are such pairs, a homomorphism $(M, z) \rightarrow (M', z')$ will mean a monoid homomorphism $M \rightarrow M'$ carrying z to z' .

Corollary 7.4. Let M be a monoid with a zero element z (an element satisfying $zx = z = xz$ for all $x \in M$), such that (M, z) is embeddable in $(\text{Rel}(\Omega), \emptyset)$, and let J be a set of cardinality $\leq \text{card}(\Omega)$. Then the complete lattice $L_{(M, z), J}^{\overline{z}}$ contains no chains with $> \text{card}(\Omega)$ jumps. \square

Proof. In $\text{Rel}(\Omega)^J$, any set of the form $S_{v=\emptyset}$ can clearly also be described as $S_{v \leq \emptyset}$, whence the assertion follows immediately from Lemma 7.2. \square

A way of getting restrictions on embeddability in $\text{Rel}(\Omega)$ without bringing in additional structure is to note, as we did in §5 for $\text{Self}(\Omega)$, that any monoid homomorphism from a group into $\text{Rel}(\Omega)$ will land in the group of invertible relations, which is again $\text{Sym}(\Omega)$. Hence our restrictions on groups embeddable in $\text{Sym}(\Omega)$ are also restrictions on the groups of invertible elements of monoids embeddable in $\text{Rel}(\Omega)$.

But in fact, we can use the group $\text{Sym}(\Omega)$ of invertible elements of $\text{Rel}(\Omega)$ in a way that brings in noninvertible elements as well. Note that the function topology on $\text{Self}(\Omega)$, and hence on $\text{Sym}(\Omega)$, is the restriction thereto both of the topology of (34), and of the weaker topology having only the sets $U_{p,q}$ as subbasis of open sets. (In $\text{Self}(\Omega)$, the set ${}^cU_{p,q}$ can be written as $\bigcup_{p' \neq p} U_{p',q}$, so the topology generated by the sets $U_{p,q}$ also contains the complementary sets.) This immediately gives case (i) of the next lemma. Case (ii), the one we shall make use of, is more surprising. (We shall not use the final parenthetical strengthening of (ii).)

Lemma 7.5. The restrictions of the monoid multiplication of $\text{Rel}(\Omega)$ to maps

$$(44) \quad \text{Sym}(\Omega) \times \text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega) \quad \text{and} \quad \text{Rel}(\Omega) \times \text{Sym}(\Omega) \rightarrow \text{Rel}(\Omega)$$

are continuous if we put the function topology on $\text{Sym}(\Omega)$, and put on $\text{Rel}(\Omega)$ the topology with subbasis of open sets consisting either of

- (i) the sets $U_{p,q}$, or
- (ii) the sets $U_{p,q}$ and ${}^cU_{p,q}$. (This remains true if we replace $\text{Sym}(\Omega)$ by $\text{Self}(\Omega)$ in the second map of (44), though not in the first.)

Proof. As noted, continuity in the topology determined by (i) follows from the continuity of the multiplication of $\text{Rel}(\Omega)$ in that topology. This also gives half of continuity in the topology specified in (ii), namely openness of the inverse images of the sets $U_{p,q}$. We shall prove the corresponding statement for ${}^cU_{p,q}$ for the second map in (44) with $\text{Sym}(\Omega)$ replaced by $\text{Self}(\Omega)$. The case where $\text{Sym}(\Omega)$ is left unchanged follows immediately, and the first statement of (44) then follows by reversing coordinates in ordered pairs.

Note that a necessary and sufficient condition for a composite xa ($x \in \text{Rel}(\Omega)$, $a \in \text{Self}(\Omega)$) to lie in ${}^cU_{p,q}$, i.e., not to contain (p,q) , is that for the unique $r \in \Omega$ such that $(r,q) \in a$, we have $(p,r) \notin x$. Hence the inverse image of ${}^cU_{p,q}$ under the above composition map is the union $\bigcup_{r \in \Omega} ({}^cU_{p,r} \times (U_{r,q} \cap \text{Self}(\Omega)))$. This set is open in $\text{Rel}(\Omega) \times \text{Self}(\Omega)$, as claimed.

To get the negative part of the final parenthetical assertion, assume without loss of generality that $\omega \subseteq \Omega$. For each $n \in \omega$, let $x_n \in \text{Rel}(\Omega)$ be the partial function which sends 0 to n and does nothing else, and let $a \in \text{Self}(\Omega)$ be the function sending all elements to 0. Then for all n , $a x_n = x_0$, hence $\lim_{n \rightarrow \infty} a x_n = x_0$. On the other hand, $\lim_{n \rightarrow \infty} x_n = \emptyset$, the empty relation, which when left multiplied by a gives $\emptyset \neq x_0$. So composition is not continuous in the Hausdorff topology of (ii). \square

From the operations (44) and the group operations of $\text{Sym}(\Omega)$, we can form words in a mixture of $\text{Sym}(\Omega)$ - and $\text{Rel}(\Omega)$ -valued elements; but any such word can involve, at most, *either* one occurrence of a $\text{Rel}(\Omega)$ -valued variable *or* one occurrence of a non- $\text{Sym}(\Omega)$ -valued constant, since the operations (44) do not allow the multiplication of two non- $\text{Sym}(\Omega)$ -valued elements. (Here by a “ $\text{Rel}(\Omega)$ -valued variable” I mean a variable that is allowed to range over all of $\text{Rel}(\Omega)$, taking on both invertible and noninvertible values. Constants, on the other hand, have specific values rather than ranges, so for these the relevant concept is that of a non- $\text{Sym}(\Omega)$ -valued constant, i.e., a noninvertible element of $\text{Rel}(\Omega)$.) By Lemma 7.5, if we take a family of such words, possibly involving many $\text{Rel}(\Omega)$ -valued variables and non- $\text{Sym}(\Omega)$ -valued constants altogether (though with at most one of these per word), and a set of equations in these words, then its solution set is closed in the Hausdorff topology defined by the subbasis (ii). Hence we can apply Lemma 5.1 and get restrictions on the lattice of such solution sets, which imply the same restrictions on the corresponding lattice obtained from any monoid embeddable in $\text{Rel}(\Omega)$.

It is not clear to me whether so allowing more than one $\text{Rel}(\Omega)$ -valued variable or constant in our system of equations actually contributes to the generality of this result. Note that if we have a system of equations in several such variables and constants, then any equation involving different $\text{Rel}(\Omega)$ -valued variables on the two sides, say x on the left and y on the right, will allow us to solve for y in terms of x and the $\text{Sym}(\Omega)$ -valued variables and constants (since the latter can all be inverted and brought to the left side of the equation); and we can then substitute the resulting expression for all occurrences of y in the remaining equations, and so eliminate y from the system. On the other hand, our interest is not in the solution set of a *single* system of equations but in the relation between solution sets of many such systems, and a variable that can be eliminated from one of these will not in general be eliminable from all of them. So if the consequences of our conditions on lattices of solution sets can indeed be reduced to the case where there is only one $\text{Rel}(\Omega)$ -valued variable, the argument by which this reduction is done may be nontrivial.

Leaving it to others to determine whether such a reduction is possible, I will, for simplicity, record here only the statements for systems with at most one $\text{Rel}(\Omega)$ -valued-variable.

Theorem 7.6. *Let J be any set of cardinality $\leq \text{card}(\Omega)$.*

Let L denote the lattice of solution sets in $\text{Sym}(\Omega)^J \times \text{Rel}(\Omega)$ of systems of equations in a J -tuple of $\text{Sym}(\Omega)$ -valued variables, and a single $\text{Rel}(\Omega)$ -valued variable which appears at most once on each side of any equation, together with arbitrary $\text{Sym}(\Omega)$ -valued constants.

Let L' similarly denote the lattice of solution sets in $\text{Sym}(\Omega)^J$ of systems of equations having on each side a word in a J -tuple of $\text{Sym}(\Omega)$ -valued variables, arbitrary $\text{Sym}(\Omega)$ -valued constants, and at most one occurrence of a $\text{Rel}(\Omega)$ -valued constant.

Then neither of these lattices contains a chain with $> \text{card}(\Omega)$ jumps. Hence for any monoid M embeddable in $\text{Rel}(\Omega)$, the corresponding restrictions hold, with the group $U(M)$ of invertible elements of M taking the place of $\text{Sym}(\Omega)$. \square

Simple examples of equations of the sort arising above are stabilizer relations $gx = x$ and $xg = x$ ($x \in M$, $g \in U(M)$), and more generally, $g x g' = x$ ($x \in M$, $g, g' \in U(M)$). Here is an example based on relations $gx = x$.

Recall that a *left zero element* in a monoid M means an element z satisfying $zx = z$ for all $x \in M$.

Corollary 7.7. *Let I be a set, let G be the restricted direct product of an I -tuple of copies of the group Z_2 , with generators g_i ($i \in I$) (i.e., the additive group of the vector space with basis $\{g_i \mid i \in I\}$ over the field of two elements), and let M be the monoid whose group of invertible elements is this group G , and whose other elements, denoted z_i, z'_i ($i \in I$), are left zero elements whose behavior under the left action of G is described by*

$$(45) \quad \text{Left multiplication by } g_i \text{ interchanges } z_i \text{ and } z'_i, \text{ and fixes all } z_j \text{ and } z'_j \text{ with } j \neq i.$$

Then M is embeddable in $\text{Rel}(\Omega)$ if and only if $\text{card}(I) \leq \text{card}(\Omega)$.

Proof. It is straightforward to verify that the above operations define a monoid, which has cardinality $\leq \max(\text{card}(I), \aleph_0)$, and so is embeddable in $\text{Self}(\Omega) \subseteq \text{Rel}(\Omega)$ if $\text{card}(I) \leq \text{card}(\Omega)$. To prove the “only if” part of the conclusion, assume I is a cardinal $\kappa > \text{card}(\Omega)$. For each $\alpha \in \kappa$, let $S_\alpha = \{x \in M \mid (\forall \beta > \alpha) g_\beta x = x\}$. Note that $z_\beta \in S_\alpha$ if and only if $\beta \leq \alpha$, so the S_α form a well-ordered chain of cardinality $\kappa > \text{card}(\Omega)$. By the “ L ” case of Theorem 7.6 (with $J = 0$), this precludes embeddability in $\text{Rel}(\Omega)$. \square

In proving the above corollary we could, alternatively, have used the z_α as monoid-valued constants, and let the g in the relations $gz_\alpha = z_\alpha$ be an invertible-element-valued variable, getting a system to which the “ L ” case of Theorem 7.6 with $J = 1$ applied.

Question 7.8. *Can one give stronger necessary conditions for embeddability of a monoid M of cardinality $\leq 2^{\text{card}(\Omega)}$ in $\text{Rel}(\Omega)$ than those of Theorem 7.6?*

In particular, suppose we define the left, right, and 2-sided stabilizers of an element x of a monoid M as $\{y \in M \mid yx = x\}$, $\{y \in M \mid xy = x\}$, and $\{(y, y') \in M^2 \mid yxy' = x\}$. Does $\text{Rel}(\Omega)$ have any chains of intersections of such stabilizers with $> \text{card}(\Omega)$ jumps? What if we restrict y and y' here to left invertible or to right invertible elements?

Are there any monoids of cardinality $\leq 2^{\text{card}(\Omega)}$ having no invertible elements other than 1 (or having $\leq \text{card}(\Omega)$ invertible elements) which are not embeddable in $\text{Rel}(\Omega)$?

For $\text{card}(I) > \text{card}(\Omega)$, is the monoid of Corollary 7.7 ever embeddable as a semigroup in $\text{Rel}(\Omega)$?

One kind of noninvertible elements one might look at in approaching these questions are the *idempotents*, since for e idempotent, the solution sets of $xe = x$ and $ex = x$ are particularly natural objects. Idempotents in $\text{Rel}(\Omega)$ come in more forms than one might expect. Obvious examples are equivalence relations, subsets of the identity relation, graphs of retractions of Ω onto subsets, and the opposites of such graphs; these four constructions can also be mixed in fairly natural ways. For less obvious examples, note that (i) for any partial ordering, \leq , on Ω with no jumps (e.g., the ordinary ordering on the set of rational numbers), the graph of the relation “ $<$ ” is idempotent, though it has trivial intersection with the identity relation; and (ii) for any nondisjoint subsets X and Y of Ω , the set $X \times Y \subseteq \Omega \times \Omega$ is an idempotent relation.

We have not written down the positive embeddability result analogous to Theorem 3.1, i.e., the embeddability in $\text{Rel}(\Omega)$ of monoids $F(2^{\text{card}(\Omega)})$ for appropriate functors F , since this follows from Theorem 3.1 and the inclusion $\text{Self}(\Omega) \subseteq \text{Rel}(\Omega)$. It would, of course, be of interest if one could get *stronger* results of this sort for $\text{Rel}(\Omega)$ than for $\text{Self}(\Omega)$. I also don’t know the answer to

Question 7.9. *Is $\text{Rel}(\Omega) \amalg \text{Rel}(\Omega)$ embeddable as a monoid in $\text{Rel}(\Omega)$?*

(If so, then by the usual argument, $\text{Rel}(\Omega)$ in fact contains a coproduct of $2^{\text{card}(\Omega)}$ copies of itself.)

Here are some partial positive results on that question.

Let $1_\Omega \in \text{Rel}(\Omega)$ denote the identity element, that is, the diagonal subset of $\Omega \times \Omega$. Let us say that elements $g, h \in \text{Rel}(\Omega)$ “differ off the diagonal” if $g - 1_\Omega \neq h - 1_\Omega$ (where “ $-$ ” denotes set-theoretic difference); and for a subset $X \subseteq \text{Rel}(\Omega)$, let us say “the members of X are distinguishable off the diagonal” if every pair of distinct elements of X differs off the diagonal.

Lemma 7.10. *There exists a monoid homomorphism $f : \text{Rel}(\Omega) \amalg \text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega)$ with the property that whenever M and N are submonoids of $\text{Rel}(\Omega)$, each of which has the property that its members are distinguishable off the diagonal, then the restriction of f to a homomorphism $M \amalg N \rightarrow \text{Rel}(\Omega)$ is an embedding.*

Proof. Let us understand a “relational action” of a monoid M on a set X to mean a homomorphism $M \rightarrow \text{Rel}(X)$. Paralleling the proof of Lemma 4.1(ii), we shall construct a relational action of $\text{Rel}(\Omega) \amalg \text{Rel}(\Omega)$ on the disjoint union of a family of copies of $\Omega \times \omega$, where the family is again indexed by the group $\text{Sym}_{\text{fin}}(\Omega \times \omega)$, such that the restriction of this action to any submonoid $M \amalg N$ as in the statement of the lemma is faithful.

Let the “natural relational action” of $\text{Rel}(\Omega)$ on $\Omega \times \omega$ be defined to take each $g \in \text{Rel}(\Omega)$ to the relation on $\Omega \times \omega$ consisting of all pairs $((q, k), (p, k))$ with $(q, p) \in g$ and $k \in \omega$. Let α, β be the two coprojections $\text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega) \amalg \text{Rel}(\Omega)$. For each $g \in \text{Rel}(\Omega)$, let us send $\alpha(g)$ to the natural relational action of g on each copy of $\Omega \times \omega$, while letting $\beta(g)$ act on the copy of $\Omega \times \omega$ indexed by each $t \in \text{Sym}_{\text{fin}}(\Omega \times \omega)$ via the conjugated relation tgt^{-1} . To complete our proof, we need to show that for M and N as in the statement of the lemma, if (13) and (14) are distinct elements of $M \amalg N \subseteq \text{Rel}(\Omega) \amalg \text{Rel}(\Omega)$, then there exists $t \in \text{Sym}_{\text{fin}}(\Omega \times \omega)$ such that the two relations (15) on $\Omega \times \omega$ are distinct.

As in our previous argument, we assume that the length of (13), which we will call n , is at least the length of (14). Moreover, we assume that if these lengths are equal, and if α s and β s occur in the same positions in both expressions, then at the first position k from the right where (13) and (14) differ, g_k contains some *nondiagonal* ordered pair which does not lie in h_k . Since g_k and h_k differ off the diagonal, we can achieve this by interchanging (13) and (14) if necessary. We now choose, for each $k \in \{1, \dots, n\}$, a nondiagonal pair $(q_k, p_k) \in g_k$ (which must exist, because g_k is not the identity, hence is distinguishable from the identity off the diagonal), using, when possible, a pair not also contained in h_k ; and we define t as in (18), except that wherever (18) shows an element $g_k(p_k)$, we now use q_k . The same considerations as in the proof of Lemma 4.1 show that the first product in (17), but not the second, contains the pair $((q_n, n+1), (p_0, 0))$; so these two products are distinct relations on $\Omega \times \omega$, completing the proof. \square

Here are some consequences.

Corollary 7.11. (i) $\text{Self}(\Omega) \amalg \text{Self}(\Omega)^{\text{op}}$ is embeddable in $\text{Rel}(\Omega)$.

(ii) Let $\text{Rel}(\Omega)_{\geq 1} \subseteq \text{Rel}(\Omega)$ denote the submonoid of relations that contain the diagonal (the reflexive relations). Then $\text{Rel}(\Omega)_{\geq 1} \amalg \text{Rel}(\Omega)_{\geq 1}$ is embeddable in $\text{Rel}(\Omega)_{\geq 1}$. Hence the coproduct of $2^{\text{card}(\Omega)}$ copies of $\text{Rel}(\Omega)_{\geq 1}$ is embeddable in $\text{Rel}(\Omega)_{\geq 1}$.

(iii) Let $\text{Rel}^{\text{semi}}(\Omega)$ denote the underlying semigroup of $\text{Rel}(\Omega)$. Then the semigroup coproduct $\text{Rel}^{\text{semi}}(\Omega) \amalg \text{Rel}^{\text{semi}}(\Omega)$ is embeddable in $\text{Rel}^{\text{semi}}(\Omega)$. Hence the semigroup coproduct of $2^{\text{card}(\Omega)}$ copies of $\text{Rel}^{\text{semi}}(\Omega)$ is embeddable in $\text{Rel}^{\text{semi}}(\Omega)$.

Proof. (i) is immediate from the lemma, since both $\text{Self}(\Omega)$ and the natural copy of $\text{Self}(\Omega)^{\text{op}}$ in $\text{Rel}(\Omega)$ (gotten by taking the opposite relations to all members of $\text{Self}(\Omega)$) have the property that their members are distinguishable off the diagonal.

In (ii), Lemma 7.10 immediately gives embeddability of $\text{Rel}(\Omega)_{\geq 1} \amalg \text{Rel}(\Omega)_{\geq 1}$ in $\text{Rel}(\Omega)$. Moreover, we see that the construction of that lemma takes reflexive relations to reflexive relations, so the embedding lands in $\text{Rel}(\Omega)_{\geq 1}$. The method of proof of Theorem 4.2 now allows one to work one’s way up to the coproduct of $2^{\text{card}(\Omega)}$ copies.

To prove (iii) let us begin by constructing an embedding of $\text{Rel}^{\text{semi}}(\Omega)$ in the semigroup of relations on a set of the same cardinality as Ω , by relations that are distinguishable from each other and from the identity relation off the diagonal.

We first have to declaw the empty relation; so let z be an element not in Ω , and let us embed $\text{Rel}(\Omega)$ in $\text{Rel}(\Omega \cup \{z\})$ by sending each relation g to $g \cup \{(z, z)\}$; this is an embedding of monoids whose image consists of nonempty relations.

We now map $\text{Rel}^{\text{semi}}(\Omega \cup \{z\})$ into $\text{Rel}^{\text{semi}}((\Omega \cup \{z\}) \times 2)$ by

$$(46) \quad g \mapsto \{((q, i), (p, j)) \mid (q, p) \in g; i, j \in 2\}.$$

This construction is easily seen to respect composition, and to take distinct nonempty relations to relations that differ off the diagonal both from each other and from $1_{\Omega \cup \{z\}}$. The image of the composite map $\text{Rel}^{\text{semi}}(\Omega) \rightarrow \text{Rel}^{\text{semi}}(\Omega \cup \{z\}) \rightarrow \text{Rel}^{\text{semi}}((\Omega \cup \{z\}) \times 2)$ is in particular a subsemigroup $S \subseteq \text{Rel}^{\text{semi}}((\Omega \cup \{z\}) \times 2)$ isomorphic to $\text{Rel}^{\text{semi}}(\Omega)$. Applying Lemma 7.10 with $S \cup \{1\}$ in the role of both M and N , we get an embedding of monoids $(S \cup \{1\}) \amalg (S \cup \{1\}) \rightarrow \text{Rel}(\Omega)$, which, restricted to the

subsemigroup generated by the two copies of S , gives the desired embedding of semigroups. As before, the method of §4 allows us to push this up to an embedding of a $2^{\text{card}(\Omega)}$ -fold coproduct of copies of $\text{Rel}^{\text{semi}}(\Omega)$ in $\text{Rel}^{\text{semi}}(\Omega)$. \square

It would be nice if we could carry the idea of part (iii) of the above corollary further. If we could find a *monoid* embedding $\varphi : \text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega')$ with $\text{card}(\Omega') = \text{card}(\Omega)$, such that images of distinct elements were distinguishable off the diagonal, then an application of Lemma 7.10 would give us our desired embedding $\text{Rel}(\Omega) \amalg \text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega)$. But when we attempt to construct such a φ , we run into difficulty trying to simultaneously

- (a) make it carry the identity relation 1_Ω to $1_{\Omega'}$,
- (b) handle relations which are properly contained in 1_Ω , and
- (c) handle relations which are both infinitely-many-to-one and one-to-infinitely-many.

Part (iii) of the above corollary showed that we could get an embedding if we dropped the requirement (a). The following result (strengthening part (ii) of that corollary) shows the same if we instead drop (b), and the result after that will do the same for (c).

Lemma 7.12. *There exists a monoid homomorphism $f' : \text{Rel}(\Omega) \amalg \text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega)$ with the property that whenever M and N are submonoids of $\text{Rel}(\Omega)$, neither of which contains a proper subrelation of 1_Ω , then the restriction of f' to a homomorphism $M \amalg N \rightarrow \text{Rel}(\Omega)$ is an embedding.*

Proof. Let $\varphi : \text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega^2)$ take each $g \in \text{Rel}(\Omega)$ to $\{(p, p'), (q, q') \mid (p, q), (p', q') \in g\}$. It is straightforward to verify that this is a monoid homomorphism. We shall show that if $g, h \in \text{Rel}(\Omega)$ are distinct elements which are not both subrelations of 1_Ω , then $\varphi(g)$ and $\varphi(h)$ are distinguishable off the diagonal. This, together with Lemma 7.10, yields the desired result.

First, suppose g and h are themselves distinguishable off the diagonal. Then without loss of generality we may assume that (p, q) is contained in g but not h for some $p \neq q$, and we see that $((p, p), (q, q))$ belongs to $\varphi(g)$ but not $\varphi(h)$. On the other hand, if g and h are not distinguishable off the diagonal, then since they are not both subrelations of 1_Ω , there must be some (p, q) with $p \neq q$ that is contained in both of them. Also, since $g \neq h$, some (r, r) ($r \in \Omega$) will belong to one of them but not the other; say to g but not h . Then we see that $((p, r), (q, r))$ is a nondiagonal element belonging to $\varphi(g)$ but not $\varphi(h)$. \square

Finally, let us see what we can get if we sacrifice (c).

Given $g \in \text{Rel}(\Omega)$ and $X \subseteq \Omega$, let us define the “image” set

$$(47) \quad gX = \{q \in \Omega \mid (\exists p \in X) (q, p) \in g\},$$

and let

$$(48) \quad \text{Rel}_{\text{fin}\leftarrow 1}(\Omega) = \{g \in \text{Rel}(\Omega) \mid (\forall p \in \Omega) g\{p\} \text{ is finite}\}.$$

The arrow points to the left to show that we are defining this set in terms of the left action (47). We can clearly also describe (48) as the set of g such that for every finite $X \subseteq \Omega$, the set gX is again finite. This shows $\text{Rel}_{\text{fin}\leftarrow 1}(\Omega)$ to be a submonoid of $\text{Rel}(\Omega)$, and it clearly acts faithfully – by functions, not relations – on $\mathbf{P}_{\text{fin}}(\Omega)$. This gives us an embedding $\text{Rel}_{\text{fin}\leftarrow 1}(\Omega) \rightarrow \text{Self}(\mathbf{P}_{\text{fin}}(\Omega)) \cong \text{Self}(\Omega)$. Of course, we also have $\text{Self}(\Omega) \subseteq \text{Rel}_{\text{fin}\leftarrow 1}(\Omega)$; so embeddability of a monoid in $\text{Rel}_{\text{fin}\leftarrow 1}(\Omega)$ and in $\text{Self}(\Omega)$ are equivalent. This is the first assertion of the next lemma, and by the results of §4 it implies the second.

Lemma 7.13. *$\text{Rel}_{\text{fin}\leftarrow 1}(\Omega)$ and $\text{Self}(\Omega)$ are each embeddable in the other.*

Hence $\text{Rel}_{\text{fin}\leftarrow 1}(\Omega)$ contains a coproduct of $2^{\text{card}(\Omega)}$ copies of itself as a monoid. \square

Let me record here a curious construction which I thought, at one point, would give a more elegant proof of our semigroup-embedding result, Corollary 7.11(iii). This did not quite work; but perhaps it is nonetheless of interest.

Consider the map $\theta : \text{Rel}^{\text{semi}}(\Omega) \rightarrow \text{Rel}^{\text{semi}}(\mathbf{P}_{\text{fin}}(\Omega))$ defined by

$$(49) \quad \theta(g) = \{(t, s) \in (\mathbf{P}_{\text{fin}}(\Omega))^2 \mid t \subseteq gs\}.$$

It is easy to verify that θ is a semigroup homomorphism. Moreover, every $\theta(g)$ contains all pairs (\emptyset, t) with $t \in \mathbf{P}_{\text{fin}}(\Omega)$, and hence differs from $1_{\mathbf{P}_{\text{fin}}(\Omega)}$ off the diagonal. For most pairs of distinct $g, h \in \text{Rel}(\Omega)$, one finds that $\theta(g)$ and $\theta(h)$ differ in the nondiagonal pairs of one of the forms $(\{p\}, \{q\})$, $(\{p\}, \{p, q\})$, or $(\{p, q\}, \{p\})$ that they contain. There are exceptions, however; for instance, if $p \in \Omega$ and we take $g = \{p\} \times \Omega$, $h = \{p\} \times (\Omega - \{p\})$; then $\theta(g)$ and $\theta(h)$ differ only with regard to the diagonal pair $(\{p\}, \{p\})$. Hence this construction is not itself a substitute for the one used in proving Corollary 7.11(iii). We can cure the problem using the “ $\text{Rel}(\Omega) \rightarrow \text{Rel}(\Omega \cup \{z\})$ ” trick that was used (to cure a different problem) in the existing proof of that statement (and after this is done, the images of any two relations in fact differ in the pairs $(\{p\}, \{q, z\})$ that they contain); but I wouldn’t call the resulting proof more elegant than the one we used.

We end this section with a different sort of self-embeddability question for $\text{Rel}(\Omega)$.

Question 7.14. *If $e \in \text{Rel}(\Omega)$ is an idempotent, and we regard $eRe = \{x \in \text{Rel}(\Omega) \mid exe = x\}$ as a monoid with identity element e , is this monoid always embeddable in $\text{Rel}(\Omega)$? (Cf. remarks on idempotents following Question 7.8.)*

8 Restrictions on K -algebras embeddable in $\text{End}_K(V)$.

Recall that V denotes a vector space with basis Ω over a field K . Since the endomorphism algebra $\text{End}_K(V)$ is a K -linear analog of the monoid $\text{Self}(\Omega)$, we would hope to get restrictions on associative unital K -algebras embeddable in $\text{End}_K(V)$ parallel to our restrictions on monoids embeddable in $\text{Self}(\Omega)$. We can do this – except that, where we would like to bound the number of jumps in a chain of solution sets by $\text{card}(\Omega) = \dim_K(V)$, I only know how to bound it by $\text{card}(V) = \max(\text{card}(\Omega), \text{card}(K))$. The following theorem is proved exactly like Theorem 5.4, using the fact that addition and composition of members of $\text{End}_K(V)$, and multiplication of these maps by members of K , are continuous in the function topology on $\text{End}_K(V)$, regarded as a subset of $\text{Self}(V)$.

Theorem 8.1. *Let J be any set of cardinality $\leq \text{card}(V)$. Then $L_{\text{End}_K(V), J}^{\bar{=}}$ contains no chains with $> \text{card}(V)$ jumps. Hence for any K -algebra A embeddable in $\text{End}_K(V)$, the lattice $L_{A, J}^{\bar{=}}$ contains no such chains; in particular, it contains no well-ordered or reverse-well-ordered chains of cardinality $> \text{card}(V)$. \square*

In the hope of reducing the bound $\text{card}(V)$ to $\text{card}(\Omega)$, we might try replacing the function topology on $\text{End}_K(V)$ by some topology with a smaller basis of open sets; say one that defines its subbasic open sets not by considering the values of elements of $\text{End}_K(V)$ at arbitrary elements of V , but only at the elements of our basis Ω . Unfortunately, each of these images still has $\text{card}(V)$ possible values. However, a linear restriction on these images corresponds to a proper subspace of the space V of possible values, suggesting that the vector space dimension should still bound lengths of chains. On the other hand, the conditions on the coordinates of our elements induced by ring-theoretic relations are not necessarily linear. Perhaps one should seek bounds on the lengths of chains of solution sets by methods of algebraic geometry. Or perhaps one can get stronger results for relations that are multilinear, such as centralizer and annihilator relations, than for general relations.

In another direction, if K is a topological field such as the real or complex numbers, perhaps we could use the topology of that field instead of the discrete topology on our coordinates, and replace $\text{card}(V)$ in the above theorem by $\max(\text{card}(\Omega), \kappa)$ where κ is the least cardinality of a basis for the topology of K . We record these problems as

Question 8.2. *In Theorem 8.1, can the bound $\text{card}(V)$ be lowered to $\text{card}(\Omega)$? If not in general, what if we restrict attention to the lattice determined by K -linear or K -affine relations? Can one at least improve the bound $\text{card}(V)$ if K admits a structure of topological field with a basis of $< 2^{\text{card}(\Omega)}$ open sets?*

For one sort of system of relations, we can indeed get the expected bound.

Lemma 8.3. *Any set of nonzero pairwise orthogonal idempotent elements of $\text{End}_K(V)$ has cardinality $\leq \text{card}(\Omega)$. Hence the same is true in any K -algebra embeddable in $\text{End}_K(V)$.*

In particular, $\text{End}_K(V)$ does not contain a direct product of $> \text{card}(\Omega)$ nontrivial K -algebras.

Proof. Given an infinite family S of nonzero pairwise orthogonal idempotents, the images of these as endomaps of V form a set of subspaces of V whose sum is direct. Hence the dimension of that sum is at least $\text{card}(S)$; so $\text{card}(S) \leq \dim(V) = \Omega$.

To see the final sentence, note that in a direct product of K -algebras $\prod_{i \in I} A_i$, if e_i denotes the element with i -component 1 and all other components 0, then these are pairwise orthogonal idempotents. \square

On the other hand, some ways in which the behavior of $\text{End}_K(V)$ for K large is indeed different from that of $\text{Self}(\Omega)$ will be noted at the end of §10.

Wehrung, paralleling his result mentioned earlier that $\text{Self}(\Omega)$ cannot be embedded in $\text{Self}(\Omega)^{\text{op}}$, also shows in [19] that $\text{End}_K(V)$ cannot be embedded in $\text{End}_K(V)^{\text{op}}$. In fact, he shows that $\text{Self}(\Omega)$ cannot be embedded in the underlying multiplicative semigroup of $\text{End}_K(V)^{\text{op}}$, yielding both results!

9 Other directions for generalization.

In this note, we have concentrated on questions of embeddability in a small number of objects: $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$, $\text{End}_K(V)$, $\text{Equiv}(\Omega)$ and $\text{Rel}(\Omega)$, with brief observations on a few more: $\text{Sym}(\Omega)/\text{Sym}_{<}(\Omega)$, $(\text{Rel}(\Omega), \subseteq)$, $(\text{Rel}(\Omega), \emptyset)$, $\text{Rel}^{\text{semi}}(\Omega)$, and objects which we could write $\text{Equiv}^0(\Omega)$, $\text{Equiv}^1(\Omega)$ and $\text{Equiv}^{0,1}(\Omega)$, i.e., $\text{Equiv}(\Omega)$ regarded as a member of the variety of lattices with least and/or greatest element. Similar questions for other objects of the same flavor, for example the groups of automorphisms of various structures considered in [9], [10] and [11], the lattices of congruences of these objects, etc., would also be of interest.

Above, I quoted results from McKenzie [16] only in the forms in which they were relevant to the questions considered here; but that paper in fact considers the group $\text{Sym}(\Omega, \beta)$ of all permutations of Ω moving $< \beta$ elements, for a fixed cardinal β , and many of the restrictions proved there are in terms of β , rather than $\text{card}(\Omega)$. This, too, represents a direction in which the present results might be generalized.

The variant of the technique of §3 illustrated in Theorem 3.4, based on considering functors on the category **T.ord** of totally ordered sets, rather than on **Set**, also admits wide generalization. Note that any algebra-valued functor F on **T.ord** can be extended to the category **Poset** of *partially* ordered sets, though usually not uniquely. (For instance, by taking each partially ordered set P to the colimit of the algebras $F(C)$ as C ranges over all chains in P ; or to the limit of the algebras $F(C)$ as C ranges over all quotient-sets of P given with total orderings that make the quotient-map isotone.) Hence in stating Theorem 3.4 and seeking generalizations, we could just as well let F be a functor on **Poset**. Still more generally, why not let it be defined on arbitrary preordered sets? Or on sets with an arbitrary binary relation? Or several binary relations? And for such generalizations, what would be the “best” analog of (9)? Not knowing what the useful generalizations would be, I have merely given a sample result.

The referee notes that de Bruijn’s groups with presentation (22)-(24) are the values of a functor on the category whose objects are sets I , and whose morphisms are *one-to-one* set-maps; this also applies to several other examples for which we proved nonembeddability results. Hence that domain category is *not* good from the above point of view. A repeated use that we made of non-one-to-one maps in the arguments of §3 was in getting left inverses to one-to-one maps with nonempty domains. The fact that such inverses exist has the consequence (also pointed out by the referee) that our key condition (9) is equivalent to the condition that F respect direct limits.

A question I have not thought much about, but to which some of the techniques we have introduced above should be applicable, is

Question 9.1. *What can be said about groups A embeddable in $(\prod_{n \in \omega} \text{Sym}(n))^\Omega$, monoids A embeddable in $(\prod_{n \in \omega} \text{Self}(n))^\Omega$, K -algebras A embeddable in $(\prod_{n \in \omega} \text{End}_K(K^n))^\Omega$, and lattices A embeddable in $(\prod_{n \in \omega} \text{Equiv}(n))^\Omega$?*

Clearly, such an A must be residually finite(-dimensional), and embeddable in $\text{Sym}(\Omega)$, $\text{Self}(\Omega)$, $\text{End}_K(V)$, respectively $\text{Equiv}(\Omega)$. Are these conditions on an algebra A sufficient for it to be embeddable in the indicated object?

I record next a question which it would have been natural to give in [4], [5], or [6], but which occurred to me too late to include in those papers. (It is an instance of the direction for investigation suggested in the

second sentence of [6, §10].) There are obvious variants and strengthenings, but for concreteness, I pose the question here for the object that has been most studied.

Question 9.2. *Suppose G and H are subgroups of $\text{Sym}(\omega)$, which together generate that group. Must $\text{Sym}(\omega)$ be finitely generated over one of these subgroups?*

We end this note with a few items that we have postponed.

10 Appendix: $\text{End}_K(V) \amalg \text{End}_K(V)$ can be embedded in $\text{End}_K(V)$.

We shall now give the postponed proof of the above embeddability statement, Lemma 4.1(iii).

Suppose S and T are two nonzero K -algebras (as always, associative and unital), and we form their coproduct $S \amalg T$ in the variety of such algebras, calling the coprojection maps $\alpha : S \rightarrow S \amalg T$ and $\beta : T \rightarrow S \amalg T$. Recall [2, Corollary 8.2] that if B_S, B_T are K -vector-space bases for S and T , containing 1_S and 1_T respectively, then a K -vector-space basis for $S \amalg T$ is given by the set of finite products

$$(50) \quad \dots \alpha(b_i) \beta(b_{i-1}) \alpha(b_{i-2}) \beta(b_{i-3}) \dots$$

where, as in (13), those b_j that are arguments of α (i.e., in (50), those with subscript j having the same parity as i) are taken from $B_S - \{1_S\}$, those that are arguments of β are taken from $B_T - \{1_T\}$, and the empty product is understood to give the identity element $1 = \alpha(1_S) = \beta(1_T)$. As in (13), I have not shown the first and last terms, since each may be either an α -term or a β -term.

Letting $S = T = \text{End}_K(V)$, and letting B be a K -basis for this algebra containing 1, a basis for $\text{End}_K(V) \amalg \text{End}_K(V)$ is thus given by the words (50) with all b_j taken from $B - \{1\}$. To show $\text{End}_K(V) \amalg \text{End}_K(V)$ embeddable in $\text{End}_K(V)$, it will suffice to find a representation of $\text{End}_K(V) \amalg \text{End}_K(V)$ by K -linear endomorphisms on a vector space of the same dimension as V , such that the images of the elements (50) are linearly independent. For this, in turn, it will suffice to find a family of $\leq \text{card}(\Omega)$ representations of $\text{End}_K(V) \amalg \text{End}_K(V)$ on such spaces, such that every nontrivial linear relation among elements (50) fails to hold in at least one of these representations, since then all such relations will fail in the direct sum of the representations.

The representations we use will each be on $\bigoplus_{\omega} V$, a direct sum of countably many copies of V , with basis $\Omega \times \omega$. For each $k \in \Omega$ we shall call the k th copy of V , i.e., the span of $\Omega \times \{k\}$, the “ k th level” of $\bigoplus_{\omega} V$, and, extending the notation we are using on its basis, we shall denote the element at the k th level corresponding to any $v \in V$ by (v, k) . We define the *natural action* of each $f \in \text{End}_K(V)$ on the k th level of $\bigoplus_{\omega} V$ to be given by $f(v, k) = (f(v), k)$, i.e., to mimic its action on V . The natural action on $\bigoplus_{\omega} V$ will mean the direct sum of these actions.

We now define our $\text{card}(\Omega)$ actions of $\text{End}_K(V) \amalg \text{End}_K(V)$ on $\bigoplus_{\omega} V$. They will be indexed by the set of those vector space automorphisms t of $\bigoplus_{\omega} V$ which have order 2, fix all but finitely many members of the basis $\Omega \times \omega$, and take the remaining members of that basis to linear combinations of members of that basis with integer coefficients. The last two conditions insure there are only $\text{card}(\Omega)$ such t . For each such choice of t , we map $\text{End}_K(V) \amalg \text{End}_K(V)$ to $\text{End}_K(\bigoplus_{\omega} V)$ by sending each element $\alpha(f)$ ($f \in \text{End}_K(V)$) to the natural action of f on $\bigoplus_{\omega} V$, which we denote by the same symbol f , while we send $\beta(f)$ to $t f t^{-1}$.

For the remainder of the proof, let us fix a nonzero $x \in \text{End}_K(V) \amalg \text{End}_K(V)$. If $x \in K$, then clearly x has nonzero action under all of our representations; so let us assume $x \notin K$, and show how to construct a t such that the action of x on $\bigoplus_{\omega} V$ under the representation indexed by t is nonzero.

To do this, let us fix an arbitrary element $r \in \Omega$, and note that $\text{End}_K(V)$ will be the direct sum of the 1-dimensional subspace spanned by the identity map, and the subspace $\text{End}_K(V)_0$ of consisting of those maps that take r to a linear combination of elements of $\Omega - \{r\}$. Choosing, temporarily, an arbitrary basis B' of $\text{End}_K(V)$ consisting of 1 and elements of $\text{End}_K(V)_0$, let us express x as a linear combination of words of the form (50) with the b_j taken from $B' - \{1\}$. Let n be the maximum of the lengths of the words occurring with nonzero coefficient in this expression. Let $S \subseteq B'$ be the set consisting of 1 and all those elements of $B' - \{1\}$ that occur (as arguments of α or β) in the expression for x . From the fact that the span of S is a finite dimensional subspace of $\text{End}_K(V)$, it is not hard to see that there will exist a finite subset $\Sigma \subseteq \Omega$ containing the element r , such that, if we write $P_{\Sigma} \in \text{End}_K(V)$ for the element that fixes all members of Σ and annihilates all members of $\Omega - \Sigma$, then the linear operator on $\text{End}_K(V)$ given by

$$(51) \quad f \mapsto P_\Sigma f P_\Sigma$$

is one-to-one on the span of S . Let us fix such a set $\Sigma \subseteq \Omega$.

We can now describe the basis B of $\text{End}_K(V)$ in terms of which we will work for the rest of the proof. For all $p, q \in \Omega$, let $E(q, p) \in \text{End}_K(V)$ be the linear map that takes p to q , and all other members of Ω to 0. Since the typical member of $\text{End}_K(V)$ has infinite-dimensional range, the elements $E(q, p)$ do not span $\text{End}_K(V)$; but their linear combinations do give all possible behaviors on our finite set Σ , which is what we will need. Let us choose B to consist of the identity operator 1, all the operators $E(q, p)$ with $p \in \Sigma$ (and q unrestricted) except for $E(r, r)$, and the members of any basis of the space of those endomorphisms that annihilate Σ .

To see that an arbitrary $f \in \text{End}_K(V)$ may be represented by a linear expression in members of B , first set the coefficient of 1 in this expression to be the coefficient of r in $f(r)$. Subtracting from f that multiple of 1 gives a member of $\text{End}_K(V)_0$, whose behavior on the elements of Σ can be represented by a finite linear combination of the operators $E(q, p)$ with $p \in \Sigma$ and $(p, q) \neq (r, r)$. Subtracting this off, we are left with an operator annihilating Σ , which can be uniquely represented using the elements introduced in the last part of our definition of B . Clearly, this expression for f is unique.

We now take our earlier expression for x as a linear combination of words (50) with all $b_j \in B' - \{1\}$, and substitute for the b_j their expressions as linear combinations of elements of $B - \{1\}$, getting an expression for x , again as a combination of words (50), with the b_j now in $B - \{1\}$. Clearly, these words still have length $\leq n$. I claim, moreover, that by our choice of Σ , the expression contains, with nonzero coefficient, at least one product (50) of length n in which all b_j have the form $E(q, p)$ with p and q both in Σ (and by definition of B , with $(p, q) \neq (r, r)$).

To see this, note that a consequence of our normal form for coproducts of associative K -algebras is that $\text{End}_K(V) \amalg \text{End}_K(V)$ can be identified with a direct sum of iterated tensor products $\text{End}_K(V)_0 \otimes_K \cdots \otimes_K \text{End}_K(V)_0$, where the 0-fold tensor product, i.e., K , occurs once, and each higher tensor product occurs twice, corresponding to the two ways of labeling the tensor factors alternately with α and β . This identification maps any element of one of these direct summands (e.g., an element $v_1 \otimes v_2 \otimes v_3$ in the summand labeled with α, β, α) to the corresponding sum of products of elements of $\alpha(\text{End}_K(V)_0)$ and $\beta(\text{End}_K(V)_0)$ (e.g., $\alpha(v_1)\beta(v_2)\alpha(v_3)$ in that example), and is therefore independent of choice of basis. (It does depend on our choice of linear complement $\text{End}_K(V)_0$ for K in $\text{End}_K(V)$, which depends on our choice of r , but we made that choice at the start and have not changed it.) I claim that the condition by which we chose the finite set Σ implies that if the K -linear map (51) is applied simultaneously to every tensor factor $\text{End}_K(V)_0$ in every summand in the above expression for $\text{End}_K(V) \amalg \text{End}_K(V)$, the element x continues to have nonzero components in all the degrees where x had them; in particular, in degree n . Indeed, the linear relations holding among a set of expressions in a tensor product of vector spaces depend only on the linear relations holding among the elements of the given spaces that occur in these expressions; and Σ was chosen so that (51) creates no new linear relations among the elements occurring in our original expression for x . Now in terms of our new basis $B - \{1\}$ of $\text{End}_K(V)_0$, the map (51) acts by throwing out all basis elements other than the $E(q, p)$ with $p, q \in \Sigma$. So since x continues to have nonzero component in degree n after the application of (51), the expression for x using the basis $B - \{1\}$ does indeed involve at least one length- n word in such elements $E(q, p)$ alone.

We now choose, subject to a restriction to be given in a moment, a particular length- n word of this sort occurring with nonzero coefficient in x ,

$$(52) \quad \dots \alpha(E(q_i, p_i)) \beta(E(q_{i-1}, p_{i-1})) \alpha(E(q_{i-2}, p_{i-2})) \beta(E(q_{i-3}, p_{i-3})) \dots,$$

where $(q_j, p_j) \in \Sigma \times \Sigma - \{(r, r)\}$ ($j = 1, \dots, n$).

To do this, let us work from the right, first choosing an $\alpha(E(q_1, p_1))$ or $\beta(E(q_1, p_1))$ that occurs as a rightmost factor in some length- n word of the form (52) in our expression for x ; then choosing for $\beta(E(q_2, p_2))$ or $\alpha(E(q_2, p_2))$, as the case may be, an element of this form that occurs in second position from the right, immediately to the left of our first chosen factor, in at least one length- n word of that form; and so on. The one restriction we impose is that for each $j = 1, \dots, n$, (q_j, p_j) should satisfy $q_j \neq p_j$ if this is possible, i.e., if there is a factor satisfying $q_j \neq p_j$ which occurs, followed by the terms chosen so far, in the j th position of a length- n word occurring in x . Henceforth, (52) will denote the particular word so chosen.

The reason we avoid choices with $q_k = p_k$ whenever possible is that it will not be as easy to make use of the fact that elements of the form $E(p, p)$ are nonscalar as it will for other elements; but we will be able

to do so if there are no elements $E(q, p)$ with $q \neq p$ “in the vicinity”. The distinction between these cases is used in the next definition.

For $k = 1, \dots, n$, let us define elements $p'_k \in V$, by

$$(53) \quad \begin{aligned} p'_k &= p_k \in \Omega && \text{if } q_k \neq p_k, \\ p'_k &= p_k + r \in V && \text{if } q_k = p_k. \end{aligned}$$

Note that in the second line above, the summands p_k and r are distinct members of Ω , for if $q_k = p_k$, we cannot have $p_k = r$, since $E(r, r) \notin B$. For those values of k such that $q_k = p_k$, let us temporarily form a new basis of the k th level (the k th direct summand) of $\bigoplus_{\omega} V$, by deleting from $\Omega \times \{k\}$ the basis element (r, k) and inserting (p'_k, k) in its place, while for those k such that $q_k \neq p_k$, let us keep the basis $\Omega \times \{k\}$. Thus, the union over k of these bases is a new basis of $\bigoplus_{\omega} V$. Note that for *every* k , (p'_k, k) and (q_k, k) are distinct elements of our new basis.

We now define our automorphism t of $\bigoplus_{\omega} V$ to fix all elements of our new basis, except for the following $2(n+1)$ elements, which we let it transpose in pairs as shown:

$$(54) \quad (p_1, 0) \leftrightarrow (p'_1, 1), \quad (q_{k-1}, k-1) \leftrightarrow (p'_k, k) \quad (1 < k \leq n), \quad (q_n, n) \leftrightarrow (q_n, n+1).$$

This is where we need the second line of (53). It insures that even if $q_k = p_k$, the basis elements (p'_k, k) and (q_k, k) are distinct, so that the rules $(q_{k-1}, k-1) \leftrightarrow (p'_k, k)$ and $(q_k, k) \leftrightarrow (p'_{k+1}, k+1)$ do not give contradictory specifications of the action of t on the same basis element at the k th level.

As promised, t has order 2 and sends members of our original basis $\Omega \times \omega$ to linear combinations of members of that basis with integer coefficients. Having noted this, we shall now go back to using $\Omega \times \omega$ as the basis in terms of which we will compute with elements of $\bigoplus_{\omega} V$, and shall think of the terms of (54) as expressions in those basis elements.

As sketched earlier, we now let $h : \text{End}_K(V) \amalg \text{End}_K(V) \rightarrow \text{End}_K(\bigoplus_{\omega} V)$ be the homomorphism which takes elements $\alpha(f)$ to f (acting by the natural action) and elements $\beta(f)$ to $t f t^{-1} = t f t$.

The remainder of the proof follows closely the concluding steps of the proof of Lemma 4.1(ii). We want to show that $h(x) \neq 0$. Clearly, this is equivalent to proving nonzero the element $h(x)'$ that we get on multiplying $h(x)$ on the right by t if the rightmost term of (52) is $\alpha(E(p_1, q_1))$, while leaving that side unchanged if that term is $\beta(E(p_1, q_1))$, and multiplying on the left by t if the leftmost term of (52) is $\alpha(E(p_n, q_n))$, while leaving that side unchanged if that term is $\beta(E(p_n, q_n))$. To avoid cumbersome language, we shall call products of $E(p, q)$'s and t s occurring with nonzero K -coefficient in our expression for $h(x)'$ the “summands” in that expression (not counting the K -coefficients as parts of these “summands”). A consequence of our definition of $h(x)'$ is that the summand therein arising from the term (52) of $h(x)$ has a t at each end, so we can now write it so as show those ends. Let us also give it a name:

$$(55) \quad u = t E(q_n, p_n) t E(q_{n-1}, p_{n-1}) t \dots t E(q_2, p_2) t E(q_1, p_1) t.$$

The general summand in the expression for $h(x)'$ will likewise be an alternating product of t and elements of $B - \{1\}$ (which we will call the “ B -factors” of the summand), with at most n of the latter. We claim now that when we apply $h(x)'$ to $(p_1, 0)$, the summand u shown in (55), and only that summand, leads to a nonzero component at the $n+1$ st level of $\bigoplus_{\omega} V$.

Since in each summand, the only factors that carry elements of $\bigoplus_{\omega} V$ from one level to the next are the factors t , and each of these moves elements by only one level, the only summands in $h(x)'$ that can possibly lead to components at the $n+1$ st level in $h(x)'(p_1, 0)$ are those which, like (55), have exactly n (rather than fewer) B -factors, and have a t at each end. Consider any such summand

$$(56) \quad w = t b_n t b_{n-1} t \dots t b_2 t b_1 t,$$

where $b_k \in B - \{1\}$ ($k = 1, \dots, n$). We shall show inductively for $k = 1, \dots, n+1$ that if we apply to $(p_1, 0)$ the substring $t b_{k-1} t b_{k-2} t \dots t b_2 t b_1 t$ of w , then the components of the result in levels higher than the k th are zero, the component in the k th level is a scalar multiple of (p'_k, k) , and the scalar factor is nonzero if and only if our substring agrees exactly with the corresponding substring of u , i.e., if and only if $b_i = E(q_i, p_i)$ for $i = 1, \dots, k-1$.

The base case $k = 1$ is immediate: we are merely applying t to $(p_1, 0)$, and by (54) the result is $(p'_1, 1)$. (This rightmost factor t was important in distinguishing the action of u from actions of summands of $h(x)'$ not ending in t ; but we have already used it to exclude such strings from consideration.)

Now let our inductive assumption hold for some k with $1 \leq k \leq n$. If we do not have $b_i = E(q_i, p_i)$ for all $i < k$, then by that inductive assumption, $t b_{k-1} t \dots t b_1 t(p_1, 0)$ has zero component at the k th and higher levels, so multiplication by b_k and then by t will not bring anything to the $k+1$ st level or higher.

If the factors of w so far *have* agreed with those of u , then by inductive hypothesis, $t b_{k-1} t \dots t b_1 t(p_1, 0)$ has at the k th level a nonzero scalar multiple of (p'_k, k) . When we apply b_k to this, if $b_k = E(q_k, p_k)$ i.e., if this too agrees with the corresponding factor of u , then we see that, whether (p'_k, k) is equal to (p_k, k) or to $(p_k, k) + (r, k)$, the factor $E(q_k, p_k)$ will send it to (q_k, k) ; and the subsequent application of t will give us a term $(p'_{k+1}, k+1)$ at the $k+1$ st level, as desired.

If $b_k \neq E(q_k, p_k)$, there are several cases to consider. First, b_k might be one of the members of B belonging to the subspace of $\text{End}_K(V)_0$ annihilating Σ . In that case, it annihilates (p'_k, k) , leaving nothing at the k th level, and the subsequent application of t brings nothing to the $k+1$ st level. Otherwise, we have $b_k = E(q, p)$ for some $(q, p) \neq (q_k, p_k)$. Clearly, the only cases in which $E(q, p)$ can fail to annihilate (p'_k, k) are

- (a) if $p = p_k$, or
- (b) if p'_k has the form $p_k + r$, and $p = r$.

In case (a), since we have assumed that $E(q, p) \neq E(q_k, p_k)$, we must have $q \neq q_k$, so $E(q, p)(p'_k, k) = (q, k) \neq (q_k, k)$, and by (54), the subsequent application of t will not bring this up to the $k+1$ st level.

In case (b), the assumption $p'_k = p_k + r$ means, by (53), that $E(q_k, p_k) = E(p_k, p_k)$. But recall that in choosing the term (52), we avoided this possibility whenever possible. A consequence is that since w was not chosen in preference to u , we must likewise have $b_k = E(p, p)$. However, case (b) assumed $p = r$, so $b_k = E(r, r)$, which is excluded by the definition of B . Hence case (b) does not occur.

We thus conclude that u is the unique summand in $h(x)'$ which, when applied to $(p_1, 0)$, gives an element having nonzero component at the $n+1$ st level. Hence $h(x)'(p_1, 0) \neq 0$, so $h(x)' \neq 0$, so $h(x) \neq 0$, completing the proof that our action of $\text{End}_K(V) \amalg \text{End}_K(V)$ is faithful. \square

Now for an unexpected bonus. We saw in §8 that when K had large cardinality, our tools for proving *nonembeddability* of algebras in $\text{End}_K(V)$ gave weaker results than we expected from our results on groups, monoids and lattices. We shall now see that we also get stronger *positive* embeddability results in that situation. We need the following fact.

Lemma 10.1 (cf. [13, Exercise 1, p.248]). *Let k be any commutative integral domain, and M the multiplicative monoid of k . Then the product k -algebra k^ω contains a subalgebra isomorphic to the monoid algebra kM . In particular, letting G denote the group of units of k , it contains a copy of the group algebra kG .*

Proof. For each $a \in k$, let $x_a \in k^\omega$ denote the sequence of powers, $(1, a, a^2, \dots)$. Clearly, $a \mapsto x_a$ is a monoid homomorphism from M into the multiplicative monoid of k^ω . Moreover, the elements x_a are k -linearly independent by the properties of the Vandermonde determinant. Hence they span a subalgebra of k^ω isomorphic to kM . The final assertion clearly follows. \square

Proposition 10.2. *$\text{End}_K(V)$ contains a coproduct of $\text{card}(K)$ copies of itself as an associative K -algebra. (Hence, in view of Theorem 4.2(iii), it contains such a coproduct of $\max(2^{\text{card}(\Omega)}, \text{card}(K))$ copies of itself.)*

Proof. For K finite, our main statement is weaker than Theorem 4.2(iii), so assume K infinite.

Since $\text{End}_K(V)$ contains a subalgebra isomorphic to K^ω , the preceding lemma shows that it contains a copy of the group algebra KG on a group G of cardinality $\text{card}(K - \{0\}) = \text{card}(K)$.

We have also just seen that it contains a copy of $\text{End}_K(V) \amalg \text{End}_K(V)$; combining these observations, we conclude that it contains a copy of $\text{End}_K(V) \amalg KG$. For notational convenience, let us identify $\text{End}_K(V)$ with the first factor in this coproduct, and G (which we regard as an abstract group, forgetting about its relation with K) with its image in the second factor. Then for each $g \in G$, we can form the conjugate algebra $g \text{End}_K(V) g^{-1}$, getting $\text{card}(K)$ isomorphic copies of $\text{End}_K(V)$. Moreover, from the linear independence of the elements of G in KG and the normal form for the coproduct $\text{End}_K(V) \amalg KG$, one sees that the subalgebra these generate will be their coproduct, so $\text{End}_K(V)$ indeed contains a coproduct of $\text{card}(K)$ copies of itself. (This is the ring-theoretic analog of the appearance of big free products inside smaller free products in the Kurosh Subgroup Theorem.) The final assertion clearly follows. \square

Corollary 10.3. *$\text{End}_K(V)$ contains a coproduct, as K -algebras, of $\max(2^{\text{card}(\Omega)}, \text{card}(K))$ copies of every K -algebra of dimension $\leq \text{card}(\Omega)$.* \square

Of course, these results lead to the questions:

Question 10.4. *Can we prove a version of Theorem 3.2(iii) with $F(2^{\text{card}(\Omega)})$ replaced by $F(\max(2^{\text{card}(\Omega)}, \text{card}(K)))$ for a wider class of functors F than the “coproduct as K -algebras of I copies of A ” functors of Corollary 10.3?*

An interesting test case would be that in which F is the functor taking I to the K -algebra presented by an I -tuple of commuting idempotents.

Remark: One can strengthen Proposition 10.2 so as to increase $\max(2^{\text{card}(\Omega)}, \text{card}(K))$ to $\text{card}(K)^{\text{card}(\Omega)}$, which occasionally exceeds the former value; e.g., when $\text{card}(K)$ has the form $\aleph_{\alpha+\omega}$ and is $> 2^{\text{card}(\Omega)}$. To do so, replace ω in Lemma 10.1 with the free abelian monoid $A(\Omega)$ on Ω , which has cardinality $\text{card}(\Omega)$, replace the maps x_a in the proof by all homomorphisms of $A(\Omega)$ into the multiplicative monoid M of k , apply the theorem on linear independence of characters to conclude that $k^{A(\Omega)}$ contains a copy of the monoid algebra kM^Ω , and use this version of the lemma to get the strengthened proposition. (One can’t raise the cardinal in the proposition higher than $\text{card}(K)^{\text{card}(\Omega)}$, since $\dim_K(\text{End}_K(V)) = \dim_K(V^\Omega) \leq \dim_K((K^\Omega)^\Omega) = \dim_K(K^{\Omega \times \Omega}) = \dim_K(K^\Omega) \leq \text{card}(K^\Omega) = \text{card}(K)^{\text{card}(\Omega)}$; cf. [13, p.247, Theorem 2].)

It is curious that the nonembeddability results we are able to prove become weaker than expected as soon as $\text{card}(K) > \text{card}(\Omega)$, but our positive results become stronger only when $\text{card}(K) > 2^{\text{card}(\Omega)}$.

11 Appendix: another embedding of $\text{Sym}(\Omega) \amalg \text{Sym}(\Omega)$ in $\text{Sym}(\Omega)$.

Here is the alternative proof of Lemma 4.1(i) mentioned shortly before the statement of Lemma 4.1.

As in the preceding appendix, we begin by recalling a structure theorem for coproducts, this time coproducts of groups. But we will make different assumptions (we won’t assume there are only two groups, but we will assume the groups are disjoint except for their identity elements, so that we do not have to write the coprojection maps explicitly), we will use the structure theorem, initially, for a different purpose (to motivate a somewhat bizarre action of the coproduct group), and we will consider for most of this section arbitrary groups (or sometimes, monoids) rather than a pair of copies of $\text{Sym}(\Omega)$, though that is the case to which we will ultimately apply our result.

Let $(G_i)_{i \in I}$ be any family of groups whose sets of nonidentity elements are pairwise disjoint. Recall that the general element of the coproduct group $\coprod_{i \in I} G_i$ can be written uniquely as

$$(57) \quad g_n \dots g_2 g_1,$$

where $n \geq 0$ and $g_n, \dots, g_1 \in \bigcup_{i \in I} G_i - \{1\}$, say with

$$(58) \quad g_r \in G_{i_r} \quad (r = 1, \dots, n),$$

and where successive indices i_r, i_{r+1} are distinct; i.e., two elements from the same group G_i never occur in immediate succession.

Suppose we are given a representation of each G_i by permutations of a nonempty set Ω_i , and we form the direct product $\prod \Omega_i$. Then starting at any point $(x_i)_{i \in I} \in \prod \Omega_i$, an expression (57) allows us to construct a “path” in $\prod \Omega_i$: At the first step, we move from (x_i) to the point agreeing with (x_i) on all but the i_1 -coordinate, which is moved by g_1 ; the next step takes us to the point whose i_2 -coordinate has also been moved by g_2 , etc.. This suggests that we make the set of “paths” in $\prod \Omega_i$ in which each step involves changing just one coordinate, and two successive steps never change the same coordinate, into a $\coprod G_i$ -set.

Some difficulties arise. Though, by assumption, no g_r in (57) is an identity element, some of these factors may lie in the stabilizers of the coordinates they are to be applied to. In such cases should we, as the above description might suggest, allow trivial “steps” in our path, where no coordinate is changed? It turns out that this would lead to difficulties; so we specify that in such cases, no step is added to our path. Also, for inverses to behave correctly, we must allow some elements to delete rather than adding steps to our paths.

The resulting construction is described in the next lemma. Note that it is not claimed that if the Ω_i are faithful G_i -sets, then the $\coprod G_i$ -set $\boxtimes_{i \in I} \Omega_i$ described there is also faithful. Rather, we shall subsequently note additional conditions that ensure faithfulness.

Up to the step of achieving faithfulness, our construction works as well for arbitrary monoids as for groups, so the lemma below is stated in that context. As usual, we understand actions to be *left* actions.

Note that in the statement below, subscripts n and r do not correspond to the subscript i above; rather, each of the steps x_r ($r = 1, \dots, n$) in our “path” is itself an I -tuple $x_r = (x_{r,i})_{i \in I}$.

Lemma 11.1. *Let $(M_i)_{i \in I}$ be a family of monoids, and for each $i \in I$ let Ω_i be an M_i -set. Let $\boxtimes_{i \in I} \Omega_i$ be the set of all finite sequences (x_1, \dots, x_n) ($n \geq 1$) where each $x_r \in \prod_{i \in I} \Omega_i$ ($r = 1, \dots, n$), every pair of successive terms x_r, x_{r+1} ($1 \leq r < n$) differs in one and only one coordinate, and the coordinate at which x_{r+1} differs from x_r is not the same as the coordinate at which x_r differs from x_{r-1} ($1 < r < n$).*

Then $\boxtimes_{i \in I} \Omega_i$ may be made a $\prod_{i \in I} M_i$ -set (which we might call the “path product” of the M_i -sets Ω_i) by defining the action of each $g \in M_j$ ($j \in I$) on $x = (x_1, \dots, x_n) \in \boxtimes_{i \in I} \Omega_i$ by the following rules:

- (59) *Let $x'_n \in \prod_{i \in I} \Omega_i$ denote the I -tuple obtained from x_n by modifying its j -coordinate via the action of g on that element of M_j . Then*
- (i) *If $x'_n = x_n$, we define $gx = x$.*
 - (ii) *If $x'_n \neq x_n$, then:*
 - (ii.a) *If either $n = 1$, or if $n > 1$ and the coordinate at which x_n differs from x_{n-1} is not the j th, we define $gx = (x_1, \dots, x_n, x'_n)$,*
 - (ii.b) *If $n > 1$ and the coordinate at which x_n differs from x_{n-1} is the j th, then:*
 - (ii.b.1) *If $x'_n \neq x_{n-1}$, we define $gx = (x_1, \dots, x_{n-1}, x'_n)$, while*
 - (ii.b.2) *If $x'_n = x_{n-1}$, we define $gx = (x_1, \dots, x_{n-1})$.*

Proof. By the universal property of the coproduct $\prod_{i \in I} M_i$, an action of that monoid on the set $\boxtimes_{i \in I} \Omega_i$ will be defined if we verify that for each $j \in I$, the above conditions define an action of M_j on that set.

This could be done by brute force, dividing into cases according to which headings of the above definition the actions of *two* successive elements of M_j come under. But there is a trick that greatly simplifies this calculation (cf. [3, proof of Proposition 3.6.5]). For each j , we shall define a bijection φ_j between $\boxtimes_{i \in I} \Omega_i$ and a set $(\boxtimes_{i \in I} \Omega_i)_{(j)}$, such that it will be easy to define an action of M_j on $(\boxtimes_{i \in I} \Omega_i)_{(j)}$, and also easy to verify that when we transport this action from $(\boxtimes_{i \in I} \Omega_i)_{(j)}$ to $\boxtimes_{i \in I} \Omega_i$ via the bijection φ_j , the resulting action is described by (59). Roughly, elements of $(\boxtimes_{i \in I} \Omega_i)_{(j)}$, like elements of $\boxtimes_{i \in I} \Omega_i$, will represent “paths” in $\prod \Omega_i$, but in $(\boxtimes_{i \in I} \Omega_i)_{(j)}$ we require every such path to have a final step involving the j -coordinate, at the price of allowing this step (and only this step) to be trivial (i.e., to satisfy $x_{n-1} = x_n$).

Here is the precise description. We take the elements of $(\boxtimes_{i \in I} \Omega_i)_{(j)}$ to be sequences $x = (x_1, \dots, x_n)$ of elements of $\prod \Omega_i$, this time with n always ≥ 2 , such that

- (i) $(x_1, \dots, x_{n-1}) \in \boxtimes_{i \in I} \Omega_i$,
- (ii) x_{n-1} and x_n are either equal, or differ in the j th coordinate only.
- (iii) If $n > 2$, the coordinate at which x_{n-2} and x_{n-1} differ is *not* the j th.

Note that none of these conditions constrains the j th coordinate of x_n . Hence a monoid action of M_j on $(\boxtimes_{i \in I} \Omega_i)_{(j)}$ may be defined by the rule

$$(60) \quad g(x_1, \dots, x_{n-1}, x_n) = (x_1, \dots, x_{n-1}, x'_n),$$

where, as in the statement of the lemma, x'_n is obtained from x_n by modifying its j th coordinate by the action of g , and leaving all other coordinates unchanged.

Let us now define $\varphi_j : \boxtimes_{i \in I} \Omega_i \rightarrow (\boxtimes_{i \in I} \Omega_i)_{(j)}$ to leave unchanged all $x = (x_1, \dots, x_n) \in \boxtimes_{i \in I} \Omega_i$ with $n > 1$ in which the coordinate at which x_{n-1} and x_n differ happens to be the j th, while appending to every other element of $\boxtimes_{i \in I} \Omega_i$ a repetition of its final term. It is straightforward that this is a bijection, and easy to verify that the action of M_j on $\boxtimes_{i \in I} \Omega_i$ induced, via this bijection, by its action (60) on $(\boxtimes_{i \in I} \Omega_i)_{(j)}$, is as described in (59), completing our proof. \square

We now want to use the above construction to get *faithful* actions. We shall see that we can do this if our monoids are right cancellative, in particular, if they are groups.

Let us call an action of a monoid M on a set Σ *strongly faithful* if for every finite family of distinct elements $g_1, \dots, g_n \in M$, there exists $y \in \Sigma$ such that $g_1 y, \dots, g_n y$ are distinct. (The $n = 0$ case of this condition says that Σ is nonempty.) It is easy to see that if Σ is a faithful M -set, then the disjoint union $1 \cup \Sigma \cup \Sigma^2 \cup \Sigma^3 \cup \dots$ (indexed by the natural numbers, with each Σ^n given the product M -set structure) is strongly faithful, and that this construction does not increase infinite cardinalities. On the other hand, we have

Lemma 11.2. *Let $(M_i)_{i \in I}$ be a family of monoids, each having the right cancellation property*

$$(61) \quad ac = bc \implies a = b \quad (a, b, c \in M_i),$$

and for each $i \in I$, let Ω_i be a strongly faithful left M_i -set. Then $\boxtimes_{i \in I} \Omega_i$, defined and made a $\prod_{i \in I} M_i$ -set as in Lemma 11.1, is a faithful $\prod_{i \in I} M_i$ -set.

Proof. Given distinct elements $g = g_m \dots g_2 g_1$ and $h = h_n \dots h_2 h_1$ in $\prod M_i$, we wish to find an $x \in \boxtimes_{i \in I} \Omega_i$ such that $gx \neq hx$. We shall construct below such an element which is of length 1, i.e., a 1-tuple (x_1) with $x_1 \in \prod \Omega_i$. To do this let us, for each $j \in I$, choose the j -coordinate of x_1 as follows. Let $r_{j,1} < \dots < r_{j,m_j}$ be the values of r for which $g_r \in G_j$ and $s_{j,1} < \dots < s_{j,n_j}$ the values s for which $h_s \in G_j$. Although the elements

$$(62) \quad 1, g_{r_{j,1}}, g_{r_{j,2}} g_{r_{j,1}}, \dots, g_{r_{j,m_j}} \dots g_{r_{j,2}} g_{r_{j,1}}, \quad \text{and} \quad 1, h_{s_{j,1}}, h_{s_{j,2}} h_{s_{j,1}}, \dots, h_{s_{j,n_j}} \dots h_{s_{j,2}} h_{s_{j,1}}$$

of M_j need not all be distinct, *successive* elements of each of these lists will be so, by right cancellation. For each $j \in I$ let us use strong faithfulness of M_j to choose, as the j -coordinate of x_1 , an element of Ω_j whose images under *distinct* elements of the combined list (62) are distinct. (For some j , the m_j and n_j in (62) may both be zero, in which case this condition is vacuous.) Thus, the images of x_1 under successive terms of each list in (62) are distinct.

It follows from this choice that each time we apply an element g_r or h_r in building up $g_m \dots g_1 x$ or $h_n \dots h_1 x$, it in fact moves the coordinate of the term of $g_{r-1} \dots g_1 x$, respectively $h_{r-1} \dots h_1 x$ to which it is applied, and successive steps move different coordinates. Thus, we are always in case (ii.a) of the definition of the action of M_i , so the final elements $g_m \dots g_1 x$ and $h_n \dots h_1 x$ reflect all the steps of this process. In particular, if $m \neq n$, the elements gx and hx have different lengths, if $m = n$ and the sequence of indices in I determined by g and h differ, then $g_m \dots g_1 x$ and $h_n \dots h_1 x$ are clearly different, while if $m = n$ and these sequences are the same, then there must be some $j \in M$ such that the sequences of elements of G_j differ; and if we look at the first terms where this occurs, then by right cancellativity, the corresponding terms of (62) will differ, and by our choice of $x_{1,j}$, the elements gx and hx will differ at that step. \square

(The above action will, in fact, be strongly faithful – we could have handled any finite set of elements of $\prod_{i \in I} M_i$ as we did $\{g, h\}$; but we only need the two-element case, and it allowed simpler notation.)

Now letting G_0, G_1 be two isomorphic copies of the group $\text{Sym}(\Omega)$, disjoint except for the identity, each represented naturally on Ω , we get as in the paragraph before Lemma 11.2 strongly faithful actions of G_0 and G_1 on $1 \cup \Omega \cup \Omega^2 \cup \Omega^3 \cup \dots$, which has cardinality $\text{card}(\Omega)$. Calling this set, regarded as a G_0 -set Ω_0 , and regarded as a G_1 -set Ω_1 , the above lemma tells us that the coproduct of G_0 and G_1 acts faithfully on $\Omega_0 \boxtimes \Omega_1$, which is also of cardinality $\text{card}(\Omega)$, completing our alternative proof of Lemma 4.1(i).

More generally, the submonoid of *surjective* endomaps of Ω satisfies (61), so the above construction embeds the coproduct of two copies of that monoid in $\text{Self}(\Omega)$.

On the other hand, if we delete the right cancellativity assumption (61) from the hypothesis of Lemma 11.2, there will in general be no choice of M_i -sets Ω_i making $\boxtimes_{i \in I} \Omega_i$ faithful. For if one of the monoids, say M_1 , has elements $a \neq b$ and c satisfying $ac = bc$, and at least one other of the monoids is nontrivial, say M_2 , with a nonidentity element d , then in $\prod M_i$ we find that adc and bdc are distinct elements having the same action on $\boxtimes_{i \in I} \Omega_i$ for any family of M_i -sets Ω_i .

12 Appendix: Some conditions on complete lattices.

We noted in the discussion preceding Question 5.9 that the conditions we had proved on chains of solution sets in $\text{Sym}(\Omega)$ and $\text{Self}(\Omega)$ were consequences of the stronger statement that the lattice of all such solution sets embeds in the system of closed sets of a topological space having a basis of $\leq \text{card}(\Omega)$ open sets. The same observation holds for the results obtained in later sections on solution sets in $\text{Equiv}(\Omega)$, etc..

The next lemma compares these and related conditions. Here a κ -generated topological space means a topological space having a basis of open sets of cardinality $\leq \kappa$; equivalently, having a *subbasis* of open sets of that cardinality; equivalently, having such a basis or subbasis of *closed* sets. An *embedding* of lattices, of complete lower semilattices, etc., means a one-to-one homomorphism of such structures; an embedding of *partially ordered sets* means a (necessarily one-to-one) map that preserves both the relations \leq and $\not\leq$.

Lemma 12.1. *Let κ be an infinite cardinal, and A a complete lattice. Then of the following conditions, each implies the next, and conditions with the same roman numeral and different suffixes are equivalent.*

- (i.a) A is embeddable as a complete lower semilattice in $\mathbf{P}(\kappa)$.
- (i.b) A is generated as a complete upper semilattice by a set of $\leq \kappa$ elements.
- (ii.a) A is embeddable as a complete lower semilattice in a complete lower semilattice generated by $\leq \kappa$ elements.
- (ii.b) A is embeddable as a partially ordered set in a complete lower semilattice generated by $\leq \kappa$ elements.
- (ii.c) A is embeddable as a partially ordered set in $\mathbf{P}(\kappa)$.
- (ii.d) A is embeddable as a complete lower semilattice in the system of closed subsets of a κ -generated topological space.
- (ii.e) A is embeddable as a partially ordered set in the system of closed subsets of a κ -generated topological space.
- (ii.a*)-(ii.e*) *The duals of (ii.a)-(ii.e); i.e., the corresponding statements with “upper semilattice” replaced by “lower semilattice”, and “closed subsets” by “open subsets”, wherever applicable. (So, no change in (ii.c)).*
- (iii) *Every chain in A has a dense subset of cardinality $\leq \kappa$.*
- (iv.a) *No chain in A has a family of $> \kappa$ disjoint intervals.*
- (iv.b) *No chain in A has $> \kappa$ jumps.*
- (v) *A contains no well-ordered or reverse-well-ordered chain of cardinality $> \kappa$.*

Moreover, if for each family of conditions whose equivalence is asserted above, we denote the common condition using the corresponding roman numeral with suffixes dropped, then the implications (i) \implies (ii) \implies (iii) are irreversible for all κ , but are both reversible if A is restricted to be a chain; the implication (iv) \implies (v) is irreversible for $\kappa = \aleph_0$, while the reversibility of (iii) \implies (iv) for $\kappa = \aleph_0$ is equivalent to Suslin’s Hypothesis, known to be independent of ZFC.

Proof. (i.a) \implies (i.b): Given a complete lower semilattice embedding $f : A \rightarrow \mathbf{P}(\kappa)$, let us associate to each $\alpha \in \kappa$ the meet $g(\alpha) \in A$ of all elements $x \in A$ satisfying $\alpha \in f(x)$. We see that $g(\alpha)$ will be the least y such that $\alpha \in f(y)$, and we deduce that every $x \in A$ is the join in A of $\{g(\alpha) \mid \alpha \in f(x)\}$. So $\{g(\alpha) \mid \alpha \in \kappa\}$ generates A as a complete upper semilattice.

(i.b) \implies (i.a): Given a generating set $\{g_\alpha \mid \alpha \in \kappa\}$ for A as a complete upper semilattice, we find that an embedding $A \rightarrow \mathbf{P}(\kappa)$ as complete lower semilattices is given by the map $x \mapsto \{\alpha \mid g_\alpha \leq x\}$.

(i.a) \implies (ii.c) is immediate.

To prove the equivalence of the versions of (ii), we shall show (ii.a) \implies (ii.d) \implies (ii.e) \implies (ii.b) \implies (ii.c) \implies (ii.a). Since (ii.c) is self-dual, it will follow that these conditions are also equivalent to their starred variants.

(ii.a) \implies (ii.d): It suffices to show that every complete lower semilattice A' generated by a set $\{x_\alpha \mid \alpha \in \kappa\}$ is embeddable as a complete lower semilattice in the system of closed subsets of a κ -generated topology. Given such an A' , define for each $x \in A'$ the “principal downset”

$$(63) \quad D(x) = \{y \in A' \mid y \leq x\} \subseteq \mathbf{P}(A').$$

We see that these sets form a complete lower semilattice isomorphic to A' ; hence if we define a topology on the underlying set of A' using the $D(x)$ as a subbasis of closed sets, A' embeds as a complete lower semilattice in the complete lattice of closed sets of that topology. Moreover, the closed sets $D(x_\alpha)$ ($\alpha \in \kappa$) also form a subbasis of closed sets for this topology, so it is κ -generated, as required.

(ii.d) \implies (ii.e) \implies (ii.b): Trivial.

(ii.b) \implies (ii.c): Suppose A is embeddable as partially ordered set in a complete lower semilattice A' as in (ii.b). Thus, A' satisfies the dual of (i.b). Hence it satisfies the dual of (i.a), hence A , being embeddable in A' as a partially ordered set, satisfies (ii.c).

(ii.c) \implies (ii.a): Note that the map D of (63) (with A in the role of A') is an embedding $A \rightarrow \mathbf{P}(A)$ as complete lower semilattices. Hence it will suffice to show that (ii.c) implies that the image of A under D lies in a complete lower subsemilattice of $\mathbf{P}(A)$ generated by $\leq \kappa$ elements. Given an embedding $f : A \rightarrow \mathbf{P}(\kappa)$ as in (ii.c), let us define for each $\alpha \in \kappa$ the set $c_\alpha = \{x \in A \mid \alpha \notin f(x)\} \in \mathbf{P}(A)$. We see that for every $x \in A$, the $D(x)$ of (63) is the intersection of those members of $\{c_\alpha \mid \alpha \in \kappa\} \cup \{A\} \subseteq \mathbf{P}(A)$ that contain x , so D carries A into the complete lower subsemilattice of $\mathbf{P}(A)$ generated by these elements.

(ii) $\not\Rightarrow$ (i): Let A be a lattice consisting of a least element 0, a greatest element 1, and 2^κ pairwise incomparable elements lying between these. Clearly it does not satisfy (i.b). To see that it satisfies (ii.d), recall that the product topology on $\mathbf{P}(\kappa)$ is κ -generated, and let us map A into the closed sets of that topology by sending 0 to \emptyset , 1 to the improper subset, and the 2^κ intermediate elements to the singletons $\{S\} \subseteq \mathbf{P}(\kappa)$ ($S \subseteq \kappa$). (Alternatively, one can show that A satisfies (ii.c) by noting that $\mathbf{P}(\kappa \times 2) \cong \mathbf{P}(\kappa)$ has an antichain of cardinality 2^κ , consisting of the sets $(s \times \{0\}) \cup (s^c \times \{1\})$ ($s \subseteq \kappa$).

(ii.c) \Rightarrow (iii): It suffices to show that every chain C in $\mathbf{P}(\kappa)$ has a dense subset of cardinality $\leq \kappa$. Let C be such a chain, and for each pair $\alpha, \beta \in \kappa$ such that some $x \in C$ contains α but not β , choose such an element, $x_{\alpha, \beta}$. This gives a family of $\leq \kappa$ elements which is easily seen to be dense.

(iii) $\not\Rightarrow$ (ii): Take A as in the example used to show (ii) $\not\Rightarrow$ (i), but this time with $> 2^\kappa$ pairwise incomparable elements. This clearly satisfies (iii), but in view of its cardinality, cannot satisfy (ii.c).

On the other hand, the assertion that if A is a chain the implications (i) \Rightarrow (ii) \Rightarrow (iii) are reversible follows from the obvious implication (iii) \Rightarrow (i.b) in this case.

(iii) \Rightarrow (iv.a): Given any chain C in A and any dense set S of $\leq \kappa$ elements of C , we see that for every interval $[x, y]$ in C , at least one element of S must belong to $[x, y]$, showing that C cannot have $> \kappa$ disjoint intervals.

Concerning the reverse implication, recall that Suslin's Hypothesis says that every totally ordered set S having no uncountable family of disjoint intervals has a countable dense subset, and that this is independent of ZFC [18]. Assuming Suslin's Hypothesis, we immediately get (iv.a) \Rightarrow (iii) for $\kappa = \aleph_0$ by applying that statement to an arbitrary chain C in A .

In proving the converse assertion, note that (iv.a) and (iii) are statements about a *complete* lattice A (first sentence of the lemma); so we need to show that if S is any counterexample to Suslin's hypothesis, we can obtain from it a complete lattice A that satisfies (iv.a) but not (iii). Given such an S , let A be its completion as a totally ordered set; I claim that A inherits the properties making S a counterexample to Suslin's conjecture, and thus gives the desired example. Indeed, for any infinite dense subset $D \subseteq A$, if we take an element of S between every two distinct elements of D (and throw in the greatest and/or least element of S if these exist), we get a dense subset D' of S of the same cardinality; so since S has *no* countable dense subset, neither does A . Likewise, if some chain $C \subseteq A$ had an uncountable family of disjoint intervals, then for each of these intervals $[x, y]_C$ we could choose $x', y' \in S$ with $x \leq x' < y' \leq y$, getting an uncountable family of disjoint intervals $[x', y']_S$ in S ; so chains in A inherit from S the nonexistence of such families.

Since (iv.a) and (iv.b) are negative statements, we will prove their equivalence in contrapositive form:

\neg (iv.a) \Rightarrow \neg (iv.b): If a chain C in A has a family of disjoint intervals $[x_\alpha, y_\alpha]$ where α ranges over some $\lambda > \kappa$, in then the subchain $C' = \{x_\alpha, y_\alpha \mid \alpha \in \lambda\}$, the pairs $x_\alpha < y_\alpha$ will be jumps.

\neg (iv.b) \Rightarrow \neg (iv.a): If C has $> \kappa$ jumps, let us associate to each jump $x < y$ the two-element interval $[x, y]$. There is the slight difficulty that distinct jumps may not yield disjoint intervals: the upper endpoint of one may equal the lower endpoint of the other. However, if we take a family of these intervals maximal for the property of being pairwise disjoint, is easy to verify that this still has cardinality $> \kappa$, giving the desired assertion.

(iv.b) \Rightarrow (v) is clear.

To show that for $\kappa = \aleph_0$, (v) $\not\Rightarrow$ (iv.b), let A be $\mathbb{R} \times 2$, lexicographically ordered. This has continuum many jumps (a jump $(r, 0) < (r, 1)$ for each real number r), but has no uncountable well-ordered or reverse well-ordered subsets. \square

Some remarks on the above lemma:

It is, of course, condition (ii) and its consequences (iii)-(v) that are directly relevant to the results of preceding sections. I have included (i) for perspective.

(i) is the only condition in the lemma that is not equivalent to its dual. To show this inequivalence, let $A \subseteq \mathbf{P}(\kappa)$ consist of \emptyset and all sets of cardinality κ . This clearly satisfies the dual of (i.a), but we claim it does not satisfy (nondualized) (i.b). Indeed, given a putative generating set $\{x_\alpha \mid \alpha \in \kappa\}$ for A as a complete upper semilattice (where we allow repetitions in the indexing in case this set has cardinality $< \kappa$), one can construct by transfinite recursion an element $y \in \mathbf{P}(\kappa)$ having κ elements, but missing at least one element from each $x_\alpha \neq \emptyset$ in our family. We see that y will belong to A , but not to the complete upper subsemilattice generated by $\{x_\alpha\}$.

We could have written (i.a*) and (i.b*) for the duals to (i.a) and (i.b), adding to the lemma their mutual equivalence and the implication (i*) \implies (ii); but so naming those conditions would have broken the convention that conditions beginning with the same roman numeral are equivalent.

Conditions (ii.d) and (ii.e) (and hence their duals), which refer to a κ -generated topological space, are equivalent to the formally stronger conditions referring to a κ -generated *Hausdorff* (and if we wish, *totally disconnected*) topological space. For given a topological space X as in one of those statements, which we may assume without loss of generality to be T_0 , and which has a subbasis of $\leq \kappa$ closed sets, we can throw in the complements of those sets to get a stronger topology on X which is still κ -generated, but is now totally disconnected and Hausdorff, and whose lower semilattice of closed sets contains the lower semilattice of sets closed in the original topology.

It is curious that the example we gave for (ii) $\not\Rightarrow$ (i) in the proof of the lemma is an instance of (37) with $\text{card}(I) > \kappa$. Thus, although the lattice A in question satisfies (ii), the lattice $L_{A^\wedge, 1}^\equiv$ does not even satisfy (v). The example given above satisfying the dual of (i.a), but not (i.b), similarly contains such an instance of (37), in view of Sierpiński's result [17] that there exists a family of $> \kappa$ subsets of κ , each having cardinality κ , but with pairwise intersections all of smaller cardinality. These observations suggest the first part of the next question. The second part is also natural, in view of the simpleminded example we used for (iii) $\not\Rightarrow$ (ii).

Question 12.2. (a) *If we add to the hypotheses of Lemma 12.1 the assumption that $L_{A^\wedge, 1}^\equiv$ satisfies (v), does this change the validity of the nonimplications shown? (If not, we might try imposing the stronger condition (iv) or (iii) on these lattices, and/or looking at $L_{A^\wedge, J}^\equiv$ for J of larger cardinality, up to κ .)*

(b) *If we add to the hypotheses of Lemma 12.1 the assumption that A has cardinality $\leq 2^\kappa$, or that it has order-dimension $\leq \kappa$ (i.e., is embeddable as a partially ordered set in a direct product of $\leq \kappa$ totally ordered sets – both conditions being implied by (ii)), does this affect the validity of the assertion (iii) $\not\Rightarrow$ (ii)?*

Our final corollary, below, answers a couple of other questions suggested by that lemma. The formulation of statement (b.2) of that corollary uses implicitly the fact that in a complete upper semilattice A' with least element, every subset has a greatest lower bound, so that A' may be regarded as a complete lattice (though if A' was obtained as a complete upper subsemilattice of a complete lattice A , the meet operation of A' , will not in general agree with that of A). Statement (b.2*) uses the dual observation.

Corollary 12.3. (a) *For the conditions of Lemma 12.1 that treat A only as a partially ordered set, namely (ii.b), (ii.c), (ii.e), (ii.b*), (ii.c*), (ii.e*) and (iii)-(v), the implications stated in that lemma (for A the underlying partially ordered set of a complete lattice) in fact hold for any partially ordered set A .*

(b) *Let C be a complete lattice, and A a nonempty subset of C . Then the following conditions are equivalent.*

(b.1) *As a partially ordered set, A forms a complete lattice.*

(b.2) *A has a greatest element, and forms a complete lower subsemilattice of a complete upper subsemilattice B of C having a least element.*

(b.2*) *A has a least element, and forms a complete upper subsemilattice of a complete lower subsemilattice B of C having a greatest element.*

(Note: in (b.2) and (b.2*), the statement that A is a complete lower or upper subsemilattice of B means that it is closed under the relevant meet or join operations of B , not under those of C , which do not in general carry B into itself.)

Sketch of proof. (a): The versions of condition (ii) listed here all involve embeddings of A as a partially ordered set in a certain complete semilattice B , which, by adjoining a greatest or least element if necessary, can be assumed a complete lattice. The mutual equivalence of these conditions for complete lattices, applied to B , gives embeddings of B ; these in turn lead to the embeddings of A as a partially ordered set that we want. The same method yields the implication (ii) \implies (iii). The proofs of the remaining implications were entirely order-theoretic, and go over unchanged.

(b): Clearly (b.2) \implies (b.1). On the other hand, if A satisfies (b.1), and one lets B denote the result of closing A in C under arbitrary joins and throwing in the least element, then one finds that arbitrary meets in A are still meets under the operation of B , so that B witnesses (b.2). Thus, (b.1) \iff (b.2), and by duality, (b.1) \iff (b.2*). \square

References

- [1] R. Baer, *Die Kompositionsreihe der Gruppe aller eineindeutigen Abbildungen einer unendlichen Menge auf sich*, *Studia Math.* **4** (1934) 15–17.
- [2] George M. Bergman, *The diamond lemma for ring theory*, *Adv. in Math.* **29** (1978) 178–218. MR **81b**:16001.
- [3] George M. Bergman, *An Invitation to General Algebra and Universal Constructions*, pub. Henry Helson, Berkeley, CA, 1998. Readable online at <http://math.berkeley.edu/~gbergman/245>. MR **99h**:18001.
- [4] George M. Bergman, *Generating infinite symmetric groups*, *J. London Math. Soc.* **38** (2006) 429–440. MR **2007e**:20004.
- [5] George M. Bergman, *Two statements about infinite products that are not quite true*, pp.35–58 of *Groups, Rings and Algebras* (Proceedings of a conference in honor of Donald S. Passman), ed. W. Chin, J. Osterburg and D. Quinn, *Contemporary Mathematics*, v.**420**, 2006. MR 2279231.
- [6] George M. Bergman and Saharon Shelah, *Closed subgroups of the infinite symmetric group*, *Algebra Universalis* **55** (2006) 137–173. MR 2280223.
- [7] N. G. de Bruijn, *Embedding theorems for infinite groups*, *Nederl. Akad. Wetensch. Proc. Ser. A.* 60 = *Indag. Math.* **19** (1957) 560–569. MR **20**#4589. (In addition to the correction to this paper made in [8], the result quoted from [14] at p.561, 6th paragraph should exclude the countable case, and the result quoted from [1] at p.566, 3rd from last paragraph, should include, when $\mathfrak{n} = \aleph_0$, the case of the infinite alternating group.)
- [8] N. G. de Bruijn, *Addendum to “Embedding theorems for infinite groups”*, *Nederl. Akad. Wetensch. Proc. Ser. A* 67 = *Indag. Math.* **26** (1964) 594–595. MR **30**#142.
- [9] R. M. Bryant and V. A. Roman’kov, *The automorphism groups of relatively free algebras*, *J. Algebra* **209** (1998) 713–723. MR **99k**:20073.
- [10] Manfred Droste and Rüdiger Göbel, *Uncountable cofinalities of permutation groups*, *J. London Math. Soc.* (2) **71** (2005) 335–344. MR **2006a**:20008.
- [11] Manfred Droste and W. Charles Holland, *Generating automorphism groups of chains*, *Forum Math.* **17** (2005) 699–710. MR **2006c**:06023.
- [12] Ulrich Felgner and Frieder Haug, *The homomorphic images of infinite symmetric groups*, *Forum Math.* **5** (1993) 505–520. MR **95e**:20007.
- [13] Nathan Jacobson, *Lectures in abstract algebra. Volume II: Linear algebra*, Van Nostrand, 1953, and Springer GTM, No. 31, 1975. MR **14**,837e and MR **51**#5614.
- [14] Bjarni Jónsson, *Universal relational systems*, *Math. Scand.* **4** (1956) 193–208. MR **20**#3091.
- [15] Roger C. Lyndon and Paul E. Schupp, *Combinatorial Group Theory*, Springer-Verlag, *Ergebnisse der Math. u. ihrer Grenzgebiete*, v.89, 1977. MR **58**#28182.
- [16] Ralph McKenzie, *A note on subgroups of infinite symmetric groups*, *Nederl. Akad. Wetensch. Proc. Ser. A74 = Indag. Math.* **33** (1971) 53–58. MR **45**#3545.
- [17] W. Sierpiński, *Sur une décomposition d’ensembles*, *Monatsh. Math.* **35** (1928) 239–248. <http://www.springerlink.com/index/G613664821643112.pdf>
- [18] R. M. Solovay and S. Tennenbaum, *Iterated Cohen extensions and Souslin’s problem*, *Ann. of Math.* (2) **94** (1971) 201–245. MR **45**#3212.
- [19] Friedrich Wehrung, *Embedding monoids of self-maps*, (in preparation, title tentative; July 2007 draft, 5 pp.).

- [20] Friedrich Wehrung, *Embedding coproducts of partition lattices*, (in preparation, title tentative; July 2007 draft, 10 pp.).
- [21] Philip M. Whitman, *Lattices, equivalence relations, and subgroups*, Bull. Amer. Math. Soc. **52** (1946) 507–522. MR [8,62b](#).

George M. Bergman
Department of Mathematics
University of California
Berkeley, CA 94720-3840
USA
gbergman@math.berkeley.edu