

# Hints

8.  $\overrightarrow{AA'} = 3\overrightarrow{AM}/2$ .
14.  $\cos^2 \theta \leq 1$ .
19. Rotate the regular  $n$ -gon through  $2\pi/n$ .
22. Take  $X = A$  first.
23.  $\overrightarrow{AB} = \overrightarrow{OB} - \overrightarrow{OA}$ .
24. If  $a + b + c + d = 0$ , then  $2(b + c)(c + d) = a^2 - b^2 + c^2 - d^2$ .
25.  $XA^2 = |\overrightarrow{OX} - \overrightarrow{OA}|^2 = 2R^2 - 2\langle \overrightarrow{OX}, \overrightarrow{OA} \rangle$  if  $O$  is the center.
26. Consider projections of the vertices to any line.
27. Project faces to an arbitrary plane and compute signed areas.
35. Compute  $(\cos \theta + i \sin \theta)^3$ .
37. Divide  $P$  by  $z - z_0$  with a remainder.
40.  $(z - z_0)(z - \bar{z}_0)$  is real.
41. Apply Vieta's formula.
45.  $e^{\pm i\theta} = \cos \theta \pm i \sin \theta$ .
46. See Figure 13.
50. Show first that  $\binom{n}{k}$  equals the number of  $k$ -element subsets in the set of  $n$  objects.
52. Show that the sum of distances to the foci of an ellipse from points on a tangent line is minimal at the tangency point; then reflect the source of light about the line.
54.  $e = |P'P|/|P'Q|$  is the slope of the secting plane.
55. Apply the forthcoming classification of quadratic curves.
56.  $AX^2 + CY^2$  is even in  $X$  and  $Y$ .
57.  $Q(-x, y) = Q(x, y)$  only if  $b = 0$ .
60.  $\cos \theta = 4/5$ .
62. Rotating the coordinate system as in Example on p. 11, don't forget to apply the change to the linear terms.

- 63.**  $Q(-x, -y) = Q(x, y)$  for a quadratic form  $Q$ .
- 65.** Reflect the plane about a suitable line.
- 68.**  $x^2 + 4xy = (x + 2y)^2 - 4y^2$ .
- 69.** Examine when the quadratic equation  $at^2 + 2bt + c = 0$  has a double root.
- 76.** Given that A and B can be transformed into C, find a way to transform A into B.
- 80.** For each of the normal forms, draw the region of the plane where the quadratic form is positive.
- 81.** Draw the region on the plane where  $xy > 0$ .
- 85.** First apply the Inertia Theorem to make  $Q = x_1^2 + x_2^2$ .
- 91.** Take  $x_1 = x$ ,  $x_2 = \dot{x}$ .
- 97.** In the “toy version,” one of the quadratic forms is  $x^2 + y^2$ , the squared Euclidean distance to the origin.
- 98.**  $4ac = (a + c)^2 - (a - c)^2$ .
- 102.**  $\frac{3n(n+1)}{2} - n^2 = \frac{n(n+3)}{2} > \frac{n^2}{2}$ .
- 103.**  $1 \neq 0$ .
- 105.** If  $ba = 1$  and  $ab' = 1$ , then  $b = bab' = b'$ .
- 110.** This is a rephrasing of the previous exercise.
- 114.** If  $(a, n) = 1$ , then  $ka + ln = 1$  for some integers  $k$  and  $l$ .
- 116.** Show that a non-zero homomorphism between two fields maps 1 to 1, and inverses to inverses.
- 118.** Construct the homomorphism  $f : \mathbb{Z} \rightarrow \mathbb{K}$  such that  $f(1) = 1$ .
- 119.** Take  $\mathbb{F} = \{0, 1, \eta, \zeta\}$ , and set  $1 + 1 = 0$ ,  $\eta + \zeta = 1$ ,  $\eta\zeta = 1$ .
- 120.**  $x^2 + x + 1 = (x + \eta)(x + \zeta)$ .
- 121.** Compute  $\mathbf{0} + \mathbf{0}'$ .
- 122.**  $-\mathbf{u} = (-1)\mathbf{u}$ .
- 123.** Redefine multiplication by scalars as  $\lambda\mathbf{u} = \mathbf{0}$  for all  $\lambda \in \mathbb{K}$  and all  $\mathbf{u} \in \mathcal{V}$ .
- 124.**  $0\mathbf{u} = (0 + 0)\mathbf{u} = 0\mathbf{u} + 0\mathbf{u}$ .
- 129.** Rotated parallelograms are still parallelograms.
- 132.** A linear function on  $\mathcal{V} \oplus \mathcal{W}$  is uniquely determined by its restrictions to  $\mathcal{V}$  and  $\mathcal{W}$ .
- 135.** Translate given linear subspaces by any vector in the intersection of the affine ones.
- 141.** Examine the kernel of the map  $\mathbb{R}[x] \rightarrow \mathbb{R}^2$  which associates to a polynomial  $P$  the pair  $(P(1), P(-1))$  of its values at  $x = \pm 1$ .
- 152.**  $\cos 4\theta + i \sin 4\theta = (\cos \theta + i \sin \theta)^4$ .

**154.** Prove that  $E$  is injective by identifying  $\mathcal{V}$  with  $\mathbb{K}^n$  and evaluating coordinate functions on a non-zero vector.

**158.** To a linear form  $f : \mathcal{V}/\mathcal{W} \rightarrow \mathbb{K}$ , associate  $\mathcal{V} \xrightarrow{\pi} \mathcal{V}/\mathcal{W} \xrightarrow{f} \mathbb{K}$ .

**160.**  $|\mathbb{F}| = p^n$ , where  $n$  is the dimension of  $\mathbb{F}$  as a  $\mathbb{Z}_p$ -vector space.

**167.** Both associate to an input  $x \in X$  the output  $w = f(g(h(x)))$ .

**176.** Pick two  $2 \times 2$ -matrices at random.

**179.** Compute  $BAC$ , where  $B$  and  $C$  are two inverses of  $A$ .

**180.** Given two isomorphisms  $B : \mathcal{U} \rightarrow \mathcal{V}$  and  $A : \mathcal{V} \rightarrow \mathcal{W}$ , find an isomorphism from  $\mathcal{W}$  to  $\mathcal{U}$ .

**181.** Consider  $1 \times 2$  and  $2 \times 1$  matrices.

**183.** Solve  $D^{-1}AC = I$  for  $D$  and  $C$ .

**187.** Which  $2 \times 2$ -matrices are anti-symmetric?

**191.** Start with any non-square matrix.

**201.** Each elementary product contains a zero factor.

**204.** There are 12 even and 12 odd permutations.

**206.** Permutation  $\begin{pmatrix} 1 & 2 & \dots & n-1 & n \\ n & n-1 & \dots & 2 & 1 \end{pmatrix}$  has length  $\binom{n}{2}$ .

**208.** Pairs of indices inverted by  $\sigma$  and  $\sigma^{-1}$  are the same.

**209.** The transformation: *logarithm*  $\rightarrow$  *lagorithm*  $\rightarrow$  *algorithm* consists of two transpositions.

**212.** Locate a pair  $(i, i+1)$  of nearby indices which  $\sigma$  inverts, compose  $\sigma$  with the transposition of these indices, and show that the length decreases by 1.

**217.**  $247 = 2 \cdot 100 + 4 \cdot 10 + 7$ .

**221.**  $\det(-A) = (-1)^n \det A$ .

**223.**  $\det(C^tQC) = (\det Q)(\det C)^2$ .

**234.** Apply the 1st of the “cool formulas.”

**235.** Compute the determinant of a  $4 \times 4$ -matrix the last two of whose rows repeat the first two.

**236.** Apply Binet–Cauchy’s formula to the  $2 \times 3$  matrix whose rows are  $\mathbf{x}^t$  and  $\mathbf{y}^t$ .

**237.** Show that  $\Delta_n = a_n\Delta_{n-1} + \Delta_{n-2}$ .

**239.** Use the defining property of **Pascal’s triangle**:  $\binom{m}{k} = \binom{m-1}{k} + \binom{m-1}{k-1}$ .

**240.** Use the fact of algebra that a polynomial in  $(x_1, \dots, x_n)$ , which vanishes when  $x_i = x_j$ , is divisible by  $x_i - x_j$ .

**241.** Divide the  $k$ th column by  $k$  and apply Vandermonde’s identity.

**243.**  $H : \mathbb{R}^7 \rightarrow \mathbb{R}^1$ .

- 244.** The space spanned by columns of  $A$  and  $B$  contains columns of  $A+B$ .
- 252.**  $(A^t \mathbf{a})(\mathbf{x}) = \mathbf{a}(A\mathbf{x})$  for all  $\mathbf{x} \in \mathbb{K}^n$  and  $\mathbf{a} \in (\mathbb{K}^m)^*$ .
- 266.** Modify the Gaussian elimination algorithm of Section 2 by permuting unknowns (instead of equations).
- 267.** Apply the  $LUP$  decomposition to  $M^t$ .
- 268.** Apply  $LPU$ ,  $LUP$ , and  $PLU$  decompositions to  $M^{-1}$ .
- 273.** When  $\mathbb{K}$  has  $q$  elements, each cell of dimension  $l$  has  $q^l$  elements.
- 277.** Consider the map  $\mathbb{R}^n \rightarrow \mathbb{R}^{p+q}$  defined by the linear forms.
- 290.**  $T = A + iB$  where both  $A$  and  $B$  are Hermitian.
- 291.**  $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{z}^\dagger \mathbf{w}$ .
- 293.**  $\langle \mathbf{z}, \mathbf{w} \rangle = \mathbf{z}^\dagger \mathbf{w}$ .
- 294.** Use the previous exercise.
- 296.** The normal forms are:  $i|Z_1|^2 - i|Z_2|^2$ ,  $|Z_1|^2 - |Z_2|^2$ ,  $|Z_1|^2$ .
- 301.** Take the linear form for one of new coordinates.
- 303.** Prove that  $-1$  is a non-square.
- 306.** There are 1 even and 3 odd non-degenerate forms.
- 310.**  $\frac{1}{3} = 1 - \frac{2}{3}$ .
- 311.** Start with inverting 2-adic units  $\cdots * * * 1$ . ( $*$  is a wild card).
- 313.** Use the previous exercise.
- 337.** Apply Vieta's formula.
- 339.** Compute  $\langle \mathbf{x}, A\mathbf{x} \rangle$  for  $\mathbf{x} = \sum t_i \mathbf{v}_i$ , where  $\{\mathbf{v}_i\}$  is an orthonormal basis of eigenvectors, and  $\sum t_i^2 = 1$ .
- 340.** If  $AB = BA$ , then eigenspaces of  $A$  are  $B$ -invariant.
- 345.** Find an eigenvector, and show that its orthogonal complement is invariant.
- 349.** Construct  $\mathbf{v}_1, \dots, \mathbf{v}_n$  by applying the Spectral Theorem to the non-negative Hermitian operator  $A^\dagger A$ , and check that  $A\mathbf{v}_i \in \mathcal{W}$  corresponding to distinct nonzero eigenvalues  $\mu_i$  of  $A^\dagger A$  are pairwise perpendicular.
- 350.** Rephrase the previous exercise in matrix form.
- 353.** The equation  $\langle \mathbf{x} + t\mathbf{y}, \mathbf{x} + t\mathbf{y} \rangle = 0$  quadratic in  $t$  has no real solutions unless  $\mathbf{x}$  is proportional to  $\mathbf{y}$ .
- 355.**  $\langle \mathbf{x}, \mathbf{x} \rangle = (\sum x_i)^2 + \sum x_i^2$ .
- 357.**  $\det(U^t U) = 1$ .
- 366.** See Example 1.
- 376.** Use the previous exercise.
- 378.** Prove first that the radius of the circle must be equal to the middle semiaxis of the ellipsoid.

**381.** For “exactly onne” claim, use the coordinate-less description of the spectrum provided by the minimax principle.

**382.** When  $x \approx 0$ ,  $\sin x \approx x$ .

**385.** If  $T$  denotes the cyclic shift of coordinates in  $\mathbb{C}^n$ , then  $C = C_0I + C_1T + \cdots + C_{n-1}T^{n-1}$ .

**398.** Use Newton’s binomial formula.

**411.** Pick  $A$  and  $B$  so that  $e^A e^B \neq e^B e^A$ .

**414.** Verify it first for a Jordan cell.

**419.** Make two of the lines the coordinate axes on the plane, and the third one the graph of an isomorphism between them.

**420.** If the first three lines are  $y = 0$ ,  $x = 0$ , and  $y = x$ , then the fourth one is  $y = \lambda x$ .

**421.** If  $d = 0$ ,  $a = \infty$ ,  $b = 1$ , then  $\lambda = c$ .

**423.** Assuming that the spaces  $\mathcal{U}_i$  have the same dimension, and the maps between them are invertible, consider the determinant (or eigenvalues) of the composition of the maps around the cycle.

**426.** Use the theory of Bruhat cells.

**427.** Use Lemma, and apply induction on the length of the shortest of the legs.

**428.** Use Sylvester’s rule together with the previous exercise.

**431.** On the unit sphere of area  $4\pi$ , the area of a triangle with the angles  $(\frac{\pi}{p}, \frac{\pi}{q}, \frac{\pi}{r})$  has the area  $\frac{\pi}{p} + \frac{\pi}{q} + \frac{\pi}{r} - \pi$ . Use this to compute the number of required triangles.

**433.** Compute  $R_{\mathbf{v}_1} R_{\mathbf{v}_2}$  to show that it is a Jordan cell of size 2 with the eigenvalue 1.

**434.** They correspond to positive roots  $\mathbf{e}_i - \mathbf{e}_j = \mathbf{v}_{i+1} + \cdots + \mathbf{v}_j$ , where  $0 \leq i < j \leq n$ .



# Answers

1.  $mg/2, mg\sqrt{3}/2$ .
2. 18 min (reloading time excluded).
6. It rotates with the same angular velocity along a circle centered at the barycenter of the triangle formed by the centers of the given circles.
10.  $3/4$ .
13.  $7\vec{OA} = \vec{OA}' + 2\vec{OB}' + 4\vec{OC}'$  for any  $O$ .
15.  $3/2$ .
17.  $2\langle \mathbf{u}, \mathbf{v} \rangle = |\mathbf{u} + \mathbf{v}|^2 - |\mathbf{u}|^2 - |\mathbf{v}|^2$ .
18. (b) No.
28. Yes; Yes (0); Yes.
29. (a)  $\frac{1+5i}{13}$ ; (b)  $\frac{1-i\sqrt{3}}{2}$ .
31.  $|z|^{-2}$ .
33. (a)  $\sqrt{2}, -\pi/4$ ; (b)  $2, -\pi/3$ .
34.  $\frac{-1+i\sqrt{3}}{2}$ .
38.  $2 \pm i; i\frac{1 \pm \sqrt{5}}{2}; 1 + i, 1 + i; 1 \pm \frac{\sqrt{6-i\sqrt{2}}}{2}$ .
39.  $-2, 1 \pm i\sqrt{3}; i, \frac{\pm\sqrt{3}-i}{2}; \pm i\sqrt{2}, \pm i\sqrt{2}; \frac{\sqrt{3 \pm i}}{2}, \frac{\sqrt{3 \pm i}}{2}; \pm i, \pm \frac{\sqrt{3 \pm i}}{2}$ .
42.  $a_k = (-1)^k \sum_{1 \leq i_1 < \dots < i_k \leq n} z_{i_1} \cdots z_{i_k}$ .
45.  $\cos \theta = (e^{i\theta} + e^{-i\theta})/2, \sin \theta = (e^{i\theta} - e^{-i\theta})/2i$ .
47. The upper half-plane  $\text{Im } w > 0$ .
49.  $\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta, \sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$ .
58. Yes,  $k(x^2 + y^2)$ .
59.  $y = \pm x$ ; level curves of (a) are ellipses, of (c) hyperbolas.
60.  $2X^2 - Y^2 = 1$ .
62. Semiaxes  $2/\sqrt{3}$  and 2.

64. (a)  $\pm(\sqrt{\alpha^2 - \beta^2}, 0)$ , (b)  $\pm(\sqrt{\alpha^2 + \beta^2}, 0)$ .
68. 2nd ellipse, 1st & 4th hyperbolas.
72. A pair of intersecting lines,  $x - 1 = \pm(2y - 1)$ .
79. For normal forms,  $X^2$  and 0 can be taken.
84.  $(n + 1)(n + 2)/2$ .
88.  $x(t) = x(0)e^{\lambda t}$ .
89.  $x(t) = x(0)e^{3t}$ ,  $y(t) = y(0)e^{-t}$ ,  $z(t) = z(0)$ .
92.  $\dot{X}_1 = iX_1$ ,  $\dot{X}_2 = -iX_2$ .
96. Yes, some non-equivalent equations represent the empty set.
101.  $\frac{n(n+1)}{2}$ .
107. Use the **Euclidean algorithm**: divide  $a$  by  $b$  with the remainder  $r$ , then divide  $b$  by  $r$  with the remainder  $r_1$ , then divide  $r$  by  $r_1$  with the remainder  $r_2$ , etc. until the remainder becomes 0.
108. Use the previous exercises.
111. n.
113. 1, 3, 5, 7.
134. Points, lines, and planes.
137.  $p^n$ ;  $p^{mn}$ .
138.  $p^{n(n-1)/2}$ .
140. (b)  $\text{Ker } D = \mathbb{K}[x^p] \subset \mathbb{K}[x]$ .
149.  $\dim = 2$ .
151.  $x^k = x_1^k L_1(x) + \cdots + x_n^k L_n(x)$ ,  $k = 0, \dots, n - 1$ .
155. (a)  $n$ ; (b)  $n(n + 1)/2$ .
156. 2, if  $n > 1$ .
161. yes, no, no, yes, no, yes, yes, no.
163.  $E = \mathbf{va}$ , i.e.  $e_{ij} = v_i a_j$ ,  $i = 1, \dots, m$ ,  $j = 1, \dots, n$ .
164.  $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ .
166. Check your answers using associativity, e.g.  $(CB)A = C(BA)$ .

$$\text{Other answers: } BAC = \begin{bmatrix} 3 & 1 \\ 3 & 3 \end{bmatrix}, \quad ACB = \begin{bmatrix} 0 & -2 \\ 3 & 6 \end{bmatrix}.$$

$$172. \begin{bmatrix} \cos 1^\circ & -\sin 1^\circ \\ \sin 1^\circ & \cos 1^\circ \end{bmatrix}.$$

173. If  $B = A^k$ , then  $b_{i,i+k} = 1$ , and all other  $b_{ij} = 0$ .
174. The answer is the same as in the previous problem.



175. (a) If  $A$  is  $m \times n$ ,  $B$  must be  $n \times m$ . (b) Both must be  $n \times n$ .
177. Exactly when  $AB = BA$ .
184. Restriction of linear functions on  $\mathcal{W}$  to the subspace  $\mathcal{V}$ .
186. I.
188.  $S = 2x_1y_1 + x_1y_2 + x_2y_1$ ,  $A = x_1y_2 - x_2y_1$ .
189. No, only if  $AB = BA$ .
192. No; yes (of  $x, y \in \mathbb{R}^1$ ); yes (in  $\mathbb{R}^2$ ).
193.  $(\sum x_i)(\sum y_i)$  and  $\sum_{i \neq j} x_i y_j / 2$ .
195.  $T^\dagger = \overline{z_2} w_1$ ;  $S_1 = (\overline{z_1} w_2 + \overline{z_2} w_1) / 2$ ,  $S_2 = (\overline{z_1} w_2 - \overline{z_2} w_1) / 2i$ ;  
 $H_1 = (\overline{z_1} z_2 + \overline{z_2} z_1) / 2$ ,  $H_2 = (\overline{z_1} z_2 - \overline{z_2} z_1) / 2i$ .
196.  $(\overline{z_1} w_2 + \cdots + \overline{z_{n-1}} w_n + \overline{z_2} w_1 + \cdots + \overline{z_n} z_{n-1}) / 2$ .
197.  $T \mapsto C^\dagger T D$ .
199. No, yes,  $n^2$ .
202. 1; 1;  $\cos(\mathbf{x} + \mathbf{y})$ .
203. 2, 1; -2.
205.  $k(k-1)/2$ .
207.  $n(n-1)/2 - l$ .
211. Transpositions  $\tau^{(i)}$  of nearby indices  $(i, i+1)$ ,  $i = 1, \dots, n-1$ .
213. E.g.  $\tau_{14}\tau_{34}\tau_{25}$ .
214. No; yes.
215. + (6 inverted pairs); + (8 inverted pairs).
216. -1 522 200; -29 400 000.
218. 0.
219. Changes sign, if  $n = 4k + 2$  or  $n = 4k + 3$  for some  $k$ , and remains unchanged otherwise.
220.  $x = a_1, \dots, a_n$ .
222. Leave it unchanged.
226. (a)  $(-1)^{n(n-1)/2} a_1 a_2 \cdots a_n$ , (b)  $(ad - bc)(eh - fg)$ .
227. (a) 9, (b) 5.
228. (a)  $\begin{bmatrix} -\frac{5}{18} & \frac{7}{18} & \frac{1}{18} \\ \frac{1}{18} & -\frac{5}{18} & \frac{7}{18} \\ \frac{7}{18} & \frac{1}{18} & -\frac{5}{18} \end{bmatrix}$ , (b)  $\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}$ .
229. (a)  $\mathbf{x} = [-\frac{2}{9}, \frac{1}{3}, \frac{1}{9}]^t$ , (b)  $\mathbf{x} = [1, -1, 1]^t$ .
231.  $(\det A)^{n-1}$ , where  $n$  is the size of  $A$ .
232. Those with determinants  $\pm 1$ .
233. (a)  $x_1 = 3, x_2 = x_3 = 1$ , (b)  $x_1 = 3, x_2 = 4, x_3 = 5$ .
234. (a)  $-x_1^2 - \cdots - x_n^2$ , (b)  $(ad - bc)^3$ .

**238.**  $\lambda^n + a_1\lambda^{n-1} + \cdots + a_{n-1}\lambda + a_n$ .

**239.** 1.

**241.**  $1!3!5! \cdots (2n-1)!$ .

**243.** 1.

**245.** The system is consistent whenever  $b_1 + b_2 + b_3 = 0$ .

**251.** (a) Yes, (b) yes, (c) yes, (d) no, (e) yes, (f) no, (g) no, (h) yes.

**255.**  $0 \leq \text{codim} \leq k$ .

**257.** Two subspaces are equivalent if and only if they have the same dimension.

**258.** The equivalence class of an ordered pair  $\mathcal{U}, \mathcal{V}$  of subspaces is determined by  $k := \dim U$ ,  $l := \dim \mathcal{V}$ , and  $r := \dim(\mathcal{U} + \mathcal{V})$ , where  $k, l \leq r \leq n$  can be arbitrary.

**259.** (a)  $q^n$ ; (b)  $(q^n - 1)(q^n - q)(q^n - q^2) \cdots (q^n - q^{n-1})$ ;

(c)  $(q^n - 1)(q^n - q)(q^n - q^2) \cdots (q^n - q^{r-1})$ ;

(d)  $(q^n - 1)(q^n - q) \cdots (q^n - q^{r-1}) / (q^r - 1)(q^r - q) \cdots (q^r - q^{r-1})$ .

**260.**  $x_1 = 3, x_2 = 1, x_3 = 1$ ; inconsistent;  $x_1 = 1, x_2 = 2, x_3 = -2$ ;

$x_1 = 2t_1 - t_2, x_2 = t_1, x_3 = t_2, x_4 = 1$ ;

$x_1 = -8, x_2 = 3 + t, x_3 = 6 + 2t, x_4 = t$ ;  $x_1 = x_2 = x_3 = x_4 = 0$ ;

$x_1 = \frac{3}{17}t_1 - \frac{13}{17}t_2, x_2 = \frac{19}{17}t_1 - \frac{20}{17}t_2, x_3 = t_1, x_4 = t_2$ ;

$x_1 = -16 + t_1 + t_2 + 5t_3, x_2 = 23 - 2t_1 - 2t_2 - 6t_3, x_3 = t_1, x_4 = t_2, x_5 = t_3$ .

**261.**  $\lambda = 5$ .

**262.** (a)  $\text{rk} = 2$ , (b)  $\text{rk} = 2$ , (c)  $\text{rk} = 3$ , (d)  $\text{rk} = 4$ , (e)  $\text{rk} = 2$ .

**263.** Inverse matrices are:

$$\begin{bmatrix} 1 & -4 & -3 \\ 1 & -5 & -3 \\ -1 & 6 & 4 \end{bmatrix}, \quad \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}, \quad \begin{bmatrix} 2 & -1 & 0 & 0 \\ -3 & 2 & 0 & 0 \\ 31 & -19 & 3 & -4 \\ -23 & 14 & -2 & 3 \end{bmatrix}.$$

**271.**  $\binom{n}{2} - l(\sigma)$ , where  $l(\sigma)$  is the length of the permutation.

**272.**  $[n]_q! := [1]_q[2]_q \cdots [n]_q$  (called **q-factorial**), where  $[k]_q := \frac{q^k - 1}{q - 1}$ .

**274.** Inertia indices  $(p, q) = (1, 1), (2, 0), (1, 2)$ .

**279.** Empty for  $p = 0$ , has 2 components for  $p = 1$ , and 1 for  $p = 2, 3, 4$ .

**283.**  $z_1^2 + z_2^2 = 1, z_1^2 + z_2^2 = 0, z_2 = z_1^2, z_1^2 = 1, z_1^2 = 0$ .

**284.** Two parallel lines.

**289.** Those all of whose entries are imaginary.

**290.**  $T(\mathbf{z}, \mathbf{w}) =$

$$\frac{1}{2} [T(\mathbf{z} + \mathbf{w}, \mathbf{z} + \mathbf{w}) + iT(\mathbf{iz} + \mathbf{w}, \mathbf{iz} + \mathbf{w}) - (1 + i)(T(\mathbf{z}, \mathbf{z}) + T(\mathbf{w}, \mathbf{w}))].$$

**297.**  $d_{ii} = \Delta_i / \Delta_{i-1}$ .

**298.**  $(p, q) = (2, 2), (3, 1)$ .

**316.** No.

**317.** The sum of the squares of all sides of a parallelogram is equal to the sum of the squares of the diagonals.

**330.** id.

**354.**  $\theta(\mathbf{e}_i, \mathbf{e}_{i+1}) = 2\pi/3$ , while all other pairs are perpendicular.

**356.** One can take  $\mathbf{f}_i = \mathbf{e}_i - \mathbf{e}_0$ , and  $\mathbf{h}_i = \mathbf{e}_i - \mathbf{e}_{i-1}$ , where  $i = 1, \dots, n$ .

**358.** Rotations about the origin, and reflections about a line passing through the origin, which have determinants 1 and  $-1$  respectively.

**364.**  $\pm 1$ .

**371.** The operator is  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ .

**397.** (a)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}$  (b)  $\begin{bmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  (c)  $\begin{bmatrix} -3 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$   
 (e)  $\begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$  (f)  $\begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$  (k)  $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$  (l)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & -i \end{bmatrix}$ .

**407.**  $(a_0 - n(a_0 - a_1/3))3^n$ .

**408.** (a)  $(1 + \sqrt{10})/3$ .

**411.**  $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$  and  $B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$  will do.

**418.** Pairs of  $m \times n$ -matrices; pairs of matrices of transposed sizes.

**422.**  $\mathbb{K}^1$ .

**424.** The restriction to the eigenline.

**429.** (a)  $(2, 2, r)$ ,  $(2, 3, 3)$ ,  $(2, 3, 4)$ ,  $(2, 3, 5)$ ; (b)  $(3, 3, 3)$ ,  $(2, 4, 4)$ ,  $(2, 3, 6)$ .

**431.** Symmetry planes of regular polyhedron (tetrahedron in case (b), octahedron or cube in case (c), and icosahedron or dodecahedron in case (d)) partition the surface of the sphere circumscribed around the polyhedron into (respectively, 24, 48, and 120) required spherical triangles.

**432.** Since  $\langle \mathbf{v}, \mathbf{v} \rangle = 2$ , we have:

$$\langle \mathbf{x} - \langle \mathbf{x}, \mathbf{v} \rangle \mathbf{v}, \mathbf{y} - \langle \mathbf{y}, \mathbf{v} \rangle \mathbf{v} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle - \langle \mathbf{x}, \mathbf{v} \rangle \langle \mathbf{v}, \mathbf{y} \rangle - \langle \mathbf{y}, \mathbf{v} \rangle \langle \mathbf{x}, \mathbf{v} \rangle + 2 \langle \mathbf{x}, \mathbf{v} \rangle \langle \mathbf{y}, \mathbf{v} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle.$$

**434.**  $\mathcal{V}^{i,j} : \dots \rightarrow 0 \rightarrow \mathbb{K}^1 \xrightarrow{\cong} \dots \xrightarrow{\cong} \mathbb{K}^1 \rightarrow 0 \rightarrow \dots$  ( $i$  zeroes followed by  $j - i$  copies of  $\mathbb{K}^1$ , followed by  $n - j$  zeroes).

**435.**  $\mathcal{V}^{0,n} \oplus \mathcal{V}^{1,n} \oplus \dots \oplus \mathcal{V}^{n-1,n}$ .

**436.**  $X(t) = e^{tA} X(0) e^{-tA}$ .

**437.** Eigenvalues are  $\lambda_i - \lambda_j$ ,  $i, j = 1, \dots, n$ . In the space of linear forms, they form the root system of type  $A_{n-1}$ . Eigenvectors are matrices  $E_{ij}$  whose all entries are 0 except a single 1 in the  $i$ th row and  $j$ th column.



# Index

- $n$ -space, 43
- $n$ -vectors, 43
  
- absolute value, 12
- addition of vectors, 39
- additivity, 7
- adjoint form, 70
- adjoint map, 65, 96
- adjoint systems, 99
- adjugate matrix, 84
- affine subspace, 45, 49
- algebraic number, 40
- algebraically closed field, 170
- annihilator, 57
- anti-Hermitian form, 71, 122
- anti-Hermitian quadratic form, 122
- anti-linear, 70
- anti-symmetric bilinear form, 68
- anti-symmetric matrix, 68
- anti-symmetric operator, 151
- argument of complex number, 13
- associative, 62
- associativity, 4
- augmented matrix, 102
- axiom, 39
- axis of symmetry, 25
  
- Bézout, 17
- Bézout's theorem, 17
- back substitution, 102
- barycenter, 4
- basis, 5, 51
- bijective, 44
- bilinear form, 67
- bilinearity, 8
- Binet, 89
- Binet–Cauchy formula, 89
- block, 79
- block triangular matrix, 79
- Bruhat cell, 113
  
- canonical form, 29
- canonical isomorphism, 56
- canonical projection, 49
- Cartesian coordinates, 9
- category of vector spaces, 44
- Cauchy, 89
- Cauchy – Schwarz inequality, 9, 135
- Cauchy's interlacing theorem, 156
- Cayley transform, 139
- characteristic, 42
- characteristic equation, 140
- characteristic polynomial, 35, 163
- Chebyshev, 55
- Chebyshev polynomials, 55
- classification theorem, 29
- codimension, 98
- cofactor, 83
- cofactor expansion, 83, 87
- column, 47
- column space, 105
- commutative square, 95, 189
- commutativity, 4
- commutator, 124
- complementary multi-index, 87
- complete flag, 111
- completing squares, 26
- complex conjugate, 11
- complex conjugation, 149
- complex multiplication, 153
- complex sphere, 120
- complex vector space, 40
- complexification, 149
- components, 47
- composition, 61
- congruent modulo  $n$ , 41

- conic section, 21  
 conics, 119  
 connected graph, 187  
 continuous time, 175  
 coordinate Euclidean space, 147  
 coordinate flag, 112  
 coordinate system, 5  
 coordinate vectors, 43, 47  
 coordinates, 5, 43, 47, 53  
 Cramer's rule, 85  
 cross product, 90  
 cross-ratio, 193  
 cylinder, 119
- Dandelin, 22  
 Dandelin's spheres, 21  
 Darboux, 130  
 Darboux basis, 130  
 decomposable representation, 190  
 degenerate bilinear form, 155  
 Descartes, 9  
 determinant, 37, 73  
 diagonal matrix, 47  
 diagonalizable matrix, 169  
 dimension, 52  
 dimension of Bruhat cell, 113  
 direct sum, 32, 44, 190  
 direct sum of representations, 190  
 directed segment, 3  
 directrix, 23  
 discrete dynamical systems, 175  
 discrete time, 175  
 discriminant, 14, 133  
 distance, 135  
 distributive law, 12, 60  
 distributivity, 4  
 dot product, 7  
 dot-product, 67  
 dual basis, 54  
 dual map, 65  
 dual space, 46
- eccentricity, 23  
 edge, 187  
 eigenspace, 141, 163  
 eigenvalue, 33, 140, 163  
 eigenvector, 140, 163  
 elementary product, 73
- elementary row operations, 101  
 ellipse, 22  
 ellipsoid, 157  
 entry, 47  
 equivalent, 29  
 equivalent conics, 119  
 equivalent linear maps, 95  
 equivalent representations, 188  
 Euclidean algorithm, 208  
 Euclidean inner product, 147  
 Euclidean space, 147  
 Euclidean structure, 147  
 Euler's formula, 18  
 evaluation, 60  
 evaluation map, 48  
 even form, 129  
 even permutation, 75  
 exponential function, 17
- Fibonacci, 175  
 Fibonacci sequence, 175  
 field, 12, 40  
 field of  $p$ -adic numbers, 132  
 finite dimensional spaces, 52  
 flag, 111  
 focus of ellipse, 22  
 focus of hyperbola, 23  
 focus of parabola, 23  
 Fourier, 160  
 Fourier basis, 160  
 Fundamental Formula of Mathematics, 19
- Gabriel, 191  
 Gaussian elimination, 101  
 Gelfand's problem, 155  
 golden ratio, 176  
 Gram matrix, 137  
 Gram-Schmidt process, 126, 137  
 graph, 45, 187  
 gravitation constant, 162  
 greatest common divisor, 41  
 group, 198
- half-linear, 70  
 Hamilton-Cayley equation, 171  
 harmonic oscillator, 159  
 Hasse, 132

- head, 3
- Hermite, 70
- Hermite polynomials, 55
- Hermitian adjoint, 138
- Hermitian adjoint form, 70
- Hermitian conjugate matrix, 70
- Hermitian form, 71
- Hermitian inner product, 135
- Hermitian isomorphic, 136
- Hermitian operator, 138
- Hermitian quadratic form, 71, 121
- Hermitian space., 135
- Hermitian-anti-symmetric form, 70
- Hermitian-symmetric form, 70
- Hilbert space, 135
- homogeneity, 7
- homogeneous system, 98
- homomorphism, 41
- homomorphism theorem, 50
- hyperbola, 22
- hyperplane, 100
- hypersurface, 119
  
- identity matrix, 63
- identity permutation, 75
- imaginary part, 11
- imaginary unit, 11
- inconsistent system, 103
- indecomposable representation, 190
- indices in inversion, 75
- induction hypothesis, 53
- inertia index, 117
- Inertia Theorem, 28
- injective, 44
- inner product, 7
- invariant subspace, 141
- inverse matrix, 63
- inverse transformation, 64
- inversion of indices, 75
- involution, 149
- irreducible representation, 191
- isometric Hermitian spaces, 136
- isomorphic spaces, 44
- isomorphism, 44
- iterations of linear maps, 175
  
- Jordan block, 35
- Jordan canonical form, 168
  
- Jordan cell, 168
- Jordan normal form, 168
- Jordan system, 35
  
- kernel, 44
- kernel of form, 118, 121
- kinetic energy, 158
  
- Lagrange, 87
- Lagrange polynomials, 55
- Lagrange's formula, 87
- law of cosines, 9
- LDU decomposition, 110
- leading coefficient, 102
- leading entry, 102
- leading minors, 124
- left inverse, 96
- length, 135
- length of permutation, 75
- linear combination, 4
- linear dynamical systems, 175
- linear form, 45, 59
- linear function, 45, 59
- linear map, 44
- linear ODE, 175
- linear recursion relation, 175
- linear subspace, 43
- linear transformation, 64
- linearly dependent, 52
- linearly independent, 52
- Linnaeus, 29
- lower triangular, 108
- lower-triangular matrix, 47
- LPU decomposition, 108
- LU decomposition, 110
- LUP decomposition, 111
  
- Möbius band, 56
- mathematical induction, 53
- matrix, 47, 59
- matrix entry, 59
- matrix product, 60, 61
- metric space, 135
- Minkowski, 132
- Minkowski–Hasse theorem, 132
- minor, 83, 87
- multi-index, 87
- multiplication by scalar, 3

- multiplication by scalars, 39  
 multiplicative, 12  
 multiplicity, 14  
  
 nilpotent operator, 165  
 non-degenerate bilinear form, 155  
 non-degenerate Hermitian form, 124  
 nontrivial linear combination, 52  
 normal form, 29  
 normal operator, 138, 151  
 null space, 44, 105  
  
 odd form, 129  
 odd permutation, 75  
 ODE, 175  
 operator, 138  
 opposite coordinate flag, 112  
 opposite vector, 39  
 oriented graph, 187  
 orthogonal, 136  
 orthogonal basis, 115  
 orthogonal complement, 141  
 orthogonal diagonalization, 145  
 Orthogonal Diagonalization Theorem, 155  
 orthogonal projection, 137  
 orthogonal projector, 139  
 orthogonal transformation, 147  
 orthogonal vectors, 9  
 orthonormal basis, 125, 136, 147  
  
 parabola, 27  
 partition, 167  
 Pascal's triangle, 203  
 pendulum, 162  
 permutation, 73  
 permutation matrix, 108  
 phase plane, 159  
 pivot, 102  
 Plücker, 90  
 Plücker coordinates, 98  
 Plücker identity, 90  
 PLU decomposition, 111  
 polar, 13  
 polar decomposition, 146  
 positive definite, 116  
 positive operator, 145  
 positivity, 7  
  
 potential energy, 158  
 power of matrix, 65  
 principal axes, 25, 157  
 principal minor, 117  
 projection, 136  
 Pythagorean theorem, 9  
  
 q-factorial, 114, 210  
 quadratic curve, 21  
 quadratic form, 23, 31, 69  
 quadratic formula, 14  
 quiver, 187  
 quotient space, 49  
  
 range, 44  
 rank, 31, 93  
 rank of linear system, 98  
 rank of matrix, 94  
 real normal operators, 148  
 real part, 11  
 real spectral theorem, 151  
 real vector space, 40  
 realification, 149  
 reduced row echelon form, 102  
 reducible representation, 191  
 reflection, 198  
 regular nilpotent, 165  
 representation, 187  
 right inverse, 96  
 root of unity, 15  
 root space, 164  
 root system, 199  
 row echelon form, 102  
 row echelon form of rank  $r$ , 102  
 row space, 105  
  
 scalar, 39, 40  
 scalar product, 7  
 semiaxes, 157  
 semiaxis of ellipse, 26  
 sesquilinear form, 70, 121  
 sign of permutation, 73  
 similarity, 163  
 similarity transformation, 64  
 simple problems, 37  
 simple quiver, 191  
 singular value decomposition, 145  
 span, 51



- spectral theorem, 140
- spectrum, 33, 155
- square matrix, 47, 63
- square root, 15, 145
- standard basis, 51
- standard coordinate flag, 111
- standard coordinate space, 43
- standard Euclidean space, 71
- standard Hermitian space, 71
- subrepresentation, 191
- subspace, 43
- surjective, 44
- Sylvester, 124
- symmetric bilinear form, 68, 115
- symmetric matrix, 68
- symmetric operator, 151
- symmetricity, 7
- system of linear equations, 61
  
- tail, 3
- tautology, 50
- time-independent dynamical system, 175
- total anti-symmetry, 77
- trace, 139
- transition matrix, 63
- transposed form, 68
- transposed map, 65
- transposed matrix, 66
- transposed partition, 167
- transposition matrix, 109
- transposition permutation, 75
- triangle inequality, 9, 135
  
- unipotent, 168
- unipotent matrix, 110
- unit coordinate vectors, 51
- unit vector, 8
- unitary rotation, 143
- unitary space, 135
- unitary transformation, 139
- upper triangular, 108
- upper-triangular matrix, 47
  
- Vandermonde, 91
- Vandermonde's identity, 91
- vector, 3, 39
- vector space, 36, 39
- vector subspace, 43
- vector sum, 3
- vertex, 187
- Vieta, 17
- Vieta's theorem, 16
  
- Young tableaux, 167
  
- zero representation, 190
- zero vector, 3, 39