

## Quiz 11. Discussion Section 103. Math 110 Fall 2014.

Name: Solution

1. Consider the following permutation matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}.$$

Let  $X = -I_3 + 2A - 2A^2$ , where  $I_3$  is the  $3 \times 3$  identity matrix. Determine the eigenvalues  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{C}$  and the corresponding eigenspaces.

**Solution:** The given matrix satisfies  $A^3 = I_3$  (notice that  $A$  is a permutation matrix). If  $\lambda$  is an eigenvalue of  $A$  and  $Av = \lambda v$ , with  $v \neq 0$ , then

$$Xv = (-I_3 + 2A - 2A^2)v = -v + 2Av - 2A^2v = (-1 + 2\lambda - 2\lambda^2)v.$$

Here we have used the fact: if  $Av = \lambda v$ , with  $v \neq 0$ , then  $A^n v = \lambda^n v$ .

Hence, if  $\lambda$  is an eigenvalue of  $A$  then  $(-1 + 2\lambda - 2\lambda^2)$  is an eigenvalue of  $X$ . You can check that  $A$  has characteristic polynomial  $-\lambda^3 + 1$ , so its eigenvalues are  $\lambda_1 = 1, \lambda_2 = e^{2\pi i/3}, \lambda_3 = e^{4\pi i/3}$  (the cube roots of unity). Also, you can show that (by row-reduction)

$$\text{nul}(A - \lambda_1 I_3) = \text{span}((1, 1, 1)^t), \quad \text{nul}(A - \lambda_2 I_3) = \text{span}((\lambda_1, \lambda_1^2, 1)^t),$$

$$\text{nul}(A - \lambda_3 I_3) = \text{span}((\lambda_2, \lambda_2^2, 1)^t).$$

Hence,  $X$  must have the eigenvalues  $-1 + 2\lambda_i - 2\lambda_i^2$ , for  $i = 1, 2, 3$ , and with the same eigenspaces as above.