

QUIZ, SEP 25

NAME:


**Directional Derivatives (8 pts).** Compute the gradient of

$$f(x, y) = x^2 - y^2$$

and use it to determine the directional derivative in the direction  $\hat{u} = \langle \frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}} \rangle$  at the point  $(1, 1)$ .

**Derivatives and Contour Plots (6 pts).**

- Mark with  $\circ$  a point on this contour plot where the gradient vector is  $\langle 1, 0 \rangle$ .<sup>1</sup>
- Mark with  $\times$  a point on this contour plot where the derivative in the  $\hat{u} = \langle 1, 0 \rangle$  direction is 0.



<sup>1</sup>Be sure to think about both the direction and magnitude of the gradient; in particular, pay attention to the values of  $f$ .

**Chain Rule.** (6 pts) Let  $x(t) = \cos(t)$  and  $y(t) = \sin(t)$ . Suppose that for all  $t$ ,  $\frac{d}{dt}f(x(t), y(t)) = 0$ . Additionally, suppose that  $|\nabla f| = 1$ . Find the gradient vector at the point  $(1, 0)$ . <sup>2</sup>

**Bonus Problem.** *Worth no points!* There is a landscape whose elevation is described by the function  $f(x, y)$ . A hiker walks along a loop described by  $\langle x(t), y(t) \rangle$ . As they walk, they record the gradient vector of this hill, and use this to produce a new vector valued function

$$\vec{\kappa}(t) := \frac{\nabla f|_{(x(t), y(t))}}{\left| \nabla f|_{(x(t), y(t))} \right|}$$

(So, think of this as the function which records at time  $t$  the direction of the gradient that the hiker encounters. ) The hiker notices that

$$\kappa(t) = \langle \cos t, \sin t \rangle.$$

Argue why the hiker must have walked around a maxima or minima of  $f$ .

---

<sup>2</sup>Hint: Expand  $\frac{d}{dt}f(x(t), y(t))$  using the chain rule. Notice that  $(x(0), y(0)) = (1, 0)$ .