

Representation Theory of Symmetric Groups

Cambridge Part III, Michaelmas 2022

Taught by Stacey Law

Notes taken by Leonard Tomczak

Contents

1	Introduction	2
1.1	Motivation	2
1.2	Background	2
1.2.1	Representations & modules	3
1.2.2	Some Linear Algebra	4
1.2.3	Character Theory	5
2	Specht Modules	6
2.1	The Symmetric Group	6
2.2	Irreducible modules	11
2.3	Standard Basis Theorem	18
3	Character Theory	23
3.1	Hook Length Formula	23
3.2	The Determinantal Form	29
3.3	Applications	42
3.3.1	Young's Rule Revisited	42
3.3.2	Branching Rule	46
3.3.3	Murnaghan-Nakayama Rule	48
4	McKay Numbers	53
4.1	James's Abacus	53
4.2	Towers	65
4.3	The McKay Conjecture	74
	Bibliography	76

1 Introduction

1.1 Motivation

- Representation theory of finite groups: active area of research
- Many open problems, e.g. Local-Global Conjectures

Definition. Let G be a finite group, p a prime. Then we let

- $\text{Irr}(G) := \{\text{irreducible characters of } G\}$,
- $\text{Irr}_{p'}(G) := \{\chi \in \text{Irr}(G) \mid p \nmid \chi(1)\}$.

Conjecture (McKay 1972). Let G be a finite group, p a prime, P a Sylow p -subgroup of G . Then

$$|\text{Irr}_{p'}(G)| = |\text{Irr}_{p'}(N_G(P))|.$$

The case $p = 2$ has been proved in 2016.

Theorem 1.1 (Olsson 1976). *The McKay Conjecture holds for all symmetric groups S_n and all primes p .*

Outline of the course:

- Chapter 1: Introduction and background
- Chapter 2: Specht modules ([Jam78])
- Chapter 3: Character theory ([JK84])
- Chapter 4: McKay numbers ([Ols94])

1.2 Background

Notation.

- $\mathbb{N} = \{1, 2, 3, \dots\}$, $\mathbb{N}_0 = \{0, 1, 2, \dots\}$.
- If $n \in \mathbb{N}$, let $[n] := \{1, 2, \dots, n\}$.
- $\text{Irr}(G)$ (or $\text{Irr}_{\mathbb{F}}(G)$ to specify the field \mathbb{F}) is a complete set of irreducible representations of G over \mathbb{F} .

1.2.1 Representations & modules

\mathbb{F} will denote an arbitrary field and G a finite group. All modules considered in this course will be finite-dimensional left modules.

A (finite-dimensional) *representation of G over \mathbb{F}* is a group homomorphism $\rho : G \rightarrow \text{GL}(V)$, where V is a (finite-dimensional) vector space over \mathbb{F} . We write $g \cdot v$ for $\rho(g)(v)$. Equivalently a representation is an $\mathbb{F}G$ -module. The *degree* or *dimension* of a representation is the dimension of the underlying vector space.

Example. The (one-dimensional) *trivial* representation of G is a one-dimensional vector space with trivial G -action. It will be denoted by $\mathbb{1}_G$.

Other concepts.

- *Subrepresentations* W of V , written $W \leq V$
- *Simple* or *irreducible* modules, i.e. those with no proper non-zero submodules.
- *Semisimple* or *completely reducible* modules, i.e. direct sums of simple modules.
- *Decomposable* modules, i.e. modules decomposing into a direct sum of proper submodules; opposite: *indecomposable*.
- *G -homomorphisms*: If V, W are G -modules, then an \mathbb{F} -linear map $\theta : V \rightarrow W$ is a G -homomorphism if $g \cdot \theta(v) = \theta(g \cdot v)$ for all $g \in G, v \in V$.

Useful results.

Lemma 1.2 (Schur's Lemma). *Let V, W be simple G -modules, $\theta : V \rightarrow W$ a G -homomorphism. Then $\theta = 0$ or θ is an isomorphism. If $\mathbb{F} = \mathbb{F}^{\text{alg}}$ and $V = W$, then $\theta = c \text{id}_V$ for some $c \in \mathbb{F}$, i.e. $\text{End}_{\mathbb{F}G}(V) \cong \mathbb{F}$.*

Example. The (left) *regular module* of G is $\mathbb{F}G$ viewed as a left module over itself. If $\text{Irr}_{\mathbb{F}}(G) = \{S_i \mid i \in I\}$ and $\text{char } \mathbb{F} = 0$, then

$$\mathbb{F}G \cong \bigoplus_{i \in I} S_i^{\oplus \dim_{\mathbb{F}} S_i}$$

as G -modules.

Theorem 1.3 (Maschke's Theorem). *Suppose $\text{char } \mathbb{F} \nmid |G|$. If $U \leq V$ are G -modules, then there is a G -submodule $W \leq V$ such that $V = U \oplus W$.*

Corollary 1.4. *Every finite-dimensional representation of a finite group G over \mathbb{F} where $\text{char } \mathbb{F} \nmid |G|$ is semisimple.*

Common constructions.

- *Tensor products*: If V, W are G -modules, then $V \otimes_{\mathbb{F}} W$ becomes a G -module via $g \cdot (v \otimes w) = (gv) \otimes (gw)$ for all $g \in G, v \in V, w \in W$.

- *Restriction:* If $H \leq G$, V is a G -module, then we can also view V as an H -module, written $V \downarrow_H^G$, $V \downarrow_H$, V_H or $\text{Res}_H^G(V)$.
- *Induction:* If $H \leq G$, U is an H -module, we can get a G -module out of it. Let $\{t_i \mid i \in I\}$ be a set of left coset representatives of H in G . Then the induction of U from H to G is the vector space direct sum

$$\bigoplus_{i \in I} (t_i \otimes U) =: U \uparrow_H^G, U \uparrow^G \text{ or } U^G,$$

where $t_i \otimes U = \{t_i \otimes u \mid u \in U\}$, and the G -action is as follows: $g \cdot (t_i \otimes u) := t_j \otimes (t_j^{-1}gt_i)u$ where given $g \in G, i \in I$, then $j \in I$ is the unique index such that $gt_i \in t_jH$. Equivalently, we can define the induction as $U \uparrow_H^G = \mathbb{F}G \otimes_{\mathbb{F}H} U$, see Example Sheet 1, Question 1.

- *Permutation modules:* A G -module with a permutation basis B , i.e. $g \cdot b \in B$ for all $g \in G, b \in B$. E.g. the left regular module $\mathbb{F}G$ is a permutation module with basis $B = G$.

Lemma 1.5. *Suppose G acts transitively on a set Ω . Let M be the corresponding permutation module. Then $M \cong \mathbb{1}_H \uparrow^G$, where $H = \text{Stab}_G(\omega)$ for any $\omega \in \Omega$.*

Proof. Special case of Example Sheet 1, Question 2. □

1.2.2 Some Linear Algebra

- Recall that if M is a (finite-dimensional) \mathbb{F} -vector space, $M^* = \text{Hom}_{\mathbb{F}}(M, \mathbb{F})$ is again an \mathbb{F} -vector space. If e_1, \dots, e_k is a basis of M , then the dual basis $\varepsilon_1, \dots, \varepsilon_k \in M^*$ is defined by $\varepsilon_i(e_j) = \delta_{ij}$.
- Let M be a G -module, the dual M^* of M carries the G action $(g \cdot \phi)(v) = \phi(g^{-1} \cdot v)$.
- Suppose we have a symmetric bilinear form $\langle \cdot, \cdot \rangle$ on some finite-dimensional \mathbb{F} -vector space M . For a vector subspace V of M define

$$V^\perp = \{m \in M \mid \langle v, m \rangle = 0 \forall v \in V\}.$$

Consider the linear map $\phi : M \rightarrow M^*, m \mapsto \langle \cdot, m \rangle$. Note even if $\langle \cdot, \cdot \rangle$ is non-singular, i.e. $\ker \phi = M^\perp = 0$, we could have $V^\perp \cap V \neq 0$.

We can describe how large this is using a basis of V . Let e_1, \dots, e_k of V . The Gram matrix of V w.r.t. this basis be the matrix A with $A_{ij} = \langle e_i, e_j \rangle$.

Lemma 1.6. *We have that $\dim_{\mathbb{F}} V/(V \cap V^\perp) = \text{rank } A$.*

Proof. Consider $\varphi : V \rightarrow V^*, v \mapsto \langle \cdot, v \rangle$. Let $\varepsilon_1, \dots, \varepsilon_k$ be the basis of V^* dual to e_1, \dots, e_k . Then $\varphi(e_i) = \sum_{j=1}^k \langle e_j, e_i \rangle \varepsilon_j$. So the Gram matrix A is the matrix of φ with respect to the basis e_1, \dots, e_k and $\varepsilon_1, \dots, \varepsilon_k$. Clearly $\ker \varphi = V \cap V^\perp$, and so $\dim V/(V \cap V^\perp) = \dim V - \dim \ker \varphi = \text{rank } A$. □

1.2.3 Character Theory

In this subsection, $\mathbb{F} = \mathbb{C}$. Let $\rho : G \rightarrow \text{GL}(V)$ be a representation of the finite group G over some finite-dimensional \mathbb{C} -vector space V . Recall that this representation *affords* the character $\chi_V : G \rightarrow \mathbb{C}, g \mapsto \text{tr } \rho(g)$.

Theorem 1.7. $\mathbb{C}G$ -modules U, V are isomorphic iff $\chi_U = \chi_V$.

Useful facts.

- There is an inner product on class functions on G given by

$$\langle \chi, \phi \rangle = \frac{1}{|G|} \sum_{g \in G} \chi(g^{-1})\phi(g) = \frac{1}{|G|} \sum_{g \in G} \overline{\chi(g)}\phi(g).$$

- $\text{Irr}(G)$ is an orthonormal basis for the space of class functions w.r.t. $\langle \cdot, \cdot \rangle$, in particular $|\text{Irr}(G)|$ is the number of conjugacy classes of G .
- Characters of the usual constructions:
 - Direct sum: $\chi_{U \oplus V} = \chi_U + \chi_V$.
 - Tensor product: $\chi_{U \otimes V} = \chi_U \chi_V$.
 - Permutation modules: If V is a permutation module with permutation basis B , then $\chi(g) = |\{b \in B \mid gb = b\}|$ is the number of fixed points of g .
 - Restriction: If $H \leq G$ is a subgroup and V a representation of G , then $\chi_V \downarrow_H := \chi_{V \downarrow_H} = \chi_V|_H$.
- *Frobenius reciprocity:* If χ is a character of G , θ a character of H , then

$$\langle \chi \downarrow_H, \theta \rangle = \langle \chi, \theta \uparrow^G \rangle.$$

- *Mackey's theorem:* For $H, K \leq G$, ϕ a character of H , we can compute $(\phi \uparrow_H^G) \downarrow_K$ by decomposing it as a sum of characters indexed by a set of double coset representations of K, H in G . (See handout for details)

2 Specht Modules

Let \mathbb{F} be an arbitrary field.

2.1 The Symmetric Group

Let Ω be a finite set. Call the symmetric group on Ω , $\text{Sym}(\Omega)$. When $\Omega = [n]$, write S_n for $\text{Sym}(\Omega)$.

Conventions:

- $(123)(12) = (13)$ (i.e. composition from right to left)
- $S_0 = \text{Sym}(\emptyset) = \text{trivial group}$

Some representations of S_n :

- *Trivial representation* of S_n , $\mathbb{1}_{S_n}$.
- *Sign representation* of S_n , $\text{sgn}_{S_n} : \rho : S_n \rightarrow \mathbb{F}^*$, $g \mapsto \text{sgn}(g)$.
- *Natural permutation module* V_n with permutation basis $[n]$.

Note $V_n \cong \mathbb{1}_{S_{n-1}} \uparrow^{S_n}$, because $\text{Stab}(n) = S_{n-1}$.

Also $V_n \downarrow_{S_{n-1}} \cong V_{n-1} \oplus \mathbb{1}_{S_{n-1}}$.

Definition. A partition λ of n , written $\lambda \vdash n$, is a non-increasing sequence of positive integers which sum to n , i.e. $\lambda = (\lambda_1, \dots, \lambda_k)$ with $\lambda_i \in \mathbb{N}$, $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$ and $\sum_{i=1}^k \lambda_i = n$. We call

- λ_i the parts of the partition,
- n the size of λ (also denoted $|\lambda|$),
- k the length of λ (also denoted $\ell(\lambda)$).

The set $\{\lambda \mid \lambda \vdash n\}$ of all partitions of n will be denoted by $\wp(n)$.

We can extend this notion to 0 by convention: the only partition of 0 is the empty sequence, i.e. $\wp(0) = \{\emptyset\}$.

Short notation: $\lambda = (4, 3, 3, 1) = (4, 3^2, 1) \vdash 11$.

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k)$ be a partition. The Young diagram of λ is

$$Y(\lambda) = \{(i, j) \in \mathbb{N} \times \mathbb{N} \mid 1 \leq i \leq k, 1 \leq j \leq \lambda_i\}.$$

Typically, Young diagrams are drawn using boxes rather than points, e.g.:

$$\wp(4) = \left\{ \begin{array}{c} (4) \\ \square \square \square \square \end{array}, \begin{array}{c} (3, 1) \\ \square \square \square \\ \square \end{array}, \begin{array}{c} (2, 2) \\ \square \square \\ \square \square \end{array}, \begin{array}{c} (2, 1^2) \\ \square \square \\ \square \square \end{array}, \begin{array}{c} (1^4) \\ \square \\ \square \\ \square \\ \square \end{array} \right\}.$$

The rows and columns are numbered as in a matrix.

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k)$ be a partition. The conjugate partition of λ is the partition λ' such that $Y(\lambda')$ is the transpose of $Y(\lambda)$. Explicitly, $\lambda' = (\mu_1, \dots, \mu_{\lambda_1})$ where $\mu_j = \#\{i \in [k] \mid \lambda_i \geq j\}$. Note $|\lambda'| = |\lambda|$ and $(\lambda')' = \lambda$.

Example. Consider $\lambda = (4, 3, 1) \vdash 8$. Then

$$Y(\lambda) = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array},$$

and so

$$Y(\lambda') = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array},$$

i.e. $\lambda' = (3, 2, 2, 1)$.

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k)$ and $\mu = (\mu_1, \dots, \mu_s)$ be two partitions of $n \in \mathbb{N}$. Then we say that λ dominates μ , written $\lambda \trianglerighteq \mu$ or $\mu \trianglelefteq \lambda$, if $\sum_{i=1}^l \lambda_i \geq \sum_{i=1}^l \mu_i$ for all $l \in \{1, 2, \dots, \min(k, s)\}$.

Example. Take $n = 4$. Then $(4) \trianglerighteq (3, 1) \trianglerighteq (2, 2) \trianglerighteq (2, 1^2) \trianglerighteq (1^4)$.

However, in general, dominance is only a partial order, for example $(4, 3, 1) \not\trianglerighteq (5, 1^3)$ and $(5, 1^3) \not\trianglerighteq (4, 3, 1)$.

Dominance can be extended to a total ordering on $\wp(n)$, e.g. the lexicographic ordering: If $\lambda \neq \mu$, we say $\lambda > \mu$ if $\lambda_i > \mu_i$ where $i = \min\{j \in \mathbb{N} \mid \lambda_j \neq \mu_j\}$.

Definition. Let λ be a partition of n . A λ -tableau, or Young tableau of shape λ , is a bijection $t : Y(\lambda) \rightarrow [n]$. The set of all λ -tableaux will be denoted by Δ^λ .

We usually write the values of a Young tableau t in the boxes of the Young diagram $Y(\lambda)$.

Example. Take $\lambda = (3, 1) \vdash 4$, so $Y(\lambda) = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}$. Consider the tableau $t : Y(\lambda) \rightarrow [4]$, $(1, 1) \mapsto 2$, $(1, 2) \mapsto 3$, $(1, 3) \mapsto 4$, $(2, 1) \mapsto 1$. Then we write this as a labelled Young diagram, namely

$$t = \begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & & \\ \hline \end{array}.$$

The natural permutation action of S_n on $[n]$ extends to a permutation action on Δ^λ :

$$(g \cdot t)(i, j) = g(t(i, j)) \text{ for } (i, j) \in Y(\lambda), t \in \Delta^\lambda,$$

i.e. we just apply g to each entry of t .

To continue the example above, take $g = (123) \in S_4$. Then

$$g \cdot t = g \cdot \begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & & \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline 3 & 1 & 4 \\ \hline 2 & & \\ \hline \end{array}.$$

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k)$ be a partition and $t \in \Delta^\lambda$. For each $1 \leq i \leq k$, define

$$R_i(t) := \{t(i, j) \mid 1 \leq j \leq \lambda_i\}$$

and for each $1 \leq j \leq \lambda_1$, define

$$C_j(t) = \{t(i, j) \mid 1 \leq i \leq (\lambda')_j\},$$

i.e. $R_i(t)$, $C_j(t)$ are the sets of entries in the i -th row, resp. j -th column of t .

Definition. Let $\lambda \vdash n$ and $t, s \in \Delta^\lambda$. We say that t and s are row-equivalent, written $t \sim_R s$, if $R_i(t) = R_i(s)$ for all i . Note that \sim_R is an equivalence relation on Δ^λ , we will denote the equivalence classes by $\Omega^\lambda := \Delta^\lambda / \sim_R$. Each element of Ω^λ (i.e. equivalence class) will be called a λ -tabloid. We write $\{t\}$ for the equivalence class containing $t \in \Delta^\lambda$.

Example. Consider $\lambda = (3, 2) \vdash 5$ and $t = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array}$, $s = \begin{array}{|c|c|c|} \hline 2 & 3 & 1 \\ \hline 5 & 4 & \\ \hline \end{array}$. Clearly $\{t\} = \{s\}$.

To denote tabloids, we omit the vertical bars, i.e. we write

$$\{t\} = \overline{\begin{array}{ccc} 1 & 2 & 3 \\ 4 & 5 & \end{array}} = \overline{\begin{array}{ccc} 2 & 3 & 1 \\ 5 & 4 & \end{array}} = \{s\}.$$

The natural permutation of S_n on Δ^λ descends to a well-defined action on Ω^λ .

Definition. Let $\lambda \vdash n$. The λ -Young permutation module M^λ is the S_n -module with permutation basis Ω^λ .

Lemma 2.1. Let $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$. Then $M^\lambda \cong \mathbb{1}_{S_\lambda} \uparrow_{S_\lambda}^{S_n}$ where $S_\lambda \cong S_{\lambda_1} \times \dots \times S_{\lambda_k}$.

Proof. S_n acts transitively on $[n]$ and so acts transitively on Ω^λ . For $t \in \Delta^\lambda$,

$$\begin{aligned} S_\lambda &:= \text{Stab}_{S_n}(\{t\}) = \{g \in S_n \mid gR_i(t) = R_i(t) \forall i\} = \text{Sym}(R_1(t)) \times \dots \times \text{Sym}(R_k(t)) \\ &\cong S_{\lambda_1} \times \dots \times S_{\lambda_k}. \end{aligned}$$

The claim then follows from Lemma 1.5. □

Remark. The subgroup S_λ of S_n above is called a *Young subgroup of type λ* . There is a Young subgroup of type λ for each set partitions of $[n]$ into subsets of sizes $\lambda_1, \dots, \lambda_k$ and for fixed λ they are all conjugate to each other in S_n , and all isomorphic to $S_{\lambda_1} \times \dots \times S_{\lambda_k}$.

Example. Take $n = 9$, $\lambda = (4, 3, 2)$. There are $\binom{9}{4} \binom{5}{3} \binom{2}{2} = 1260$ many Young subgroups of type λ .

Examples.

(a) Let $\lambda = (n)$. Then

$$\Omega^\lambda = \{\overline{1 \ 2 \ \dots \ n}\},$$

and S_n acts trivially on this single λ -tabloid. Then $S_\lambda = S_n$ and $M^{(n)} \cong \mathbb{1}_{S_n}$.

(b) Let $\lambda = (n-1, 1) \vdash n$, for $n \geq 2$. Then

$$\Omega^\lambda = \left\{ \overline{\begin{array}{cccccc} 1 & 2 & \dots & i-1 & i+1 & \dots & n \\ j & & & & & & \end{array}} \mid 1 \leq i \leq n \right\}.$$

Then $S_\lambda \cong S_{n-1} \times S_1 \cong S_{n-1}$, hence $M^{(n-1,1)} \cong \mathbb{1}_{S_{n-1}} \uparrow^{S_n} \cong V_n$, the natural permutation representation.

(c) Let $\lambda = (1^n) \vdash n$. Then $\{t\} = \{s\}$ iff $t = s$ for $t, s \in \Delta^\lambda$. So S_λ is trivial and so $M^{(1^n)} \cong \mathbb{1}_1 \uparrow^{S_n}$ is the regular module $\mathbb{F}S_n$.

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$ and $t \in \Delta^\lambda$.

(i) The row stabiliser of t is

$$R(t) := \{g \in S_n \mid gR_i(t) = R_i(t) \forall i\},$$

and similarly define the column stabiliser $C(t)$.

(ii) The column symmetriser of t is

$$\mathbf{b}_t := \sum_{g \in C(t)} \text{sgn}(g)g \in \mathbb{F}S_n.$$

(iii) The polytabloid corresponding to t , or t -polytabloid, is

$$e(t) := \mathbf{b}_t \cdot \{t\} = \sum_{g \in C(t)} \text{sgn}(g)g \cdot \{t\} \in M^\lambda.$$

Note that $e(t)$ depends on the tableau t , not just the tabloid $\{t\}$.

Example. Let $\lambda = (2, 1) \vdash 3$. Then

$$e\left(\overline{\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \end{array}}\right) = \frac{\overline{1 \ 2}}{3} - \frac{\overline{3 \ 2}}{1} \neq \frac{\overline{2 \ 1}}{3} - \frac{\overline{3 \ 1}}{2} = e\left(\overline{\begin{array}{|c|c|} \hline 2 & 1 \\ \hline 3 & \end{array}}\right).$$

Definition. Let $\lambda \vdash n$. The λ -Specht module is defined as

$$\mathcal{S}^\lambda := \langle e(t) \mid t \in \Delta^\lambda \rangle_{\mathbb{F}} \subseteq M^\lambda,$$

i.e. \mathcal{S}^λ is the \mathbb{F} -vector space spanned by polytabloids corresponding to tableaux of shape λ .

The next lemma shows that \mathcal{S}^λ is indeed a module over S_n .

Lemma 2.2. *Let $\lambda \vdash n$ and $t \in \Delta^\lambda$.*

- (1) $e(t) \neq 0$
- (2) $\forall g \in S_n, g \cdot e(t) = e(g \cdot t)$
- (3) $\forall g \in C(t), g \cdot e(t) = \text{sgn}(g)e(t)$
- (4) \mathcal{S}^λ is a cyclic submodule of M^λ , in particular $\mathcal{S}^\lambda = \mathbb{F}S_n \cdot e(u)$ for any $u \in \Delta^\lambda$.

Proof.

- (1) Observe that $R(t) \cap C(t) = 1$, and so if $g \in C(t)$ and $g \cdot \{t\} = \{t\}$, then $g = 1$.

It follows that the coefficient of $\{t\}$ in $e(t)$ is $\text{sgn}(1) = 1 \neq 0$, hence $e(t) \neq 0$.

[In fact, $R(t) \cap C(t) = 1$ implies that $e(t)$ is a signed sum of $|C(t)|$ distinct λ -tabloids]

- (2) Observe that $C(g \cdot t) = gC(t)g^{-1}$, and so

$$\begin{aligned}
g \cdot e(t) &= g \sum_{h \in C(t)} \text{sgn}(h)h \cdot \{t\} \\
&= \sum_{h \in C(t)} \text{sgn}(h)\{gh \cdot t\} \\
&= \sum_{h \in C(t)} \text{sgn}(ghg^{-1})ghg^{-1} \cdot \{g \cdot t\} \\
&= \sum_{x \in C(g \cdot t)} \text{sgn}(x)x \cdot \{g \cdot t\} = e(g \cdot t).
\end{aligned}$$

- (3) If $g \in C(t)$, then

$$g \cdot e(t) = \sum_{h \in C(t)} \text{sgn}(h)\{gh \cdot t\} = \sum_{y \in C(t)} \text{sgn}(g^{-1}y)\{y \cdot t\} = \text{sgn}(g)e(t).$$

- (4) That \mathcal{S}^λ is an S_n -submodule of M^λ follows from (2)

That \mathcal{S}^λ can be generated as an $\mathbb{F}S_n$ -module by $e(u)$ for any $u \in \Delta^\lambda$ also follows from (2) and the fact that S_n acts transitively on Δ^λ .

□

Examples.

- (a) Let $\lambda = (n)$. We have by (1) and (4) of the lemma that $0 \neq \mathcal{S}^\lambda \leq M^\lambda$. But in a previous example we showed that $M^{(n)} \cong \mathbb{1}_{S_n}$. Hence $\mathcal{S}^{(n)} \cong \mathbb{1}_{S_n}$ also.

(b) Let $\lambda = (1^n) \vdash n$. Then $C(t) = S_n$ and thus by the lemma, $g \cdot e(t) = \text{sgn}(g)e(t)$ for all $g \in S_n$, for any $t \in \Delta^\lambda$. Thus, $\dim_{\mathbb{F}}(\mathcal{S}^{(1^n)}) = 1$ and $\mathcal{S}^{(1^n)} = \mathbb{F}S_n \cdot e(t) \cong \text{sgn}_{S_n}$.

(c) Let $\lambda = (2, 1) \vdash 3$. Then

$$\mathcal{S}^\lambda = \left\langle e\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}\right), e\left(\begin{array}{|c|c|} \hline 2 & 1 \\ \hline 3 & \\ \hline \end{array}\right), e\left(\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}\right), e\left(\begin{array}{|c|c|} \hline 3 & 1 \\ \hline 2 & \\ \hline \end{array}\right), e\left(\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & \\ \hline \end{array}\right), e\left(\begin{array}{|c|c|} \hline 3 & 2 \\ \hline 1 & \\ \hline \end{array}\right) \right\rangle_{\mathbb{F}}.$$

By (iii) of the lemma,

$$\mathcal{S}^\lambda = \left\langle \alpha := e\left(\begin{array}{|c|c|} \hline 2 & 1 \\ \hline 3 & \\ \hline \end{array}\right), \beta := e\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}\right), \gamma := e\left(\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}\right) \right\rangle_{\mathbb{F}},$$

since e.g. $e\left(\begin{array}{|c|c|} \hline 3 & 2 \\ \hline 1 & \\ \hline \end{array}\right) = -\alpha$. Moreover,

$$\begin{aligned} \alpha &= \frac{\overline{1 \ 2}}{3} - \frac{\overline{3 \ 2}}{1}, \\ \beta &= \frac{\overline{2 \ 1}}{3} - \frac{\overline{3 \ 1}}{2}, \\ \gamma &= \frac{\overline{1 \ 3}}{2} - \frac{\overline{2 \ 3}}{1}, \end{aligned}$$

so $\alpha = \beta + \gamma$. Since β, γ are linearly independent, $\dim \mathcal{S}^\lambda = 2$ for all fields \mathbb{F} . See Exercise Sheet 1, Question 4 for more.

2.2 Irreducible modules

Goal: If $\text{char } \mathbb{F} = 0$, then $\{\mathcal{S}^\lambda \mid \lambda \vdash n\}$ is a full set of irreducible $\mathbb{F}S_n$ -modules.

Definition. Let $\lambda \vdash n$. Define a symmetric bilinear form $\langle \cdot, \cdot \rangle$ on M^λ via

$$\langle \{t\}, \{s\} \rangle = \begin{cases} 1 & \text{if } \{t\} = \{s\}, \\ 0 & \text{otherwise,} \end{cases}$$

for $t, s \in \Delta^\lambda$ and then extend linearly, i.e. we take the tabloids to be an “orthonormal basis”.

We will always take the orthogonal complement U^\perp of a subspace U with respect to this bilinear form.

Lemma 2.3. Let $\lambda \vdash n$.

(1) The form $\langle \cdot, \cdot \rangle$ is S_n -invariant, i.e. $\langle gx, gy \rangle = \langle x, y \rangle$ for all $x, y \in M^\lambda, g \in S_n$.

(2) If U is an S_n -submodule of M^λ , then so is U^\perp .

Proof.

(1) This is clearly true for $x = \{t\}, y = \{s\}$, where $t, s \in \Delta^\lambda$, then follows by bilinearity.

- (2) This follows from (1): For $x \in U^\perp, g \in S_n$ we have $\langle gx, u \rangle = \langle x, g^{-1}u \rangle = 0$ for all $u \in U$, so $gx \in U^\perp$.

□

Plan:

- *James's Submodule Theorem:* If $U \leq M^\lambda$, then $U \geq \mathcal{S}^\lambda$ or $U \leq (\mathcal{S}^\lambda)^\perp$.
- JST \implies certain quotients of \mathcal{S}^λ are irreducible.

This will give us the first part of our goal: \mathcal{S}^λ is irreducible when $\text{char } \mathbb{F} = 0$.

Then the second part will be to show that they are pairwise non-isomorphic.

Proposition 2.4. *Let $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$. Suppose $t, u \in \Delta^\lambda$ satisfy $\mathfrak{b}_t \cdot \{u\} \neq 0$. Then*

- (1) $\exists h \in C(t)$ such that $h \cdot \{t\} = \{u\}$,
- (2) $\mathfrak{b}_t \cdot \{u\} = \pm e(t)$,
- (3) $\mathfrak{b}_t \cdot M^\lambda = \mathbb{F}e(t)$.

Proof.

- (1) We want to construct $h \in C(t)$ such that $R_i(h \cdot t) = R_i(u)$ for all i .

Claim: $\mathfrak{b}_t \cdot \{u\} \neq 0 \implies$ if $x \neq y$ are the numbers appearing in the same row of u , then they appear in different columns of t .

Proof of claim: Suppose not, so $(xy) \in C(t)$. Take Z to be a set of left coset representatives of $\langle (xy) \rangle$ in $C(t)$, i.e. $C(t) = Z \dot{\cup} Z(xy)$.

Then $\mathfrak{b}_t = \sum_{g \in C(t)} \text{sgn}(g)g = \sum_{z \in Z} \text{sgn}(z)z(1 - (xy))$. But then

$$\mathfrak{b}_t \cdot \{u\} = \sum_{z \in Z} \text{sgn}(z)z(\{u\} - (xy) \cdot \{u\}) = 0,$$

since $(xy) \in R(u)$ as x, y belong to the same row in u . This concludes the proof of the claim.

Returning to the proof of (1), let $R_1(u) = \{x_1, x_2, \dots, x_{\lambda_1}\}$. Suppose x_r belongs to column j_r of t , for each $r \in [\lambda_1]$. By the claim the j_r are pairwise distinct. Let $y_r = t((1, j_r))$.

Define $h_1 = \prod_{\substack{r \in [\lambda_1] \\ x_r \neq y_r}} (x_r y_r) \in C(t)$. Then

$$R_1(h_1 \cdot t) = \{h_1(y_1), \dots, h_1(y_{\lambda_1})\} = \{x_1, \dots, x_{\lambda_1}\} = R_1(u).$$

Since $h_1 \in C(t)$, then $C(h_1 \cdot t) = h_1 C(t) h_1^{-1} = C(t)$. Thus $\mathfrak{b}_t = \mathfrak{b}_{h_1 t}$, and so $\mathfrak{b}_{h_1 t} \cdot \{u\} \neq 0$.

Let $R_2(u) = \{x'_1, \dots, x'_{\lambda_2}\}$. Suppose x'_r belongs to column j'_r of $t' = h_1 \cdot t$. By the claim, the j'_r are pairwise distinct. Let $y'_r = t'((2, j'_r))$. Define $h_2 = \prod_{r \in [\lambda_2]} (x'_r y'_r) \in C(t')$. Observe $R_2(h_2 \cdot t') = R_2(u)$ and $R_1(h_2 \cdot t') = R_1(t') = R_1(u)$. That is: $R_i(h_2 h_1 \cdot t) = R_i(u)$ for all $i \in \{1, 2\}$.

Iteratively, we construct for each $m \in \{3, 4, \dots, k\}$ an element $h_m \in C(t)$ such that $R_i(h_m h_{m-1} \cdots h_1 \cdot t) = R_i(u)$ for all $i \in [m]$. For $m = k$ we get what we want by taking $h = h_k \cdots h_2 h_1$.

- (2) Let h be as in (1). Then $\mathfrak{b}_t \cdot \{u\} = \mathfrak{b}_t h \cdot \{t\} = \text{sgn}(h) \mathfrak{b}_t \cdot \{t\} = \text{sgn}(h) e(t)$.
- (3) For all $\{u\} \in M^\lambda$ we have either $\mathfrak{b}_t \cdot \{u\} = 0$ or $\mathfrak{b}_t \cdot \{u\} = \pm \{u\}$ by (2), hence $\mathfrak{b}_t \cdot M^\lambda \subseteq \mathbb{F}e(t)$ and equality holds as $\mathfrak{b}_t \{t\} = e(t)$.

□

Theorem 2.5 (James's Submodule Theorem). *Let $\lambda \vdash n$, $U \leq M^\lambda$. Then either $U \geq \mathcal{S}^\lambda$ or $U \leq (\mathcal{S}^\lambda)^\perp$.*

Proof. Suppose $U \not\leq (\mathcal{S}^\lambda)^\perp$, then there exists $x \in U$ and $t \in \Delta^\lambda$ such that $\langle x, e(t) \rangle \neq 0$. Then

$$0 \neq \langle x, e(t) \rangle = \sum_{g \in C(t)} \text{sgn}(g) \langle g^{-1} x, \{t\} \rangle = \langle \mathfrak{b}_t \cdot x, \{t\} \rangle,$$

so in particular $\mathfrak{b}_t \cdot x \neq 0$. By the proposition we have $\mathfrak{b}_t \cdot x = ce(t)$ for some $c \in \mathbb{F}^*$. So from $\mathfrak{b}_t \cdot x \in U$ we get $e(t) \in U$ and thus $\mathcal{S}^\lambda = \mathbb{F}S_n e(t) \subseteq U$. □

Remark. By JST, if we decompose M^λ into a direct sum of indecomposable modules, then there is a unique summand that contains \mathcal{S}^λ . This module is denoted Y^λ , and called the *Young module* corresponding to λ (more later).

Corollary 2.6. *Let $\lambda \vdash n$. Then $\mathcal{S}^\lambda / (\mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp)$ is either 0 or irreducible.*

Proof. If $\mathcal{S}^\lambda \leq (\mathcal{S}^\lambda)^\perp$, then the quotient is zero, so now suppose $\mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp$ is a proper submodule of \mathcal{S}^λ . Let $U \leq \mathcal{S}^\lambda$. Then $U \leq M^\lambda$, so by JST we have $U = \mathcal{S}^\lambda$ or $U \leq \mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp$. This tells us that $\mathcal{S}^\lambda / (\mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp)$ is irreducible. □

Definition. *A representation $\rho : G \rightarrow \text{GL}_n(\mathbb{F})$ is absolutely irreducible if for any field extension \mathbb{K} of \mathbb{F} , the corresponding representation $\bar{\rho} : G \rightarrow \text{GL}_n(\mathbb{K})$ is irreducible.*

Example. Let $G = C_4 = \langle g \rangle$. The representation

$$\rho : G \rightarrow \text{GL}_2(\mathbb{Q}), \quad \rho(g) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

is irreducible, since it has no 1-dimensional submodules (i.e. eigenspaces of $\rho(g)$) when we work over \mathbb{Q} . However, it is not absolutely irreducible: $\bar{\rho} : G \rightarrow \text{GL}_2(\mathbb{Q}(i))$ is a direct sum of two 1-dimensional submodules (because the eigenvalues of $\bar{\rho}(g)$ are $\pm i$).

Theorem 2.7. *Let $\lambda \vdash n$. Then $\mathcal{S}^\lambda / (\mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp)$ is either 0 or absolutely irreducible.*

Proof. We can extract a basis e_1, \dots, e_k of \mathcal{S}^λ consisting of polytabloids. By Lemma 1.6,

$$\dim_{\mathbb{F}} \mathcal{S}^\lambda / (\mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp) = \text{rank } A$$

where A is the Gram matrix corresponding to e_1, \dots, e_k . But $A_{ij} = \langle e_i, e_j \rangle$ belongs to the prime subfield of \mathbb{F} (i.e. \mathbb{Q} or \mathbb{F}_p) and so the dimension of $\mathcal{S}^\lambda / (\mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp)$ doesn't change when we extend \mathbb{F} . Since over any field $\mathcal{S}^\lambda / (\mathcal{S}^\lambda \cap (\mathcal{S}^\lambda)^\perp)$ is either 0 or irreducible by our previous result, it is either 0 or absolutely irreducible. \square

Corollary 2.8. *If $\text{char } \mathbb{F} = 0$, then \mathcal{S}^λ is irreducible for all partitions λ .*

Proof. Over \mathbb{Q} the form $\langle \cdot, \cdot \rangle$ satisfies $\langle u, u \rangle \geq 0$ for all $u \in M_{\mathbb{Q}}^\lambda$, with equality iff $u = 0$. Hence $\mathcal{S}_{\mathbb{Q}}^\lambda \cap (\mathcal{S}_{\mathbb{Q}}^\lambda)^\perp = 0$. Thus $\mathcal{S}_{\mathbb{Q}}^\lambda$ is absolutely irreducible by the theorem. Hence $\mathcal{S}_{\mathbb{F}}^\lambda$ is irreducible since \mathbb{F} extends \mathbb{Q} . \square

Proposition 2.9. *Let $\lambda = (\lambda_1, \dots, \lambda_k)$ and $\mu = (\mu_1, \dots, \mu_s)$ be two partitions of n . Suppose $t \in \Delta^\lambda$ and $u \in \Delta^\mu$ with $\mathfrak{b}_t \cdot \{u\} \neq 0$. Then*

(1) $\exists h \in C(t)$ such that for all $l \in \{1, 2, \dots, \min(k, s)\}$ we have

$$\bigsqcup_{i=1}^l R_i(u) \subseteq \bigsqcup_{i=1}^l R_i(h \cdot t).$$

(2) $\lambda \supseteq \mu$.

Proof.

(1) Arguing as in the claim in the proof of Proposition 2.4 we have that if $x \neq y$ appear in the same row of u , then they appear different columns of t .

Let $R_1(u) = \{x_1, x_2, \dots, x_{\mu_1}\}$. Suppose x_r lies in column j_r of t , so the j_r are pairwise distinct. Let $y_r = t((1, j_r))$.

Define $h_1 = \prod_{\substack{r \in [\mu_1] \\ x_r \neq y_r}} (x_r y_r) \in C(t)$. Then

$$R_1(u) = \{x_1, \dots, x_{\mu_1}\} = \{h_1(y_1), \dots, h_1(y_{\mu_1})\} \subseteq R_1(h_1 \cdot t).$$

Since $C(h_1 \cdot t) = h_1 C(t) h_1^{-1} = C(t)$, so $\mathfrak{b}_{h_1 \cdot t} = \mathfrak{b}_t$, so $\mathfrak{b}_{h_1 \cdot t} \cdot \{u\} \neq 0$.

Let $R_2(u) = \{x'_1, x'_2, \dots, x'_{\mu_2}\}$ and $t' = h_1 \cdot t$. Suppose $t'((i'_r, j'_r)) = x'_r$. If $i'_r \geq 2$, then let $y'_r = t'((2, j'_r))$. Define $h_2 = \prod_{\substack{r \in [\mu_2] \\ i'_r \geq 2 \\ x'_r \neq y'_r}} (x'_r y'_r) \in C(t') = C(t)$. Then

$$R_2(u) = \{x'_1, \dots, x'_{\mu_2}\} = \{x'_r \mid i'_r \geq 2\} \sqcup \{x'_r \mid i'_r = 1\}$$

$$\begin{aligned}
&= \{h_2(y'_r)\} \sqcup \{x'_r \mid i'_r = 1\} \\
&\subseteq R_2(h_2 \cdot t') \sqcup R_1(h_1 \cdot t')
\end{aligned}$$

Also $R_1(u) \subseteq R_1(t') = R_1(h_2 \cdot t')$. Therefore $\bigsqcup_{i=1}^l R_i(u) \subseteq \bigsqcup_{i=1}^l R_i(h_2 h_1 \cdot t')$ for all $l \in \{1, 2\}$. Now induct.

(2) By (1),

$$\sum_{i=1}^l \mu_i = \sum_{i=1}^l |R_i(u)| \leq \sum_{i=1}^l |R_i(h \cdot t)| = \sum_{i=1}^l \lambda_i,$$

for all $l = 1, \dots, \min(k, s)$. □

Theorem 2.10. *Let $\lambda, \mu \vdash n$. Suppose $0 \neq \phi \in \text{Hom}_{\mathbb{F}S_n}(\mathcal{S}^\lambda, M^\mu)$. If there exists $\tilde{\phi} \in \text{Hom}_{\mathbb{F}S_n}(M^\lambda, M^\mu)$ extending ϕ , then $\lambda \trianglerighteq \mu$.*

Proof. Since $\mathcal{S}^\lambda = \mathbb{F}S_n \cdot e(t)$ for any $t \in \Delta^\lambda$, then $\phi(e(t)) \neq 0$ as $\phi \neq 0$. Fix any $t \in \Delta^\lambda$. Then $0 \neq \phi(e(t)) = \tilde{\phi}(e(t)) = \tilde{\phi}(\mathbf{b}_t \cdot \{t\}) = \mathbf{b}_t \cdot \tilde{\phi}(\{t\})$. Writing $\phi(\{t\})$ as a sum of μ -tabloids, we see that there is $u \in \Delta^\mu$ such that $\mathbf{b}_t \cdot \{u\} \neq 0$, so we are done by the proposition. □

Example. Let $\text{char } \mathbb{F} = 2$, $n = 2$, $\lambda = (1^2)$, $\mu = (2)$. Then $\mathcal{S}^{(1^2)} \cong \text{sgn}_{S_2} \cong \mathbb{1}_{S_2} \cong M^{(2)}$, and so $\text{Hom}_{\mathbb{F}S_2}(\mathcal{S}^\lambda, M^\mu) \neq 0$, in particular, it contains isomorphisms.

On the other hand, $M^\lambda = \left\langle \overline{\frac{1}{2}}, \overline{\frac{2}{1}} \right\rangle$ and if $\theta : M^\lambda \rightarrow M^\mu$ is $\mathbb{F}S_2$ linear, then $\theta(\overline{\frac{1}{2}}) = (12)\theta(\overline{\frac{2}{1}}) = \theta(\overline{\frac{2}{1}})$. In particular $\theta(e(\overline{\frac{1}{2}})) = \theta(\overline{\frac{1}{2}}) - \theta(\overline{\frac{2}{1}}) = 0$. So for any $\theta \in \text{Hom}_{\mathbb{F}S_2}(M^\lambda, M^\mu)$ we have $\theta|_{\mathcal{S}^\lambda} = 0$, in particular not all $\phi \in \text{Hom}_{\mathbb{F}S_2}(\mathcal{S}^\lambda, M^\mu)$ have extensions to M^λ .

Corollary 2.11. *If $\text{char } \mathbb{F} = 0$, $\lambda, \mu \vdash n$, then $\mathcal{S}^\lambda \cong \mathcal{S}^\mu$ iff $\lambda = \mu$.*

Proof. Suppose $\mathcal{S}^\lambda \cong \mathcal{S}^\mu$, take an isomorphism $\mathcal{S}^\lambda \rightarrow \mathcal{S}^\mu$ and compose this with the natural inclusion $\mathcal{S}^\mu \rightarrow M^\mu$ to get $0 \neq \phi \in \text{Hom}_{\mathbb{F}S_n}(\mathcal{S}^\lambda, M^\mu)$. By Maschke's Theorem there exists $V \leq M^\lambda$ such that $M^\lambda = \mathcal{S}^\lambda \oplus V$. And so we can extend ϕ to $\tilde{\phi} \in \text{Hom}_{\mathbb{F}S_n}(M^\lambda, M^\mu)$ by setting $\tilde{\phi}|_V = 0$, so $\lambda \trianglerighteq \mu$ by the theorem. By symmetry we also have $\mu \trianglerighteq \lambda$, so $\lambda = \mu$. □

So far we showed: If $\text{char } \mathbb{F} = 0$, then

- each \mathcal{S}^λ is irreducible,
- the \mathcal{S}^λ are pairwise non-isomorphic.

If $\mathbb{F} = \mathbb{C}$, then $|\text{Irr}_{\mathbb{C}}(S_n)| = \#\text{conjugacy classes of } S_n = |\varphi(n)|$, so

$$\text{Irr}_{\mathbb{C}}(S_n) = \{\mathcal{S}_{\mathbb{C}}^\lambda \mid \lambda \vdash n\}.$$

We now extend this to arbitrary fields of characteristic 0.

Theorem 2.12. *If $\text{char } \mathbb{F} = 0$, then $\text{Irr}_{\mathbb{F}}(S_n) = \{\mathcal{S}_{\mathbb{F}}^{\lambda} \mid \lambda \vdash n\}$.*

We already know that $|\text{Irr}_{\mathbb{F}}(S_n)| \geq |\wp(n)|$. We now want to prove the reverse inequality.

Definition. \mathbb{F} is a splitting field for the finite group G if every irreducible $\mathbb{F}G$ -representation is absolutely irreducible.

Fact. If $\mathbb{F} = \mathbb{F}^{\text{alg}}$, then \mathbb{F} is a splitting field. See [Isa76, Corollary 9.4]

Theorem 2.13. *If \mathbb{F} is a splitting field for G , and \mathbb{K} a field extension of \mathbb{F} , then \mathbb{K} is also a splitting field for G , and $|\text{Irr}_{\mathbb{K}}(G)| = |\text{Irr}_{\mathbb{F}}(G)|$.*

Proof. See [Isa76, Corollary 9.8]. □

Fact. Every field is a splitting field for S_n . See [JK84, Theorem 2.1.12] and [CR62].

So in particular, \mathbb{Q} is a splitting field for S_n . Hence $|\text{Irr}_{\mathbb{F}}(S_n)| = |\text{Irr}_{\mathbb{Q}}(S_n)| = |\text{Irr}_{\mathbb{C}}(S_n)| = |\wp(n)|$.

Alternatively, one can use the following:

Theorem 2.14. *Let \mathbb{K} be a field with $\text{char } \mathbb{K} \nmid |G|$. Then $|\text{Irr}_{\mathbb{K}}(G)| \leq \#\text{conjugacy classes of } G$. If $\mathbb{K} = \mathbb{K}^{\text{alg}}$, then equality holds.*

Proof. See Moodle for a sketch using the Artin-Wedderburn theorem. □

Proof of Theorem 2.12. Corollary 2.8 and Corollary 2.11 show that the \mathcal{S}^{λ} are pairwise distinct and irreducible. Then the claim follows either from Theorem 2.13 and the fact or from Theorem 2.14. □

Remarks.

- Modular representation theory: $\text{char} = p > 0$, ordinary representation theory: $\text{char} = 0$.
- If $\text{char} = p > 0$, but $p \nmid |G|$, then the situation is similar to $\text{char} = 0$.
- If $\text{char} = p \mid |G|$, the situation is very different.
- For $\text{char}(\mathbb{F}) = p > 0$:

$$\text{Irr}_{\mathbb{F}}(S_n) = \left\{ \frac{\mathcal{S}^{\lambda}}{\mathcal{S}^{\lambda} \cap (\mathcal{S}^{\lambda})^{\perp}} \mid \lambda \vdash n \text{ is "p-regular"} \right\}.$$

Theorem 2.15 (Brauer). *Suppose $\text{char } \mathbb{F} = p > 0$. Then the number of isomorphism classes of absolutely irreducible $\mathbb{F}G$ -modules is at most the number of p -regular conjugacy classes of G . If \mathbb{F} is a splitting field for G , then equality holds.*

Proof. See [CR62, pp. 82.6, 83.6] □

Definition. Let p be a prime.

- (i) A partition $\lambda = (\lambda_1, \dots, \lambda_k)$ is p -singular if it has at least p equal parts, i.e. there exists $i \in [k - p + 1]$ such that $\lambda_i = \lambda_{i+1} = \dots = \lambda_{i+p-1}$. Otherwise, λ is called p -regular.
- (ii) An element $g \in G$ is p -regular if $p \nmid \text{ord } g$. A conjugacy class of G is p -regular if its elements are p -regular.

If $g \in S_n$, then g is p -regular iff in its disjoint cycle decomposition, no cycle has length divisible by p .

Proposition 2.16. Let p be a prime, $n \in \mathbb{N}$. Then

$$\#\{p\text{-regular } \lambda \vdash n\} = \#\{\lambda \vdash n \mid p \nmid \lambda_i \forall i\}.$$

Proof. **Proof 1.** The generating function for all partitions is

$$G(x) = \sum_{n \geq 0} |\wp(n)| x^n = \prod_{i \in \mathbb{N}} (1 + x^i + x^{2i} + \dots) = \prod_{i \in \mathbb{N}} \frac{1}{1 - x^i}$$

where a partition with a_i many parts of size i corresponds to choosing the x^{ia_i} term from the i -th bracket when we multiply out. The generating function for p -regular partitions is

$$\begin{aligned} F(x) &= \sum_{n \geq 0} \#\{p\text{-regular } \lambda \vdash n\} x^n = \prod_{i \in \mathbb{N}} (1 + x^i + \dots + x^{(p-1)i}) \\ &= \prod_{i \in \mathbb{N}} \frac{1 - x^{pi}}{1 - x^i} \\ &= \prod_{i \in \mathbb{N}, p \nmid i} \frac{1}{1 - x^i} \\ &= \sum_{n \geq 0} \#\{\lambda \vdash n \mid p \nmid \lambda_i \forall i\}. \end{aligned}$$

Proof 2. Consider

$$\{p\text{-regular } \lambda \vdash n\} \stackrel{\theta}{\rightleftarrows} \{\lambda \vdash n \mid p \nmid \lambda_i \forall i\} \stackrel{\varphi}{\leftleftarrows}$$

where θ, φ are as follows:

- θ : If λ has a part of size divisible by p , break it into p equal parts; repeat until there are no more parts of size divisible by p .
- φ : For each s , suppose λ has $\sum_{i \geq 0} a_i p^i$ parts of size s where $0 \leq a_i \leq p - 1$. Glue them together to form a_i many parts of size sp^i for each i .

Then check that θ, φ are inverses. □

In fact, the proposition and both proofs hold for all $p \in \mathbb{N}$ (not necessarily prime), provided we extend the definition accordingly.

2.3 Standard Basis Theorem

We have $\mathcal{S}^\lambda = \langle e(t) \mid t \in \Delta^\lambda \rangle_{\mathbb{F}}$. Our goal for this section is to extract a basis of polytabloids for \mathcal{S}^λ , uniform over all \mathbb{F} , thereby computing $\dim \mathcal{S}^\lambda$ (independently of \mathbb{F}).

Definition. Let $\lambda \vdash n$, $t \in \Delta^\lambda$. Then we say

- t is row-standard if the entries of t increase along rows from left to right, i.e. $t((i, j)) < t((i, j + 1))$ for all $i \in [\ell(\lambda)]$, $j \in [\lambda_i - 1]$,
- t is column-standard if the entries of t increase along columns from top to bottom, i.e. $t((i, j)) < t((i + 1, j))$ for all $j \in [\lambda_1]$, $i \in [(\lambda')_j - 1]$,
- t is standard if it is both row- and column-standard.

Define $\text{std}(\lambda) = \{t \in \Delta^\lambda \mid t \text{ is standard}\}$. We say a polytabloid $e(t)$ is standard if t is.

Examples.

- Let $\lambda = (n)$, so $\dim \mathcal{S}^\lambda = 1$ and $\text{std}(\lambda) = \{\boxed{1 \mid 2 \mid \dots \mid n}\}$.
- Let $\lambda = (1^n) \vdash n$, so $\dim \mathcal{S}^\lambda = 1$ and $\text{std}(\lambda) = \left\{ \begin{array}{c} \boxed{1} \\ \boxed{2} \\ \dots \\ \boxed{n} \end{array} \right\}$.
- Let $\lambda = (2, 1)$. We have seen earlier that then $\dim \mathcal{S}^\lambda = 2$. Then

$$\text{std}(\lambda) = \left\{ \begin{array}{c} \boxed{1 \mid 2} \\ \boxed{3} \end{array}, \begin{array}{c} \boxed{1 \mid 3} \\ \boxed{2} \end{array} \right\}.$$

- More generally, let $\lambda = (n - 1, 1)$ with $n \geq 2$. Then $\dim \mathcal{S}^\lambda = n - 1$ by Example Sheet 1, Question 5, and

$$\text{std}(\lambda) = \left\{ \begin{array}{c} \boxed{1 \mid 2 \mid \dots \mid \hat{j} \mid \dots \mid n} \\ \boxed{j} \end{array} \mid 2 \leq j \leq n \right\}$$

Our aim will be to show that $\{e(t) \mid t \in \text{std}(\lambda)\}$ is an \mathbb{F} -basis for \mathcal{S}^λ .

For linear independence, we begin by putting a total order on Ω^λ , the set of all tableaux of shape λ .

Definition. Let $\lambda \vdash n$, $t, u \in \Delta^\lambda$. Let

$$\begin{aligned} A &= \{\text{numbers that don't appear in the same row of } t \text{ and } u\} \\ &= [n] \setminus \bigcup_{i=1}^{\ell(\lambda)} R_i(t) \cap R_i(u). \end{aligned}$$

If $\{t\} \neq \{u\}$, equivalently $A \neq \emptyset$, then let $y = \max(A)$. We say $\{t\} > \{u\}$ if $y \in R_i(t) \cap R_j(u)$ where $i > j$.

Remark. Note that $>$ is a total order on Ω^λ ; it is equivalent to a total order on the set of all row-standard λ -tableaux. The maximal element w.r.t. $>$ is

$$\begin{array}{c} \overline{1 \ 2 \ 3 \ \dots \ \lambda_1} \\ \overline{\lambda_1+1 \ \dots \ \lambda_1+\lambda_2} \\ \vdots \\ \overline{\dots \ n} \end{array}$$

Small example: Take $\lambda = (3, 2)$, $t = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array}$, $u = \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array}$. Then $A = \{3, 4\}$, so $y = 4$ and $\{t\} > \{u\}$.

Lemma 2.17. *Let $\lambda \vdash n$, $t \in \Delta^\lambda$ column-standard. Let $h \in C(t) \setminus \{1\}$. Then $\{h \cdot t\} < \{t\}$.*

Proof. Since $h \neq 1$ and $R(t) \cap C(t) = \{1\}$, then $\{h \cdot t\} \neq \{t\}$. Then

$$y := \max \left([n] \setminus \bigcup_{i=1}^{\ell(\lambda)} R_i(t) \cap R_i(h \cdot t) \right)$$

exists. Suppose $y = t((i, j))$. Where is y in $h \cdot t$? Since $h \in C(t)$, then $y \in C_j(h \cdot t)$, say $y \in R_{i'}(h \cdot t)$. First, $i' \neq i$ by definition of y . But also, $i' \not> i$ since the entries in column j below row i must match exactly in t and $h \cdot t$ by maximality of y and column-standardness of t . Hence $i' < i$, so $\{h \cdot t\} < \{t\}$. \square

Proposition 2.18. *Let $\lambda \vdash n$. Then the $e(t)$ with $t \in \text{std}(\lambda)$ are linearly independent.*

Proof. Suppose not. Then there exists $\emptyset \neq \Delta \subseteq \text{std}(\lambda)$ such that $\sum_{t \in \Delta} \alpha_t e(t) = 0$ where $\alpha_t \in \mathbb{F}^\times$. For $t, u \in \text{std}(\lambda)$, we have $\{t\} = \{u\}$ iff $t = u$. So there is a unique $m \in \Delta$ such that $\{m\} > \{t\}$ for all $t \in \Delta, t \neq m$. For $t \in \Delta^\lambda$, recall $e(t) = \sum_{g \in C(t)} \text{sgn}(g)g \cdot \{t\}$, so by the lemma,

$$e(t) = \{t\} + (\text{a signed sum of tabloids } < \{t\}).$$

Therefore,

$$0 = \alpha_m e(m) + \sum_{\substack{t \in \Delta \\ t \neq m}} \alpha_t e(t) = \alpha_m \{m\} + X \in M^\lambda,$$

where X is a linear combination of tabloids $< \{m\}$. Hence $\alpha_m = 0$, a contradiction. \square

To show that the $e(t)$ for $t \in \text{std}(\lambda)$ span \mathcal{S}^λ , we want to find elements of $\mathbb{F}S_n$ that annihilate a given $e(t)$.

Definition. *Let $\lambda \vdash n$, $t \in \Delta^\lambda$. Let $X \subseteq C_j(t)$ and $Y \subseteq C_{j+1}(t)$ for some $j \in [\lambda_1 - 1]$. Then choose T a set of left coset representatives for $S_X \times S_Y$ in $S_{X \sqcup Y}$ where we abbreviate $\text{Sym}(X) =: S_X$, etc. Define the Garnir element $G_{X,Y} := \sum_{g \in T} \text{sgn}(g)g \in \mathbb{F}S_n$.*

Example. Let $\lambda = (2, 1)$, $t = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}$, $j = 1$, $X = \{1, 3\}$, $Y = \{2\}$. Then choose $T = \{1, (12), (23)\}$ for $S_X \times S_Y = \langle (13) \rangle \times 1$ in S_3 . Then $G_{X,Y} = 1 - (12) - (23)$. Observe

$$\begin{aligned} G_{X,Y}e(t) &= (1 - (12) - (23)) \left(\frac{1 \ 2}{3} - \frac{3 \ 2}{1} \right) \\ &= \left(\frac{1 \ 2}{3} - \frac{3 \ 2}{1} \right) - \left(\frac{2 \ 1}{3} - \frac{3 \ 1}{2} \right) - \left(\frac{1 \ 3}{2} - \frac{2 \ 3}{1} \right) \\ &= 0. \end{aligned}$$

Proposition 2.19. *Let $\lambda \vdash n$, $t \in \Delta^\lambda$, $j \in [\lambda_1 - 1]$, $X \subseteq C_j(t)$, $Y \subseteq C_{j+1}(t)$. Choose a set T of left coset representatives for $S_X \times S_Y$ in $S_{X \sqcup Y}$. Then if $|X| + |Y| > (\lambda')_j$, the length of the j -th column of $Y(\lambda)$, then $G_{X,Y} \cdot e(t) = 0$*

Proof. Consider $G_{X \sqcup Y} := \sum_{\rho \in S_{X \sqcup Y}} \text{sgn}(\rho) \rho \in \mathbb{F}S_n$. Then

$$G_{X \sqcup Y} = \sum_{g \in T} \sum_{h \in S_X} \sum_{k \in S_Y} \text{sgn}(ghk)ghk = \underbrace{\left(\sum_{g \in T} \text{sgn}(g)g \right)}_{=G_{X,Y}} \left(\sum_{h \in S_X} \text{sgn}(h)h \right) \left(\sum_{k \in S_Y} \text{sgn}(k)k \right).$$

Recall from Lemma 2.2: For $\sigma \in C(t)$, $\sigma \cdot e(t) = \text{sgn}(\sigma)e(t)$, and note $S_X, S_Y \subseteq C(t)$, so

$$G_{X \sqcup Y} \cdot e(t) = G_{X,Y} |S_X| |S_Y| e(t) = |X|! |Y|! (G_{X,Y} \cdot e(t)).$$

We will show $G_{X \sqcup Y} \cdot e(t) = 0$. If $\text{char } \mathbb{F} = 0$, then we immediately deduce $G_{X,Y} \cdot e(t) = 0$, but in positive characteristic we could have $|X|! |Y|! = 0$. But once we have that $G_{X,Y} \cdot e(t) = 0$ holds in characteristic 0, then $G_{X,Y} \cdot e(t)$ is just an integer linear combination of tabloids, so we can reduce the coefficients mod p to obtain $G_{X,Y} \cdot e(t) = 0$, viewed as an \mathbb{F}_p -linear combination. Hence we have $G_{X,Y} \cdot e(t) = 0$ for all fields.

It remains to show $G_{X \sqcup Y} \cdot e(t) = 0$. For $\sigma \in C(t)$, since $|X| + |Y| > (\lambda')_j$, there exist $x_\sigma \in X, y_\sigma \in Y$ such that x_σ, y_σ lie in the same row of $\sigma \cdot t$, i.e. $(x_\sigma y_\sigma) \cdot \{\sigma \cdot t\} = \{\sigma \cdot t\}$. Let Z be a set of left coset representatives for $\langle (x_\sigma y_\sigma) \rangle$ in $S_{X \sqcup Y}$, i.e. $S_{X \sqcup Y} = Z \sqcup Z(x_\sigma y_\sigma)$. Then

$$G_{X \sqcup Y} \cdot \{\sigma \cdot t\} = \sum_{z \in Z} \text{sgn}(z) z (1 - (x_\sigma y_\sigma)) \cdot \{\sigma \cdot t\} = 0.$$

Thus

$$G_{X \sqcup Y} \cdot e(t) = \sum_{\sigma \in C(t)} \text{sgn}(\sigma) G_{X \sqcup Y} \cdot \{\sigma \cdot t\} = 0.$$

□

Definition. *Let $\lambda \vdash n$, $t, u \in \Delta^\lambda$ column-standard. Let*

$$B = \{\text{numbers not in the same column of } t \text{ and } u\}$$

$$= [n] \setminus \bigcup_{j=1}^{\lambda_1} C_j(t) \cap C_j(u).$$

If for all $\sigma \in C(t)$, $\sigma \cdot t \neq u$, then $B \neq \emptyset$, so $\max B =: x$ exists. In this case, we say $t \gg u$ if $x \in C_i(t) \cap C_j(u)$ where $i > j$.

Remark. This is the column analogue of the ordering $>$ defined earlier, except we defined it on tabloids earlier. The maximal column standard tableau w.r.t. \gg is

1	λ'_1+1	\dots	
2	λ'_1+2	\dots	n
\vdots	\vdots		
λ'_1			

Note that this tableau is standard.

Proposition 2.20. Let $\lambda \vdash n$, $v \in \Delta^\lambda$ column-standard. Then $e(v) \in \langle e(t) \mid t \in \text{std}(\lambda) \rangle_{\mathbb{F}}$.

Proof. Let $W = \langle e(t) \mid t \in \text{std}(\lambda) \rangle_{\mathbb{F}}$. Let the column-standard λ -tableaux be $t_1 \gg t_2 \gg t_3 \gg \dots$. We prove by induction on r that $e(t_r) \in W$.

Base case $r = 1$: t_1 is standard, see the remark above, so $e(t_1) \in W$.

Inductive step: Suppose $t = t_r$ where we have already shown that $e(t_s) \in W$ for all $s < r$, i.e. whenever u is column-standard and $u \gg t$, then $e(u) \in W$. Then we want to show $e(t) \in W$. If t is row-standard, then t is standard and so $e(t) \in W$. Otherwise, $t((i, j)) > t((i, j+1))$ for some $i \in [\ell(\lambda)]$, $j \in [\lambda_i - 1]$. Define $X = \{t((l, j)) \mid i \leq j \leq (\lambda')_j\}$ and $Y = \{t((l, j+1)) \mid 1 \leq l \leq i\}$. Then $G_{X,Y} \cdot e(t) = 0$ by Proposition 2.19, where $G_{X,Y}$ is defined w.r.t. any set T of coset representatives of $S_X \times S_Y$ in $S_{X \sqcup Y}$. Choose $1 \in T$. Then

$$0 = G_{X,Y} \cdot e(t) = e(t) + \sum_{g \in T \setminus \{1\}} \text{sgn}(g) g \cdot e(t).$$

We will prove that $e(g \cdot t) \in W$ for all $g \in T \setminus \{1\}$. Then we also get $e(t) \in W$ from this relation. Fix $g \in T \setminus \{1\}$. Since $g \notin S_X \times S_Y$, we must have some $y \in Y$ such that $g(y) \in X$. Hence $A := \{g(y) \mid y \in Y, g(y) \in X\} \neq \emptyset$. It is easy to see that $A = X \cap C_{j+1}(g \cdot t)$.

Consider $B := [n] \setminus \bigcup_{i=1}^{\lambda_1} C_i(t) \cap C_i(g \cdot t) \subseteq X \sqcup Y$. Moreover,

$$\begin{aligned} B &= \{x \in X \mid x \in C_{j+1}(g \cdot t)\} \sqcup \{y \in Y \mid y \in C_j(g \cdot t)\} \\ &= \underbrace{(X \cap C_{j+1}(g \cdot t))}_{=A \neq \emptyset} \sqcup (Y \cap C_j(g \cdot t)) \end{aligned}$$

Therefore $\max(B) = \max(A) \in X \cap C_{j+1}(g \cdot t)$ (using that t is column-standard and $t((i, j)) > t((i, j+1))$). Let u be the unique column-standard λ -tableau such that $C_i(u) =$

$C_l(g \cdot t)$ for all l . Then $B = [n] \setminus \bigcup_{l=1}^{\lambda_1} C_l(t) \cap C_l(u)$. We have shown that $\max(B) \in X \cap C_{j+1}(g \cdot t) \subseteq C_j(t) \cap C_{j+1}(u)$, hence $u \gg t$, so $e(u) \in W$ by inductive hypothesis. There exists $\sigma \in C(u)$ such that $\sigma \cdot u = g \cdot t$, and so $e(g \cdot t) = e(\sigma \cdot u) = \sigma \cdot e(u) = \pm e(u)$. Therefore, $e(g \cdot t) \in W$ as desired. \square

Theorem 2.21 (Standard Basis Theorem). *Let $\lambda \vdash n$, \mathbb{F} any field. Then $\{e(t) \mid t \in \text{std}(\lambda)\}$ is a basis for \mathcal{S}^λ , called the standard basis.*

Proof. Linear independence holds by Proposition 2.18. For span, let $v \in \Delta^\lambda$. Then there is a $g \in C(v)$ such that $u := g \cdot v$ is column standard. By Proposition 2.20, $e(u) \in \langle e(t) \mid t \in \text{std}(\lambda) \rangle_{\mathbb{F}}$. But $e(u) = \pm e(v)$, so we are done. \square

Note that the standard basis is not a permutation basis in general: $g \cdot e(t) = e(g \cdot t)$ for all $g \in S_n, t \in \Delta^\lambda$. But there are many g, t such that $t \in \text{std}(\lambda)$, but $g \cdot t$ is not.

Corollary 2.22. *For $\lambda \vdash n$, any field \mathbb{F} ,*

$$\dim_{\mathbb{F}} \mathcal{S}^\lambda = \#\text{standard } \lambda\text{-tableaux.}$$

3 Character Theory

From now on, $\mathbb{F} = \mathbb{C}$, unless otherwise stated.

Notation. Let $\lambda \vdash n$. We will let χ^λ denote the character of the irreducible λ -Specht module.

3.1 Hook Length Formula

Goal. Prove the hook length formula, a closed formula for calculating $\dim \mathcal{S}^\lambda = \chi^\lambda(1)$.

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$. Write $\lambda' = (\mu_1, \dots, \mu_{\lambda_1})$.

(i) For a box $(i, j) \in Y(\lambda)$, the (i, j) -hook of λ is

$$H_{i,j}(\lambda) := \{(i, j)\} \sqcup \underbrace{\{(i, y) \mid j < y \leq \lambda_i\}}_{\text{arm}} \sqcup \underbrace{\{(x, j) \mid i < x \leq \mu_j\}}_{\text{leg}}.$$


(ii) The arm of $H_{i,j}(\lambda)$ is $\{(i, j) \mid j < y \leq \lambda_i\}$, the leg is $\{(x, j) \mid i < x \leq \mu_j\}$.

(iii) The hand of $H_{i,j}(\lambda)$ is the box (i, λ_i) , the foot is (μ_i, j) .

(iv) The hook length corresponding to (i, j) is $|H_{i,j}(\lambda)| =: h_{i,j}(\lambda)$.

(v) Let $\mathcal{H}(\lambda) = \{h_{i,j}(\lambda) \mid (i, j) \in Y(\lambda)\}$ be the multiset of hook lengths of λ (i.e. we also count repetitions of the same hook length).

Example. Take $\lambda = (8, 6, 5, 4, 2, 1) \vdash 26$, $(i, j) = (2, 3)$. Then the hook is $\{\bullet\} \sqcup \text{arm} \sqcup \text{leg}$ as indicated in the diagram.



Theorem 3.1 (Hook Length Formula). Let $\lambda \vdash n$. Then

$$\chi^\lambda(1) = \frac{n!}{\prod_{h \in \mathcal{H}(\lambda)} h}.$$

Examples.

(a) Let $\lambda = (n)$. List the hook lengths in $Y(\lambda)$: $\boxed{n \cdots 2 \ 1}$. So $\chi^\lambda(1) = \frac{n!}{n!} = 1$. This is not unexpected as we already knew that $\mathcal{S}^\lambda \cong \mathbb{1}_{S_n}$.

(b) Let $\lambda = (3, 2) \vdash 5$. Then

$$\text{std}(\lambda) = \left\{ \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline \end{array} \right\},$$

so $\chi^\lambda(1) = 5$ by the standard basis theorem. This is consistent with the hook length formula. Indeed, the hook lengths are $\begin{array}{|c|c|c|} \hline 4 & 3 & 1 \\ \hline 2 & 1 & \\ \hline \end{array}$, so $\chi^\lambda(1) = \frac{5!}{4 \cdot 3 \cdot 2} = 5$.

(c) Let $\lambda = (6, 4, 4, 3, 2, 1, 1) \vdash 21$. Then the hook lengths are

12	9	7	5	2	1
9	6	4	2		
8	5	3	1		
6	3	1			
4	1				
2					
1					

Therefore

$$\chi^\lambda(1) = \frac{21!}{\prod_{h \in \mathcal{H}(\lambda)} h} = 905304400.$$

We give a probabilistic proof of the hook length formula due to Greene, Nijenhuis and Wilf (1979). Another proof will be on the example sheets. The proof will be by induction on n .

Definition. By a composition of n , we mean a sequence of non-negative integers which sum to n , written $\lambda \models n$.

Define a function F on $\{\lambda \mid \lambda \models n\}$ as follows:

$$F(\lambda) = \begin{cases} \frac{n!}{\prod_{h \in \mathcal{H}(\lambda)} h} & \text{if } \lambda \vdash n, \\ 0 & \text{otherwise.} \end{cases}$$

If $\lambda = (\lambda_1, \dots, \lambda_k) \models n$, we want the inductive step to look like

$$F(\lambda) = \sum_{i=1}^k F(\underbrace{(\lambda_1, \dots, \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, \dots, \lambda_k)}_{\models n-1 \text{ if } \lambda_i \geq 1}).$$

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$. Define

$$\begin{aligned} \lambda^- &:= \{\mu \vdash n-1 \mid Y(\mu) \text{ can be obtained from } Y(\lambda) \text{ by removing one box}\} \\ &= \{(\lambda_1, \dots, \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, \dots, \lambda_k) \mid i \in [k] \text{ such that } \lambda_i - 1 \geq \lambda_{i+1}\}. \end{aligned}$$

(Here we treat $\lambda_{k+1} = 0$.)

We say the box (i, j) of $Y(\lambda)$ is removable if $Y(\lambda) \setminus \{(i, j)\} = Y(\mu)$ for some $\mu \in \lambda^-$.

Example. Let $\lambda = (3, 3, 1) \vdash 7$, so $Y(\lambda) =$

. Then

$$\lambda^- = \left\{ \begin{array}{c} (3, 3) \\ \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \end{array} , \begin{array}{c} (3, 2, 1) \\ \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \end{array} \right\}.$$

Observe that $\chi^\lambda(1) = \sum_{\mu \in \lambda^-} \chi^\mu(1)$. This follows from the Standard Basis Theorem. Indeed, $\chi^\lambda(1) = |\text{std}(\lambda)|$ and in a standard λ -tableau, $\lambda \vdash n$, the number n must appear in a removable box, which when removed, leaves a standard μ -tableau for some $\mu \in \lambda^-$.

We would be able to prove Theorem 3.1 by induction on n if we can show

$$F(\lambda) = \sum_{\mu \in \lambda^-} F(\mu),$$

because we would have $\sum_{\mu \in \lambda^-} F(\mu) = \sum_{\mu \in \lambda^-} \chi^\mu(1)$ by the inductive hypothesis.

We will in fact show that $1 = \sum_{\mu \in \lambda^-} \frac{F(\mu)}{F(\lambda)}$ by interpreting $\frac{F(\mu)}{F(\lambda)}$ as probabilities. For the rest of this section, fix $\lambda \vdash n$, and abbreviate $H_{i,j}(\lambda) = H_{i,j}$ and $h_{i,j}(\lambda) = h_{i,j}$.

Consider the following probabilistic process on $Y(\lambda)$:

- **Step 1.** Pick a box of $Y(\lambda)$ uniformly at random (probability = $\frac{1}{n}$).
- **Step 2.** Suppose that (i, j) is the currently chosen box. If (i, j) is removable, equivalently $h_{i,j} = 1$, then terminate the process. Otherwise, choose $(i', j') \in H_{i,j} \setminus \{(i, j)\}$ (probability = $\frac{1}{h_{i,j}-1}$).
- **Step 3.** Repeat Step 2 until we terminate.

We will call each run of the process a *trial*.

Definition. For $(\alpha, \beta) \in Y(\lambda)$, let $\mathbb{P}(\alpha, \beta)$ be the probability that a trial terminates at (α, β) .

Our aim is to show that $\mathbb{P}(\alpha, \beta) = \frac{F(\mu)}{F(\lambda)}$ where $\mu \in \lambda^-$ and $Y(\mu) = Y(\lambda) \setminus \{(\alpha, \beta)\}$ (note that if a trial terminates at (α, β) , then this is necessarily a removable box, so this makes sense).

Definition. Let $\pi : (a_1, b_1) \rightarrow (a_2, b_2) \rightarrow \dots \rightarrow (a_m, b_m)$ be a trial of the process. Define $A_\pi = \{a_1, \dots, a_m\}$, the set of horizontal projections of π . Analogously, let $B_\pi = \{b_1, \dots, b_m\}$, the set of vertical projections of π .

Example. Let $\lambda = (4, 4, 3, 3, 2)$. We could have the trial

1	2		
	3	4	

where we indicate the box we are in at time t by t . So $\pi : (2, 1) \rightarrow (2, 2) \rightarrow (4, 2) \rightarrow (4, 3)$. Then

$$A_\pi = \{2, 4\}, \quad B_\pi = \{1, 2, 3\}.$$

Observe that for $\pi : (a_1, b_1) \rightarrow \cdots \rightarrow (a_m, b_m)$,

- the starting box (a_1, b_1) must equal $(\min A_\pi, \min B_\pi)$.
- the last box (a_m, b_m) must equal $(\max A_\pi, \max B_\pi)$.
- for each $i \in [m-1]$, either $a_i < a_{i+1}$ and $b_i = b_{i+1}$ (step down), or $a_i = a_{i+1}$ and $b_i < b_{i+1}$ (step right). So $m = |A_\pi| + |B_\pi| - 1$.

Definition. Given $(a, b) \in Y(\lambda)$, $A, B \subseteq \mathbb{N}$, define $\mathbb{P}(A, B \mid a, b)$ to be the probability that a trial π starting at box (a, b) has $A_\pi = A, B_\pi = B$.

Outline of proof of the hook length formula:

- We will calculate $\mathbb{P}(A, B \mid a, b)$ in terms of $\frac{1}{h_{ij}-1}$ for various i, j .
- For $\mu \in \lambda^-$, we will calculate $\frac{F(\mu)}{F(\lambda)}$ as a product of terms of the form $\frac{1}{h_{i,j}-1}$, and interpret the terms in the expansion as probabilities of the form $\mathbb{P}(A, B \mid a, b)$.
- We will show $\mathbb{P}(\alpha, \beta)$, the probability that a trial terminates at (α, β) , is

$$\sum_{\substack{\text{possible} \\ A, B}} \sum_{\substack{\text{projections starting} \\ \text{box} \\ (a, b)}} \mathbb{P}(A, B \mid a, b)$$

to conclude $\mathbb{P}(\alpha, \beta) = \frac{F(\mu)}{F(\lambda)}$, where $\mu \in \lambda^-$ satisfies $Y(\mu) = Y(\lambda) \setminus \{(\alpha, \beta)\}$.

Lemma 3.2. Let $(\alpha, \beta) \in Y(\lambda)$ be removable. Let $A = \{a_1, \dots, a_t\}, B = \{b_1, \dots, b_u\} \subseteq \mathbb{N}$, where $a_1 < a_2 < \cdots < a_t = \alpha$, $b_1 < b_2 < \cdots < b_u = \beta$. Then

$$\mathbb{P}(A, B \mid a_1, b_1) = \prod_{x \in A \setminus \{\alpha\}} \frac{1}{h_{x, \beta} - 1} \prod_{y \in B \setminus \{\beta\}} \frac{1}{h_{\alpha, y} - 1}.$$

Proof. Induct on $t + u = |A| + |B|$. Base case $t + u = 2$, then $A = \{a_1 = \alpha\}$ and $B = \{b_1 = \beta\}$. Then $\mathbb{P}(A, B \mid a, b) = 1$ which is also the value of the RHS which is an empty product. For the inductive step now suppose $t + u > 2$, and so $(a_1, b_1) \neq (\alpha, \beta)$. Condition on the second box in the trial:

$$\mathbb{P}(A, B \mid a_1, b_1) = \sum_{\substack{(a', b') \\ \in H_{a_1, b_1} \setminus \{(a_1, b_1)\}}} \left[\mathbb{P} \left(\begin{array}{l} \text{proj. sets} \\ = A, B \end{array} \mid \begin{array}{l} \text{first box is } (a_1, b_1) \text{ and} \\ \text{second box is } (a', b') \end{array} \right) \right. \\ \left. \cdot \mathbb{P}(\text{second box is } (a', b') \mid \text{first box is } (a_1, b_1)) \right]$$

$$\begin{aligned}
&= \sum_{(a',b') \in \text{arm of } H_{a_1,b_1}} + \sum_{(a',b') \in \text{leg of } H_{a_1,b_1}} \\
&= \sum_{b_1 < b' \leq \lambda_{a_1}} \mathbb{P}(A, B \setminus \{b_1\} \mid a_1, b') \mathbb{P}\left(\begin{array}{c} \text{second box} \\ = (a_1, b') \end{array} \mid \begin{array}{c} \text{first box} \\ = (a_1, b_1) \end{array} \right) \\
&\quad + \sum_{a_1 < a' \leq (\lambda')_{b_1}} \mathbb{P}(A \setminus \{a_1\}, B \mid a', b_1) \mathbb{P}\left(\begin{array}{c} \text{second box} \\ = (a', b_1) \end{array} \mid \begin{array}{c} \text{first box} \\ = (a_1, b_1) \end{array} \right) \\
&= \frac{1}{h_{a_1,b_1} - 1} \left(\sum_{b_1 < b' \leq \lambda_{a_1}} \mathbb{P}(A, B \setminus \{b_1\} \mid a_1, b') \right. \\
&\quad \left. + \sum_{a_1 < a' \leq (\lambda')_{b_1}} \mathbb{P}(A \setminus \{a_1\}, B \mid a', b_1) \right)
\end{aligned}$$

Note that $\mathbb{P}(A, B \setminus \{b_1\} \mid a_1, b') = 0$ unless $b' = b_2$. Indeed, if $b' \neq b_2$ in a trial, then

- either $b_1 < b' < b_2$: b' is in the vertical projection set, but $b' \notin B \setminus \{b_1\}$.
- or $b' > b_2$: b' is in the vertical projection set, but b_2 is not in the vertical projection set.

Similarly, $\mathbb{P}(A \setminus \{a_1\}, B \mid a', b_1) = 0$ unless $a' = a_2$. Therefore,

$$\mathbb{P}(A, B \mid a_1, b_1) = \frac{1}{h_{a_1,b_1} - 1} \left(\mathbb{P}(A, B \setminus \{b_1\} \mid a_1, b_2) + \mathbb{P}(A \setminus \{a_1\}, B \mid a_2, b_1) \right).$$

If one of u, t is 1, we simply omit the corresponding term. By the induction hypothesis, this is

$$\begin{aligned}
&\frac{1}{h_{a_1,b_1} - 1} \left(\prod_{x \in A \setminus \{\alpha\}} \frac{1}{h_{x,\beta} - 1} \prod_{y \in B \setminus \{\beta, b_1\}} \frac{1}{h_{\alpha,y} - 1} + \prod_{x \in A \setminus \{\alpha, a_1\}} \frac{1}{h_{x,\beta} - 1} \prod_{y \in B \setminus \{\beta\}} \frac{1}{h_{\alpha,y} - 1} \right) \\
&= \frac{(h_{\alpha,b_1} - 1) + (h_{a_1,\beta} - 1)}{h_{a_1,b_1} - 1} \prod_{x \in A \setminus \{\alpha\}} \frac{1}{h_{x,\beta} - 1} \prod_{y \in B \setminus \{\beta\}} \frac{1}{h_{\alpha,y} - 1}
\end{aligned}$$

Now draw a picture to see why $(h_{\alpha,b_1} - 1) + (h_{a_1,\beta} - 1) = h_{a_1,b_1} - 1$, so the first term disappears and we are done. \square

Proposition 3.3. *Let $(\alpha, \beta) \in Y(\lambda)$ be a removable box. Suppose $\mu \in \lambda^-$ is such that $Y(\mu) = Y(\lambda) \setminus \{(\alpha, \beta)\}$. Then*

$$\mathbb{P}(\alpha, \beta) = \frac{F(\mu)}{F(\lambda)}.$$

Proof. Observe that

- $h_{x,y}(\mu) = h_{x,y}(\lambda)$ if $x \neq \alpha$ and $y \neq \beta$,

- $h_{\alpha,y}(\mu) = h_{\alpha,y}(\lambda) - 1$ if $y \neq \beta$,
- $h_{x,\beta}(\mu) = h_{x,\beta}(\lambda) - 1$ if $x \neq \alpha$.

Thus,

$$\begin{aligned}
\frac{F(\mu)}{F(\lambda)} &= \frac{\prod_{h \in \mathcal{H}(\lambda)} h}{n!} \frac{(n-1)!}{\prod_{h \in \mathcal{H}(\mu)} h} \\
&= \frac{1}{n} \prod_{1 \leq x < \alpha} \frac{h_{x,\beta}}{h_{x,\beta} - 1} \prod_{1 \leq y < \beta} \frac{h_{\alpha,y}}{h_{\alpha,y} - 1} \\
&= \frac{1}{n} \prod_{1 \leq x < \alpha} \left(1 + \frac{1}{h_{x,\beta} - 1}\right) \prod_{1 \leq y < \beta} \left(1 + \frac{1}{h_{\alpha,y} - 1}\right)
\end{aligned}$$

We want to interpret the terms in the expansion as the probabilities that a trial terminating at (α, β) has certain horizontal and vertical projections. We have

$$\begin{aligned}
\prod_{1 \leq x < \alpha} \left(1 + \frac{1}{h_{x,\beta} - 1}\right) &= \left(1 + \frac{1}{h_{1,\beta} - 1}\right) \left(1 + \frac{1}{h_{2,\beta} - 1}\right) \cdots \left(1 + \frac{1}{h_{\alpha-1,\beta} - 1}\right) \\
&= \sum_{\substack{A \subseteq [\alpha] \\ \alpha \in A}} \prod_{x \in A \setminus \{\alpha\}} \frac{1}{h_{x,\beta} - 1}
\end{aligned}$$

and similarly

$$\prod_{1 \leq y < \beta} \left(1 + \frac{1}{h_{\alpha,y} - 1}\right) = \sum_{\substack{B \subseteq [\beta] \\ \beta \in B}} \prod_{y \in B \setminus \{\beta\}} \frac{1}{h_{\alpha,y} - 1}.$$

Then

$$\begin{aligned}
\frac{F(\mu)}{F(\lambda)} &= \frac{1}{n} \sum_{\substack{A \subseteq [\alpha], \alpha \in A \\ B \subseteq [\beta], \beta \in B}} \prod_{x \in A \setminus \{\alpha\}} \frac{1}{h_{x,\beta} - 1} \prod_{y \in B \setminus \{\beta\}} \frac{1}{h_{\alpha,y} - 1} \\
&= \frac{1}{n} \sum_{\substack{A \subseteq [\alpha], \alpha \in A \\ B \subseteq [\beta], \beta \in B}} \mathbb{P}(A, B \mid \min(A), \min(B)).
\end{aligned}$$

Also, $\mathbb{P}(\alpha, \beta)$, the probability of terminating at (α, β) , is

$$\begin{aligned}
&\sum_{(a,b) \in Y(\lambda)} \mathbb{P}\left(\begin{array}{c} \text{terminate at} \\ (\alpha, \beta) \end{array} \middle| \begin{array}{c} \text{start at} \\ (a, b) \end{array} \right) \cdot \mathbb{P}(\text{start at } (a, b)) \\
&= \frac{1}{n} \sum_{(a,b) \in Y(\lambda)} \mathbb{P}\left(\begin{array}{c} \text{terminate at} \\ (\alpha, \beta) \end{array} \middle| \begin{array}{c} \text{start at} \\ (a, b) \end{array} \right) \\
&= \frac{1}{n} \sum_{(a,b) \in Y(\lambda)} \sum_{A', B'} \mathbb{P}(A', B' \mid a, b)
\end{aligned}$$

where the second sum runs over $A' \subseteq [\alpha], B' \subseteq [\beta]$ such that $\alpha = \max A', a = \min A', \beta = \max B', b = \min B'$. We conclude $\mathbb{P}(\alpha, \beta) = \frac{F(\mu)}{F(\lambda)}$. \square

Proof of Theorem 3.1. Since a trial must terminate at a removable box,

$$1 = \sum_{(\alpha, \beta) \text{ removable}} \mathbb{P}(\alpha, \beta) = \sum_{\mu \in \lambda^-} \frac{F(\mu)}{F(\lambda)}.$$

So we are done by induction on n , as previously described. \square

3.2 The Determinantal Form

Applications.

- Recall the permutation module $M^\lambda \cong \mathbb{1}_{S_\lambda} \uparrow^{S_n}$, see Lemma 2.1. In a direct sum decomposition of M^λ into irreducibles, how many times do we get S^μ ? \rightsquigarrow Young's Rule, Theorem 3.11 and Corollary 3.19.
- We have a nested structure: $S_1 < S_2 < \dots < S_{n-1} < S_n < \dots$. How do S_n -modules relate to S_{n-1} -modules?

E.g. $V_n \downarrow_{S_{n-1}}^{S_n} \cong V_{n-1} \oplus \mathbb{1}_{S_{n-1}}$ where V_n is the natural permutation module of S_n .
What is $\mathcal{S}^\lambda \downarrow_{S_{n-1}}^{S_n}$? \rightsquigarrow Branching Rule, Theorem 3.22.

- What is $\chi^\lambda(g)$ for all $g \in S_n$? \rightsquigarrow Murnaghan-Nakayama Rule, Theorem 3.25.
- And more:
 - e.g. Branching Rule describes $\mathcal{S}^\lambda \downarrow_{S_{n-1} \times S_1}^{S_n}$. What about $\mathcal{S}^\lambda \downarrow_{S_{n-m} \times S_m}^{S_n}$? \rightsquigarrow Littlewood-Richardson Rule.
 - e.g. another proof of the hook length formula, see Example Sheet 2.

Notation.

- S_n is the symmetric group, S_λ Young subgroups
- Before: \mathcal{S}^μ were Specht modules. For the rest of this chapter we use $[\mu]$ to replace \mathcal{S}^μ to denote the μ -Specht module. When it is clear from context, for $\mu = (m)$, we abbreviate $[\mu] = [(m)]$ to $[m]$.
- Let ξ^λ be the character of M^λ .

Definition.

- Let G, H be finite groups, V a G -module, W an H -module. Then $V \otimes W$ can be into a $(G \times H)$ -module via

$$(g, h) \cdot (v \otimes w) = (gv) \otimes (hw)$$

for all $g \in G, h \in H, v \in V, w \in W$. The resulting $(G \times H)$ -module is the (outer) tensor product of V and W , which we will denote by $V \# W$. If V affords χ , and W affords ϕ , then $V \# W$ affords $\chi \# \phi$ where

$$(\chi \# \phi)((g, h)) = \chi(g)\phi(h)$$

for all $g \in G, h \in H$.

- Let $m, n \in \mathbb{N}, \alpha \vdash m, \beta \vdash n$. Then $\chi^\alpha \# \chi^\beta \in \text{Irr}(S_m \times S_n)$ since $\chi^\alpha \in \text{Irr}(S_m), \chi^\beta \in \text{Irr}(S_n)$. Note that $S_m \times S_n$ naturally embeds inside S_{m+n} as $\text{Sym}\{1, 2, \dots, m\} \times \text{Sym}\{m+1, \dots, m+n\}$. Then the outer product of $[\alpha]$ and $[\beta]$ is defined as

$$[\alpha][\beta] = [\alpha] \# [\beta] \uparrow_{S_m \times S_n}^{S_{m+n}}$$

Remarks.

- The outer product is associative and commutative.
- Let $H \leq G, x \in G$. Then $\mathbb{1}_H \uparrow^G \cong \mathbb{1}_{xHx^{-1}} \uparrow^G$. Suppose that $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$. Recall $S_\lambda \cong S_{\lambda_1} \times \dots \times S_{\lambda_k}$. We may fix $S_\lambda = \text{Sym}\{1, \dots, \lambda_1\} \times \text{Sym}\{\lambda_1+1, \dots, \lambda_1+\lambda_2\} \times \dots \times \text{Sym}\{\sum_{i=1}^{k-1} \lambda_i+1, \dots, n\}$ when we consider $M^\lambda \cong \mathbb{1}_{S_\lambda} \uparrow^{S_n}$, since all Young subgroups of type λ are conjugate to this one.

Also,

$$M^\lambda \cong \mathbb{1}_{S_\lambda} \uparrow^{S_n} = \mathbb{1}_{S_{\lambda_1}} \# \mathbb{1}_{S_{\lambda_2}} \# \dots \# \mathbb{1}_{S_{\lambda_k}} \uparrow^{S_n} = [\lambda_1][\lambda_2] \dots [\lambda_k],$$

and so $[\lambda_1][\lambda_2] \dots [\lambda_k]$ has character ξ^λ .

Example. For $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$, consider the $k \times k$ -matrix \mathcal{D}_λ whose (i, j) -entry is the module $[\lambda_i - i + j]$ where we interpret $[l]$ as the zero module when $l < 0$.

- Let $\lambda = (n-1, 1)$. Then, using the outer product to multiply modules,

$$\det \mathcal{D}_\lambda = \det \begin{pmatrix} [n-1] & [n] \\ [0] & [1] \end{pmatrix} = [n-1][n] - [n][0].$$

This (virtual) representation has (virtual) character

$$\xi^{(n-1,1)} - \xi^{(n)} = \xi^{(n-1,1)} - \chi^{(n)} = \chi^{(n-1,1)} = \chi^\lambda$$

by Example Sheet 1, Question 5.

- Let $\lambda = (3, 1^2) \vdash 5$. Then

$$\begin{aligned} \det \mathcal{D}_\lambda &= \det \begin{pmatrix} [3] & [4] & [5] \\ [0] & [1] & [2] \\ 0 & [0] & [1] \end{pmatrix} = [3] \det \begin{pmatrix} [1] & [2] \\ [0] & [1] \end{pmatrix} - [0] \det \begin{pmatrix} [4] & [5] \\ [0] & [1] \end{pmatrix} \\ &= [3][1][1] - [3][2][0] - [4][1][0] + [5][0][0] \end{aligned}$$

which has (virtual) character

$$\xi^{(3,1^2)} - \xi^{(3,2)} - \xi^{(4,1)} + \xi^{(5)} = \xi^{(3,1^2)} = \xi^\lambda.$$

Definition. A virtual character of G is a \mathbb{Z} -linear combination of irreducible characters of G .

Definition. Let $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$. Let \mathcal{D}_λ be the $k \times k$ matrix whose (i, j) -entry is the module $[\lambda_i - i + j]$ (i.e. as in the example above).

We could in fact have defined $\mathcal{D}_\lambda = [\lambda_i - i + j]_{ij}$ to be $k' \times k'$ for any $k' \geq k$, and the determinant remains unchanged. E.g. for $\lambda = (3, 1, 1)$,

$$\det \begin{pmatrix} [3] & [4] & [5] \\ [0] & [1] & [2] \\ 0 & [0] & [1] \end{pmatrix} = \det \begin{pmatrix} [3] & [4] & [5] & [6] \\ [0] & [1] & [2] & [3] \\ 0 & [0] & [1] & [2] \\ 0 & 0 & 0 & [0] \end{pmatrix},$$

viewing $(3, 1, 1) = (3, 1, 1, 0, 0, \dots)$.

Goal. Prove that $\det \mathcal{D}_\lambda$ has character χ^λ for all $\lambda \vdash n$.

For the rest of this chapter, we will work with $\mathbb{Z}^{\mathbb{N}}$, the set of sequences with integer entries, under pointwise addition.

Let $n \in \mathbb{N}$. Summary:

Term	Notation	Def.: $\lambda = (\lambda_1, \lambda_2, \dots) \in \mathbb{Z}^{\mathbb{N}}$ s.t. $\sum_i \lambda_i = n$ and
partition of n	$\lambda \vdash n$	$\lambda_1 \geq \lambda_2 \geq \dots$ and $\lambda_i \in \mathbb{N}_0$ for all i
composition of n	$\lambda \models n$	$\lambda_i \in \mathbb{N}_0$ for all i
integer composition of n	$\lambda \vDash n$	only finitely many λ_i are non-zero

(a) Recall $S_n = \text{Sym}\{1, 2, \dots, n\}$. Define $S_{\mathbb{N}} = \bigcup_{n \in \mathbb{N}} S_n$.

- For $\pi \in S_{\mathbb{N}}$, we can view it as an element of $\mathbb{Z}^{\mathbb{N}}$ via $\pi = (\pi^{-1}(1), \pi^{-1}(2), \dots)$. Note that π does not have finite support, but $\pi^{-1}(i) = i$ for all sufficiently large i . In particular, the identity of $S_{\mathbb{N}}$ is $\text{id} = (1, 2, 3, \dots)$.
- For $\pi \in S_{\mathbb{N}}$ and $\lambda \in \mathbb{Z}^{\mathbb{N}}$, we define $\pi \cdot \lambda := (\lambda_{\pi^{-1}(1)}, \lambda_{\pi^{-1}(2)}, \dots)$. Then $\pi \cdot \text{id} = \text{id} \cdot \pi = \pi$, $\pi \cdot \pi^{-1} = \pi^{-1} \cdot \pi = \text{id}$, and $\tau \cdot (\pi \cdot \lambda) = (\tau\pi) \cdot \lambda$.
- For $\pi \in S_{\mathbb{N}}$ and $\lambda \vDash n$, observe that $\pi \cdot \lambda \vDash n$. Also $\lambda - \text{id} + \pi = (\lambda_1 - 1 + \pi^{-1}(1), \lambda_2 - 2 + \pi^{-1}(2), \dots) \vDash n$.

(b) In the above, we let λ_j be the j -th entry of λ as usual. If λ has finite support, we can define $\ell(\lambda) = \max\{i \in \mathbb{N} \mid \lambda_i \neq 0\}$. We may write $(\lambda_1, \dots, \lambda_{\ell(\lambda)})$ and $(\lambda_1, \dots, \lambda_{\ell(\lambda)}, 0, 0, \dots)$ interchangeably.

(c) We can extend Young subgroups to have type given by compositions, not just partitions; these will be conjugate to Young subgroups of type given by partitions. E.g. $S_{(1,0,0,2,0,0,\dots)} = S_{(1,2)} = \text{Sym}\{1\} \times \text{Sym}\{2, 3\}$ is conjugate to $S_{(2,1)} = \text{Sym}\{1, 2\} \times \text{Sym}\{3\}$.

(d) We can extend ξ^λ to be defined for all integer compositions $\lambda \equiv n$ by:

$$\xi^\lambda = \begin{cases} \mathbb{1}_{S_\lambda} \uparrow^{S_n} & \text{if } \lambda \vdash n, \\ 0 & \text{otherwise.} \end{cases}$$

So for all $\lambda \equiv n$, $[\lambda_1][\lambda_2] \dots [\lambda_{\ell(\lambda)}]$ has character ξ^λ , since $[r] = 0$ if $r < 0$.

(e) We could e.g. dominance partial ordering to $\lambda \equiv n$, Young diagrams, \mathcal{D}_λ , etc.

Definition. For $\lambda \equiv n$, define

$$\psi^\lambda = \sum_{\pi \in S_{\mathbb{N}}} \text{sgn}(\pi) \xi^{\lambda - \text{id} + \pi},$$

it is a virtual character of S_n .

Lemma 3.4. Let $\lambda \equiv n$.

- (i) Only finitely many terms in the sum defining ψ^λ are non-zero.
- (ii) The virtual character afforded by $\det \mathcal{D}_\lambda = \det([\lambda_i - i + j]_{ij})$ is ψ^λ .

Proof.

- (i) Since λ has finite support, $k = \ell(\lambda)$ is defined. Let $\pi \in S_{\mathbb{N}} \setminus S_k$. We claim that $\lambda - \text{id} + \pi$ has a negative entry. Indeed, let $m := \max\{i \mid \pi^{-1}(i) \neq i\}$. Since $\pi \notin S_k$, we must have $m > k$. By maximality of m , we must have $\pi^{-1}(m) < m$. Then $(\lambda - \text{id} + \pi)_m = \lambda_m - m + \pi^{-1}(m) = \pi^{-1}(m) - m < 0$. So $\xi^{\lambda - \text{id} + \pi} = 0$ for such π , and so $\psi^\lambda = \sum_{\pi \in S_k} \text{sgn}(\pi) \xi^{\lambda - \text{id} + \pi}$ is a finite sum.
- (ii) Recall that the determinant of a $k \times k$ matrix D is given by

$$\det D = \sum_{\pi \in S_k} \text{sgn} \pi \prod_{i=1}^k D_{i, \pi(i)}.$$

The claim follows since $[\alpha_1][\alpha_2] \dots [\alpha_{\ell(\alpha)}]$ has character ξ^α for all $\alpha \equiv n$. □

So our goal is to show $\psi^\lambda = \chi^\lambda$ for all $\lambda \vdash n$.

Lemma 3.5. Let $\lambda \equiv n$. Let $i \in \mathbb{N}$ and suppose that $\mu \equiv n$ satisfies $\mu - \text{id} = (i \ i + 1) \cdot (\lambda - \text{id})$, i.e.

$$\mu = (\lambda_1, \dots, \lambda_{i-1}, \lambda_{i+1} - 1, \lambda_i + 1, \lambda_{i+2}, \dots).$$

Then $\psi^\mu = -\psi^\lambda$. In particular, if $\lambda_i - i = \lambda_{i+1} - (i + 1)$ for some $i \in \mathbb{N}$, then $\psi^\lambda = 0$.

Proof. Let $\tau = (i \ i+1)$. Then $\mu - \text{id} = \tau \cdot (\lambda - \text{id})$, so $\mu - \text{id} + \tau\pi = \tau \cdot (\lambda - \text{id} + \pi)$ for any $\pi \in S_{\mathbb{N}}$. Hence

$$\begin{aligned}\psi^\lambda &= \sum_{\pi \in S_{\mathbb{N}}} \text{sgn}(\pi) \xi^{\lambda - \text{id} + \pi} = \sum_{\pi \in S_{\mathbb{N}}} \text{sgn}(\pi) \xi^{\tau \cdot (\lambda - \text{id} + \pi)} \\ &= \sum_{\pi \in S_{\mathbb{N}}} \text{sgn}(\pi) \xi^{\mu - \text{id} + \tau\pi} = - \sum_{\pi \in S_{\mathbb{N}}} \text{sgn}(\tau\pi) \xi^{\mu - \text{id} + \tau\pi} = -\psi^\mu.\end{aligned}$$

If $\lambda_i - i = \lambda_{i+1} - (i+1)$, then $\mu = \lambda$, and so $\psi^\lambda = -\psi^\lambda$, so $\psi^\lambda = 0$. \square

Next we look at $\xi^\lambda \downarrow_{S_m \times S_k}$ where $\lambda \equiv n = m + k$. Note that $\xi^\lambda \downarrow_{S_m \times S_k} = \mathbb{1}_{S_\lambda} \uparrow^{S_n} \downarrow_{S_m \times S_k}$, so we will use Mackey's theorem. For this we will need to know the double cosets of S_λ , $S_m \times S_k$ in S_n .

Proposition 3.6. *Let $\lambda, \mu \vdash n$. There is a bijection between the set of double cosets of S_λ and S_μ in S_n , and the set of $\ell(\lambda) \times \ell(\mu)$ -matrices with entries in \mathbb{N}_0 whose row sums are λ , and column sums are μ .*

Proof. Write $S_\lambda = S_{A_1} \times S_{A_1} \times \cdots \times S_{A_{\ell(\lambda)}}$ where $A_1 = [\lambda_1], A_2 = \{\lambda_1 + 1, \dots, \lambda_1 + \lambda_2\}, \dots$, and $S_\mu = S_{B_1} \times S_{B_2} \times \cdots \times S_{B_{\ell(\mu)}}$ similarly.

For each $\sigma \in S_n$, define a matrix $Z(\sigma)$ via $Z(\sigma)_{ij} := |A_i \cap \sigma(B_j)|$ for all i, j . Note that the i -th row sum is

$$\sum_j |A_i \cap \sigma(B_j)| = |A_i \cap \bigcup_j \sigma(B_j)| = |A_i \cap [n]| = |A_i| = \lambda_i,$$

and similarly the j -th column sum is

$$\sum_i |A_i \cap \sigma(B_j)| = |\sigma(B_j)| = |B_j| = \mu_j.$$

Conversely, any matrix in the set described in the proposition is $Z(\sigma)$ for some $\sigma \in S_n$ (exercise).

Now we claim that for $\sigma, \tau \in S_n$, we have $Z(\sigma) = Z(\tau)$ iff $S_\lambda \sigma S_\mu = S_\lambda \tau S_\mu$. First, suppose $\tau = h\sigma k$ for some $h \in S_\lambda, k \in S_\mu$. Then $Z(\tau)_{ij} = |A_i \cap \tau(B_j)| = |A_i \cap h\sigma k(B_j)| = |A_i \cap h\sigma(B_j)|$ since $k \in S_\mu$, so that $k(B_j) = B_j$ for all j . Similarly, $h^{-1}(A_i) = A_i$, so $Z(\tau)_{ij} = |h^{-1}(A_i) \cap \sigma(B_j)| = |A_i \cap \sigma(B_j)| = Z(\sigma)_{ij}$. Conversely, suppose that $|A_i \cap \sigma(B_j)| = |A_i \cap \tau(B_j)|$ for all i, j . For each fixed i , $\{A_i \cap \sigma(B_j)\}_j$ and $\{A_i \cap \tau(B_j)\}_j$ are both partitions of the set A_i . But $|A_i \cap \sigma(B_j)| = |A_i \cap \tau(B_j)|$ for all j , so there exists $h_i \in S_{A_i}$ such that $h_i(A_i \cap \sigma(B_j)) = A_i \cap \tau(B_j)$ for all j . Then $h := h_1 \cdot h_2 \cdots h_{\ell(\lambda)} \in S_{A_1} \times \cdots \times S_{A_{\ell(\lambda)}} = S_\lambda$ satisfies $h(\sigma(B_j)) = \tau(B_j)$ for all j . Therefore $\tau^{-1}h\sigma(B_j) = B_j$ for all j , and so $\tau^{-1}h\sigma \in S_\mu$. Say $\tau^{-1}h\sigma = k^{-1}$, then $\tau = h\sigma k$ where $h \in S_\lambda, k \in S_\mu$.

Thus $S_\lambda \sigma S_\mu \mapsto Z(\sigma)$ is a well-defined bijection between the two sets in the proposition. \square

Lemma 3.7. *Let $\lambda \models n = m + k$, $m, k \in \mathbb{N}_0$. Then*

$$(i) \quad \xi^\lambda \downarrow_{S_m \times S_k} = \sum_{\mu \models k} \xi^{\lambda - \mu} \# \xi^\mu,$$

$$(ii) \quad \psi^\lambda \downarrow_{S_m \times S_k} = \sum_{\mu \models k} \psi^{\lambda - \mu} \# \xi^\mu.$$

Proof.

- (i) Both sides of (i) are equal to zero if $\lambda \not\models n$. So we may now assume that $\lambda \models n$. Also, note that the sum over $\mu \models k$ is finite since $\xi^{\lambda - \mu} = 0$ unless $\lambda - \mu \models m$, meaning we need $0 \leq \mu_i \leq \lambda_i - m$ for all i .

By Mackey:

$$\begin{aligned} \xi^\lambda \downarrow_{S_m \times S_k} &= \mathbb{1}_{S_\lambda} \uparrow^{S_n} \downarrow_{S_m \times S_k} \\ &= \sum_{\sigma \in S_m \times S_k \backslash S_n / S_\lambda} \mathbb{1} \uparrow_{\sigma S_\lambda \sigma^{-1} \cap (S_m \times S_k)}^{S_m \times S_k}. \end{aligned}$$

By Proposition 3.6 there is a bijection between $(S_m \times S_k) - S_\lambda$ double cosets in S_n and $2 \times \ell(\lambda)$ matrices over \mathbb{N}_0 with row sums (m, k) and column sums λ . Specifically, if $A_1 = [m], A_2 = [m + 1, \dots, m + k]$, $B_1 = [\lambda_1], B_2 = \{\lambda_1 + 1, \dots, \lambda_1 + \lambda_2\}$, etc., then the double coset $(S_m \times S_k)\sigma S_\lambda$ corresponds to $Z(\sigma)$ where $Z(\sigma)_{1j} = |A_i \cap \sigma(B_j)|$ and $Z(\sigma)_{2j} = |A_2 \cap \sigma(B_j)|$. Since $Z(\sigma)_{1j} + Z(\sigma)_{2j} = \lambda_j$ for all j , this matrix is in fact determined by just its second row, say, which we will call $\mu := (|A_2 \cap \sigma(B_1)|, \dots, |A_2 \cap \sigma(B_{\ell(\lambda)}|) \models k$. In particular, the first row is then $\lambda - \mu$ and note $0 \leq \mu_i \leq \lambda_i$ for all i .

Observe

$$\sigma S_\lambda \sigma^{-1} = \sigma(S_{B_1} \times \dots \times S_{B_{\ell(\lambda)}})\sigma^{-1} = S_{\sigma(B_1)} \times \dots \times S_{\sigma(B_{\ell(\lambda)})}$$

and hence $\sigma S_\lambda \sigma^{-1} \cap (S_m \times S_k)$ is conjugate to $S_{\lambda - \mu} \times S_\mu$. Then

$$\begin{aligned} \mathbb{1} \uparrow_{\sigma S_\lambda \sigma^{-1} \cap (S_m \times S_k)}^{S_m \times S_k} &= \mathbb{1}_{S_{\lambda - \mu}} \uparrow^{S_m \times S_k} \\ &= \mathbb{1}_{S_{\lambda - \mu}} \uparrow^{S_m} \# \mathbb{1}_{S_\mu} \uparrow^{S_k} \\ &= \xi^{\lambda - \mu} \# \xi^\mu. \end{aligned}$$

This finishes the proof of (i).

- (ii) We have by (i),

$$\begin{aligned} \psi^\lambda \downarrow_{S_m \times S_k} &= \sum_{\pi \in S_{\mathbb{N}}} \text{sgn}(\pi) \xi^{\lambda - \text{id} + \pi} \downarrow_{S_m \times S_k} \\ &= \sum_{\pi \in S_{\mathbb{N}}} \text{sgn}(\pi) \sum_{\mu \models k} \xi^{\lambda - \text{id} + \pi - \mu} \# \xi^\mu \end{aligned}$$

$$\begin{aligned}
&= \sum_{\mu \vdash k} \left(\sum_{\pi \in S_{\mathbb{N}}} \operatorname{sgn}(\pi) \xi^{(\lambda-\mu)-\operatorname{id}+\pi} \right) \# \xi^{\mu} \\
&= \sum_{\mu \vdash k} \psi^{\lambda-\mu} \# \xi^{\mu}.
\end{aligned}$$

□

Definition. Let $0 \leq k \leq n$, $\lambda \equiv n$, $\mu \vdash k$. Define

$$\psi^{\lambda \setminus \mu} := \sum_{\pi \in S_{\mathbb{N}}} \operatorname{sgn}(\pi) \xi^{\lambda-\operatorname{id}-\pi \cdot (\mu-\operatorname{id})},$$

it is a virtual character of S_{n-k} .

Note. If $k = 0$, then $\mu = (0, 0, \dots)$; and $\psi^{\lambda \setminus \mu} = \psi^{\lambda}$.

We will informally call the $\psi^{\lambda \setminus \mu}$ skew characters, one can also define skew diagrams, etc.

We have the following analogue of Lemma 3.4

Lemma 3.8. Let $0 \leq k \leq n$, $\lambda \equiv n$, $\mu \vdash k$.

- (i) Only finitely many terms in the sum defining $\psi^{\lambda \setminus \mu}$ are non-zero.
- (ii) The virtual character afforded by the determinant $\det([\lambda_i - i - (\mu_j - j)])_{ij}$ is $\psi^{\lambda \setminus \mu}$.

Proof. Very similar as the proof of Lemma 3.4, see Example Sheet 2, Question 5. □

Lemma 3.9. Let $\lambda \equiv m + k$, $m, k \in \mathbb{N}_0$. Then

$$\psi^{\lambda} \downarrow_{S_m \times S_k} = \sum_{\beta \vdash k} \psi^{\lambda \setminus \beta} \# \psi^{\beta}.$$

Proof. All sums involved will be finite. First, from Lemma 3.7 we have

$$\begin{aligned}
\psi^{\lambda} \downarrow_{S_m \times S_k} &= \sum_{\pi \in S_{\mathbb{N}}} \operatorname{sgn}(\pi) \sum_{\mu \vdash k} \xi^{\lambda-\operatorname{id}+\pi-\mu} \# \xi^{\mu} \\
&= \sum_{\pi \in S_{\mathbb{N}}} \operatorname{sgn}(\pi) \sum_{\nu \vdash k} \xi^{\lambda-\operatorname{id}+\pi-\pi \circ \nu} \# \xi^{\pi \circ \nu} && \nu = \pi^{-1} \circ \mu \\
&= \sum_{\pi \in S_{\mathbb{N}}} \operatorname{sgn}(\pi) \sum_{\nu \vdash k} \xi^{\lambda-\operatorname{id}-\pi \circ (\nu-\operatorname{id})} \# \xi^{\nu} && (*)
\end{aligned}$$

On the other hand,

$$\sum_{\beta \vdash k} \psi^{\lambda \setminus \beta} \# \psi^{\beta} = \sum_{\beta \vdash k} \left(\sum_{\pi \in S_{\mathbb{N}}} \operatorname{sgn}(\pi) \xi^{\lambda-\operatorname{id}-\pi \circ (\beta-\operatorname{id})} \right) \# \left(\sum_{\tau \in S_k} \operatorname{sgn}(\tau) \xi^{\beta-\operatorname{id}+\tau} \right)$$

Note that if $\beta \vdash k$, then $\ell(\beta) \leq k$, so in the last sum we can sum over $\tau \in S_k$ instead of $S_{\mathbb{N}}$. Then

$$\begin{aligned}
\sum_{\beta \vdash k} \psi^{\lambda \setminus \beta} \# \psi^\beta &= \sum_{\beta \vdash k} \sum_{\pi \in S_{\mathbb{N}}} \sum_{\tau \in S_k} \operatorname{sgn}(\pi\tau) \xi^{\lambda - \operatorname{id} - \pi \circ (\beta - \operatorname{id})} \# \xi^{\tau^{-1} \circ (\beta - \operatorname{id}) + \tau} \\
&= \sum_{\beta \vdash k} \sum_{\tau \in S_k} \sum_{\rho \in S_{\mathbb{N}}} \operatorname{sgn}(\rho) \xi^{\lambda - \operatorname{id} - \rho \tau^{-1} \circ (\beta - \operatorname{id})} \# \xi^{\overbrace{\tau^{-1} \circ (\beta - \operatorname{id}) + \operatorname{id}}^{=: \mu}} \quad \rho = \pi\tau \\
&= \sum_{\rho \in S_{\mathbb{N}}} \sum_{\substack{\mu \models k \text{ such that} \\ \mu = \tau^{-1} \circ (\beta - \operatorname{id}) + \operatorname{id} \\ \text{for some } \tau \in S_k, \beta \vdash k}} \operatorname{sgn}(\rho) \xi^{\lambda - \operatorname{id} - \rho \circ (\mu - \operatorname{id})} \# \xi^\mu \\
&= \sum_{\rho \in S_{\mathbb{N}}} \sum_{\substack{\mu \models k \text{ such that} \\ \mu = \tau^{-1} \circ (\beta - \operatorname{id}) + \operatorname{id} \\ \text{for some } \tau \in S_k, \beta \vdash k}} \operatorname{sgn}(\rho) \xi^{\lambda - \operatorname{id} - \rho \circ (\mu - \operatorname{id})} \# \xi^\mu \quad (**).
\end{aligned}$$

Note that we may replace $\sum_{\beta} \sum_{\tau}$ by $\sum_{\mu \text{ s.t. } \dots}$ because: if $\tau^{-1} \circ (\beta - \operatorname{id}) + \operatorname{id} = \tilde{\tau}^{-1} \circ (\tilde{\beta} - \operatorname{id}) + \operatorname{id}$ for some $\tau, \tilde{\tau} \in S_k$, $\beta, \tilde{\beta} \vdash k$, then $\beta - \operatorname{id} = (\tau \circ \tilde{\tau}^{-1}) \circ (\tilde{\beta} - \operatorname{id})$. Since $\beta \vdash k$, $\beta_i \geq \beta_{i+1}$ for all i . But then $\beta_i - i > \beta_{i+1} - (i+1)$ for all i , i.e. $\beta - \operatorname{id}$ is strictly decreasing. Similarly for $\tilde{\beta} - \operatorname{id}$. Therefore, $\tau \circ \tilde{\tau}^{-1} = 1$, i.e. $\tau = \tilde{\tau}$ and then also $\beta = \tilde{\beta}$.

We want to show $(*) = (**)$. For this we have to show that the restriction in $(**)$ can be removed.

First, we claim that

$$\begin{aligned}
&\{\mu \models k \mid \mu = \tau^{-1} \circ (\beta - \operatorname{id}) + \operatorname{id} \text{ for some } \tau \in S_k, \beta \vdash k\} \\
&= \{\mu \models k \mid \mu_i = 0 \text{ for all } i > k, \mu_i - i \text{ are distinct for all } i\}
\end{aligned}$$

To see \subseteq : Take τ, β , define $\mu = \tau^{-1} \circ (\beta - \operatorname{id}) + \operatorname{id}$. Then

- $|\mu| = |\beta| = k$,
- since $\beta \vdash k$, then $\beta - \operatorname{id}$ is strictly increasing, and so the $\mu_i - i$ are distinct for all i .
- since $\tau \in S_k$ and $\beta_i = 0$ for all $i > k$, then $\mu_i = 0$ for all $i > k$.

To see \supseteq : given $\mu \models k$ such that $\mu_i = 0$ for all $i > k$, $\mu_i - i$ are distinct for all i , we will construct $\tau \in S_k, \beta \vdash k$ as follows:

Since $\mu_i = 0$ for all $i > k$, $\mu_i - i = -i$ for all $i > k$. Since the $\mu_i - i$ are distinct, $\mu_i - i \geq -k$ for all $i \leq k$. Moreover, we can order the $\mu_i - i$ and then uniquely define $\tau \in S_k$ by

$$\mu_{\tau^{-1}(1)} - \tau^{-1}(1) > \mu_{\tau^{-1}(2)} - \tau^{-1}(2) > \dots > \mu_{\tau^{-1}(k)} - \tau^{-1}(k) > -(k+1) > -(k+2) > \dots$$

Then define $\beta := \tau \circ (\mu - \operatorname{id}) + \operatorname{id}$. Then we get $\mu = \tau^{-1} \circ (\beta - \operatorname{id}) + \operatorname{id}$, so we only have to check that $\beta \vdash k$. We have $|\beta| = |\mu| = k$. By construction, $\beta - \operatorname{id}$ is strictly decreasing,

therefore $\beta_i \geq \beta_{i+1}$ for all i . Since $\tau \in S_k$, $\mu_i = 0$ for all $i > k$, then $\beta_i = 0$ for all $i > k$. Hence $\beta \vdash k$.

Second, we claim that

$$\begin{aligned} & \{\mu \models k \mid \mu_i = 0 \text{ for all } i > k, \text{ the } \mu_i - i \text{ are distinct for all } i\} \\ &= \{\mu \models k \mid \mu_i - i \text{ are distinct for all } i\} \end{aligned}$$

See Example Sheet 2.

Hence (**) becomes

$$\sum_{\beta \vdash k} \psi^{\lambda \setminus \beta} \# \psi^\beta = \sum_{\rho \in S_{\mathbb{N}}} \sum_{\substack{\mu \models k \text{ such that} \\ \mu_i - i \text{ distinct } \forall i}} \text{sgn}(\rho) \xi^{\lambda - \text{id} - \rho \circ (\mu - \text{id})} \# \xi^\mu$$

Finally, if $\mu \models k$ is such that $\mu_i - i = \mu_j - j$ for some $i \neq j$, then

$$\begin{aligned} & \sum_{\rho \in S_{\mathbb{N}}} \text{sgn}(\rho) \xi^{\lambda - \text{id} - \rho(\mu - \text{id})} \# \xi^\mu \\ &= \frac{1}{2} \sum_{\sigma \in S_{\mathbb{N}}} \left[\text{sgn}(\sigma) \xi^{\lambda - \text{id} - \sigma \circ (\mu - \text{id})} \# \xi^\mu + \text{sgn}(\sigma \circ (ij)) \xi^{\lambda - \text{id} - \sigma \circ \overbrace{(ij) \circ (\mu - \text{id})}^{\mu - \text{id}}} \# \xi^\mu \right] \\ &= \frac{1}{2} \sum_{\sigma \in S_{\mathbb{N}}} \left[\text{sgn}(\sigma) \xi^{\lambda - \text{id} - \sigma \circ (\mu - \text{id})} \# \xi^\mu - \text{sgn}(\sigma) \xi^{\lambda - \text{id} - \sigma \circ (\mu - \text{id})} \# \xi^\mu \right] \\ &= 0 \end{aligned}$$

Then

$$\sum_{\beta \vdash k} \psi^{\lambda \setminus \beta} \# \psi^\beta = \sum_{\rho \in S_{\mathbb{N}}} \sum_{\mu \models k} \text{sgn}(\rho) \xi^{\lambda - \text{id} - \rho \circ (\mu - \text{id})} \# \xi^\mu = (*) = \psi^\lambda \downarrow_{S_m \times S_k}.$$

□

Theorem 3.10. *Let $0 \leq k \leq n$, $\alpha \vdash n$, $\beta \vdash k$.*

- (i) *If $\psi^{\alpha \setminus \beta} \neq 0$, then $\alpha_i \geq \beta_i$ for all i ,*
- (ii) $\langle \psi^{\alpha \setminus \beta}, \xi^{(n-k)} \rangle = \begin{cases} 1 & \text{if } \alpha_1 \geq \beta_1 \geq \alpha_2 \geq \beta_2 \geq \dots, \\ 0 & \text{otherwise.} \end{cases}$

If $\alpha_1 \geq \beta_1 \geq \alpha_2 \geq \beta_2 \geq \dots$, we say that α and β *intertwine*.

Proof.

- (i) Recall from Lemma 3.8 that $\psi^{\alpha \setminus \beta}$ is the character of the determinant of the matrix A where $A_{ij} = [\alpha_i - i - (\beta_j - j)]$. Note that since α, β are partitions, $\alpha - \text{id}, \beta - \text{id}$ are strictly decreasing. If A_{ij} is zero (in other words, $\alpha_i - i - (\beta_j - j) < 0$), then all entries to its left and below are zero. Thus the determinant vanishes if a diagonal entry is zero. So if $\psi^{\alpha \setminus \beta} \neq 0$, we must have $\alpha_i - i - (\beta_i - i) \geq 0$, i.e. $\alpha_i \geq \beta_i$ for all i .

(ii) For $\lambda \equiv n - k$, recall $\xi^\lambda = 0$ if $\lambda \not\equiv n - k$. If $\lambda \equiv n - k$, then

$$\langle \xi^\lambda, \xi^{(n-k)} \rangle = \langle \mathbb{1}_{S_\lambda} \uparrow^{S_{n-k}}, \mathbb{1}_{S_{n-k}} \rangle \stackrel{\text{Frobenius reciprocity}}{=} \langle \mathbb{1}_{S_\lambda}, \mathbb{1}_{S_\lambda} \rangle = 1.$$

Thus

$$\langle \psi^{\alpha \setminus \beta}, \xi^{(n-k)} \rangle = \sum_{\pi \in S_{\mathbb{N}}} \text{sgn } \pi \langle \xi^{\alpha - \text{id} - \pi \circ (\beta - \text{id})}, \xi^{(n-k)} \rangle = \sum_{\pi \in S_{\mathbb{N}}} (\text{sgn } \pi) \delta_{\{\alpha - \text{id} - \pi \circ (\beta - \text{id}) \equiv n - k\}}$$

This is the determinant of M where $M_{ij} = \delta_{\{\alpha_i - i - (\beta_j - j) \geq 0\}}$. Note if $M_{ij} = 0$, then all entries to its left and below are also zero. Also, M only has 0–1 entries. If $\alpha_1 \geq \beta_1 \geq \alpha_2 \geq \beta_2 \geq \dots$, then $M_{ii} = 1$ and $M_{i+1, i} = 0$ for all i , and so $\det M = 1$. Otherwise, $M_{ii} = 0$ for some i , or $M_{i+1, i} = 1$ for some i , but then M must have a column of all 0's, or have two equal columns; and therefore $\det M = 0$. □

Theorem 3.11 (Young's Rule). *Let $\lambda \equiv n$ with $\ell(\lambda) \leq n$. Let $\alpha \vdash n$. Then $\langle \psi^\alpha, \xi^\lambda \rangle$ is equal to the number of tuples of partitions $(\beta^{(1)}, \beta^{(2)}, \dots, \beta^{(n-1)})$ satisfying*

- (i) $\beta^{(i)} \vdash \sum_{j=1}^i \lambda_j$ for all $i \in [n-1]$,
- (ii) $0 \leq \beta_j^{(1)} \leq \beta_j^{(2)} \leq \dots \leq \beta_j^{(n-1)} \leq \alpha_j$ for all $j \in [n]$,
- (iii) $\beta_j^{(i)} \leq \beta_{j-1}^{(i-1)}$ for all $j > 1, i \geq 1$, where we treat $\beta^{(0)} = (0, 0, \dots)$ and $\beta^{(n)} = \alpha$.

Once we have proved $\psi^\alpha = \chi^\alpha$, then Young's Rule will tell us the multiplicity of the Spect module $[\alpha]$ in a direct sum decomposition of M^λ into irreducibles.

Example. Let $n = 5, \alpha = (3, 2)$.

(i) Let $\lambda = (2, 0, 1, 2) \equiv 5$. Then

$$\begin{array}{l} \beta^{(0)} = (0, 0, 0, \dots) \\ \quad \quad \quad \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(1)} = (\beta_1^{(1)}, \beta_2^{(1)}, \beta_3^{(1)}, \dots) \vdash 2 \\ \quad \quad \quad \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(2)} = (\beta_1^{(2)}, \beta_2^{(2)}, \beta_3^{(2)}, \dots) \vdash 2 + 0 = 2 \\ \quad \quad \quad \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(3)} = (\beta_1^{(3)}, \beta_2^{(3)}, \beta_3^{(3)}, \dots) \vdash 2 + 0 + 1 = 3 \\ \quad \quad \quad \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(4)} = (\beta_1^{(4)}, \beta_2^{(4)}, \beta_3^{(4)}, \dots) \vdash 2 + 0 + 1 + 2 = 5 \\ \quad \quad \quad \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \alpha = \beta^{(5)} = (3, 2, 0, \dots) \end{array}$$

where an arrow $a \rightarrow b$ indicates that $a \leq b$. The yellow arrows are by (ii) and the violet arrows by condition (iii). We see that most entries are uniquely determined as follows:

$$\begin{array}{rcl}
\beta^{(0)} & = & (0, \quad 0, \quad 0, \quad \dots) \\
& & \begin{array}{c} \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(1)} = (2, \quad 0, \quad 0, \quad \dots) \vdash 2 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(2)} = (2, \quad 0, \quad 0, \quad \dots) \vdash 2 + 0 = 2 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(3)} = (\beta_1^{(3)}, \beta_2^{(3)}, 0, \quad \dots) \vdash 2 + 0 + 1 = 3 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(4)} = (3, \quad 2, \quad 0, \quad \dots) \vdash 2 + 0 + 1 + 2 = 5 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \alpha = \beta^{(5)} = (3, \quad 2, \quad 0, \quad \dots) \end{array}
\end{array}$$

We can have $\beta^{(3)} = (3, 0, \dots)$ or $(2, 1, 0, \dots)$. Therefore $\langle \psi^\alpha, \xi^\lambda \rangle = 2$.

(ii) Let $\lambda = (0, 2, 2, 0, 1) \models 5$. [Since $\xi^{(2,0,1,2)} = \xi^{(0,2,2,0,1)}$, we expect again two tuples]

$$\begin{array}{rcl}
\beta^{(0)} & = & (0, \quad 0, \quad 0, \quad \dots) \\
& & \begin{array}{c} \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(1)} = (\beta_1^{(1)}, \beta_2^{(1)}, \beta_3^{(1)}, \quad \dots) \vdash 0 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(2)} = (\beta_1^{(2)}, \beta_2^{(2)}, \beta_3^{(2)}, \quad \dots) \vdash 0 + 2 = 2 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(3)} = (\beta_1^{(3)}, \beta_2^{(3)}, \beta_3^{(3)}, \quad \dots) \vdash 0 + 2 + 2 = 4 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \beta^{(4)} = (\beta_1^{(4)}, \beta_2^{(4)}, \beta_3^{(4)}, \quad \dots) \vdash 0 + 2 + 2 + 0 = 4 \\ \downarrow \quad \swarrow \quad \downarrow \quad \swarrow \quad \downarrow \\ \alpha = \beta^{(5)} = (3, \quad 2, \quad 0, \quad \dots) \end{array}
\end{array}$$

Again we see that most entries are uniquely determined:

$$\begin{aligned}
\beta^{(0)} &= (0, 0, 0, \dots) \\
&\quad \begin{array}{cccc} & & & \\ & & & \\ & & & \\ & & & \\ & & & \end{array} \\
\beta^{(1)} &= (0, 0, 0, \dots) \vdash 0 \\
&\quad \begin{array}{cccc} & & & \\ & & & \\ & & & \\ & & & \\ & & & \end{array} \\
\beta^{(2)} &= (2, 0, 0, \dots) \vdash 0 + 2 = 2 \\
&\quad \begin{array}{cccc} & & & \\ & & & \\ & & & \\ & & & \\ & & & \end{array} \\
\beta^{(3)} &= (2/3, 2/1, 0, \dots) \vdash 0 + 2 + 2 = 4 \\
&\quad \begin{array}{cccc} & & & \\ & & & \\ & & & \\ & & & \\ & & & \end{array} \\
\beta^{(4)} &= (2/3, 2/1, 0, \dots) \vdash 0 + 2 + 2 + 0 = 4 \\
&\quad \begin{array}{cccc} & & & \\ & & & \\ & & & \\ & & & \\ & & & \end{array} \\
\alpha = \beta^{(5)} &= (3, 2, 0, \dots) \\
&\quad \begin{array}{cccc} & & & \\ & & & \\ & & & \\ & & & \\ & & & \end{array}
\end{aligned}$$

So we can have either $\beta^{(3)} = \beta^{(4)} = (2, 2, 0, \dots)$ or $\beta^{(3)} = \beta^{(4)} = (3, 1, 0, \dots)$. So we again get $\langle \psi^\alpha, \xi^\lambda \rangle = 2$.

Proof of Theorem 3.11. We have

$$\begin{aligned}
\langle \psi^\alpha, \xi^\lambda \rangle &= \langle \psi^\alpha, \mathbb{1}_{S_\lambda} \uparrow^{S_n} \rangle \stackrel{\text{F.R.}}{=} \langle \psi^\alpha \downarrow_{S_\lambda}^{S_n}, \mathbb{1}_{S_\lambda} \rangle \\
&= \langle \left(\psi^\alpha \downarrow_{S_{\lambda_n} \times S_{\lambda_{n-1} + \dots + \lambda_1}}^{S_n} \right) \downarrow_{S_{\lambda_n} \times S_{\lambda_{n-1}} \times \dots \times S_{\lambda_1}}, \xi^{(\lambda_n)} \# \dots \# \xi^{(\lambda_1)} \rangle \\
&\stackrel{\text{Lemma 3.9}}{=} \left\langle \sum_{\beta^{(n-1)} \vdash \sum_{j=1}^{n-1} \lambda_j} \psi^{\alpha \setminus \beta^{(n-1)}} \# (\psi^{\beta^{(n-1)}}) \downarrow_{S_{\lambda_{n-1}} \times \dots \times S_{\lambda_1}}^{S_{\lambda_{n-1} + \dots + \lambda_1}}, \xi^{(\lambda_n)} \# \dots \# \xi^{(\lambda_1)} \right\rangle \\
&= \sum_{\beta^{(n-1)} \vdash \sum_{j=1}^{n-1} \lambda_j} \langle \psi^{\alpha \setminus \beta^{(n-1)}}, \xi^{(\lambda_n)} \rangle \cdot \langle \psi^{\beta^{(n-1)}} \downarrow_{S_{\lambda_{n-1}} \times \dots \times S_{\lambda_1}}^{S_{\lambda_{n-1} + \dots + \lambda_1}}, \xi^{(\lambda_{n-1})} \# \dots \# \xi^{(\lambda_1)} \rangle \\
&\stackrel{\text{Theorem 3.10}}{=} \sum_{\beta^{(n-1)}} \langle \psi^{\beta^{(n-1)}} \downarrow_{S_{\lambda_{n-1}} \times \dots \times S_{\lambda_1}}^{S_{\lambda_{n-1} + \dots + \lambda_1}}, \xi^{(\lambda_{n-1})} \# \dots \# \xi^{(\lambda_1)} \rangle
\end{aligned}$$

where we sum over $\beta^{(n-1)} \vdash \sum_{j=1}^{n-1} \lambda_j$ such that α and $\beta^{(n-1)}$ intertwine. Iteratively applying Lemma 3.9 and Theorem 3.10, we get

$$\begin{aligned}
\langle \psi^\alpha, \xi^\lambda \rangle &= \sum_{\beta^{(n-1)}} \sum_{\beta^{(n-2)}} \dots \sum_{\beta^{(1)}} \langle \psi^{\beta^{(1)}} \downarrow_{S_{\lambda_1}}^{S_{\lambda_1}}, \xi^{(\lambda_1)} \rangle \\
&= \sum_{\beta^{(n-1)}, \beta^{(n-2)}, \dots, \beta^{(1)}} 1
\end{aligned}$$

where we sum over $\beta^{(i)} \vdash \sum_{j=1}^i \lambda_j$ (this is condition (i)) such that $\beta^{(i)}$ and $\beta^{(i-1)}$ intertwine for all $i \in [n]$ (this gives conditions (ii) and (iii)). \square

Lemma 3.12. *Let $\alpha, \beta \vdash n$. If $\langle \xi^\alpha, \chi^\beta \rangle > 0$, then $\beta \triangleright \alpha$.*

Proof. Since $\langle \xi^\alpha, \chi^\beta \rangle > 0$, we have $\text{Hom}_{\mathbb{C}S_n}([\beta], M^\alpha) \neq 0$. Since $\text{char } \mathbb{C} = 0$, Maschke's theorem gives a complement of $[\beta]$ in M^β , so we can extend any $\phi \in \text{Hom}_{\mathbb{C}S_n}([\beta], M^\alpha)$ to $\tilde{\phi} \in \text{Hom}_{\mathbb{C}S_n}(M^\beta, M^\alpha)$. Then $\beta \succeq \alpha$ by Theorem 2.10. \square

Remarks.

- We can't use Theorem 3.11 to prove the lemma, since we don't have $\chi^\beta = \psi^\beta$ yet.
- The converse holds, see Lemma 3.20.

Theorem 3.13. *Let $\alpha \vdash n$. Then $\psi^\alpha = \chi^\alpha$. In particular, the irreducible representation $[\alpha]$ has determinantal form $\det([\alpha_i - i + j])_{ij}$.*

Proof.

- **Step 1.** We first show that if $\lambda \models n$ with $\ell(\lambda) \leq n$ and $\langle \xi^\lambda, \psi^\alpha \rangle > 0$, then $\alpha \succeq \lambda$.

Proof. Suppose $\langle \xi^\lambda, \psi^\alpha \rangle > 0$. Then there exists $(\beta^{(1)}, \dots, \beta^{(n-1)})$ satisfying Theorem 3.11 (i), (ii), (iii). By (iii),

$$0 = \beta_1^{(0)} \geq \beta_2^{(1)} \geq \beta_3^{(2)} \geq \dots \geq 0,$$

so $\ell(\beta^{(i)}) \leq i$ for all i . Now $\beta^{(i)} = (\beta_1^{(i)}, \beta_2^{(i)}, \dots, \beta_i^{(i)}, 0, \dots) \vdash \sum_{j=1}^i \lambda_j$ by (i), and $\alpha_j \geq \beta_j^{(i)}$ for all j by (ii). So

$$\alpha_1 + \alpha_2 + \dots + \alpha_i \geq \beta_1^{(i)} + \beta_2^{(i)} + \dots + \beta_i^{(i)} = \lambda_1 + \lambda_2 + \dots + \lambda_i,$$

for all i , in other words, $\alpha \succeq \lambda$. \square

- **Step 2.** We show $\langle \psi^\alpha, \xi^\alpha \rangle = 1$.

Proof. Observe that $(\beta^{(1)}, \dots, \beta^{(n-1)})$ with $\beta^{(i)} = (\alpha_1, \alpha_2, \dots, \alpha_i)$ satisfies the conditions in Theorem 3.11, and so $\langle \psi^\alpha, \xi^\alpha \rangle \geq 1$. Conversely, suppose $(\beta^{(1)}, \dots, \beta^{(n-1)})$ satisfies (i), (ii), (iii) in Theorem 3.11 with $\lambda = \alpha$. Then, as in Step 1, we obtain $\ell(\beta^{(i)}) \leq i$, $\alpha_j \geq \beta_j^{(i)}$ for all j , and $\alpha_1 + \dots + \alpha_i \geq \beta_1^{(i)} + \dots + \beta_i^{(i)} = \alpha_1 + \dots + \alpha_i$ for all i . Hence, we must have equality in $\alpha_j \geq \beta_j^{(i)}$ for $j = 1, \dots, i$, so $\beta^{(i)} = (\alpha_1, \dots, \alpha_i)$. So there is only one such tuple $(\beta^{(1)}, \dots, \beta^{(n-1)})$ and therefore $\langle \psi^\alpha, \xi^\alpha \rangle = 1$. \square

- **Step 3.** We show $\langle \psi^\alpha, \psi^\alpha \rangle = 1$.

Proof. First, for any $\pi \in S_{\mathbb{N}}$, $\alpha - \text{id} + \pi \succeq \alpha$ since for all i ,

$$\pi^{-1}(1) + \pi^{-1}(2) + \dots + \pi^{-1}(i) \geq 1 + 2 + \dots + i,$$

so

$$(\alpha_1 - 1 + \pi^{-1}(1)) + (\alpha_2 - 2 + \pi^{-1}(2)) + \dots + (\alpha_i - i + \pi^{-1}(i)) \geq \alpha_1 + \alpha_2 + \dots + \alpha_i.$$

On the other hand, if $\pi \in S_n$ and $\langle \xi^{\alpha - \text{id} + \pi}, \psi^\alpha \rangle > 0$, then $\alpha \triangleright \alpha - \text{id} + \pi$ by Step 1 (because $\alpha - \text{id} + \pi \models n$, else $\xi^{\alpha - \text{id} + \pi} = 0$, and $\alpha \vdash n$, so $\ell(\alpha) \leq n$, so $\ell(\alpha - \text{id} + \pi) \leq n$). Hence $\alpha \triangleright \alpha - \text{id} + \pi \triangleright \alpha$, so $\pi = \text{id}$. Thus,

$$\begin{aligned} \langle \psi^\alpha, \psi^\alpha \rangle &= \sum_{\pi \in S_n} \text{sgn } \pi \langle \xi^{\alpha - \text{id} + \pi}, \psi^\alpha \rangle \\ &= \langle \xi^\alpha, \psi^\alpha \rangle \\ &= 1. \end{aligned}$$

□

We can now prove $\psi^\alpha = \chi^\alpha$. Since $\langle \psi^\alpha, \psi^\alpha \rangle = 1$, we have $\psi^\alpha = \pm \phi$ for some $\phi \in \text{Irr}(S_n)$. Since also $\chi^\alpha \in \text{Irr}(S_n)$, it thus suffices to prove $\langle \psi^\alpha, \chi^\alpha \rangle > 0$. Next, if $\lambda \models n$ such that $\langle \xi^\lambda, \chi^\alpha \rangle > 0$, then $\langle \xi^\beta, \chi^\alpha \rangle > 0$ where $\beta \vdash n$ is obtained from λ permuting its parts. By Lemma 3.12, $\alpha \triangleright \beta$, but also clearly $\beta \triangleright \lambda$. Therefore $\alpha \triangleright \lambda$. So if $\pi \in S_n$, and $\langle \xi^{\alpha - \text{id} + \pi}, \chi^\alpha \rangle > 0$, then $\alpha \triangleright \alpha - \text{id} + \pi \triangleright \alpha$, i.e. $\pi = \text{id}$. Thus

$$\langle \psi^\alpha, \chi^\alpha \rangle = \sum_{\pi \in S_n} \text{sgn } \pi \langle \xi^{\alpha - \text{id} + \pi}, \chi^\alpha \rangle = \langle \xi^\alpha, \chi^\alpha \rangle.$$

This is > 0 , since $[\alpha] \leq M^\alpha$.

Therefore $\langle \psi^\alpha, \psi^\alpha \rangle = 1$ and $\langle \psi^\alpha, \chi^\alpha \rangle > 0$. These imply $\psi^\alpha = \chi^\alpha$. □

3.3 Applications

3.3.1 Young's Rule Revisited

Corollary 3.14. *Let $\alpha \vdash n$. Then $\langle \chi^\alpha, \xi^\alpha \rangle = 1$.*

Proof.

- Either from James Submodule Theorem and complete reducibility in char 0,
- or use Theorem 3.13 and Step 2 in its proof.

□

Corollary 3.15. *The permutation characters $\{\xi^\alpha \mid \alpha \vdash n\}$ gives a basis of the \mathbb{C} -vector space of class functions of S_n . In particular, the change of basis matrix to $\text{Irr}(S_n) = \{\chi^\beta \mid \beta \vdash n\}$ is \mathbb{Z} -valued, and unitriangular if we order the partitions in a way that extends the dominance partial ordering.*

Proof. From the definition of ψ^β and the fact that $\psi^\beta = \chi^\beta$, it is clear that the χ^β are \mathbb{Z} -linear combinations of the permutation characters. Conversely, it is clear that the permutation characters are \mathbb{Z} -linear combinations of the χ^α . From Lemma 3.12 it follows that the matrix is triangular and Corollary 3.14 gives that the diagonal entries are 1. □

Remark. Young's Rule tells us the multiplicity of $[\alpha]_{\mathbb{C}}$ in a direct sum decomposition of $M_{\mathbb{C}}^{\lambda}$ into irreducibles. Over an arbitrary field \mathbb{F} , $M_{\mathbb{F}}^{\lambda}$ decomposes as a direct sum of indecomposables: We saw from James' Submodule Theorem that there is a unique summand containing $[\lambda]_{\mathbb{F}}$, which we called the Young module $Y_{\mathbb{F}}^{\lambda}$.

In general, Young modules for S_n are defined as the indecomposable summands of $M_{\mathbb{F}}^{\lambda}$ for some $\lambda \vdash n$. It turns out that isomorphism classes are indexed by $\wp(n)$.

Fact. $M_{\mathbb{F}}^{\lambda}$ can be decomposed as a direct sum of S_n -modules each of which is isomorphic to $Y_{\mathbb{F}}^{\mu}$ for some $\mu \supseteq \lambda$, and $Y_{\mathbb{F}}^{\lambda}$ appears exactly once.

If $\text{char } \mathbb{F} = 0$, then indecomposable = irreducible, and we have proven this fact (then $Y_{\mathbb{C}}^{\lambda} = [\lambda]_{\mathbb{C}}$).

In general, $Y_{\mathbb{F}}^{\lambda} \not\cong [\lambda]_{\mathbb{F}}$, e.g. in Example Sheet, Question 5, we saw that $[(n-1, 1)]_{\mathbb{F}}$ was a submodule, but not a direct summand of $M_{\mathbb{F}}^{(n-1, 1)}$ in the case $\text{char } \mathbb{F} \mid n$.

If $\text{char } \mathbb{F} > 2$, then it is known that Specht modules are always indecomposable. In $\text{char } \mathbb{F} = 2$, this is still an open problem.

Next, we work towards another combinatorial way to interpret Young's Rule.

Lemma 3.16. *Let $m, k \in \mathbb{N}$, let $\alpha \vdash m+k, \beta \vdash k, \gamma \vdash m$. Then*

$$\langle \psi^{\alpha \setminus \beta}, \chi^{\gamma} \rangle = \langle \chi^{\alpha} \downarrow_{S_m \times S_k}, \chi^{\gamma} \# \chi^{\beta} \rangle.$$

Moreover, $\langle \psi^{\alpha \setminus \beta}, \chi^{\gamma} \rangle = \langle \psi^{\alpha \setminus \gamma}, \chi^{\beta} \rangle$.

Letting γ vary, this shows that $\psi^{\alpha \setminus \beta}$ is a genuine character.

Proof. We have

$$\begin{aligned} \langle \chi^{\alpha} \downarrow_{S_m \times S_k}, \chi^{\gamma} \# \chi^{\beta} \rangle &= \langle \psi^{\alpha} \downarrow_{S_m \times S_k}, \chi^{\gamma} \# \chi^{\beta} \rangle \\ &= \sum_{\delta \vdash k} \langle \psi^{\alpha \setminus \delta} \# \psi^{\delta}, \chi^{\gamma} \# \chi^{\beta} \rangle \\ &= \sum_{\delta \vdash k} \langle \psi^{\alpha \setminus \delta}, \chi^{\gamma} \rangle \cdot \langle \psi^{\delta}, \chi^{\beta} \rangle \\ &= \sum_{\delta \vdash k} \langle \psi^{\alpha \setminus \delta}, \chi^{\gamma} \rangle \cdot \langle \chi^{\delta}, \chi^{\beta} \rangle \\ &= \langle \psi^{\alpha \setminus \beta}, \chi^{\gamma} \rangle. \end{aligned}$$

The last part follows from $\langle \chi^{\alpha} \downarrow_{S_m \times S_k}, \chi^{\gamma} \# \chi^{\beta} \rangle = \langle \chi^{\alpha} \downarrow_{S_k \times S_m}, \chi^{\beta} \# \chi^{\gamma} \rangle$. \square

Remark. Multiplicities of the form $\langle \chi^{\alpha} \downarrow_{S_m \times S_k}, \chi^{\gamma} \# \chi^{\beta} \rangle$ are called *Littlewood-Richardson coefficients*, also denoted $c_{\gamma, \beta}^{\alpha}$, and they occur in many different contexts, e.g. symmetric functions and algebraic combinatorics, representation theory of algebraic groups, etc.

Lemma 3.17. *Let $m, k \in \mathbb{N}$, $\alpha \vdash m$. Then*

$$\chi^\alpha \# \chi^{(k)} \uparrow_{S_m \times S_k}^{S_{m+k}} = \sum \chi^\gamma$$

where the sum runs over $\gamma \vdash m+k$ such that $\alpha_i \leq \gamma_i \leq \alpha_{i-1}$ for all i , treating $\alpha_0 = \infty$.

Proof. Let $\gamma \vdash m+k$. Then


$$\begin{aligned} \langle \chi^\gamma, \chi^\alpha \# \chi^{(k)} \uparrow_{S_m \times S_k}^{S_{m+k}} \rangle &= \langle \chi^\gamma \downarrow_{S_m \times S_k}, \chi^\alpha \# \chi^{(k)} \rangle \\ &= \langle \psi^{\gamma \setminus \alpha}, \chi^{(k)} \rangle \\ &= \langle \psi^{\gamma \setminus \alpha}, \xi^{(k)} \rangle \\ &\stackrel{\text{Theorem 3.10}}{=} \begin{cases} 1 & \text{if } \gamma_1 \geq \alpha_1 \geq \gamma_2 \geq \alpha_2 \geq \dots, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

□

Corollary 3.18. *Notation as in Lemma 3.17. Then the Young diagrams $Y(\gamma)$ can be obtained from $Y(\alpha)$ by adding k many boxes in all possible ways such that no two of the newly added boxes lie in the same column*

Proof. Since $\gamma_i \geq \alpha_i$ for all i , we can certainly view $Y(\gamma)$ as a superset of $Y(\alpha)$. The condition $\gamma_i \leq \alpha_{i-1}$ corresponds to the assertion that no two boxes in $Y(\gamma) \setminus Y(\alpha)$ lie in the same column. □

Example. Let $\alpha = (3, 2, 2) \vdash 7$, $k = 2$. Then $Y(\alpha) = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}$. We have the following possible $Y(\gamma)$:



Therefore

$$\begin{aligned} [\alpha][k] &= [5, 2^2] \oplus [4, 3, 2] \oplus [4, 2^2, 1] \oplus [3^2, 2, 1] \oplus [3, 2^3] \\ \xi^{(\alpha, k)} &= \chi^{(5, 2^2)} + \chi^{(4, 3, 2)} + \chi^{(4, 2^2, 1)} + \chi^{(3^2, 2, 1)} + \chi^{(3, 2^3)} \end{aligned}$$

We can use Corollary 3.18 repeatedly to decompose $M^\alpha \cong [\alpha_1][\alpha_2] \cdots [\alpha_{\ell(\alpha)}]$ into irreducibles.

¹Remark by L.T.: I believe on the LHS it should not be $\xi^{(\alpha, k)}$. Correct would be the character of $[\alpha][k]$ and this module does not coincide with $M^{(\alpha, k)} = [\alpha_1][\alpha_2][\alpha_3][k]$. E.g. we have $\dim M^{(\alpha, k)} = \frac{9!}{3!^2 2! 2!} = 7560$, but using the hook length formula we calculate $\dim[\alpha][k] = [S_9 : S_7 \times S_2] \dim[\alpha] \# [k] = \frac{9!}{7! 2!} \dim[\alpha] \dim[k] = \frac{9!}{7! 2!} \cdot \frac{7!}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 2} \cdot 1 = 756$.

Example. Let $\alpha = (3, 2, 1) \vdash 6$. First,

$$[3][2] = \begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 2 & \\ \hline & & & 2 & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline \end{array}$$

Here we label the original boxes with 1 and the new boxes with 2. Then,

$$\begin{aligned} [3][2][1] &= \begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 2 & 3 \\ \hline & & & 3 & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 2 \\ \hline 3 & & & & \\ \hline \end{array} \\ &\oplus \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 3 \\ \hline 2 & & & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 3 & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & & & 3 \\ \hline \end{array} \\ &\oplus \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 3 \\ \hline 2 & 2 & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & \\ \hline 2 & 2 & 3 & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & \\ \hline 2 & 2 & & 3 \\ \hline \end{array} \end{aligned}$$

So

$$\xi^{(3,2,1)} = \chi^{(6)} + 2\chi^{(5,1)} + 2\chi^{(4,2)} + \chi^{(4,1,1)} + \chi^{(3,3)} + \chi^{(3,2,1)}.$$

Definition.

- (i) A generalised Young tableau of shape $\alpha \vdash n$ and content (or weight, type) $\lambda \models n$ is a filling of $Y(\alpha)$ with positive integers such that i appears exactly λ_i many times for all i .
- (ii) A generalised Young tableau is semistandard if its entries weakly increase left to right along rows, but strictly increase down columns.

We will abbreviate semistandard tableaux to SSYT.

Example. $\begin{array}{|c|c|c|c|c|c|} \hline 2 & 1 & 1 & 4 & 2 & 2 \\ \hline 4 & 1 & & & & \\ \hline \end{array}$ has shape $(6, 2)$, content $(3, 3, 0, 2)$. The semistandard Young tableaux of shape this shape and content are

$$\begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 2 & 2 \\ \hline 4 & 4 & & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 2 & 4 \\ \hline 2 & 4 & & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 4 & 4 \\ \hline 2 & 2 & & & & \\ \hline \end{array}$$

Young tableaux from before are just generalised Young tableaux of content (1^n) .

Using SSYT we can generalise the above example, determining $\xi^{(3,2,1)}$, and reformulate Young's Rule.

Corollary 3.19. *Let $\alpha \vdash n, \lambda \models n$. Then $\langle \xi^\lambda, \chi^\alpha \rangle$ is the number of SSYT of shape α and content λ .*

Note that unlike in Theorem 3.11 we don't require $\ell(\lambda) \leq n$.

Proof. Apply Corollary 3.18 and note that $M^\lambda \cong [\lambda_1][\lambda_2] \dots [\lambda_{\ell(\lambda)}]$ has character ξ^λ . \square

Example 1. We revisit the example after Theorem 3.11 where we showed that $\langle \chi^\alpha, \xi^\lambda \rangle = \langle \psi^\alpha, \xi^\lambda \rangle = 2$ for $\alpha = (3, 2)$ and $\lambda = (2, 0, 1, 2)$ or $\lambda = (0, 2, 2, 0, 1)$. The SSYT of shape α and content λ are:

$$\lambda = (2, 0, 1, 2) \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 3 \\ \hline 4 & 4 & \\ \hline \end{array}, \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 4 \\ \hline 3 & 4 & \\ \hline \end{array}$$

$$\lambda = (0, 2, 2, 0, 1) \quad \begin{array}{|c|c|c|} \hline 2 & 2 & 3 \\ \hline 3 & 5 & \\ \hline \end{array}, \quad \begin{array}{|c|c|c|} \hline 2 & 2 & 5 \\ \hline 3 & 3 & \\ \hline \end{array}$$

Recall Lemma 3.12: If $\alpha, \beta \vdash n$ with $\langle \xi^\alpha, \chi^\beta \rangle > 0$, then $\alpha \preceq \beta$. The converse also holds.

Lemma 3.20. *Suppose $\alpha, \beta \vdash n$ with $\alpha \preceq \beta$. Then $\langle \xi^\alpha, \chi^\beta \rangle > 0$.*

Proof. Example Sheet 3. □

Remark. The number of SSYT of shape α and content λ is often denoted by $K_{\alpha, \lambda}$. Such quantities are known as *Kostka numbers*.

3.3.2 Branching Rule

We investigate restriction from S_n to $S_{n-1} \cong S_{n-1} \times S_1$. Note that this is a special case of $S_m \times S_k$.

Definition. Let $\lambda \vDash n$, $i \in \mathbb{N}$. Define $\lambda^{i-} \vDash n-1$ and $\lambda^{i+} \vDash n+1$ via $\lambda^{i-} = (\lambda_1, \dots, \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, \dots)$ and $\lambda^{i+} = (\lambda_1, \dots, \lambda_{i-1}, \lambda_i + 1, \lambda_{i+1}, \dots)$.

Lemma 3.21. *Let $\lambda \vDash n$. Then $\xi^\lambda \downarrow_{S_{n-1}} = \sum_{i=1}^{\infty} \xi^{\lambda^{i-}}$.*

Proof. First note that the RHS sum is finite since $\lambda^{i-} \vDash n-1$ for all $i > \ell(\lambda)$, whence $\xi^{\lambda^{i-}} = 0$. Now, by Lemma 3.7,

$$\xi^\lambda \downarrow_{S_{n-1}} = \xi^\lambda \downarrow_{S_{n-1} \times S_1} = \sum_{\mu \vdash 1} \xi^{\lambda - \mu} \# \xi^\mu.$$

But $\xi^\mu = \mathbb{1}_{S_1}$ and $\mu = (0, \dots, 0, 1, 0, \dots)$ where the 1 is in the i -th position, so $\lambda - \mu = \lambda^{i-}$. □

Recall we defined α^- , where $\alpha \vdash n$, and removable boxes in Section 3.1. Observe

$$\alpha^- = \{\beta \vdash n-1 \mid \beta = \alpha^{i-} \text{ for some } i \in \mathbb{N}\} = \{\alpha^{i-} \mid \alpha_i > \alpha_{i+1}\}.$$

Definition. Let $\alpha \vdash n$. We define

$$\alpha^+ := \{\beta \vdash n+1 \mid \beta = \alpha^{i+} \text{ for some } i \in \mathbb{N}\} = \{\alpha^{i+} \mid \alpha_i < \alpha_{i+1}\},$$

where we treat $\alpha_0 = \infty$. In other words, α^+ is the set of all partitions β such that $Y(\beta)$ can be obtained from $Y(\alpha)$ by adding a single box.

We will call (i, j) addable to α if $(i, j) \notin Y(\alpha)$ and $Y(\alpha) \cup \{(i, j)\} = Y(\beta)$ for some $\beta \in \alpha^+$.

Example. Let $\alpha = (4, 2, 2, 1) \vdash 9$. Then $Y(\alpha) = \begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \end{array} \begin{array}{l} + \\ + \\ - \\ + \end{array}$ where the removable resp. addable boxes are marked with a $-$ resp. $+$. So

$$\begin{aligned} \alpha^- &= \{(3, 2^2, 1), (4, 2, 1^1), (4, 2^2)\}, \\ \alpha^+ &= \{(5, 2^2, 1), (4, 3, 2, 1), (4, 2^3), (4, 2^2, 1^2)\}. \end{aligned}$$

Theorem 3.22 (Branching Rule - restriction). *Let $\alpha \vdash n$. Then $\chi^\alpha \downarrow_{S_{n-1}} = \sum_{\beta \in \alpha^-} \chi^\beta$.*

Proof. We have

$$\begin{aligned} \chi^\alpha \downarrow_{S_{n-1}} &= \psi^\alpha \downarrow_{S_{n-1}} = \sum_{\pi} \text{sgn } \pi \xi^{\alpha - \text{id} + \pi} \downarrow_{S_{n-1}} \\ &= \sum_{\pi} \text{sgn } \pi \sum_{i \in \mathbb{N}} \xi^{(\alpha - \text{id} + \pi)^{i-}} \\ &= \sum_{i \in \mathbb{N}} \sum_{\pi} (\text{sgn } \pi) \xi^{\alpha^{i-} - \text{id} + \pi} \\ &= \sum_{i \in \mathbb{N}} \psi^{\alpha^{i-}} \end{aligned}$$

Now if $\psi^{\alpha^{i-}} \neq 0$, then $\alpha_i^{i-} - i \neq \alpha_{i+1}^{i-} - (i+1)$ by Lemma 3.5, so $\alpha_i - 1 - i \neq \alpha_{i+1} - (i+1)$ and so $\alpha_i \neq \alpha_{i+1}$, then $\alpha^{i-} \in \alpha^-$. \square

Corollary 3.23 (Branching Rule - induction). *Let $\alpha \vdash n$. Then $\chi^\alpha \uparrow^{S_{n+1}} = \sum_{\beta \in \alpha^+} \chi^\beta$.*

Proof. This follows from Theorem 3.22 and Frobenius reciprocity noting that $\beta \in \alpha^+$ iff $\alpha \in \beta^-$. \square

Example. Let $\alpha = (4, 2^2, 1) \vdash 9$. Then


$$\begin{aligned} \chi^\alpha \downarrow_{S_8} &= \chi^{(3, 2^2, 1)} + \chi^{(4, 2, 1^2)} + \chi^{(4, 2^2)}, \\ \chi^\alpha \uparrow^{S_{10}} &= \chi^{(5, 2^2, 1)} + \chi^{(4, 3, 2, 1)} + \chi^{(4, 2^3)} + \chi^{(4, 2^2, 1^2)}. \end{aligned}$$

Definition. *The Young (branching) graph \mathbb{Y} is the graph with*

- vertex set $\bigcup_{n \in \mathbb{N}_0} \wp(n)$,
- edge set $\{(\lambda, \mu) \mid \mu \in \lambda^-\}$.

We will call $\wp(n)$ th n -th layer or level of \mathbb{Y} .

Here are the first five layers of \mathbb{Y} :



For each partition λ , there is a natural bijection between $\text{std}(\lambda)$ and the set of upwards-directed paths from \emptyset to λ in \mathbb{Y} . Indeed, given such a path, we construct the standard λ -tableau by putting in the layer number in each newly added box in the path. E.g. consider $\lambda = (3, 1)$ and the path

$$\emptyset \rightarrow \square \rightarrow \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \rightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}.$$

Then we get the sequence of tableaux

$$\emptyset \rightarrow \begin{array}{|c|} \hline 1 \\ \hline \end{array} \rightarrow \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \end{array} \rightarrow \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 2 \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 2 & 2 \\ \hline \end{array}.$$

Now $\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 2 & 2 \\ \hline \end{array}$ is the standard tableau corresponding to this path.


3.3.3 Murnaghan-Nakayama Rule

Definition. Let $\lambda \vdash n$, $(i, j) \in Y(\lambda)$.


- (i) The rim of λ is $R(\lambda) := \{(x, y) \in Y(\lambda) \mid (x + 1, y + 1) \notin Y(\lambda)\}$.
- (ii) The (i, j) -rim hook of λ is $R_{i,j}(\lambda) = \{(x, y) \in R(\lambda) \mid x \geq i \text{ and } y \geq j\}$. Its hand is (i, λ_i) and its foot is (λ'_j, j) , the same as for $H_{i,j}(\lambda)$.
- (iii) The leg length of $R_{i,j}(\lambda)$ is $\lambda'_i - j$, and arm length $\lambda_i - j$, same as for $H_{i,j}(\lambda)$.

Note that for both the hook and the rim hook the leg (resp. arm) length is the number of rows (resp. columns) occupied by the hook, minus one.


Example. Let $\lambda = (7, 5^3, 3, 1) \vdash 26$. Then the boxes in the rim $R(\lambda)$ are highlighted green.




Take $(i, j) = (2, 2)$. The rim hook is highlighted red.



Removing $H_{2,2}(\lambda)$ (resp. $R_{2,2}(\lambda)$) leaves



$(7, 1^5), (3, 3, 2)$



$(7, 4^2, 2, 1^2)$

Note that if we merge the two components obtained by removing $H_{2,2}(\lambda)$ get precisely what is left after removing $R_{2,2}(\lambda)$.

Lemma 3.24. Let $\lambda \vdash n$, $(i, j) \in Y(\lambda)$.

- (i) $|R_{i,j}(\lambda)| = |H_{i,j}(\lambda)| = h_{i,j}(\lambda)$,
- (ii) Removing $H_{i,j}(\lambda)$ from $Y(\lambda)$, and then sliding the lower-right component (if the result was disconnected) up and left one unit each, gives $Y(\lambda) \setminus R_{i,j}(\lambda)$.

Proof. Consider a walk along each hook, one box at a time traversing from the hand to the foot. Then the claims follow since

- $H_{i,j}(\lambda)$ and $R_{i,j}(\lambda)$ have the same hands and feet,
- we only move left or down at each step,
- we use the same number of leftward steps (namely the common arm length $\lambda_i - j$), and downward steps (by length $\lambda'_j - i$).

□

Definition. Let $\lambda \vdash n$, $(i, j) \in Y(\lambda)$. Define $\lambda \setminus H_{i,j}(\lambda)$ to be the partition of $|\lambda| - h_{i,j}(\lambda)$ such that $Y(\lambda \setminus H_{i,j}(\lambda)) = Y(\lambda) \setminus R_{i,j}(\lambda)$. Explicitly, letting $a = \lambda_i - j$ be the arm length, and $b = \lambda'_j - i$ be the leg length, then

$$\lambda \setminus H_{i,j}(\lambda) = (\lambda_1, \lambda_2, \dots, \lambda_{i-1}, \lambda_{i+1} - 1, \lambda_{i+2} - 1, \dots, \lambda_{i+b} - 1, j - 1, \lambda_{i+b+1}, \lambda_{i+b+2}, \dots).$$

Note that $h_{i,j}(\lambda) = 1 + a + b$, so $j - 1 = \lambda_i - a - 1 = \lambda_i - h_{i,j}(\lambda) + b$.

- From now on, when we remove a hook from λ , we mean to get $\lambda \setminus H_{i,j}(\lambda)$ for some $(i, j) \in Y(\lambda)$.
- If μ is obtained from λ by removing a hook, then we let $LL(\lambda \setminus \mu)$ denote the leg length of the removed hook. That is, $\mu = \lambda \setminus H_{i,j}(\lambda)$ for some $(i, j) \in Y(\lambda)$, and $LL(\lambda \setminus \mu)$ is the leg length of $H_{i,j}(\lambda)$, equivalently of $R_{i,j}(\lambda)$.

Theorem 3.25 (Murnaghan-Nakayama Rule). *Let $\alpha \vdash n$, $k \in [n]$. Let $\pi \in S_n$, and suppose that it has a k -cycle in its disjoint cycle decomposition. Let $\rho \in S_{n-k}$ have the same cycle type as π but with one fewer k -cycle. Then*

$$\chi^\alpha(\pi) = \sum_{\beta} (-1)^{LL(\alpha \setminus \beta)} \chi^\beta(\rho),$$

where the sum runs over partitions β obtained from α by removing a hook of size k .

Example. Let $\alpha = (3^3) \vdash 9$, $\pi = (1234)(56)(789)$. We take $k = 3$ and $\rho = (1234)(56)$. What are the possible hooks of size 3 we can remove?

$$\begin{array}{ccc} \begin{array}{|c|c|c|} \hline & & \times \\ \hline & & \times \\ \hline & & \times \\ \hline \end{array} & \begin{array}{|c|c|c|} \hline & & \\ \hline & & \times \\ \hline & \times & \times \\ \hline \end{array} & \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \times & \times & \times \\ \hline \end{array} \\ LL = 2 & LL = 1 & LL = 0 \end{array}$$

Then

$$\chi^\alpha(\pi) = (\chi^{(2^3)} - \chi^{(3,2,1)} + \chi^{(3^2)})(\mu)$$

We repeat this with $n = 6, k = 2$. So this is

$$\begin{array}{l} \begin{array}{|c|c|} \hline & \\ \hline & \times \\ \hline & \times \\ \hline \end{array} & \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \times & \times \\ \hline \end{array} & \text{no removable} & \begin{array}{|c|c|c|} \hline & & \times \\ \hline & & \\ \hline & & \\ \hline \end{array} & \begin{array}{|c|c|c|} \hline & & \\ \hline & & \times \\ \hline & \times & \times \\ \hline \end{array} \\ LL = 1 & LL = 0 & \text{hooks of size 2} & LL = 1 & LL = 0 \\ = \left(-\chi^{(2,1^2)} + \chi^{(2^2)} + 0 - \chi^{(2^2)} + \chi^{(3,1)} \right) ((1234)) \\ = (-\chi^{(2,1^2)} + \chi^{(3,1)})((1234)) \\ \begin{array}{|c|c|} \hline \times & \times \\ \hline \times & \\ \hline \times & \\ \hline \end{array} & \begin{array}{|c|c|c|} \hline \times & \times & \times \\ \hline \times & & \\ \hline \times & & \\ \hline \end{array} \\ LL = 2 & LL = 2 \\ = \left(-\chi^\emptyset - \chi^\emptyset \right) (e) \\ = -1 - 1 = -2 \end{array}$$

Proof of Theorem 3.25. Since characters are class functions, we may assume that $\pi = \rho\sigma$ for some k -cycle σ disjoint from ρ . For $\mu \models k$, recall $\xi^\mu = \mathbb{1}_{S_\mu} \uparrow^{S_k}$, so $\xi^\mu(\sigma) = \frac{1}{|S_\mu|} \sum_{\substack{x \in S_k \\ x\sigma x^{-1} \in S_\mu}} 1$. But since σ is a k -cycle, σ belongs to a conjugate of S_μ if and only if

$\mu = (\dots, 0, k, 0, \dots)$. So if μ is not of this form, then $\xi^\mu(\sigma) = 0$. On the other hand, if $\mu = (\dots, 0, k, 0, \dots)$, then $\xi^\mu = \mathbb{1}_{S_{(k)}} \uparrow^{S_k} = \mathbb{1}_{S_k}$, and so $\xi^\mu(\sigma) = 1$. Therefore

$$\begin{aligned}
\chi^\alpha(\pi) &= \psi^\alpha(\pi) = \psi^\alpha \downarrow_{S_{n-k} \times S_k}(\rho\sigma) \\
&\stackrel{\text{Lemma 3.7}}{=} \sum_{\mu \models k} \left(\psi^{\alpha-\mu} \# \xi^\mu \right) (\rho\sigma) \\
&= \sum_{\mu \models k} \psi^{\alpha-\mu}(\rho) \xi^\mu(\sigma) \\
&= \sum_{i=1}^{\infty} \psi^{(\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_i - k, \alpha_{i+1}, \dots)}(\rho) \\
&= \sum_{i \in \mathbb{N}} \psi^{\beta_{i,0}}(\rho) \tag{*}
\end{aligned}$$

where we let $\beta_{i,0} = (\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_i - k, \alpha_{i+1}, \dots) \models n - k$.

Recall from Lemma 3.5 that if $\gamma - \text{id} = (j \ j+1) \circ (\lambda - \text{id})$, then $\psi^\gamma = -\psi^\lambda$. Fix $i \in \mathbb{N}$, define $\beta_{i,m} \models n - k$ via $\beta_{i,m} - \text{id} = (i+m \ i+m-1 \ \dots \ i+2 \ i+1 \ i) \circ (\beta_{i,0} - \text{id})$, for each $m \in \mathbb{N}_0$. Explicitly, $\beta_{i,m} = (\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_{i+1} - 1, \dots, \alpha_{i+m} - 1, \alpha_i - k + m, \alpha_{i+m+1}, \dots)$. Since $(i+m \ i+m-1 \ \dots \ i+2 \ i+1 \ i) = (i+m \ i+m-1) \cdots (i+2 \ i+1)(i+1 \ i)$, we can apply Lemma 3.5 repeatedly to get

$$\psi^{\beta_{i,0}} = (-1)^m \psi^{\beta_{i,m}}.$$

We will see that

- if there exists $m \in \mathbb{N}_0$ such that $\beta_{i,m}$ is a partition, then m is unique, and we will relate $\beta_{i,m}$ to α by removing an appropriate hook,

† while if there does not exist such an m , then we will show that $\psi^{i,m} = 0$.

For $(i, j) \in Y(\alpha)$, letting b be the leg length of $H_{i,j}(\alpha)$, we recall that $\alpha \setminus H_{i,j}(\alpha) = (\alpha_1, \dots, \alpha_{i-1}, \alpha_{i+1} - 1, \alpha_{i+2} - 1, \dots, \alpha_{i+k} - 1, \alpha_i - h_{i,j}(\alpha) + b, \alpha_{i+k+1}, \alpha_{i+k+2}, \dots)$. Compare this with $\beta_{i,m}$. For any given i , we see that the following are equivalent:

- the existence of an $m \in \mathbb{N}_0$ such that $\beta_{i,m}$ is a partition,
- the existence of a rim hook $R_{i,j}(\alpha)$ of size k , for some $j \in [\lambda_i]$.

The highest row occupied by this hook is row i . In particular, $i \leq \ell(\alpha)$. A rim hook is uniquely determined by its highest row and size. In particular, there is at most one m for each i , and when this exists, m is uniquely determined as the leg length of the hook.

Notice if $i > \ell(\alpha)$, then for all $m \in \mathbb{N}_0$, $\beta_{i,m}$ has a negative part:

$$\begin{aligned}
\beta_{i,0} &= (\alpha_1, \alpha_2, \dots, \alpha_{\ell(\alpha)}, 0, \dots, 0, -k, 0 \dots) \\
\beta_{i,1} &= (\alpha_1, \alpha_2, \dots, \alpha_{\ell(\alpha)}, 0, \dots, 0, -1, -k+1, 0 \dots) \\
\beta_{i,2} &= (\alpha_1, \alpha_2, \dots, \alpha_{\ell(\alpha)}, 0, \dots, 0, -1, -1, -k+2, 0 \dots)
\end{aligned}$$

So we never talk about a hook length in row i unless i is a genuine row in $Y(\alpha)$.

Once we prove the claim †, then (*) gives

$$\begin{aligned}
\chi^\alpha(\pi) &= \sum_{i \in \mathbb{N}} \psi^{\beta_{i,0}}(\rho) \\
&= \sum_{\substack{i \in \mathbb{N} \text{ such that} \\ \exists m \in \mathbb{N}_0: \beta_{i,m} \text{ is a partition}}} (-1)^m \psi^{\beta_{i,m}}(\rho) \\
&= \sum_{\substack{\beta \vdash n-k \\ \text{obtained from } \alpha \\ \text{by removing a hook} \\ \text{of size } k}} (-1)^{LL(\alpha \setminus \beta)} \psi^\beta(\rho) \\
&= \sum_{\beta} \chi^\beta(\rho).
\end{aligned}$$

So it remains to prove †. Fix $i \in \mathbb{N}$ and suppose $\beta_{i,m} \not\vdash n - k$ for all $m \in \mathbb{N}_0$. Observe

$$\begin{aligned}
\beta_{i,m} - \text{id} &= (\alpha_1 - 1, \alpha_2 - 2, \dots, \alpha_{i-1} - (i-1), \alpha_{i+1} - (i+1), \dots, \alpha_{i+m} - (i+m), \\
&\quad \alpha_i - i - k, \alpha_{i+m+1} - (i+m+1), \dots)
\end{aligned}$$

Since α is a partition, $\alpha - \text{id}$ is strictly decreasing. Since $\alpha_i - i \geq \alpha_i - i - k \geq \alpha_{i+k} - (i+k)$, there exists a unique $t \in \mathbb{N}_0$ such that $\alpha_{i+t} - (i+t) \geq \alpha_i - i - k > \alpha_{i+t+1} - (i+t+1)$. If $\alpha_{i+t} - (i+t) = \alpha_i - i - k$, then $\beta_{i,t} - \text{id}$ has two adjacent terms equal. But then $\psi^{\beta_{i,t}} = 0$ by Lemma 3.5, hence $\psi^{\beta_{i,0}} = (-1)^t \psi^{\beta_{i,t}} = 0$. Otherwise, $\alpha_{i+t} - (i+t) > \alpha_i - i - k$. But that means $\beta_{i,t}$ is weakly decreasing. Also $(\beta_{i,t})_j = \alpha_j$ for all $j \geq i+t+1$ and $\alpha_j \geq 0$ for all $j \in \mathbb{N}$. Also α has finite support, hence so does $\beta_{i,t}$, thus $\beta_{i,t}$ is a partition, contradicting our assumption. This proves † and hence the proof of the theorem. \square


4 McKay Numbers

In this chapter we go back to partitions, and continue with $\mathbb{F} = \mathbb{C}$.

Main goal. Describe $\text{Irr}_{p'}(S_n)$ and work towards understanding the techniques in Olsson's proof of the McKay Conjecture for symmetric groups.

4.1 James's Abacus

Example. Let $\lambda = (7, 5^3, 3, 1) \vdash 26$, and consider $H_{2,2}(\lambda), R_{2,2}(\lambda)$. Write $1, 2, \dots, h_{2,2}(\lambda)$ into $R_{2,2}(\lambda)$ from hand to foot. For those numbers in boxes at the bottom of their column, write them in $H_{2,2}(\lambda)$ in the same column. For the rest, write them in to $H_{2,2}(\lambda)$ in the row below.



Observe

$$\begin{aligned} 1 &= 7 - 6 = h_{2,2}(\lambda) - h_{3,2}(\lambda) \\ 2 &= 7 - 5 = h_{2,2}(\lambda) - h_{4,2}(\lambda) \\ 5 &= 7 - 2 = h_{2,2}(\lambda) - h_{5,2}(\lambda) \end{aligned}$$

Lemma 4.1. Let $\lambda \vdash n, (i, j) \in Y(\lambda)$. Then

$$\{1, 2, \dots, h_{i,j}(\lambda)\} = \{h_{i,y}(\lambda) \mid j \leq y \leq \lambda_i\} \sqcup \{h_{i,j}(\lambda) - h_{x,j}(\lambda) \mid i < x \leq \lambda'_j\}$$

Proof. We omit (λ) from the notation. Let $A = \{(u, v) \in R_{i,j} \mid u = \lambda'_v\} = \{(\lambda'_y, y) \mid j \leq y \leq \lambda_i\}$ and $B = \{(u, v) \in R_{i,j} \mid u \neq \lambda'_v\} = \{(x-1, \lambda_x) \mid i < x \leq \lambda'_j\}$.

By Lemma 3.24, $|R_{i,j}| = h_{i,j}$, so we may fill the numbers $1, 2, \dots, h_{i,j}$ into $R_{i,j}$ one number in each box from hand to foot. We claim that A is filled with $\{h_{i,y} \mid j \leq y \leq \lambda_i\}$ and B with $\{h_{i,j} - h_{x,j} \mid i < x \leq \lambda'_j\}$, whence the lemma follows. Consider (λ'_y, y) in A . It is filled with

$$\begin{aligned} &1 + \# \text{left steps} + \# \text{down steps} \\ &= 1 + \text{arm length of } H_{i,y} + \text{leg length of } H_{i,y} = h_{i,y} \end{aligned}$$

Consider $(x-1, \lambda_x) \in B$. It is filled with

$$\begin{aligned} & 1 + \#\text{left steps} + \#\text{down steps} \\ &= 1 + (\lambda_i - \lambda_x) + (x-1-i) = (1 + \lambda_i - j + \lambda'_j - i) - (1 + \lambda_x - j + \lambda'_j - x) \\ &= h_{i,j} - h_{x,j}. \end{aligned}$$

□

Definition. Let $\lambda \vdash n$, $m = \ell(\lambda)$.

- (i) Let $X_\lambda = \{h_{1,1}(\lambda), h_{2,1}(\lambda), \dots, h_{m,1}(\lambda)\}$ be the set of first column hook lengths of λ .
- (ii) For each $i \in [m]$, let $\mathcal{H}_i(\lambda) = \{h_{i,j}(\lambda) \mid j \in [\lambda_i]\}$ be the set of row i hook lengths of λ .

Note that $\mathcal{H}_i(\lambda) = \{1, 2, \dots, h_{i,1}(\lambda)\} \setminus \{h_{1,1}(\lambda) - h_{x,1}(\lambda) \mid i < x \leq m\}$ by Lemma 4.1.

Convention: If $i > m$, then $\mathcal{H}_i(\lambda) = \emptyset$.

Notice that X_λ determines λ : If we know that $\{h_1, h_2, \dots, h_m\}$ where $h_1 > h_2 > \dots > h_m$, is the set of first column hook lengths for some partition λ , then λ must be $\lambda = (h_1 - (m-1), \dots, h_{m-1} - 1, h_m - 0)$.

Idea. We represent partitions using beads on an abacus.

- Info about hook lengths is encoded into the bead positions
- given an arrangement of beads, we will be able to reconstruct the partition using observations like the above.
- advantages: operations on partitions (e.g. hook removal) are easy to describe.

Definition. A β -set X is a finite subset $\{h_1, \dots, h_m\}$ of \mathbb{N}_0 . Convention: $h_1 > h_2 > \dots > h_m$.

For a β -set $X = \{h_1, \dots, h_m\}$ and $l \in \mathbb{N}_0$, we define X^{+l} , the l -shift of X , as follows:

- $X^{+0} = X$,
- if $l > 0$, then $X^{+l} = \{h_1 + l, h_2 + l, \dots, h_m + l\} \cup \{l-1, l-2, \dots, 1, 0\}$.

We define the partition corresponding to X to be $\mathcal{P}(X) = (h_1 - (m-1), h_2 - (m-2), \dots, h_{m-1} - 1, h_m - 0)$. This expression for $\mathcal{P}(X)$ may have trailing zeros, which can be removed.

Example. Let $X = \{4, 2\}$. Then $\mathcal{P}(X) = (4-1, 2-0) = (3, 2)$. And $X^{+2} = \{6, 4, 1, 0\}$ and $\mathcal{P}(X^{+2}) = (6-3, 4-2, 1-1, 0-0) = (3, 2)$.

Lemma 4.2. Let $\lambda \vdash n$ and X a β -set. Then X is a β -set for λ , meaning $\mathcal{P}(X) = \lambda$, if and only if $X \in \{X_\lambda^{+l} \mid l \in \mathbb{N}_0\}$.

Proof. Let $X = \{h_1, h_2, \dots, h_m\}$ and $t = \ell(\lambda)$. Then

$$\begin{aligned}
\mathcal{P}(X) = \lambda &\iff (h_1 - (m - 1), \dots, h_{m-1} - 1, h_m - 0) = (\lambda_1, \lambda_2, \dots, \lambda_t) \\
&\iff m \geq t \text{ and } \begin{cases} h_j - (m - j) = \lambda_j & \text{if } j \leq t, \\ h_j - (m - j) = 0 & \text{if } j > t \end{cases} \\
&\iff m \geq t \text{ and } h_j = \begin{cases} \lambda_j + (t - j) + (m - t) & \text{if } j \leq t, \\ m - j & \text{if } j > t \end{cases} \\
&\iff m - t \in \mathbb{N}_0 \text{ and } X = X_\lambda^{+(m-t)}
\end{aligned}$$

□

Definition. Let $e \in \mathbb{N}$. James's e -abacus consists of e runners (drawn as columns) labelled $0, 1, 2, \dots, e - 1$ from left to right, with rows labelled by \mathbb{N}_0 increasing downwards. The positions are labelled by \mathbb{N}_0 , with that in row a and runner i labelled by $ae + i$.

Given a β -set X , the e -abacus configuration corresponding to X has beads precisely in positions given by the elements of X . We call the configuration A_X . Conversely, given an e -abacus configuration A , i.e. a finite set of beads in the e -abacus, define the corresponding β -set X_A to be the set of position labels of the beads. We define the corresponding partition to be $\mathcal{P}(A) := \mathcal{P}(X_A)$.

Also, if $X = X_\lambda$, then abbreviate $A_{X_\lambda} = A_\lambda$.

Clearly,

$$\begin{aligned}
\{e\text{-abacus configurations}\} &\xleftrightarrow{1-1} \{\beta\text{-sets}\} \\
A_X &\longleftarrow X \\
A &\longmapsto X_A \\
\text{bead positions} &\longleftrightarrow \{h_1, \dots, h_m\}
\end{aligned}$$

e -abacus:

	0	1	2	\dots	$e - 1$
0	0	1	2	\dots	$e - 1$
1	e	$e + 1$	$e + 2$	\dots	$2e - 1$
2	$2e$	$2e + 1$	$2e + 2$	\dots	$3e - 1$
\vdots			\vdots		

Examples.

(i) Let $e \geq 2$, $X = \{2e, e + 1, 2\}$. On an e -abacus we have

	0	1	2	\dots	$e - 1$
0	0	1	②	\dots	$e - 1$
1	e	①	$e + 2$	\dots	$2e - 1$
2	②	$2e + 1$	$2e + 2$	\dots	$3e - 1$
\vdots			\vdots		

and $\mathcal{P}(X) = (2e - 2, e, 2)$.

(ii) Consider the 3-abacus configuration A given by

	0	1	2
0	0	①	②
1	③	4	⑤
2	6	7	8
3	⑨	10	⑪

so $X_A = \{11, 9, 5, 3, 2, 1\}$ and $\mathcal{P}(A) = \mathcal{P}(X_A) = (6, 5, 2, 1^3) \vdash 16$.

Let $X = \{6, 4, 1, 0\}$, $e = 3$ or $e = 4$. Then

	0	1	2			0	1	2	3
0	①	②	2	0	0	①	②	2	3
1	3	④	5	1	④	5	⑥	7	
2	⑥	7	8						

We have $\mathcal{P}(X) = (3, 2)$.

Lemma 4.3. *Let $e \in \mathbb{N}$. Given an e -abacus configuration A , with beads at $h_1 > h_2 > \dots > h_m$, then $\mathcal{P}(A) = (a_1, a_2, \dots, a_m)$ where a_j is the number of gaps, i.e. empty positions, i such that $0 \leq i < h_j$.*

Proof. By definition, $\mathcal{P}(A) = (h_1 - (m - 1), \dots, h_m - 0)$. But there are h_j positions before h_j , of which $m - j$ have beads, namely h_{j+1}, \dots, h_m . \square

Definition. *Let $X = \{h_1, \dots, h_m\}$ be a β -set. For $i \in [m]$, define $\mathcal{H}_i(X) = \{1, 2, \dots, h_i\} \setminus \{h_i - h_j \mid i < j \leq m\}$.*

Lemma 4.4. *Let $\lambda \vdash n$ and X a β -set for λ . If $X = \{h_1, \dots, h_m\}$, then $\mathcal{H}_i(X) = \mathcal{H}_i(\lambda)$ for all $i \in [m]$.*

Proof. We have $X = X_\lambda^{+(m-\ell(\lambda))}$ from the proof of Lemma 4.2. If $i > \ell(\lambda)$, then $|\mathcal{H}_i(X)| = h_i - (m - i) = 0$, so $\mathcal{H}_i(X) = \mathcal{H}_i(\lambda) = \emptyset$. If $i \leq \ell(\lambda)$, then $\mathcal{H}_i(\lambda) = \{1, 2, \dots, h_{i,1}(\lambda)\} \setminus$

$\{h_{1,1}(\lambda) - h_{j,1}(\lambda) \mid i < j \leq \ell(\lambda)\}$ and so clearly $\mathcal{H}_i(\lambda) = \mathcal{H}_i(X_\lambda)$. So it remains to check $\mathcal{H}_i(X) = \mathcal{H}_i(X^{+1})$. We have $X^{+1} = \{h_1 + 1, h_2 + 1, \dots, h_m + 1, 0\}$, so

$$\begin{aligned} \mathcal{H}_i(X^{+1}) &= \{1, 2, \dots, h_i + 1\} \setminus (\{(h_i + 1) - (h_j + 1) \mid i < j \leq m\} \cup \{(h_i + 1) - 0\}) \\ &= \{1, 2, \dots, h_i\} \setminus \{h_i - h_j \mid i < j \leq m\} = \mathcal{H}_i(X). \end{aligned}$$

□

Corollary 4.5. *Let $\lambda \vdash n$ and $X = \{h_1, \dots, h_m\}$ be a β -set for λ . Let $h \in \mathbb{N}_0$. Then $h \in \mathcal{H}_i(\lambda)$ iff $h_i - h \geq 0$ and $h_i - h \notin X$, for any $i \in [m]$.*

Proof. The claim is clear if $i > \ell(\lambda)$ (since then $\mathcal{H}_i(\lambda) = \emptyset$), or if $h = 0$. So we may assume that $i \leq \ell(\lambda)$ and $h > 0$. If $h > h_i$, then $h > \max \mathcal{H}_i(X) = \max \mathcal{H}_i(\lambda)$, so $h \in \mathcal{H}_i(\lambda)$. Otherwise, $h \leq h_i$. Recall $\mathcal{H}_i(\lambda) = \mathcal{H}_i(X) = \{1, 2, \dots, h_i\} \setminus \{h_i - h_j \mid i < j \leq m\}$. So

$$\begin{aligned} h \notin \mathcal{H}_i(\lambda) &\iff h = h_i - h_j \text{ for some } i < j \leq m \\ &\iff h_i - h \in X \end{aligned}$$

□

Corollary 4.6. *Let $\lambda \vdash n$ and suppose $ef \in \mathcal{H}(\lambda)$ for some $e, f \in \mathbb{N}$. Then $e \in \mathcal{H}(\lambda)$.*

Proof. Let $X = X_\lambda = \{h_1, h_2, \dots, h_m\}$. Since $ef \in \mathcal{H}(\lambda)$, then $ef \in \mathcal{H}_i(\lambda)$ for some $i \in [m]$. By Corollary 4.5, $0 \leq h_i - ef \notin X$. But $h_i \in X$, so there exists $l \in \{0, 1, \dots, f-1\}$ such that $0 \leq h_i - e(l+1) \notin X$, but $h_i - el \in X$. This means $h_i - el = h_k$ for some $i \leq k \leq m$. But then $0 \leq h_k - e = h_i - e(l+1) \notin X$, hence by Corollary 4.5 again, $e \in \mathcal{H}_k(\lambda)$. □

Example. Let $\lambda = (7, 5^2, 3, 1) \vdash 21$. So:

11						
8	×	×	×	×		
7	×					
4	×					
1						

$H_{2,2}(\lambda)$

11						
7				×		
4		×	×	×		
2	×	×				
1						

$R_{2,2}(\lambda)$

$$X_\lambda = \{11, 8, 7, 4, 1\}$$

$$X_{\lambda \setminus H_{2,2}(\lambda)} = \{11, 7, 4, 2, 1\}$$

Note that $X_{\lambda \setminus H_{2,2}(\lambda)} = (X_\lambda \setminus \{8\}) \sqcup \{8 - h_{2,2}(\lambda)\}$ and 8 is the second element in X_λ .

Proposition 4.7. *Let $\lambda \vdash n$, $X = \{h_1, h_2, \dots, h_m\}$ be a β -set for λ . Let $(i, j) \in Y(\lambda)$. Then*

(i) $0 \leq h_i - h_{i,j}(\lambda) \notin X$,

(ii) $Z := (X \setminus \{h_i\}) \sqcup \{h_i - h_{i,j}(\lambda)\}$ is a β -set for $\lambda \setminus H_{i,j}(\lambda)$

Proof.

- (i) Immediate from Corollary 4.5.
- (ii) Since β -sets are determined up to shift, and $Z^{+l} = (X^{+l} \setminus \{h_i + l\}) \sqcup \{(h_i + l) - h_{i,j}(\lambda)\}$, then it is enough to prove (ii) for $X = X_\lambda$. So now assume $X = X_\lambda$, $m = \ell(\lambda)$, $h_i = h_{i,j}(\lambda)$. Let $\mu = \lambda \setminus H_{i,j}(\lambda)$. Recall that if b is the leg length of $H_{i,j}(\lambda)$, then $\mu = (\lambda_1, \lambda_2, \dots, \lambda_{i-1}, \lambda_{i+1} - 1, \dots, \lambda_{i+b} - 1, j - 1, \lambda_{i+b+1}, \dots)$. Let Z' be the β -set for μ such that $|Z'| = m$. This does exist, since $\ell(\mu) \leq \ell(\lambda) = m$, so in particular, Z' is just $X_\mu^{+(m-\ell(\mu))}$. Let $Z' = \{k_1, \dots, k_m\}$. We compute Z' :

- For $s < i$, then $k_s = \mu_s + (\ell(\mu) - s) + (m - \ell(\mu)) = \lambda_s + m - s = h_{s,1}(\lambda) = h_s$.
- For $s \in \{0, 1, \dots, b-1\}$, $k_{i+s} = \mu_{i+s} + (\ell(\mu) - (i+s)) + (m - \ell(\mu)) = (\lambda_{i+s+1} - 1) + m - (i+s) = \lambda_{i+s+1} + m - (i+s+1) = h_{i+s+1}$.
- $k_{i+b} = \mu_{i+b} + (\ell(\mu) - (i+b)) + (m - \ell(\mu)) = j - 1 + m - i - b$.
- For $s \geq i+b+1$, $k_s = \mu_s + m - s = \lambda_s + m - s = h_s$.

So $Z' = (X \setminus \{h_i\}) \sqcup \{j - 1 + m - i - b\}$. But $h_i - h_{i,j} = h_{i,1}(\lambda) - h_{i,j}(\lambda) = (\lambda_i + m - i) - (1 + \lambda_i - j + b) = j - 1 + m - i - b$. So $Z' = Z$.

□

Corollary 4.8. *Let $e \in \mathbb{N}$, $\lambda \vdash n$, $X = \{h_1, \dots, h_m\}$ a β -set for λ , $i \in [m]$. Write $h_i = ae + j$ for $a \in \mathbb{N}_0$ and $j \in \{0, 1, \dots, e-1\}$. Then the following are equivalent:*

- *There exists $y \in [\lambda_i]$ such that $h_{i,y}(\lambda) = e$.*
- *$a \geq 1$ and $(a-1)e$ is an empty position in the e -abacus configuration A_X .*

When these hold, y is unique.

Moreover, the e -abacus configuration A' obtained from A_X by sliding the bead in position h_i to position $h_i - e$ has $\mathcal{P}(A') = \lambda \setminus H_{i,y}(\lambda)$.

In other words, removing a hook of size e is the same as sliding a bead up one row on an e -abacus.

Proof. By Corollary 4.5,

$$\begin{aligned} e \in \mathcal{H}_i(\lambda) &\iff 0 \leq h_i - e \notin X \\ &\iff a \geq 1, \text{ and } (a-1)e + j \notin X. \end{aligned}$$

Hence the equivalence. For the second part, clearly $X_{A'} = (X \setminus \{h_i\}) \cup \{h_i - e\}$, but this is a β -set for $\lambda \setminus H_{i,y}(\lambda)$ by Proposition 4.7. □

Remark. Recall the proof of Corollary 4.6 - we had $0 \leq h_i - ef \notin X$, $h_i \in X$. The existence of l in the proof is equivalent to there being a bead immediately below a gap


somewhere on this runner between h_i and $h_i - ef$. By Corollary 4.8, this corresponds to a hook of length e .

Just as we have the division algorithm for integers, giving quotients and remainders when we divide by e , we can do something similar for partitions, giving “ e -quotients”, and “ e -cores”.

Definition. Let $e \in \mathbb{N}$, $\lambda \vdash n$. We say that λ is an e -core partition if $e \notin \mathcal{H}(\lambda)$. The empty partition \emptyset is always an e -core for any e .

Example.

(i) Suppose $|\lambda| < e$. Then λ is an e -core partition.

(ii)  We can see that $(4, 3, 3)$ is not a 5 core, but $(2, 2, 1)$ is.

(iii) Let $e = 2$. Hooks of size 2 are always “dominoes” (i.e. 2×1 or 1×2 rectangles). So the 2-core partitions are precisely

$$\emptyset, \quad \square, \quad \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \quad \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \quad \dots$$

i.e. \emptyset and $(t, t - 1, \dots, 2, 1)$ for $t \in \mathbb{N}$.

Definition. Let $e \in \mathbb{N}$, $\lambda \vdash n$, X a β -set for λ .

(i) For $i \in \{0, 1, \dots, e - 1\}$, define $X_i^{(e)} = \{a \in \mathbb{N}_0 \mid ae + i \in X\}$. That is, $X_i^{(e)}$ is the set of row labels of beads on runner i of the e -abacus configuration A_X .

(ii) The e -quotient of λ is $Q_e(\lambda) := (\lambda^{(0)}, \lambda^{(1)}, \dots, \lambda^{(e-1)})$ where $\lambda^{(i)} = \mathcal{P}(X_i^{(e)})$. That is, $\lambda^{(i)}$ is the partition corresponding to the runner i of A_X viewed as a 1-abacus.

(iii) Define $X_{(e)} = \bigsqcup_{i=0}^{e-1} \{ae + i \mid 0 \leq a \leq |X_i^{(e)}| - 1\}$.

(iv) The e -core of λ is $C_e(\lambda) := \mathcal{P}(X_{(e)})$.

The e -abacus configuration $A_{X_{(e)}}$ is obtained from A_X by sliding beads up as high as possible. The description of $A_{X_{(e)}}$ and Corollary 4.8 imply that $C_e(\lambda)$ is indeed an e -core partition.

Lemma 4.9. Let $e \in \mathbb{N}$, $\lambda \vdash n$, X a β -set for λ .

(i) For $i \in \{1, 2, \dots, e - 1\}$, $(X^{+1})_i^{(e)} = X_{i-1}^{(e)}$.

(ii) $(X^{+1})_0^{(e)} = (X_{e-1}^{(e)})^{+1}$.

(iii) For $i \in \{0, 1, \dots, e - 1\}$, $\mathcal{P}((X^{+1})_i^{(e)}) = \mathcal{P}(X_{i-1}^{(e)})$ where $i - 1$ is taken mod e

(iv) $(X^{+1})_{(e)} = (X_{(e)})^{+1}$

(v) $\mathcal{P}((X^{+1})_{(e)}) = \mathcal{P}(X_{(e)})$

Proof. Example Sheet 3. □

Remarks.

- Lemma 4.9 (iv), (v) and Lemma 4.2 show that $C_e(\lambda)$ just depends only on e and λ , but not the choice of β -set X for λ .
- Lemma 4.9 (i), (ii) and (iii) show that if we shift X to X^{+1} , we induce a cyclic shift of the components of $Q_e(\lambda) = (\lambda^{(0)}, \lambda^{(1)}, \dots, \lambda^{(e-1)})$. So far, $Q_e(\lambda)$ therefore still depends on the choice of X . But X and X^{+e} give the same cyclic shift of $\lambda^{(i)}$, and $|X^{+l}| = |X| + l$, so to fix an ordering of the components of $Q_e(\lambda)$ and thereby specifying $Q_e(\lambda)$ uniquely from now on, we will always choose β -sets X such that $|X|$ is a multiple of e when calculating e -quotients.

Example. Let $e = 3$, $\lambda = (6, 5, 2, 1^3) \vdash 16$. Then $X_\lambda = \{11, 9, 5, 3, 2, 1\}$. Note that $3 \mid |X_\lambda|$. Let $X = X_\lambda$. Then

A_λ			
	0	1	2
0	0	①	②
1	③	4	⑤
2	6	7	8
3	⑨	10	⑪


$A_{X^{(3)}}$			
	0	1	2
0	①	②	③
1	④	5	⑥
2	7	8	⑩
3	9	10	11

So $C_3(\lambda) = (3, 1)$,

$$\begin{aligned} X_0^{(3)} &= \{3, 1\}, \\ X_1^{(3)} &= \{0\}, \\ X_2^{(3)} &= \{3, 1, 0\} \end{aligned}$$

and $Q_3(\lambda) = ((2, 1), \emptyset, (1))$.

Note that in total we moved four beads up when going from A_X to $A_{X^{(3)}}$. This could correspond to removing rim hooks as follows (order indicated by number)



Definition. Let $e \in \mathbb{N}$. An e -hook is a hook of size exactly e .

Theorem 4.10. Let $e \in \mathbb{N}$, $\lambda \vdash n$. Then $C_e(\lambda)$ is the unique e -core partition we obtain by successively removing e -hooks from λ until we cannot remove any more. In particular, this is independent of the order in which we removed the hooks.

Proof. Let X be a β -set for λ . Let γ be an e -core partition obtained from λ by removing some e -hooks. By Corollary 4.8, there exists a β -set Z for γ such that the e -abacus configuration A_Z is obtained from A_X by sliding all beads up as far as possible. But then clearly $Z = X_{(e)}$, and so $\gamma = \mathcal{P}(Z) = \mathcal{P}(X_{(e)}) = C_e(\lambda)$. \square

Definition. Let $e \in \mathbb{N}$, $\lambda \vdash n$. Consider $Q_e(\lambda) = (\lambda^{(0)}, \lambda^{(1)}, \dots, \lambda^{(e-1)})$. We say that H is a hook of $Q_e(\lambda)$ if $H = H_{i,j}(\lambda^{(s)})$ for some $s = 0, \dots, e-1$ and $(i, j) \in Y(\lambda^{(s)})$. Moreover, we define $Q_e(\lambda) \setminus H := (\lambda^{(0)}, \dots, \lambda^{(s-1)}, \lambda^{(s)} \setminus H, \lambda^{(s+1)}, \dots, \lambda^{(e-1)})$. When we refer to a hook H of $Q_e(\lambda)$, it is considered to carry both the information of which component $\lambda^{(s)}$ it came from, as well as the box (i, j) .

Theorem 4.11. Let $e \in \mathbb{N}$, $\lambda \vdash n$. There is a bijection

$$f : \{H_{i,j}(\lambda) \text{ s.t. } e \mid h_{i,j}(\lambda)\} \rightarrow \{\text{hooks of } Q_e(\lambda)\}$$

such that if $H = H_{i,j}(\lambda)$ with $e \mid h_{i,j}(\lambda)$, then $|H| = e|f(H)|$ and $Q_e(\lambda \setminus H) = Q_e(\lambda) \setminus f(H)$.

Proof. Let $X = \{h_1, h_2, \dots, h_m\}$ be a β -set for λ with $e \mid m$. Recall from Corollary 4.5 that for $i \in [m]$ and $h \in \mathbb{N}_0$,

$$h \in \mathcal{H}_i(\lambda) \iff 0 \leq h_i - h \notin X.$$

So we get a bijection

$$\{H_{i,j}(\lambda) \text{ s.t. } (i, j) \in Y(\lambda)\} \rightarrow \{(b, g) \in \mathbb{N}_0^2 \mid b > g, b \in X, g \notin X\},$$

i.e. pairs of positions (b, g) in the e -abacus configuration A_X such that b is a bead, g is a gap and $b > g$. If $H_{i,j}(\lambda) \mapsto (b, g)$, then $h_i = b$ and $h_i - h_{i,j}(\lambda) = g$. In particular, $h_{i,j}(\lambda) = b - g$. So this restricts to a bijection

$$F : \{H_{i,j}(\lambda) \text{ s.t. } e \mid h_{i,j}(\lambda)\} \rightarrow \{(b, g) \in \mathbb{N}_0^2 \mid b > g, b \in X, g \notin X, b \equiv g \pmod{e}\}$$

If $b \equiv g \pmod{e}$, then $b = b'e + s$ and $g = g'e + s$ for some $s \in \{0, 1, \dots, e-1\}$ and some $b' > g' \in \mathbb{N}_0$. Again by Corollary 4.5, since $Q_e(\lambda) = (\lambda^{(0)}, \dots, \lambda^{(e-1)})$ has $\lambda^{(s)} = \mathcal{P}(X_s^{(e)})$, and $X_s^{(e)} = \{a \in \mathbb{N}_0 \mid ae + s \in X\}$, we have bijections

$$f_s : \{H_{i,j}(\lambda^{(s)}) \text{ s.t. } (i, j) \in Y(\lambda^{(s)})\} \rightarrow \{(b', g') \in \mathbb{N}_0^2 \mid b' > g', b' \in X_s^{(e)}, g' \notin X_s^{(e)}\}$$

And as before, if $H_{i,j}(\lambda^{(s)}) \mapsto (b', g')$, then $h_{i,j}(\lambda^{(s)}) = b' - g'$. The bijection f that we seek follows from composing F with the inverses of f_0, f_1, \dots, f_{e-1} , noting that

$$\{(b, g) \mid b > g, b \in X, g \notin X, b \equiv g \pmod{e}\} \xrightarrow{1^{-1}} \bigsqcup_{s=0}^{e-1} \{(b', g') \mid b' > g', b' \in X_s^{(e)}, g' \notin X_s^{(e)}\}.$$

Moreover, $b - g = e(b' - g')$ gives $|H| = e|f(H)|$.

To see that $Q_e(\lambda \setminus H) = Q_e(\lambda) \setminus f(H)$ when $H = H_{i,j}(\lambda)$ with $e \mid h_{i,j}(\lambda)$: from Proposition 4.7, we know that Z is a β -set for $\lambda \setminus H$, where

$$Z = (X \setminus \{h_i\}) \sqcup \{h_i - h_{i,j}(\lambda)\} = (X \setminus \{b'e + s\}) \sqcup \{g'e + s\}$$

Note $e \mid |X| = |Z|$, so we can use Z to calculate $Q_e(\lambda \setminus H)$: $Z_t^{(e)} = X_t^{(e)}$ for all $t \in \{0, 1, \dots, e-1\} \setminus \{s\}$, and $Z_s^{(e)} = (X_s^{(e)} \setminus \{b'\}) \sqcup \{g'\}$. So $Z_s^{(e)}$ is a β -set for $\lambda^{(s)} \setminus f(H)$, hence $Q_e(\lambda \setminus H) = (\lambda^{(0)}, \dots, \lambda^{(s-1)}, \lambda^{(s)} \setminus f(H), \lambda^{(s+1)}, \dots, \lambda^{(e-1)}) =: Q_e(\lambda) \setminus f(H)$. \square

Example. Continue the example from before, so let $e = 3$, $\lambda = (6, 5, 2, 1^3) \vdash 16$. Then $X_\lambda = \{11, 9, 5, 3, 2, 1\}$.

hook lengths	A_λ	3-quotient																																																			
<table border="1" style="display: inline-table; vertical-align: middle;"> <tr><td>11</td><td>7</td><td>5</td><td>4</td><td>3</td><td>1</td></tr> <tr><td>9</td><td>5</td><td>3</td><td>2</td><td>1</td><td></td></tr> <tr><td>5</td><td>1</td><td></td><td></td><td></td><td></td></tr> <tr><td>3</td><td></td><td></td><td></td><td></td><td></td></tr> <tr><td>2</td><td></td><td></td><td></td><td></td><td></td></tr> <tr><td>1</td><td></td><td></td><td></td><td></td><td></td></tr> </table>	11	7	5	4	3	1	9	5	3	2	1		5	1					3						2						1						<table border="1" style="display: inline-table; vertical-align: middle;"> <tr><th>0</th><th>1</th><th>2</th></tr> <tr><td>0</td><td>①</td><td>②</td></tr> <tr><td>③</td><td>4</td><td>⑤</td></tr> <tr><td>6</td><td>7</td><td>8</td></tr> <tr><td>⑨</td><td>10</td><td>⑪</td></tr> </table>	0	1	2	0	①	②	③	4	⑤	6	7	8	⑨	10	⑪	$Q_3(\lambda) = ((2, 1), \emptyset, (1))$ $= \left(\begin{array}{ c c } \hline 3 & 1 \\ \hline 1 & \\ \hline \end{array}, \emptyset, \begin{array}{ c } \hline 1 \\ \hline \end{array} \right)$
11	7	5	4	3	1																																																
9	5	3	2	1																																																	
5	1																																																				
3																																																					
2																																																					
1																																																					
0	1	2																																																			
0	①	②																																																			
③	4	⑤																																																			
6	7	8																																																			
⑨	10	⑪																																																			
$h_{1,5}(\lambda) = 3$ row 1	$\textcircled{11} \rightarrow 8$ $h_1 \rightarrow h_1 - h_{1,5}(\lambda)$	$H_{1,2}(\lambda) \xrightarrow{f} H_{1,1}(\lambda^{(2)})$ $11 \equiv 2 \pmod{3}$																																																			
$h_{2,1}(\lambda) = 9$	$\textcircled{9} \rightarrow 0$	$H_{2,1}(\lambda) \xrightarrow{f} H_{1,1}(\lambda^{(0)})$																																																			
$h_{2,3}(\lambda) = 3$	$\textcircled{9} \rightarrow 6$	$H_{2,3}(\lambda) \xrightarrow{f} H_{1,2}(\lambda^{(0)})$																																																			
$h_{4,1}(\lambda) = 3$	$\textcircled{3} \rightarrow 0$	$H_{4,1}(\lambda) \xrightarrow{f} H_{2,1}(\lambda^{(0)})$																																																			

To see that e.g. $H_{2,3}(\lambda) \xrightarrow{f} H_{1,2}(\lambda^{(0)})$ note that the runner 0 of the abacus goes from

$$\begin{array}{c} 0 \\ \hline 0 \\ \textcircled{3} \\ 6 \\ \textcircled{9} \end{array} \quad \text{to} \quad \begin{array}{c} 0 \\ \hline 0 \\ \textcircled{3} \\ \textcircled{6} \\ 9 \end{array}$$

which has partition $(1, 1) = \begin{array}{|c|} \hline \\ \hline \end{array} = \lambda^{(0)} \setminus H_{1,2}(\lambda^{(0)})$ by Lemma 4.3.

Definition. Let $e \in \mathbb{N}$, $\lambda \vdash n$. Then the e -weight of λ is $w_e(\lambda) := |Q_e(\lambda)| := \sum_{i=0}^{e-1} |\lambda^{(i)}|$.

Proposition 4.12. Let $e \in \mathbb{N}$, $\lambda \vdash n$. Then

- (i) $w_e(\lambda)$ is the number of e -hooks we need to remove to get from λ to $C_e(\lambda)$.
- (ii) $|\lambda| = |C_e(\lambda)| + e|Q_e(\lambda)|$.

(iii) $w_e(\lambda)$ is the number of hooks of λ of size divisible by e .

Proof.

(i) Induct on $w_e(\lambda)$. If $w_e(\lambda) = |Q_e(\lambda)| = 0$, then by Theorem 4.11, λ has no e -hooks, and so $\lambda = C_e(\lambda)$. Now suppose $w_e(\lambda) > 0$. Then by the same theorem, λ has a hook length divisible by e . So there also exists a hook H of λ of size exactly e , by Corollary 4.6 or also Theorem 4.11. Recall $Q_e(\lambda \setminus H) = Q_e(\lambda) \setminus f(H)$ and $|f(H)| = 1$, so $w_e(\lambda) = |Q_e(\lambda)| = 1 + |Q_e(\lambda) \setminus f(H)| = 1 + |Q_e(\lambda \setminus H)| = 1 + w_e(\lambda \setminus H)$, so the claim follows from the inductive hypothesis since we removed one e -hook to get from λ to $\lambda \setminus H$ and $C_e(\lambda) = C_e(\lambda \setminus H)$.

(ii) Immediate from (i).

(iii) Follows from Theorem 4.11 as $|Q_e(\lambda)|$ is the number of hooks of $Q_e(\lambda)$. □

Theorem 4.13. *Let $e \in \mathbb{N}$, $n \in \mathbb{N}_0$, and define*

$$B(n) := \left\{ (\gamma; \rho^0, \rho^1, \dots, \rho^{e-1}) \mid \begin{array}{l} \gamma \text{ is an } e\text{-core partition, } \rho^i \text{ is a partition for all } i \\ \text{and } |\gamma| + e \sum_{i=0}^{e-1} |\rho^i| = n \end{array} \right\}.$$

Then

$$\begin{aligned} g : \wp(n) &\longrightarrow B(n), \\ \lambda &\longmapsto (C_e(\lambda); Q_e(\lambda)) \end{aligned}$$

is a bijection. In other words, a partition is uniquely determined by its e -core and e -quotient.

Proof.

- By Proposition 4.12, $n = |\lambda| = |C_e(\lambda)| + e|Q_e(\lambda)|$, so $g(\lambda) \in B(n)$ and g is well-defined.
- g is surjective: Let $(\gamma; \underline{\rho}) \in B(n)$, where $\underline{\rho} = (\rho^0, \rho^1, \dots, \rho^{e-1})$. Let X be a β -set for γ such that $e \mid |X|$ and $|X_i^{(e)}| \geq \ell(\rho^i)$ for all i . Then define Z_i to be the β -set for ρ^i such that $|Z_i| = |X_i^{(e)}|$ for all i , and set $Z := \bigsqcup_{i=0}^{e-1} \{ae + i \mid a \in Z_i\}$. Let $\lambda = \mathcal{P}(Z)$. Since γ is an e -core, $X = X_{(e)}$ and so we have $Z_{(e)} = X_{(e)} = X$. Hence $C_e(\lambda) = \mathcal{P}(Z_{(e)}) = \mathcal{P}(X) = \gamma$. Next, $e \mid |X| = |Z|$, so $Q_e(\lambda) = (\lambda^{(0)}, \dots, \lambda^{(e-1)})$ with $\lambda^{(i)} = \mathcal{P}(Z_i^{(e)}) = \mathcal{P}(Z_i) = \rho^i$. Finally, by Proposition 4.12, $|\lambda| = |C_e(\lambda)| + e|Q_e(\lambda)| = n$ since $(\gamma; \underline{\rho}) \in B(n)$. So $g(\lambda) = (\gamma; \underline{\rho})$ with $\lambda \vdash n$.
- g is injective: notation as above, suppose $g(\mu) = (\gamma; \underline{\rho})$, for some $\mu \vdash n$. Since $C_e(\mu) = \gamma$, there exists a unique β -set W for μ such that $|W| = |X|$. Now $|W_{(e)}| = |W| = |X|$, and $\mathcal{P}(W_{(e)}) = \gamma = \mathcal{P}(X)$. Hence $W_{(e)} = X$ by Lemma 4.2. Also,

$|W_i^{(e)}| = |(W_{(e)})_i^{(e)}| = |X_i^{(e)}| = |Z_i^{(e)}|$, and $\mathcal{P}(W_i^{(e)}) = \rho^i$ since $g(\mu) = (\gamma; \underline{\rho})$ noting that $e \mid |X| = |W|$. But also $\rho^i = \mathcal{P}(Z_i^{(e)})$, hence $W_i^{(e)} = Z_i^{(e)}$ for all i again from Lemma 4.2.

Thus $W_{(e)} = X = Z_{(e)}$ and $W_i^{(e)} = Z_i^{(e)}$ for all i , so $W = Z$, so $\mu = \mathcal{P}(W) = \mathcal{P}(Z) = \lambda$.

□

Example. How do we reconstruct λ , given $C_e(\lambda)$ and $Q_e(\lambda)$? Let $e = 3$ and $(\gamma; \underline{\rho}) = ((3, 1); (2, 1), \emptyset, (1)) \in B(16)$. We expect $\lambda = (6, 5, 2, 1^3) \vdash 16$.

- **Step 1.** Start with A_γ ,

	0	1	2
0	0	①	2
1	3	④	5
2	6	7	8
		⋮	


- **Step 2.** Shift to get $e \mid |X|$,

	0	1	2
0	①	1	②
1	3		⑤
2	6	7	8
		⋮	

- **Step 3.** Add enough full rows of beads, i.e. shift enough by multiples of e , to get $|X_i^{(e)}| \geq \ell(\rho^i)$ for all i ,

	0	1	2
0	①	①	②
⋮		⋮	
#	#	#	#
#	#	#	#
#	#	#	#
#	#	#	#
⋮		⋮	

The 3-cores are




Definition. Let $e \in \mathbb{N}$. An e -tower is an infinite sequence $T = (T_0, T_1, T_2, \dots)$ such that each T_j is a sequence of e^j many partitions, $T_j = (\lambda_0^j, \lambda_1^j, \dots, \lambda_{e^j-1}^j)$.

- The T_j are the layers or rows of T , define $|T_j| := \sum_{i=0}^{e^j-1} |\lambda_i^j|$.
- The depth of T is $\text{depth}(T) = \sup\{k \in \mathbb{N}_0 \mid |T_k| \neq \emptyset\}$. We will call the depth of the empty tower -1 .
- We say T is an e -core tower if $\text{depth}(T) < \infty$ and λ_i^j is an e -core partition for all i, j .

As we saw in the example above, we can visualise e -towers using graphs.

- vertices: λ_i^j ,
- edges: μ, ν are joined if $\mu = \lambda_i^j$ and $\nu = \lambda_{ie+t}^{j+1}$ for some $j \in \mathbb{N}_0, i \in \{0, 1, \dots, e^j - 1\}, t \in \{0, 1, \dots, e - 1\}$.

e.g. for $e = 2$,



These graphs are rooted, ordered, full e -ary trees. When we use graphs to describe e -towers, we always mean trees like this.

Notation. Let $e \in \mathbb{N}$

- $[\bar{e}] := \{0, 1, \dots, e - 1\}$ (residues mod e)
- For each $x \in [\bar{e}]$, write $Q_e(\lambda^{(x)}) = (\lambda^{(x,0)}, \lambda^{(x,1)}, \dots, \lambda^{(x,e-1)})$, instead of $\lambda^{(x)(0)}, \lambda^{(x)(1)}$, etc.
- Similarly, for all $r \in \mathbb{N}$, and for all $\underline{i} = (i_1, i_2, \dots, i_r) \in [\bar{e}]^r$, will write $Q_e(\lambda^{\underline{i}}) = (\lambda^{(i_1, i_2, \dots, i_r, 0)}, \dots, \lambda^{(i_1, \dots, i_r, e-1)})$.

Definition. Let $e \in \mathbb{N}$, $\lambda \vdash n$. The e -quotient tower of λ is the e -tower $T^Q(\lambda)$ with

- $T^Q(\lambda)_0 = (\lambda)$

- $T^Q(\lambda)_1 = Q_e(\lambda) = (\lambda^{(0)}, \lambda^{(1)}, \dots, \lambda^{(e-1)})$.
- For all $j \in \mathbb{N}$, $T^Q(\lambda)_j = (\lambda^{\dot{i}})_{i \in [\bar{e}]^j}$, lexicographically ordered.

Lemma 4.14. Let $e \in \mathbb{N}$, $\lambda \vdash n$, $T^Q(\lambda)$ the e -quotient tower. Suppose $e \geq 2$, then $\text{depth}(T^Q(\lambda)) < \infty$.

Proof. From Proposition 4.12, $|\lambda| = |C_e(\lambda)| + e|Q_e(\lambda)| \geq |Q_e(\lambda)|$ with equality iff $|C_e(\lambda)| = |Q_e(\lambda)| = 0$, since $e \geq 2$. By Theorem 4.13, equality holds iff $\lambda = \emptyset$. Hence $|T^Q(\lambda)_j| > |T^Q(\lambda)_{j+1}|$ unless $T^Q(\lambda_j) = (\emptyset, \dots, \emptyset)$. \square

Remark. $Q_1(\lambda) = (\lambda^{(0)}) = (\lambda)$, so the 1-quotient tower $T^Q(\lambda)$ has all layers equal to (λ) . So its depth is -1 if $\lambda = \emptyset$, and ∞ otherwise.

Definition. Let $e \in \mathbb{N}$, $\lambda \vdash n$. The e -core tower of λ is the e -tower $T^C(\lambda)$ obtain from the e -quotient tower $T^Q(\lambda)$ by replacing every vertex with its e -core. That is, $T^C(\lambda)_j = (C_e(\lambda^{\dot{i}}))_{i \in [\bar{e}]^j}$, lexicographically ordered.

When $e \geq 2$, $\text{depth}(T^C(\lambda)) < \infty$ since $\text{depth}(T^Q(\lambda)) < \infty$. When $e = 1$, $T^C(\lambda)$ is empty, so also $\text{depth}(T^C(\lambda)) < \infty$. So $T^C(\lambda)$ is indeed an e -core tower.

Lemma 4.15. Let $e \in \mathbb{N}$, $\lambda \vdash n$. For $x \in [\bar{e}]$, the subtree of $T^C(\lambda)$ rooted at $C_e(\lambda^{(x)})$ is the e -core tower of $\lambda^{(x)}$, so the $(j+1)$ -th layer of $T^C(\lambda)$ is the concatenation of the j -th layers of $T^C(\lambda^{(0)}), T^C(\lambda^{(1)}), \dots, T^C(\lambda^{(e-1)})$. That is, $T^C(\lambda^{(x)})_j = (C_e(\lambda^{(x;\dot{i})}))_{i \in [\bar{e}]^j}$ and $T^C(\lambda)_{j+1} = (T^C(\lambda^{(0)})_j, T^C(\lambda^{(1)})_j, \dots, T^C(\lambda^{(e-1)})_j)$.

Proof. The subtree of $T^Q(\lambda)$ rooted at $\lambda^{(x)}$ is $T^Q(\lambda^{(x)})$. \square

Theorem 4.16. Let $e \in \mathbb{N}$, $e \geq 2$, let $n \in \mathbb{N}_0$. Define

$$\theta(n) := \{e\text{-core towers } T \text{ such that } \sum_{j=0}^{\infty} |T_j| e^j = n\}.$$

Then

$$\begin{aligned} h : \wp(n) &\longrightarrow \theta(n), \\ \lambda &\longmapsto e\text{-core tower } T^C(\lambda) \end{aligned}$$

is a bijection.

Proof. First, we check $\sum_{j=0}^{\infty} |T^C(\lambda)_j| e^j = n$, by induction on n . The base case $n = 0$ is clear since then $\lambda = \emptyset$. Now suppose $n > 0$. Then $n = |\lambda| = |C_e(\lambda)| + e \sum_{x=0}^{e-1} |\lambda^{(x)}| = |T^C(\lambda)_0| + e \sum_{x=0}^{e-1} \sum_{j=0}^{\infty} |T^C(\lambda^{(x)})_j| e^j$ by the inductive hypothesis, since ≥ 2 means $|\lambda^{(x)}| < |\lambda|$. This is

$$|T^C(\lambda)_0| + \sum_{j=0}^{\infty} \left(\sum_{x=0}^{e-1} |T^C(\lambda_j^{(x)})| \right) e^{j+1} = |T^C(\lambda)_0| + \sum_{j=0}^{\infty} |T^C(\lambda)_{j+1}| e^{j+1} = \sum_{j=0}^{\infty} |T^C(\lambda)_j| e^j.$$


Next, to prove that h is a bijection, we show for all $T \in \theta(n)$ that there exists a unique $\lambda \vdash n$ such that $T^C(\lambda) = T$. For $x \in [\bar{e}]$, let $S(x)$ be the subtree of T rooted at λ_x^1 , where $T = (T_0, T_1, T_2, \dots)$, $T_j = (\lambda_0^j, \lambda_1^j, \dots, \lambda_{e^j-1}^j)$. Since T is an e -core tower, so is $S(x)$. Then since $n_x := \sum_{j=0}^{\infty} |S(x)_j|e^j \leq \sum_{j=0}^{\infty} |T_{j+1}|e^j < |T_0| + \sum_{j=0}^{\infty} |T_{j+1}|e^{j+1} = n$, we can use inductive hypothesis to see that there is a unique $\mu_x \vdash n_x$ such that $T^C(\mu_x) = S(x)$. By Theorem 4.13 there is a unique partition λ such that $C_e(\lambda) = \lambda_0^0$ and $Q_e(\lambda) = (\mu_0, \mu_1, \dots, \mu_{e-1})$. Observe $T^C(\lambda) = T$ since $T^C(\mu_x) = S(x) = T^C(\lambda^{(x)})$, i.e. $T^C(\mu_x)$ is the subtree of $T^C(\lambda)$ rooted at $\lambda_x^1 = C_e(\lambda^{(x)}) = C_e(\mu_x)$. To check $|\lambda| = n$:

$$\begin{aligned} |\lambda| &= |C_e(\lambda)| + e \sum_{x=0}^{e-1} |\mu_x| \\ &= |T^C(\lambda)_0| + e \sum_{x=0}^{e-1} \sum_{j=0}^{\infty} |S(x)_j|e^j \\ &= |\lambda_0^0| + \sum_{j=0}^{\infty} \left(\sum_{x=0}^{e-1} |S(x)_j| \right) e^{j+1} \\ &= \sum_{j=0}^{\infty} |T_j|e^j = n \end{aligned}$$

Uniqueness of λ is also clear from this argument. □

Remark. This is not a bijection when $e = 1$ since then $T^C(\lambda)$ is empty for all λ .


Example. Given $T \in \theta(n)$, how to compute $\lambda = h^{-1}(T)$? Let $e = 3$, $T =$




where $\underline{\emptyset}$ means that from that vertex onwards there are only empty partitions. We have $n = 1 \cdot 3^0 + 2 \cdot 3^1 + 1 \cdot 3^2 = 16$.

- When we see a subtree rooted at an e -core partition γ with all empty below, this subtree is the e -core tower of γ because $C_e(\gamma) = \gamma$, $Q_e(\gamma) = (\emptyset, \dots, \emptyset)$.
- We will draw boxes to replace subtrees by the partition whose e -core tower is that


subtree.



- Work up the layers: What μ has $T^C(\mu) = \emptyset$? It is the partition μ with




$C_3(\mu) = \emptyset$ and $Q_3(\mu) = (\emptyset, (1), \emptyset)$. We showed how to find this in the example after Theorem 4.13. In this case we get $\mu = (2, 1)$. Then we get



So $T = T^C(\lambda)$ where $C_3(\lambda) = (1)$ and $Q_3(\lambda) = ((2, 1), (2), \emptyset)$. We find that $\lambda = (7, 6, 3)$.


Example. How does hook removal interact with core towers? Let $e = 3$, $\lambda = (7, 6, 3)$,



- (a) Remove the 3-hook marked in

x	x	x				

. So let $\mu = \lambda \setminus H$, where $H = H_{3,1}(\lambda)$, $h_{3,1}(\lambda) = 3$. On the abacus:



where $T^C(\lambda)$ is the p -core tower of λ .

Proof. Recall the hook length formula, Theorem 3.1: We get

$$v_p(\chi^\lambda(1)) = v_p\left(\frac{n!}{\prod_{h \in \mathcal{H}(\lambda)} h}\right) = v_p(n!) - v_p\left(\prod_{h \in \mathcal{H}(\lambda)} h\right).$$

- **Step 1.** We compute $v_p(n!)$. Observe that

$$\begin{aligned} v_p(n!) &= \sum_{r=1}^{\infty} \lfloor \frac{n}{p^r} \rfloor = \sum_{r=1}^{\infty} (\alpha_r + \alpha_{r+1}p + \alpha_{r+2}p^2 + \dots) \\ &= \sum_{r=1}^{\infty} \alpha_r (1 + p + p^2 + \dots + p^{r-1}) \\ &= \sum_{r=1}^{\infty} \alpha_r \frac{p^r - 1}{p - 1} \\ &= \frac{1}{p - 1} \left(\sum_{r=1}^{\infty} \alpha_r p^r - \sum_{r=1}^{\infty} \alpha_r \right) \\ &= \frac{1}{p - 1} \left(\sum_{r=0}^{\infty} \alpha_r p^r - \sum_{r=0}^{\infty} \alpha_r \right) \\ &= \frac{n - \sum_{r=0}^{\infty} \alpha_r}{p - 1}. \end{aligned}$$

- **Step 2.** We claim that $v_p(\prod_{h \in \mathcal{H}(\lambda)} h) = \sum_{r=1}^{\infty} |T^Q(\lambda)_r|$, where $T^Q(\lambda)$ is the p -quotient tower of λ . We prove this by induction on n . The base case $n = 0$ is clear since $v_p(1) = 0$. Now suppose $n > 0$. We write $\mathcal{H}(Q_p(\lambda))$ for the multiset of hook lengths of $Q_p(\lambda)$. Then

$$\begin{aligned} v_p\left(\prod_{h \in \mathcal{H}(\lambda)} h\right) &= v_p\left(\prod_{\substack{h \in \mathcal{H}(\lambda) \\ p|h}} hv\right) \\ &\stackrel{\text{Theorem 4.11}}{=} v_p\left(\prod_{h \in \mathcal{H}(Q_p(\lambda))} ph\right) \\ &= |Q_p(\lambda)| + v_p\left(\prod_{h \in \mathcal{H}(Q_p(\lambda))} h\right) \\ &= |Q_p(\lambda)| + v_p\left(\prod_{x=0}^{p-1} \prod_{h \in \mathcal{H}(\lambda^{(x)})} h\right) \\ &= |T^Q(\lambda)_1| + \sum_{x=0}^{p-1} v_p\left(\prod_{h \in \mathcal{H}(\lambda^{(x)})} h\right) \end{aligned}$$

$$\begin{aligned}
& \text{ind. hypothesis} \quad |T^Q(\lambda)_1| + \sum_{x=0}^{p-1} \sum_{r=1}^{\infty} |T^Q(\lambda^{(x)})_r| \\
&= |T^Q(\lambda)_1| + \sum_{r=1}^{\infty} \sum_{x=0}^{p-1} |T^Q(\lambda^{(x)})_r| \\
&= |T^Q(\lambda)_1| + \sum_{r=1}^{\infty} |T^Q(\lambda)_{r+1}| \\
&= \sum_{r=1}^{\infty} |T^Q(\lambda)_r|
\end{aligned}$$

- **Step 3.** By Proposition 4.12 for all $r \in \mathbb{N}_0$, $\underline{i} \in [\bar{p}]^r$,

$$|\lambda^{\underline{i}}| = |C_p(\lambda^{\underline{i}})| + p|Q_p(\lambda^{\underline{i}})|.$$

Summing over $\underline{i} \in [\bar{p}]^r$, we get

$$|T^Q(\lambda)_r| = |T^C(\lambda)_r| + p|T^Q(\lambda)_{r+1}|.$$

Therefore,

$$\begin{aligned}
n = |\lambda| &= |T^Q(\lambda)_0| \\
&= \sum_{r=0}^{\infty} |T^Q(\lambda)_r| - \sum_{r=1}^{\infty} |T^Q(\lambda)_r| \\
&= \left(\sum_{r=0}^{\infty} |T^C(\lambda)_r| - p \sum_{r=0}^{\infty} |T^Q(\lambda)_{r+1}| \right) - \sum_{r=1}^{\infty} |T^Q(\lambda)_r| \\
&= \sum_{r=0}^{\infty} |T^C(\lambda)_r| + (p-1) \sum_{r=1}^{\infty} |T^Q(\lambda)_r|.
\end{aligned}$$

Hence

$$\begin{aligned}
v_p(\chi^\lambda(1)) &= v_p(n!) - v_p\left(\prod_{h \in \mathcal{H}(\lambda)} h\right) \\
&= \frac{1}{p-1} \left(n - \sum_{r=0}^{\infty} \alpha_r \right) - \sum_{r=1}^{\infty} |T^Q(\lambda)_r| \\
&= \frac{1}{p-1} \left(\sum_{r=0}^{\infty} |T^C(\lambda)_r| + (p-1) \sum_{r=1}^{\infty} |T^Q(\lambda)_r| - \sum_{r=0}^{\infty} \alpha_r \right) - \sum_{r=1}^{\infty} |T^Q(\lambda)_r| \\
&= \frac{1}{p-1} \left(\sum_{r=0}^{\infty} |T^C(\lambda)_r| - \sum_{r=0}^{\infty} \alpha_r \right).
\end{aligned}$$

□


Corollary 4.19. *Let p be prime, $n \in \mathbb{N}_0$ with p -adic expansion $n = \sum_{r=0}^{\infty} \alpha_r p^r$. Let $\lambda \vdash n$. Then $v_p(\chi^\lambda(1)) = 0$ iff $|T^C(\lambda)_r| = \alpha_r$ for all $r \in \mathbb{N}_0$, where $T^C(\lambda)$ is the p -core tower of λ .*

Proof. “if” is clear from the theorem. For “only if” the theorem gives us $\sum_{r=0}^{\infty} |T^C(\lambda)_r| = \sum_{r=0}^{\infty} \alpha_r$. Also note that $\sum_{r=0}^{\infty} |T^C(\lambda)_r| p^r = n$. Let $\beta_r = |T^C(\lambda)_r|$. So we have


$$\begin{aligned} \sum_{r \geq 0} \alpha_r &= \sum_{r \geq 0} \beta_r \\ \sum_{r \geq 0} \alpha_r p^r &= \sum_{r \geq 0} \beta_r p^r \end{aligned}$$

We show that $\alpha_r = \beta_r$ for all $r \in \mathbb{N}_0$. First, $\beta_0 \equiv \alpha_0 \pmod{p}$. Hence we can write $\beta_0 = \alpha_0 + m_1 p$, for some $m_1 \in \mathbb{N}_0$. Since $\beta_0 \in \mathbb{N}_0$ and $\alpha_0 \in \{0, 1, \dots, p-1\}$. Thus $\sum_{r \geq 0} \beta_r p^r = \sum_{r \geq 2} \beta_r p^r + (\beta_1 + m_1)p + \alpha_0 = \sum_{r \geq 0} \alpha_r p^r$. Then $\beta_1 + m_1 \equiv \alpha_1 \pmod{p}$, so $\beta_1 + m_1 = \alpha_1 + m_2 p$ for some $m_2 \in \mathbb{N}_0$. Iterating, $\beta_r + m_r = \alpha_r + m_{r+1} p$ for all $r \in \mathbb{N}_0$ where $m_r \in \mathbb{N}_0$ and $m_0 = 0$. Then $\sum_{r \geq 0} \alpha_r = \sum_{r \geq 0} \beta_r = \sum_{r \geq 0} \alpha_r + (p-1) \sum_{r \geq 0} m_r$, hence $m_r = 0$ for all r and so $\alpha_r = \beta_r$. \square

Example. We compute $\text{Irr}_{2'}(S_4)$. By Theorem 4.16 there is a bijection between partitions and 2-core towers. By the corollary, for $\lambda \vdash 4 = 1 \cdot 2^2$, we have $\chi^\lambda \in \text{Irr}_{2'}(S_4)$ iff $|T^C(\lambda)_2| = 1$ and $|T^C(\lambda)_r| = 0$ for all $r \neq 2$. So we already see that $|\text{Irr}_{2'}(S_4)| = 4$. The towers are:



As in the example after Theorem 4.16 we compute:



Hence we see that

$$\text{Irr}_{2'}(S_4) = \{\chi^\lambda \text{ s.t. } \lambda \in \{(4), (3, 1), (2, 1^2), (1^4)\}\}.$$

Note that these partitions are exactly the hooks of size 4.

4.3 The McKay Conjecture

Recall the McKay Conjecture: Let G be a finite group, p a prime, P a Sylow p -subgroup of G . Then

$$|\text{Irr}_{p'}(G)| = |\text{Irr}_{p'}(N_G(P))|.$$

Definition. Let G be a finite group, p a prime. The McKay numbers of G are

$$m_i(p, G) = |\{\chi \in \text{Irr}(G) \text{ s.t. } v_p(\chi(1)) = i\}|,$$

for $i \in \mathbb{N}_0$.

So we are interested in $m_0(p, G)$ for $G = S_n$ (and $G = N_{S_n}(P)$).

Corollary 4.20. Let $n \in \mathbb{N}$ with binary expansion $n = 2^{n_1} + 2^{n_2} + \dots + 2^{n_t}$, i.e. $t \in \mathbb{N}$, and the $n_i \in \mathbb{N}_0$ are distinct. Then

$$m_0(2, S_n) = |\text{Irr}_{2'}(S_n)| = 2^{n_1 + n_2 + \dots + n_t}.$$

Proof. By Theorem 4.16, we have a bijection

$$\begin{aligned} h : \wp(n) &\longrightarrow \theta(n) \\ \lambda &\longmapsto \text{2-core tower } T^C(\lambda) \end{aligned}$$

By Corollary 4.19, for $\lambda \vdash n$ we have $\chi^\lambda \in \text{Irr}_{2'}(S_n)$ iff

$$|T^C(\lambda)_r| = \begin{cases} 1 & \text{if } r \in \{n_1, \dots, n_t\}, \\ 0 & \text{otherwise.} \end{cases}$$

But $|T^C(\lambda)_r| = 1$ means $T^C(\lambda)_r$ is a sequence of 2^r many partitions, exactly one of which is (1), the rest \emptyset . So the number of such 2-core towers is $2^{n_1} \cdot 2^{n_2} \dots 2^{n_t}$. \square

Corollary 4.21. Let p be a prime, $n \in \mathbb{N}$ with p -adic expansion $n = \sum_{r \geq 0} \alpha_r p^r$. Then

$$m_0(p, S_n) = |\text{Irr}_{p'}(S_n)| = \prod_{r \geq 0} k_p(p^r, \alpha_r),$$

where $k_p(l, m)$ is the number of tuples of partitions $(\gamma^1, \dots, \gamma^l)$ such that each γ^i is a p -core partition and $\sum_{i=1}^l |\gamma^i| = m$.

Proof. The same as the previous corollary, use Theorem 4.16 and Corollary 4.19. \square

Sketch towards the McKay conjecture. We need some group theoretic facts.

Suppose $p = 2$.

- Let $P_n \in \text{Syl}_2(S_n)$ be a Sylow 2-subgroup of S_n . Then $N_{S_n}(P_n) = P_n$.
- For $n = 2^k$, $\text{Irr}_{2'}(N_{S_n}(P_n)) = \text{Irr}_{2'}(P_n) = \{\text{degree 1 characters of } P_n\}$ as the degree of any irreducible character divides the group order. But now the degree 1 characters of any group H are in bijection with $\text{Irr}(H/H')$ where H' is the commutator subgroup. If $H = P_{2^k}$, then $H/H' \cong C_2^{\times k}$, hence $|\text{Irr}_{2'}(P_{2^k})| = |\text{Irr}(C_2^{\times k})| = 2^k$.
- For general $n = 2^{n_1} + 2^{n_2} + \dots + 2^{n_t}$, count the number of factors of $p = 2$ in $|S_n| = n!$ to see that

$$P_n \cong P_{2^{n_1}} \times P_{2^{n_2}} \times \dots \times P_{2^{n_t}}.$$

Then

$$|\text{Irr}_{2'}(N_{S_n}(P_n))| = |\text{Irr}_{2'}(P_n)| = \prod_{i=1}^s |\text{Irr}_{2'}(P_{2^{n_i}})| = \prod_{i=1}^b 2^{n_i} = m_0(2, S_n).$$

For $p > 2$ the first point need no longer be true.

Bibliography

- [CR62] Charles W Curtis and I. Reiner. *Representation theory of finite groups and associative algebras*. 1962.
- [Isa76] I. Martin Isaacs. *Character theory of finite groups*. Pure and applied mathematics (Academic Press) ; 69. New York: Academic Press, 1976.
- [Jam78] G. D James. *The Representation Theory of the Symmetric Groups*. Vol. 682. Lecture Notes in Mathematics. Berlin, Heidelberg: Springer Berlin / Heidelberg, 1978.
- [JK84] G James and A Kerber. *The representation theory of the symmetric group*. Encyclopedia of mathematics and its applications volume 16. 1984.
- [Ols94] J. Olsson. *Combinatorics and Representations of Finite Groups*. Vorlesungen aus dem Fachbereich Mathematik der Universität GH Essen, Heft 20. 1994.