

1. AFFINE MORPHISMS

1.1.  $S$ -preschemes and  $\mathcal{O}_S$ -algebras.

(1.1.1). Given an  $S$ -prescheme  $f: X \rightarrow S$ ,  $\mathcal{A}(X)$  denotes the sheaf of  $\mathcal{O}_S$  algebras  $f_*\mathcal{O}_X$ . Given a sheaf of  $\mathcal{O}_X$  modules (or  $\mathcal{O}_X$  algebras)  $\mathcal{F}$ ,  $\mathcal{A}(\mathcal{F})$  denotes the sheaf of  $\mathcal{A}(X)$ -modules (or  $\mathcal{A}(X)$  algebras)  $f_*(\mathcal{F})$ .

(1.1.2–3).  $X \mapsto \mathcal{A}(X)$  is a contravariant functor from  $S$ -preschemes to sheaves of  $\mathcal{O}_S$  algebras. More generally, there is a contravariant functor  $(X, \mathcal{F}) \mapsto (\mathcal{A}(X), \mathcal{A}(\mathcal{F}))$  from pairs consisting of an  $S$ -prescheme  $X$  and sheaf of  $\mathcal{O}_X$  modules  $\mathcal{F}$  to pairs consisting of a sheaf of  $\mathcal{O}_S$  algebras and a sheaf of modules over it.

1.2. Preschemes affine over a prescheme.

*Definition* (1.2.1). — An  $S$ -prescheme  $f: X \rightarrow S$  is *affine over  $S$*  if  $S$  has an affine open covering  $(S_\alpha)$  such that each  $f^{-1}(S_\alpha)$  is affine.

*Example* (1.2.2). — By (I, 4.2.3-4) any closed sub-prescheme of  $S$  is affine over  $S$ .

*Remark* (1.2.3). — A prescheme affine over  $S$  need not be affine, e.g.,  $X = S$ . An affine scheme  $X$  that is a prescheme over  $S$  need not be affine over  $S$  (see (1.3.3)), but if  $S$  is a *scheme* [i.e., a separated prescheme] then any  $S$ -prescheme which is an affine scheme is affine over  $S$  (I, 5.5.10).

*Proposition* (1.2.4). — *Every prescheme affine over  $S$  is separated over  $S$ , i.e., it is a scheme over  $S$ .*

*Proposition* (1.2.5). — *If  $f: X \rightarrow S$  is affine, then for every open  $U \subseteq S$ ,  $f^{-1}(U)$  is affine over  $U$ .*

*Proposition* (1.2.6). — *If  $f: X \rightarrow S$  is affine, then for every quasi-coherent sheaf of  $\mathcal{O}_X$  modules  $\mathcal{F}$ ,  $f_*(\mathcal{F})$  is quasi coherent.*

In particular,  $\mathcal{A}(X)$  is a quasi-coherent sheaf of  $\mathcal{O}_S$  algebras.

*Proposition* (1.2.7). — *Let  $X$  be affine over  $S$ . For every  $S$ -prescheme  $Y$ , the canonical map  $\text{Hom}_S(Y, X) \rightarrow \text{Hom}_{\mathcal{O}_S\text{-Alg}}(\mathcal{A}(X), \mathcal{A}(Y))$  is bijective.*

*Corollary* (1.2.8). — *If  $X$  and  $Y$  are affine over  $S$ , then an  $S$ -morphism  $h: X \rightarrow Y$  is an isomorphism iff it induces an isomorphism  $\mathcal{A}(X) \cong \mathcal{A}(Y)$ .*

1.3. Prescheme affine over  $S$  associated to an  $\mathcal{O}_S$  algebra.

*Proposition* (1.3.1). — *Given any quasi-coherent sheaf of  $\mathcal{O}_S$  algebra  $\mathcal{B}$ , there exists a prescheme  $X$  affine over  $S$ , unique up to canonical isomorphism, such that  $\mathcal{A}(X) = \mathcal{B}$ .*

The prescheme  $X$  in the proposition is denoted  $\text{Spec}(\mathcal{B})$ .

*Corollary* (1.3.2). — *Let  $f: X \rightarrow S$  be affine. For every affine  $U \subseteq S$ ,  $f^{-1}(U)$  is an affine scheme  $\text{Spec}(\Gamma(U, \mathcal{A}(X)))$ .*

*Example (1.3.3).* — Let  $K$  be a field,  $S$  the affine plane with the origin doubled, so  $S = Y_1 \cup Y_2$ , where each  $Y_i \cong \mathbb{A}_K^2$ . Let  $f$  be the open immersion  $Y_1 \hookrightarrow S$ . Then  $f^{-1}(Y_2)$  is not affine, so  $Y_1$  is not affine over  $S$ , even though  $Y_1$  is an affine scheme.

*Corollary (1.3.4).* — *Let  $S$  be an affine scheme. Then an  $S$ -prescheme  $X$  is affine over  $S$  iff  $X$  is an affine scheme.*

*Corollary (1.3.5).* — *Let  $X$  be affine over  $S$  and let  $Y$  be an  $X$ -prescheme. Then  $Y$  is affine over  $X$  iff  $Y$  is affine over  $S$ .*

(1.3.6). Let  $X$  be affine over  $S$ . To give an  $S$ -prescheme  $Y$  affine over  $X$ , it is equivalent to give a quasi-coherent sheaf of  $\mathcal{O}_S$  algebras  $\mathcal{B}$  and a homomorphism  $\mathcal{A}(X) \rightarrow \mathcal{B}$ ; that is, to give a quasi-coherent sheaf of  $\mathcal{A}(X)$  algebra on  $S$ .

*Corollary (1.3.7).* — *Let  $X$  be affine over  $S$ . Then  $X$  is of finite type over  $S$  iff  $\mathcal{A}(X)$  is of finite type as a sheaf of  $\mathcal{O}_S$  algebras (I, 9.6.2).*

*Corollary (1.3.8).* — *A prescheme  $X$  affine over  $S$  is reduced iff  $\mathcal{A}(X)$  is reduced (0, 4.1.4).*

#### 1.4. Quasi-coherent sheaves on a prescheme affine over $S$ .

*Proposition (1.4.1).* — *Let  $X$  be affine over  $S$ ,  $Y$  any  $S$ -prescheme,  $\mathcal{F}, \mathcal{G}$  quasi-coherent sheaves of  $\mathcal{O}_X, \mathcal{O}_Y$  modules. The functorial correspondence from morphisms  $(h, u): (Y, \mathcal{G}) \rightarrow (X, \mathcal{F})$  to di-homomorphisms  $(\mathcal{A}(h), \mathcal{A}(u)): (\mathcal{A}(X), \mathcal{A}(\mathcal{F})) \rightarrow (\mathcal{A}(Y), \mathcal{A}(\mathcal{G}))$  is bijective.*

*Corollary (1.4.2).* — *In (1.4.1), suppose  $Y$  is also affine over  $S$ . Then  $(h, u)$  is an isomorphism iff  $(\mathcal{A}(h), \mathcal{A}(u))$  is an isomorphism.*

*Proposition (1.4.3).* — *Given quasi-coherent sheaves of  $\mathcal{O}_X$  algebras  $\mathcal{B}$  and  $\mathcal{B}$  modules  $\mathcal{M}$ , there exists a prescheme  $X$  affine over  $S$  and a quasi-coherent sheaf  $\mathcal{F}$  of  $\mathcal{O}_X$  modules, unique up to canonical isomorphism, such that  $(\mathcal{A}(X), \mathcal{A}(\mathcal{F})) \cong (\mathcal{B}, \mathcal{M})$ .*

The sheaf  $\mathcal{F}$  in the proposition is denoted  $\widetilde{\mathcal{M}}$ .

*Corollary (1.4.4).* —  $\mathcal{M} \mapsto \widetilde{\mathcal{M}}$  is a covariant exact functor, which commutes with direct limits and direct sums.

*Corollary (1.4.5).* — *Under the hypotheses of (1.4.3),  $\widetilde{\mathcal{M}}$  is an  $\mathcal{O}_X$  module of finite type iff  $\mathcal{M}$  is a  $\mathcal{B}$  module of finite type.*

*Proposition (1.4.6).* — *Let  $Y$  be affine over  $S$  and  $X, X'$  affine over  $Y$  (hence over  $S$  (1.3.5)). Then  $X \times_Y X' = \text{Spec}(\mathcal{A}(X) \otimes_{\mathcal{A}(Y)} \mathcal{A}(X'))$  is affine over  $Y$  (and over  $S$ ).*

*Corollary (1.4.7).* — *If  $\mathcal{F}, \mathcal{F}'$  are quasi-coherent sheaves of  $\mathcal{O}_X, \mathcal{O}_{X'}$  modules, then  $\mathcal{A}(\mathcal{F} \otimes_Y \mathcal{F}') \cong \mathcal{A}(\mathcal{F}) \otimes_{\mathcal{A}(Y)} \mathcal{A}(\mathcal{F}')$ .*

(1.4.8). In particular, taking  $X = X' = Y$  affine over  $S$ , if  $\mathcal{F}, \mathcal{G}$  are quasi-coherent sheaves of  $\mathcal{O}_X$  modules, then

$$(1.4.8.1) \quad \mathcal{A}(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}) = \mathcal{A}(\mathcal{F}) \otimes_{\mathcal{A}(X)} \mathcal{A}(\mathcal{G}).$$

If  $\mathcal{F}$  is finitely presented, then (I, 1.6.3 and 1.3.12) imply

$$(1.4.8.2) \quad \mathcal{A}(\mathcal{H}om(\mathcal{F}, \mathcal{G})) = \mathcal{H}om_{\mathcal{A}(X)}(\mathcal{A}(\mathcal{F}), \mathcal{A}(\mathcal{G})),$$

up to canonical isomorphism.

*Remark (1.4.9).* — If  $X, X'$  are affine over  $S$ , then so is  $X \amalg X'$ .

*Proposition (1.4.10).* — Let  $\mathcal{B}$  be a quasi-coherent sheaf of  $\mathcal{O}_S$  algebras,  $X = \text{Spec}(\mathcal{B})$ . If  $\mathcal{I} \subseteq \mathcal{B}$  is a quasi-coherent sheaf of ideals, then  $\widetilde{\mathcal{I}}$  is a quasi-coherent sheaf of ideals in  $\mathcal{O}_X$ , and the closed subscheme  $Y \subseteq X$  which it defines is canonically isomorphic to  $\text{Spec}(\mathcal{B}/\mathcal{I})$ .

Put another way, if  $h: \mathcal{B} \rightarrow \mathcal{B}'$  is a surjective homomorphism of quasi-coherent sheaves of  $\mathcal{O}_S$  algebras, then the induced morphism  $\text{Spec}(\mathcal{B}') \rightarrow \text{Spec}(\mathcal{B})$  is a closed immersion.

*Proposition (1.4.11).* — Let  $\mathcal{B}$  be a quasi-coherent sheaf of  $\mathcal{O}_S$  algebras,  $X = \text{Spec}(\mathcal{B})$ ,  $f: X \rightarrow S$  the structure morphism. If  $\mathcal{J} \subseteq \mathcal{O}_S$  is a quasi-coherent sheaf of ideals, then  $f^*(\mathcal{J})\mathcal{O}_X \cong (\mathcal{J}\mathcal{B})^\sim$ , canonically.

## 1.5. Change of base prescheme.

*Proposition (1.5.1).* — If  $X$  is affine over  $S$ , then any base change  $X_{(S')}$  is affine over  $S'$ .

*Corollary (1.5.2).* — Let  $f: X \rightarrow S$  be affine,  $g: S' \rightarrow S$  any  $S$ -prescheme,  $X' = X_{(S')}$ ,  $f': X' \rightarrow S'$ ,  $g': X' \rightarrow X$  the projections (note  $g \circ f' = f \circ g'$ ). For every quasi-coherent  $\mathcal{O}_X$ -module, there is a canonical isomorphism

$$(1.5.2.1) \quad u: g^*(f_*(\mathcal{F})) \cong f'_*(g'^*(\mathcal{F})).$$

In particular,  $\mathcal{A}(X') \cong g^*(\mathcal{A}(X))$ .

*Remark (1.5.3).* — Although (1.5.2) fails if  $X$  is not affine over  $S$ , a weaker version is valid for coherent sheaves on  $X$  when  $f$  is proper and  $S$  is Noetherian (III, 4.2.4).

*Corollary (1.5.4).* — For  $f: X \rightarrow S$  affine and  $s \in S$ , the fiber  $f^{-1}(s)$  is an affine scheme.

*Corollary (1.5.5).* — If  $X$  is an  $S$ -prescheme via  $f: X \rightarrow S$ , and  $S'$  is affine over  $S$ , then  $X' = X_{(S')}$  is affine over  $X$ . Moreover  $\mathcal{A}(X') \cong f^*(\mathcal{A}(S'))$  and for every quasi-coherent  $\mathcal{A}(S')$ -module  $\mathcal{M}$ ,  $f^*(\mathcal{M}) \cong \mathcal{A}(f'^*(\widetilde{\mathcal{M}}))$ , where  $f' = f_{(S')}$ .

(1.5.6). Let  $q: S' \rightarrow S$  be a morphism,  $\mathcal{B}, \mathcal{B}'$  quasi-coherent sheaves of  $\mathcal{O}_S, \mathcal{O}_{S'}$  algebras,  $u: \mathcal{B} \rightarrow \mathcal{B}'$  a  $q$ -morphism (i.e. an  $\mathcal{O}_S$  algebra homomorphism  $\mathcal{B} \rightarrow q_*(\mathcal{B}')$ ). Then  $u$  induces a morphism

$$v = \text{Spec}(u): X' = \text{Spec}(\mathcal{B}') \rightarrow \text{Spec}(\mathcal{B}) = X,$$

such that the following diagram commutes

$$(1.5.6.1) \quad \begin{array}{ccc} X' & \xrightarrow{v} & X \\ \downarrow & & \downarrow \\ S' & \xrightarrow{q} & S \end{array}.$$

(1.5.7). Moreover, if  $\mathcal{M}$  is a quasi-coherent  $\mathcal{B}$ -module, then

$$(1.5.7.1) \quad v^*(\widetilde{\mathcal{M}}) \cong (q^*(\mathcal{M}) \otimes_{q^*(\mathcal{B})} \mathcal{B}')^\sim.$$

## 1.6. Affine morphisms.

(1.6.1). A morphism  $f: X \rightarrow Y$  is *affine* if it makes  $X$  affine over  $Y$ .

*Proposition (1.6.2).* — (i) A closed immersion is affine.

(ii) The composite of affine morphisms is affine.

(iii) If  $f$  is affine, so is any base change  $f_{(S')}$ .

(iv) If  $f, g$  are affine, so is  $f \times_S g$ .

(v) If  $g \circ f$  is affine and  $g$  is separated, then  $f$  is affine.

(vi) If  $f$  is affine, then  $f_{\text{red}}$  is affine.

*Corollary (1.6.3).* — If  $X$  is an affine scheme and  $Y$  is a [separated] scheme, then any morphism  $X \rightarrow Y$  is affine.

*Proposition (1.6.4).* — Let  $Y$  be locally Noetherian and  $f: X \rightarrow Y$  a morphism of finite type. Then  $f$  is affine iff  $f_{\text{red}}$  is affine.

## 1.7. Vector bundle associated to a sheaf of modules.

(1.7.1). The *symmetric algebra*  $\mathbf{S}(E)$  of an  $A$ -module  $E$  is the quotient of the tensor algebra  $\mathbf{T}(E)$  by the relations  $x \otimes y - y \otimes x$  for  $x, y \in E$ . It has the universal property that any  $A$ -linear map  $E \rightarrow B$ , where  $B$  is a commutative  $A$ -algebra, factors uniquely as  $E \rightarrow \mathbf{S}(E) \rightarrow B$ .  $\mathbf{S}(-)$  is a functor from  $A$ -modules to commutative  $A$ -algebras; it commutes with direct limits and has  $\mathbf{S}(E \oplus F) = \mathbf{S}(E) \otimes_A \mathbf{S}(F)$ .  $\mathbf{S}(E)$  is graded, with  $\mathbf{S}_n(E)$  [the  $n$ -th symmetric power of  $E$ ] the  $A$ -linear span of products of  $n$  elements of  $E$ . We have  $\mathbf{S}(A^m) \cong A[t_1, \dots, t_m]$ .

(1.7.2). Let  $\phi: A \rightarrow B$  be a ring homomorphism,  $F$  a  $B$ -module.  $F_{[\phi]}$  denotes  $F$  regarded as an  $A$ -module. The inclusion  $F_{[\phi]} \rightarrow \mathbf{S}(F)_{[\phi]}$  and the universal property induce a canonical  $A$ -algebra homomorphism  $\mathbf{S}(F_{[\phi]}) \rightarrow \mathbf{S}(F)_{[\phi]}$ . Any  $A$ -module homomorphism  $E \rightarrow F_{[\phi]}$  induces  $\mathbf{S}(E) \rightarrow \mathbf{S}(F)_{[\phi]}$ . We also have  $\mathbf{S}(E \otimes_A B) = \mathbf{S}(E) \otimes_A B$ .

(1.7.3). Let  $R \subseteq A$  be a multiplicative set, and  $B = R^{-1}A$ . Then  $\mathbf{S}(R^{-1}E) = R^{-1}\mathbf{S}(E)$ , and if  $R \subseteq R'$ , then  $R^{-1}E \rightarrow R'^{-1}E$  commutes with  $\mathbf{S}(R^{-1}E) \rightarrow \mathbf{S}(R'^{-1}E)$ .

(1.7.4). Given a ringed space  $(S, \mathcal{A})$  and an  $\mathcal{A}$ -module  $\mathcal{E}$ , we have a presheaf of  $\mathcal{A}$ -algebras  $U \mapsto \mathbf{S}(\mathcal{E}(U))$ . Its associated sheaf is the *symmetric algebra of  $\mathcal{E}$* , denoted  $\mathbf{S}(\mathcal{E})$  or  $\mathbf{S}_{\mathcal{A}}(\mathcal{E})$ . It is functorial and has the corresponding universal property as for the symmetric algebra of a module.

We have  $\mathbf{S}(\mathcal{E})_s = \mathbf{S}(\mathcal{E}_s)$  (because  $\mathbf{S}$  commutes with direct limits) and  $\mathbf{S}(\mathcal{E} \oplus \mathcal{F}) = \mathbf{S}(\mathcal{E}) \otimes_{\mathcal{A}} \mathbf{S}(\mathcal{F})$ .  $\mathbf{S}(\mathcal{E})$  is graded, and  $\mathbf{S}(\mathcal{A}) = \mathcal{A}[t] = \mathcal{A} \otimes_{\mathbb{Z}} \mathbb{Z}[t]$  (regarding  $\mathbb{Z}, \mathbb{Z}[t]$  as constant sheaves on  $S$ ).

(1.7.5). Given a morphism of ringed spaces  $f: (S, \mathcal{A}) \rightarrow (T, \mathcal{B})$  and a  $\mathcal{B}$ -module  $\mathcal{F}$ , we have  $\mathbf{S}(f^*\mathcal{F}) \cong f^*\mathbf{S}(\mathcal{F})$ , canonically.

*Proposition (1.7.6).* — Let  $S = \text{Spec}(A)$ ,  $\mathcal{E} = \widetilde{M}$ . Then  $\mathbf{S}(\mathcal{E}) = \mathbf{S}(M)^\sim$ .

*Corollary (1.7.7).* — *If  $\mathcal{E}$  is a quasi-coherent sheaf of  $\mathcal{O}_S$  modules on a prescheme  $S$ , then  $\mathbf{S}(\mathcal{E})$  is a quasi-coherent sheaf of  $\mathcal{O}_S$  algebras. If  $\mathcal{E}$  is of finite type, then each  $\mathbf{S}_n(\mathcal{E})$  is of finite type.*

*Definition (1.7.8).* —  $\mathbf{V}(\mathcal{E}) = \text{Spec}(\mathbf{S}(\mathcal{E}))$  is the vector bundle over  $S$  associated to the quasi-coherent sheaf  $\mathcal{E}$ .

[It is more conventional to use the term ‘vector bundle’ only in the special case when  $\mathcal{E}$  is locally free of finite rank.]

Note that  $S$ -morphisms  $X \rightarrow \mathbf{V}(\mathcal{E})$  correspond bijectively to  $\mathcal{O}_S$ -algebra homomorphisms  $\mathbf{S}(\mathcal{E}) \rightarrow \mathcal{A}(X)$ , and in turn to  $\mathcal{O}_S$ -module homomorphisms  $\mathcal{E} \rightarrow \mathcal{A}(X)$  [that is, the  $S$ -prescheme  $\mathbf{V}(\mathcal{E})$  represents the functor  $X \rightarrow \text{Hom}_{\mathcal{O}_S}(\mathcal{E}, \mathcal{A}(X))$  from  $S$ -preschemes to sets].

(1.7.9). Taking  $X$  above to be an open subscheme  $U \subseteq S$ , we see that the sheaf  $U \mapsto \text{Hom}_S(U, \mathbf{V}(\mathcal{E}))$  of sections of the  $S$ -scheme  $\mathbf{V}(\mathcal{E})$  is canonically identified with the dual  $\mathcal{E}^\vee = \mathcal{H}om(\mathcal{E}, \mathcal{O}_S)$  of  $\mathcal{E}$ . In particular, there is a canonical global  $S$ -section  $S \rightarrow \mathbf{V}(\mathcal{E})$ , the zero section.

(1.7.10). Now let  $K$  be a field and take  $X = \text{Spec}(K) = \{\xi\}$ , with  $f: X \rightarrow S$  corresponding to a field extension  $k(s) \rightarrow K$  for  $s \in S$ , so the  $S$ -morphisms  $\{\xi\} \rightarrow \mathbf{V}(\mathcal{E})$  are the geometric points of  $\mathbf{V}(\mathcal{E})$  with values in the extension  $K$  of  $k(s)$ . They are identified with  $\mathcal{O}_S$ -module homomorphisms  $\mathcal{E} \rightarrow f_*(\mathcal{O}_X)$ , or equivalently with  $\mathcal{O}_X$ -module (i.e.,  $K$ -vector space) homomorphisms  $f^*(\mathcal{E}) \rightarrow K$  (0, 4.4.3). By definition,  $f^*(\mathcal{E}) = \mathcal{E}_s \otimes_{\mathcal{O}_s} K = \mathcal{E}^s \otimes_{k(s)} K$ , where we put  $\mathcal{E}^s = \mathcal{E}_s / \mathfrak{m}_s \mathcal{E}_s$ . So the geometric fiber of  $\mathbf{V}(\mathcal{E})$  rational over  $K$  at the point  $s$  is identified with the dual to the  $K$ -vector space  $\mathcal{E}^s \otimes_{k(s)} K$ , or equivalently with  $(\mathcal{E}^s)^\vee \otimes_{k(s)} K$ , where  $(\mathcal{E}^s)$  is the dual of the  $k(s)$ -vector space  $\mathcal{E}^s$ .

*Proposition (1.7.11).* — (i)  $\mathbf{V}(-)$  is a contravariant functor from quasi-coherent sheaves of  $\mathcal{O}_S$  modules to affine  $S$ -schemes.

(ii) If  $\mathcal{E}$  is of finite type, then  $\mathbf{V}(\mathcal{E})$  is a scheme of finite type over  $S$ .

(iii)  $\mathbf{V}(\mathcal{E} \oplus \mathcal{F}) = \mathbf{V}(\mathcal{E}) \times_S \mathbf{V}(\mathcal{F})$ .

(iv) For any  $g: S' \rightarrow S$ ,  $\mathbf{V}(g^*(\mathcal{E})) \cong \mathbf{V}(\mathcal{E})_{(S')} = \mathbf{V}(\mathcal{E}) \times_S S'$ .

(v) If  $\mathcal{E} \rightarrow \mathcal{F}$  is surjective, then  $\mathbf{V}(\mathcal{F}) \rightarrow \mathbf{V}(\mathcal{E})$  is a closed immersion.

(1.7.12). Taking  $\mathcal{E} = \mathcal{O}_S$ , we have  $\mathbf{S}(\mathcal{E}) = \mathcal{O}_S[t]$ , and  $\mathbf{V}(\mathcal{E}) = S \times_{\mathbb{Z}} \text{Spec}(\mathbb{Z}[t])$ . We denote it  $S[t]$  [or, more standardly these days,  $\mathbb{A}_S^1$ ]. The sheaf of  $S$ -sections of  $S[t]$  is identified with  $\mathcal{O}_S$ , by (1.7.9).

(1.7.13). For any  $S$ -prescheme  $X$ , we have  $\text{Hom}_S(X, S[t]) \cong \Gamma(S, \mathcal{A}(X))$ , which is a ring. So the functor  $S[t]$  from  $S$ -preschemes to sets factors through commutative rings. Similarly,  $\text{Hom}_S(X, \mathbf{V}(\mathcal{E}))$  is a module over  $S[t](X)$ . This can be interpreted as saying that  $S[t]$  is a commutative ring scheme over  $S$ , and  $\mathbf{V}(\mathcal{E})$  is an  $S[t]$ -module scheme over  $S$ .

(1.7.14). From the structure of  $S[t]$ -module scheme on  $\mathbf{V}(\mathcal{E})$ , we can recover  $\mathcal{E}$ , up to canonical isomorphism. First, we recover  $\mathbf{S}(\mathcal{E}) = \mathcal{A}(\mathbf{V}(\mathcal{E}))$ . For any  $S$ -prescheme  $X$ , the  $S[t]$ -module scheme structure on  $\mathbf{V}(\mathcal{E})$  identifies the the set of  $\mathcal{O}_S$  algebra homomorphisms  $\text{Hom}_{\mathcal{O}_S\text{-Alg}}(\mathbf{S}(\mathcal{E}), \mathcal{A}(X))$  with  $\mathcal{O}_S$  module homomorphisms  $\text{Hom}_{\mathcal{O}_S}(\mathcal{E}, \mathcal{A}(X))$ . In particular,

this set is naturally an  $\mathcal{A}(X)$ -module. Now  $\mathcal{E}$  is canonically identified with the sub- $\mathcal{O}_S$ -module of  $\mathbf{S}(\mathcal{E})$  whose sections  $z$  on an open set  $U$  have the following property: for every  $S$ -prescheme  $X$ , the evaluation map  $h \rightarrow h(z)$  from  $\mathrm{Hom}_{(\mathcal{O}_S|U)\text{-Alg}}(\mathbf{S}(\mathcal{E})|U, \mathcal{A}(X)|U)$  to  $\Gamma(U, \mathcal{A}(X))$  is a homomorphism of  $\Gamma(U, \mathcal{A}(X))$ -modules.

*Proposition (1.7.15). — Let  $Y$  be a quasi-compact scheme, or a prescheme whose underlying space is Noetherian. Every prescheme  $X$  affine and of finite type over  $Y$  is  $Y$ -isomorphic to a closed sub- $Y$ -scheme of a  $Y$ -scheme of the form  $\mathbf{V}(\mathcal{E})$ , where  $\mathcal{E}$  is a quasi-coherent  $\mathcal{O}_Y$ -module of finite type.*