

Practice Final, MATH 110, Linear Algebra, Fall 2013

Name (Last, First): _____

Student ID: _____

Circle your GSI and section:

| | | | | | |
|----------|------|-----------------|----------|------|-----------------|
| Scerbo | 8am | 200 Wheeler | Forman | 2pm | 3109 Etcheverry |
| Scerbo | 9am | 3109 Etcheverry | Forman | 4pm | 3105 Etcheverry |
| McIvor | 12pm | 3107 Etcheverry | Melvin | 5pm | 24 Wheeler |
| McIvor | 11am | 3102 Etcheverry | Melvin | 4pm | 151 Barrows |
| Mannisto | 12pm | 3 Evans | Mannisto | 11am | 3113 Etcheverry |
| Wayman | 1pm | 179 Stanley | McIvor | 2pm | 179 Stanley |
| Wayman | 2pm | 81 Evans | | | |

If none of the above, please explain: _____

This exam consists of 10 problems, each worth 10 points, of which you must complete 8. **Choose two problems not to be graded by crossing them out in the box below.** You must justify every one of your answers unless otherwise directed.

| Problem | Maximum Score | Your Score |
|----------------|---------------|------------|
| 1 | 10 | |
| 2 | 10 | |
| 3 | 10 | |
| 4 | 10 | |
| 5 | 10 | |
| 6 | 10 | |
| 7 | 10 | |
| 8 | 10 | |
| 9 | 10 | |
| 10 | 10 | |
| Total Possible | 80 | |

Name (Last, First): _____

1. Let V be a nonzero finite-dimensional real vector space. Suppose $T : V \rightarrow V$ is a linear transformation.

Decide if the following assertions are ALWAYS TRUE or SOMETIMES FALSE. You need not justify your answer.

- i. There exists an eigenvalue of T .
- ii. There exists a basis of V such that T is upper-triangular.
- iii. $\dim V = \dim \text{null}(T) + \dim \text{range}(T)$
- iv. If v and w are colinear, then Tv and Tw are colinear.
- v. If v and w are linearly independent, then Tv and Tw are linearly independent.
- vi. If T is invertible and λ is an eigenvalue of T , then λ^{-1} is an eigenvalue of T^{-1} .
- vii. If T is invertible and v is an eigenvector of T , then v is an eigenvector of T^{-1} .
- viii. If $T^2 = 1$, then T has an eigenvalue.
- ix. If $T^3 = T^2$, then T has an eigenvalue.
- x. If $T^3 = T^2$, then $\text{null}(T) \neq \{0\}$.

Name (Last, First): _____

2. Let V be an inner product space and v_1, \dots, v_n a list of vectors in V .

(a) State what it means for v_1, \dots, v_n to be linearly independent. State what it means for v_1, \dots, v_n to be orthonormal.

(b) Prove that if v_1, \dots, v_n is orthonormal, then v_1, \dots, v_n is linearly independent.

Name (Last, First): _____

3. Let $A \in M_{n \times n}(\mathbb{C})$ be a complex matrix. Consider the subspace $W \subset M_{n \times n}(\mathbb{C})$ given by

$$W = \text{span}\{I, A, A^2, A^3, \dots, A^k, \dots\}$$

Prove that

$$\dim W \leq n.$$

Name (Last, First): _____

4. Consider \mathbb{C}^3 with the standard Euclidean inner product. Determine whether each of the following operators $T : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ is self-adjoint, normal, or neither. You need not justify your answer.

a. T has eigenvectors $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$ with respective eigenvalues $0, 1 + i, 1 - i$.

b. T has eigenvectors $(1, i, 0)$, $(1, -i, 0)$, $(0, 0, 1)$ with respective eigenvalues $1, -1, 0$.

c. T has eigenvectors $(1, 0, 0)$, $(0, i, -i)$, $(1, 1, 1)$ with respective eigenvalues $1, -1, 1$.

d. $\dim \text{null}(T^2) = 3$, $\dim \text{range}(T) = 1$.

e. $\dim \text{null}(T - i) = 2$, $\dim \text{null}(T) = 1$ with $\text{null}(T - i) \perp \text{null}(T)$.

Name (Last, First): _____

5. Find a basis for \mathbb{C}^3 that puts the operator given by the matrix

$$T = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}$$

into Jordan canonical form. What is the Jordan canonical form?

Name (Last, First): _____

6. Consider \mathbb{R}^2 with the inner product

$$\langle (x_1, x_2), (y_1, y_2) \rangle = 2x_1y_1 + x_1y_2 + x_2y_1 + x_2y_2$$

a. Find an orthonormal basis for \mathbb{R}^2 with respect to the above inner product.

b. Find the vector $v = (a, b)$ closest to $(1, 0)$ satisfying $a + b = 0$.

Name (Last, First): _____

7. Find the Jordan form of an operator $T : \mathbb{C}^5 \rightarrow \mathbb{C}^5$ given the following information:

$$\dim \text{null}(T^2) = 2 \quad \dim \text{null}(T^3) = 3 \quad \dim \text{null}((T-1)^2) = 2 \quad \dim \text{range}(T-1) = 4$$

Be sure to justify your answer.

Name (Last, First): _____

8. Consider the following matrices:

$$T_1 = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix} \quad T_2 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad T_3 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$T_4 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad T_5 = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad T_6 = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}$$

Which of the matrices has minimal polynomial $m(z) = z^3 + z$? Be sure to justify your answer.

Name (Last, First): _____

9. Consider the matrix

$$T = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$$

Calculate T^{100} applied to the vector $(3, 2)$.

Name (Last, First): _____

10. Let V be a complex vector space of dimension n . Suppose $T : V \rightarrow V$ satisfies $T^n = 0$ but $T^{n-1} \neq 0$. Show that there is a vector $v \in V$ such that the list $v, Tv, T^2v, \dots, T^{n-1}v$ is a basis.