


PRACTICE FINAL (AGOL) - SOLUTIONS

PEYAM RYAN TABRIZIAN

1. 1) Picture:

1A/Practice Exams/Ladder.png



- 2) We want to find $\frac{d\theta}{dt}$ when $x = 6$
- 3) We know $\cos(\theta) = \frac{x}{10}$
- 4) Differentiating, we get: $-\sin(\theta)\frac{d\theta}{dt} = \frac{1}{10}\frac{dx}{dt}$.
- 5) However, $\frac{dx}{dt} = 2$, and if $x = 6$, then we get our usual 6 – 8 – 10-triangle, so $\sin(\theta) = \frac{8}{10} = \frac{4}{5}$. Putting everything together, we get $-\frac{8}{10}\frac{d\theta}{dt} = \frac{6}{10}$, so $\frac{d\theta}{dt} = \frac{6}{8} = \boxed{\frac{3}{4} \text{ ft/s}}$


2. **Note:** I think the problem should say $x > 1$

Define $f(x) = x \ln(x) - (x - 1)$, by the Mean Value Theorem applied to f at $(1, x)$, we get that there is a $c \in (1, x)$ such that $\frac{f(x) - f(1)}{x - 1} = f'(c)$. But $f(1) = 0$, and $f'(c) = \ln(c) + 1 - 1 = \ln(c) > 0$ (since $c > 1$), so we get: $\frac{f(x)}{x - 1} > 0$, so multiplying by $x - 1 > 0$, we get $f(x) > 0$, so $x \ln(x) - (x - 1) > 0$, so


Date: Monday, May 9th, 2011.

$$x \ln(x) > x - 1.$$

3. 1) The circular piece of paper looks as follows:
1A/Practice Exams/Drinking cup.png



- And when you join the edges, your drinking cup should look as follows:
1A/Practice Exams/Drinking cup 2.png



- 2) You want to maximize the volume $V = \frac{\pi}{3}r^2h$, where r is the radius of the cup, and h is the height. But by the Pythagorean theorem, you know that $r^2 + h^2 = R^2$, so $r^2 = R^2 - h^2$, so $V(h) = \frac{\pi}{3}(R^2 - h^2)h$
- 3) The only constraint is $0 \leq h \leq R$ ($h \geq 0$ is clear, and $h \leq R$ is because we want the side of the triangle to be smaller than its hypotenuse)
- 4) $V'(h) = \frac{\pi}{3}(-2h)h + \frac{\pi}{3}(R^2 - h^2) = -\pi h^2 + \frac{\pi}{3}R^2 = 0 \Leftrightarrow h^2 = \frac{R^2}{3} \Leftrightarrow h = \frac{R}{\sqrt{3}}$. Now $V(0) = V(R) = 0$, and $V\left(\frac{R}{\sqrt{3}}\right) = \frac{2\pi R^3}{9\sqrt{3}} > 0$. Hence, by the closed interval method, the maximum volume is $\frac{2\pi R^3}{9\sqrt{3}}$

4. Let F be an antiderivative of $\ln(x)$. Then $\int_{x^2}^{1+x^2} \ln(t)dt = F(x^2) - F(1+x^2)$, so:

$$\frac{d}{dx} \int_{x^2}^{1+x^2} \ln(t)dt = F'(x^2)(2x) - F'(1+x^2)(2x) = \ln(x^2)(2x) - \ln(1+x^2)(2x)$$

5. The slope of the tangent line to $x = y^3$ is $\frac{dy}{dx}$, where $1 = 3y^2 \frac{dy}{dx}$, so $\frac{dy}{dx} = \frac{1}{3y^2}$. And the slope of the tangent line to $y^2 + 3x^2 = 5$ is $\frac{dy}{dx}$, where $2y \frac{dy}{dx} + 6x = 0$, so $\frac{dy}{dx} = -\frac{3x}{y}$. Now when the two curves intersect, the slope of the tangent line to the first curve, $\frac{dy}{dx} = \frac{1}{3y^2} = \frac{1}{3\frac{x}{y}} = \frac{y}{3x}$ (here, I used the fact that $x = y^3$, so dividing by y on both sides, you get $\frac{x}{y} = y^2$), is the negative reciprocal of the slope of the tangent line to the second curve, $\frac{dy}{dx} = -\frac{3x}{y}$! Hence the tangent lines are perpendicular when the two curves intersect!

6. (a) First of all, using the substitution $u = \tan^{-1}(x)$, with $du = \frac{1}{1+x^2}dx$, we get

$$\int_0^1 \frac{\tan^{-1}(x)}{1+x^2} dx = \int_0^{\frac{\pi}{4}} u du = \left[\frac{u^2}{2} \right]_0^{\frac{\pi}{4}} = \frac{\pi^2}{32}. \text{ Hence:}$$

$$\int_0^1 x - \frac{\tan^{-1}(x)}{1+x^2} dx = \left[\frac{x^2}{2} \right]_0^1 - \frac{\pi^2}{32} = \frac{1}{2} - \frac{\pi^2}{32}$$

- (b) $\int_0^2 \sqrt{4-x^2} dx = \frac{\pi(2)^2}{4} = \pi$ (because the integral is the area of the quarter of circle of radius 2)

- (c) Let $u = 1 + e^x$, then $du = e^x dx$, so:

$$\int e^x \sqrt{1+e^x} dx = \int \sqrt{u} du = \frac{2}{3} u^{\frac{3}{2}} + C = \frac{2}{3} (1+e^x)^{\frac{3}{2}} + C$$

7. (a) Domain: $x \neq 0$
 (b) Intercepts: None
 (c) Symmetry: None
 (d) Asymptotes: $y = 0$ is a H.A. at $-\infty$ (by l'Hopital's rule), $x = 0$ is a V.A. (since $\lim_{x \rightarrow 0^+} \frac{e^x}{x} = \infty$ and $\lim_{x \rightarrow 0^+} \frac{e^x}{x} = -\infty$ by direct computation)
 (e) Intervals of increase or decrease: $f'(x) = \frac{e^x x - e^x}{x^2} = \frac{e^x(x-1)}{x^2}$. f is decreasing on $(-\infty, 0) \cup (0, 1)$ and increasing on $(1, \infty)$
 (f) Local maximum and minimum values: $(1, e)$ is a local minimum by the first derivative test!
 (g) Concavity and points of inflection:

$$f''(x) = \frac{(e^x(x-1) + e^x)x^2 - e^x(x-1)(2x)}{x^4} = \frac{x^3 e^x - 2x^2 e^x + 2x e^x}{x^4} = \frac{e^x(x^2 - 2x + 2)}{x^3}$$

But notice that $x^2 - 2x + 1 = (x-1)^2 + 1 > 0$, so the sign of f'' only depends on the sign of x^3 . In particular, f is concave down on $(-\infty, 0)$ and concave up on $(0, \infty)$. There are no inflection points!

- (h) Again, whip out your calculator and look at your rough sketch!

8. (a) $f'(x) = 1 - \frac{1}{\sqrt{x}} > 0$ when $x > 1$, so f is increasing when $x > 1$.
 (b) Let $y = x - 2\sqrt{x}$, all we need to do is to solve for x in terms of y . The trick is: Let $X = \sqrt{x}$, then $y = X^2 - 2X$, so $X^2 - 2X - y = 0$, so $(X-1)^2 + 1 - y = 0$, so $(X-1)^2 = y-1$, so $X-1 = \sqrt{y-1}$ (and this is legitimate, because since f is increasing when $x > 1$ by (a), we have $y = f(x) \geq f(1) = 1$), so $X = 1 + \sqrt{y-1}$, so $\sqrt{x} = 1 + \sqrt{y-1}$, so $x = (1 + \sqrt{y-1})^2$. Hence $f^{-1}(x) = (1 + \sqrt{x-1})^2$
 (c) We need to show that for all $M > 0$ there exists an N large enough such that when $x > N$, then $f(x) > M$.
 Let $M > 0$ be given, we want $x - 2\sqrt{x} > M$. But notice $x - 2\sqrt{x} = (\sqrt{x})(\sqrt{x} - 2) > \sqrt{x}\sqrt{x} = x$.
 Now choose $N = M$, then if $x > N$, $x - 2\sqrt{x} > x > N = M$, and we're done!

9. The picture is on page 437!

Shell method: $K = 0$, $|x| = x$, Outer = $2\sqrt{R^2 - x^2}$ (use the fact that $x^2 + y^2 = R^2$), Inner = 0,

$$\int_r^R 2\pi x(2\sqrt{R^2 - x^2})dx = \frac{4\pi}{3}(R^2 - r^2)^{\frac{3}{2}} = \frac{4\pi}{3} \left(\left(\frac{h}{2} \right)^2 \right)^{\frac{3}{2}} = \frac{4\pi}{3} \frac{h^3}{8} = \frac{\pi h^3}{6}$$

(use the substitution $u = R^2 - r^2$, and the fact that $r^2 + (\frac{h}{2})^2 = R^2$ by the Pythagorean theorem)