

# Profinite completions of products

Peter J. Haine

November 28, 2022

## 0 Introduction

Write  $\mathbf{Spc}_\pi$  for the  $\infty$ -category of  $\pi$ -finite spaces. Given a set  $\Sigma$  of primes, write  $\mathbf{Spc}_\Sigma \subset \mathbf{Spc}_\pi$  for the full subcategory spanned by those  $\pi$ -finite spaces whose homotopy groups are  $\Sigma$ -groups. Write

$$(-)_\Sigma^\wedge : \mathrm{Pro}(\mathbf{Spc}) \rightarrow \mathrm{Pro}(\mathbf{Spc}_\Sigma)$$

for the  $\Sigma$ -completion functor, i.e., the left adjoint to the inclusion  $\mathrm{Pro}(\mathbf{Spc}_\Sigma) \subset \mathrm{Pro}(\mathbf{Spc})$ . One source of difficulty in profinite homotopy theory is that the  $\Sigma$ -completion functor does not preserve finite limits, or even finite products (see [SAG, Remark E.5.2.6; 2, Remark 3.10]). The purpose of this note is to explain two situations in which  $\Sigma$ -completion does preserve products.

The first is that if one is already in the setting of profinite homotopy theory, then  $\Sigma$ -completion preserves products:

**0.1 Proposition (Corollary 2.7).** *Let  $\Sigma$  be a set of primes. Then the  $\Sigma$ -completion functor restricted to profinite spaces*

$$(-)_\Sigma^\wedge : \mathrm{Pro}(\mathbf{Spc}_\pi) \rightarrow \mathrm{Pro}(\mathbf{Spc}_\Sigma)$$

*preserves products.*

The second is that in the setting of étale homotopy theory,  $\Sigma$ -completion preserves finite products. Given a scheme  $X$ , write  $\Pi_\infty^{\mathrm{ét}}(X) \in \mathrm{Pro}(\mathbf{Spc})$  for the étale homotopy type of  $X$ .

**0.2 Proposition (Corollary 2.10).** *Let  $\Sigma$  be a set of primes and let  $X$  and  $Y$  be qcqs schemes. Then the natural map of profinite spaces*

$$(\Pi_\infty^{\mathrm{ét}}(X) \times \Pi_\infty^{\mathrm{ét}}(Y))_\Sigma^\wedge \rightarrow \Pi_\infty^{\mathrm{ét}}(X)_\Sigma^\wedge \times \Pi_\infty^{\mathrm{ét}}(Y)_\Sigma^\wedge$$

*is an equivalence.*

**0.3 Remark.** **Proposition 0.2** fills a gap in the proof of [4, Theorem 5.3]. Chouh's proof cites the false claim that profinite completion preserves finite limits. While this is not generally the case, what is actually used is **Proposition 0.2** (with  $\Sigma$  the set of all primes). In particular, the conclusion of [4, Theorem 5.3] remains valid.

**0.4 Proof Strategy.** **Propositions 0.1** and **0.2** turn out to be a consequence of a more general result. To explain why this is the case, first note that since  $\Sigma$ -completion preserves cofiltered limits, to prove **Proposition 0.1** it suffices to show that  $\Sigma$ -completion preserves finite products of  $\pi$ -finite spaces. This reduction is useful because  $\pi$ -finite spaces admit very nice presentations: every  $\pi$ -finite space can be written as the geometric realization (in  $\mathbf{Spc}$ ) of a simplicial finite set [SAG, Lemma E.1.6.5].

Similarly, to prove [Proposition 0.2](#) we use that the étale homotopy type of a qcqs scheme admits a nice presentation. To see this, the first technical observation is that since protruncation preserves limits and profinite completion factors through protruncation, it suffices to replace the étale homotopy types on the left-hand side by their protruncations. Our work with Barwick and Glasman [[1](#), Theorems 10.2.3 & 12.5.1] provides a description of the protruncated étale homotopy type as the protruncated classifying space of an explicit profinite category. Said differently, the protruncated étale homotopy type can be written as a geometric realization of a simplicial profinite space computed in the larger  $\infty$ -category of protruncated spaces (see [Example A.9](#)).

Hence we're done if we can prove the more general claim that  $\Sigma$ -completion preserves products of protruncated spaces that admit such presentations. This follows once we know that geometric realizations preserve finite products in the  $\infty$ -categories of protruncated and  $\Sigma$ -profinite spaces (see [Proposition 1.12](#) and [Corollary 1.13](#)). See [Lemma 2.6](#) for the key categorical lemma that we use.

**0.5 Linear Overview.** [Section 1](#) proves that geometric realizations preserve finite products in the  $\infty$ -categories of protruncated and profinite spaces. [Section 2](#) proves [Propositions 0.1](#) and [0.2](#). It is immediate from [[1](#), Theorem 10.2.3] that the protruncated étale homotopy type can be written as the geometric realization of a simplicial profinite space. However, for ease of reference we have given a detailed explanation of this fact in [Appendix A](#).

**0.6 Conventions.** Throughout, we use the notational conventions of [[5](#), §§1 & 3].

**Acknowledgments.** We thank Chang-Yeon Chough, Tim Holzschuh, and Sebastian Wolf for many enlightening discussions around the contents of this note. We gratefully acknowledge support from the NSF Mathematical Sciences Postdoctoral Research Fellowship under Grant #DMS-2102957 and a grant from the Simons Foundation (816048, LC).

## 1 Universality of colimits

In this section, we prove that geometric realizations are universal in the  $\infty$ -categories of protruncated and  $\Sigma$ -profinite spaces. A second goal of this section is to record a more general fact: colimits over diagrams that can be computed as finite colimits when valued in an  $n$ -category (see [Definition 1.7](#)) are universal in the  $\infty$ -categories of protruncated and  $\Sigma$ -profinite spaces ([Proposition 1.12](#) and [Corollary 1.13](#)).

The first observation is that finite colimits are universal in protruncated spaces.

**1.1 Lemma.** *Let  $\mathcal{C}$  be an  $\infty$ -category with pullbacks and finite colimits. If finite colimits in  $\mathcal{C}$  are universal, then finite colimits are universal in  $\text{Pro}(\mathcal{C})$ .*

*Proof.* By (the dual of) [[HTT](#), Proposition 5.3.5.15], pullbacks, pushouts, and finite coproducts are computed ‘levelwise’ in  $\text{Pro}(\mathcal{C})$ . Thus the assumption that finite colimits in  $\mathcal{C}$  are universal implies the claim.  $\square$

**1.2 Example.** Finite colimits in  $\text{Pro}(\mathbf{Spc})$  are universal. For each integer  $n \geq 0$ , finite colimits in  $\text{Pro}(\mathbf{Spc}_{\leq n})$  are universal.

Recall that a localization  $L : \mathcal{C} \rightarrow \mathcal{D}$  is *locally cartesian* if for any cospan  $X \rightarrow Z \leftarrow Y$  such that  $X, Z \in \mathcal{D}$ , the natural map  $L(X \times_Z Y) \rightarrow X \times_Z L(Y)$  is an equivalence. The following is immediate from the definitions:

**1.3 Lemma.** *Let  $\mathcal{J}$  be an  $\infty$ -category,  $\mathcal{C}$  an  $\infty$ -category with pullbacks and  $\mathcal{J}$ -shaped colimits, and  $L : \mathcal{C} \rightarrow \mathcal{D}$  a locally cartesian localization. If  $\mathcal{J}$ -shaped colimits in  $\mathcal{C}$  are universal, the  $\mathcal{J}$ -shaped colimits in  $\mathcal{D}$  are universal.*

**1.4 Example.** Since the protruncation functor  $\tau_{<\infty} : \text{Pro}(\mathbf{Spc}) \rightarrow \text{Pro}(\mathbf{Spc}_{<\infty})$  preserves limits [5, Proposition 3.9], [Example 1.2](#) and [Lemma 1.3](#) show that finite colimits in  $\text{Pro}(\mathbf{Spc}_{<\infty})$  are universal.

Now we formulate the key property of the category  $\Delta^{\text{op}}$  that we need.

**1.5 Definition.** Let  $n \geq 0$  be an integer. A functor between  $\infty$ -categories  $c : \mathcal{J} \rightarrow \mathcal{J}$  is *n-colimit-cofinal* if for every  $n$ -category  $\mathcal{C}$  and functor  $f : \mathcal{J} \rightarrow \mathcal{C}$ , the following conditions are satisfied:

(1.5.1) The colimit  $\text{colim}_{\mathcal{J}} f$  exists if and only if the colimit  $\text{colim}_{\mathcal{J}} fc$  exists.

(1.5.2) If the colimit  $\text{colim}_{\mathcal{J}} f$  exists, then the natural map  $\text{colim}_{\mathcal{J}} fc \rightarrow \text{colim}_{\mathcal{J}} f$  is an equivalence.

**1.6 Example.** For an integer  $n \geq 0$ , write  $\Delta_{\leq n} \subset \Delta$  for the full subcategory spanned by those linearly ordered finite sets of cardinality  $\leq n+1$ . By [6, Proposition A.1], the inclusion  $\Delta_{\leq n}^{\text{op}} \subset \Delta^{\text{op}}$  is  $n$ -colimit-cofinal.

**1.7 Definition.** Let  $\mathcal{J}$  be an  $\infty$ -category. We say that  $\mathcal{J}$  is *almost finite* if for each integer  $n \geq 0$ , there exists a *finite*  $\infty$ -category  $\mathcal{J}_n$  and an  $n$ -colimit-cofinal functor  $c_n : \mathcal{J}_n \rightarrow \mathcal{J}$ .

**1.8 Example.** If  $\mathcal{J}$  is an  $\infty$ -category that admits a colimit-cofinal functor from a finite  $\infty$ -category, then  $\mathcal{J}$  is almost finite.

**1.9 Example.** The category  $\Delta^{\text{op}}$  is almost finite: the inclusion  $\Delta_{\leq n}^{\text{op}} \hookrightarrow \Delta^{\text{op}}$  is an  $n$ -colimit-cofinal functor from a finite  $\infty$ -category [3, Example 6.5.3].

Recall that a space  $X$  is *almost  $\pi$ -finite* if  $\pi_0(X)$  is finite and all homotopy groups of  $X$  are finite. Since an almost  $\pi$ -finite space admits a presentation by a Kan complex with finitely many simplices in each dimension [SAG, Lemma E.1.6.5], the following implies that almost  $\pi$ -finite spaces are also almost finite:

**1.10 Example.** Let  $K$  be a simplicial set with finitely many simplices in each dimension and let  $\mathcal{J}$  be the underlying  $\infty$ -category of  $K$ . Then  $\mathcal{J}$  is almost finite: we take  $\mathcal{J}_n$  to be the underlying  $\infty$ -category of the  $(n+1)$ -skeleton of  $\text{sk}_{n+1} K$  and  $c_n : \mathcal{J}_n \rightarrow \mathcal{J}$  the functor induced by the inclusion  $\text{sk}_{n+1} K \subset K$ .

**1.11 Definition.** Let  $\mathcal{C}$  be an  $\infty$ -category with pullbacks. We say that *almost finite colimits in  $\mathcal{C}$  are universal* if for each almost finite  $\infty$ -category  $\mathcal{J}$ , the  $\infty$ -category  $\mathcal{C}$  admits  $\mathcal{J}$ -shaped colimits and  $\mathcal{J}$ -shaped colimits in  $\mathcal{C}$  are universal.

The argument that Lurie gives in the proof of [SAG, Theorem E.6.3.1] shows that almost finite colimits in  $\text{Pro}(\mathbf{Spc}_{\Sigma})$  are universal (however, this result is only stated when  $\Sigma$  is the set of all primes). We also need to know that almost finite colimits in  $\text{Pro}(\mathbf{Spc}_{<\infty})$  are also universal. The strategy is the same as Lurie's proof: we use that equivalences are checked on truncations to reduce to the case of finite colimits.

**1.12 Proposition.** *Almost finite colimits in  $\text{Pro}(\mathbf{Spc}_{<\infty})$  are universal.*

Moreover, [Proposition 1.12](#) reproves [SAG, Theorem E.6.3.1]:

**1.13 Corollary.** *Let  $\Sigma$  be a set of primes. Then almost finite colimits in  $\text{Pro}(\mathbf{Spc}_\Sigma)$  are universal.*

*Proof of Corollary 1.13.* Since the localization  $(-)_\Sigma^\wedge : \text{Pro}(\mathbf{Spc}_{<\infty}) \rightarrow \text{Pro}(\mathbf{Spc}_\Sigma)$  is locally cartesian [5, Proposition 3.18], this follows from Lemma 1.3 and Proposition 1.12.  $\square$

*Proof of Proposition 1.12.* Let  $\mathcal{J}$  be an almost finite  $\infty$ -category, let  $f : X \rightarrow Z$  be a morphism in  $\text{Pro}(\mathbf{Spc}_{<\infty})$ , and let

$$g : \mathcal{J} \rightarrow \text{Pro}(\mathbf{Spc}_{<\infty})/Z$$

be a diagram of protruncated spaces over  $Z$ . To prove the claim, it suffices to show that for each integer  $n \geq 0$ , the induced map

$$\tau_{\leq n} \left( \text{colim}_{i \in \mathcal{J}} X \times_Z g(i) \right) \rightarrow \tau_{\leq n} \left( X \times_Z \text{colim}_{i \in \mathcal{J}} g(i) \right)$$

is an equivalence in  $\text{Pro}(\mathbf{Spc}_{\leq n})$ . Since  $\mathcal{J}$  is almost finite, there exists a finite  $\infty$ -category  $\mathcal{J}_{n+2}$  and  $(n+2)$ -colimit-cofinal functor  $c_{n+2} : \mathcal{J}_{n+2} \rightarrow \mathcal{J}$ . Consider the commutative diagram

$$\begin{array}{ccccc} \text{colim}_{j \in \mathcal{J}_{n+2}} \tau_{\leq n} \left( X \times_Z g c_{n+2}(j) \right) & \xrightarrow{\sim} & \tau_{\leq n} \left( \text{colim}_{j \in \mathcal{J}_{n+2}} X \times_Z g c_{n+2}(j) \right) & \longrightarrow & \tau_{\leq n} \left( X \times_Z \text{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j) \right) \\ \downarrow & & \downarrow & & \downarrow \\ \text{colim}_{i \in \mathcal{J}} \tau_{\leq n} \left( X \times_Z g(i) \right) & \xrightarrow{\sim} & \tau_{\leq n} \left( \text{colim}_{i \in \mathcal{J}} X \times_Z g(i) \right) & \longrightarrow & \tau_{\leq n} \left( X \times_Z \text{colim}_{i \in \mathcal{J}} g(i) \right). \end{array}$$

(Here, the colimits in the leftmost column are computed in  $\text{Pro}(\mathbf{Spc}_{\leq n})$ .) Since  $\text{Pro}(\mathbf{Spc}_{\leq n})$  is an  $(n+1)$ -category and  $c_{n+2} : \mathcal{J}_{n+2} \rightarrow \mathcal{J}$  is  $(n+2)$ -colimit-cofinal, the leftmost vertical map is an equivalence. Thus the central vertical map is also an equivalence. Since  $\mathcal{J}_{n+2}$  is finite and finite colimits in  $\text{Pro}(\mathbf{Spc}_{<\infty})$  are universal (Example 1.4), the top right-hand horizontal map is an equivalence.

To complete the proof, it suffices to show that the rightmost vertical map is an equivalence. For this, consider the commutative square

$$\begin{array}{ccc} \text{colim}_{j \in \mathcal{J}_{n+2}} \tau_{\leq n+1} g c_{n+2}(j) & \xrightarrow{\sim} & \tau_{\leq n+1} \left( \text{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j) \right) \\ \downarrow & & \downarrow \\ \text{colim}_{i \in \mathcal{J}} \tau_{\leq n+1} g(i) & \xrightarrow{\sim} & \tau_{\leq n+1} \left( \text{colim}_{i \in \mathcal{J}} g(i) \right), \end{array}$$

where the colimits in the left-hand column are computed in  $\text{Pro}(\mathbf{Spc}_{\leq n+1})$ . Since  $\text{Pro}(\mathbf{Spc}_{\leq n+1})$  is an  $(n+2)$ -category and  $c_{n+2}$  is  $(n+2)$ -colimit-cofinal, the left-hand vertical map is an equivalence. Hence the right-hand vertical map is also an equivalence. As a consequence, the map

$$\text{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j) \rightarrow \text{colim}_{i \in \mathcal{J}} g(i)$$

is  $n$ -connected. Thus the basechange

$$X \times_Z \text{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j) \rightarrow X \times_Z \text{colim}_{i \in \mathcal{J}} g(i)$$

is also  $n$ -connected. Hence the map

$$\tau_{\leq n} \left( X \times_Z \operatorname{colim}_{j \in \mathcal{J}_{n+2}} g c_{n+2}(j) \right) \rightarrow \tau_{\leq n} \left( X \times_Z \operatorname{colim}_{i \in \mathcal{J}} g(i) \right)$$

is an equivalence, as desired.  $\square$

Note that the universality of geometric realizations implies that geometric realizations preserve finite products:

**1.14 Lemma.** *Let  $\mathcal{J}$  be a sifted  $\infty$ -category and let  $\mathcal{C}$  be an  $\infty$ -category with finite limits and  $\mathcal{J}$ -shaped colimits. If  $\mathcal{J}$ -shaped colimits in  $\mathcal{C}$  are universal, then the functor*

$$\operatorname{colim}_{\mathcal{J}} : \operatorname{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \mathcal{C}$$

*preserves finite products.*

*Proof.* Let  $X_{\bullet}, Y_{\bullet} : \mathcal{J} \rightarrow \mathcal{C}$  be functors. We have natural equivalences

$$\begin{aligned} \operatorname{colim}_{i \in \mathcal{J}} X_i \times Y_i &\simeq \operatorname{colim}_{(i,j) \in \mathcal{J} \times \mathcal{J}} X_i \times Y_j && (\mathcal{J} \text{ is sifted}) \\ &\simeq \operatorname{colim}_{i \in \mathcal{J}} \operatorname{colim}_{j \in \mathcal{J}} (X_i \times Y_j) \\ &\simeq \operatorname{colim}_{i \in \mathcal{J}} \left( X_i \times \operatorname{colim}_{j \in \mathcal{J}} Y_j \right) && (\mathcal{J}\text{-shaped colimits are universal}) \\ &\simeq \left( \operatorname{colim}_{i \in \mathcal{J}} X_i \right) \times \left( \operatorname{colim}_{j \in \mathcal{J}} Y_j \right) && (\mathcal{J}\text{-shaped colimits are universal}). \quad \square \end{aligned}$$

**1.15 Example.** Let  $\Sigma$  be a set of primes. **Proposition 1.12**, **Corollary 1.13**, and **Lemma 1.14** show that geometric realizations preserve finite products in the  $\infty$ -categories  $\operatorname{Pro}(\mathbf{Spc}_{\Sigma})$  and  $\operatorname{Pro}(\mathbf{Spc}_{<\infty})$ .

## 2 Completions of products

The goal of this section is to prove **Propositions 0.1** and **0.2**. We do this by noting that in the setting of both of these results, the prospaces of interest can be written as geometric realizations of simplicial profinite spaces. We begin by axiomatizing the situation:

**2.1 Notation.** Let  $\mathcal{C}$  be an  $\infty$ -category and let  $\mathcal{D} \subset \mathcal{C}$  be a full subcategory. Given a simplicial object  $X_{\bullet} : \Delta^{\operatorname{op}} \rightarrow \mathcal{D}$ , write

$$|X_{\bullet}|_{\mathcal{D}} := \operatorname{colim} \left( \Delta^{\operatorname{op}} \xrightarrow{X_{\bullet}} \mathcal{D} \right)$$

for the geometric realization of  $X_{\bullet}$  computed in  $\mathcal{D}$ . Write

$$|X_{\bullet}|_{\mathcal{C}} := \operatorname{colim} \left( \Delta^{\operatorname{op}} \xrightarrow{X_{\bullet}} \operatorname{Pro}(\mathbf{Spc}_{\pi}) \hookrightarrow \mathcal{C} \right)$$

for the geometric realization of  $X_{\bullet}$  computed in  $\mathcal{C}$ . Importantly, if the inclusion  $\mathcal{D} \subset \mathcal{C}$  admits a left adjoint  $L : \mathcal{C} \rightarrow \mathcal{D}$ , then

$$L |X_{\bullet}|_{\mathcal{C}} \simeq |X_{\bullet}|_{\mathcal{D}} .$$

**2.2 Definition.** Let  $\mathcal{C}$  be an  $\infty$ -category and  $\mathcal{D} \subset \mathcal{C}$  a full subcategory. We say that an object  $X \in \mathcal{C}$  admits a  $\mathcal{D}$ -presentation if there exists a simplicial object  $X_\bullet : \Delta^{\text{op}} \rightarrow \mathcal{D}$  and an equivalence

$$|X_\bullet|_{\mathcal{C}} \simeq X.$$

**2.3 Notation.** Write  $\mathbf{Set}_{\text{fin}}$  for the category of a finite sets and all maps.

**2.4 Example.** Let  $X$  be a space which is finite<sup>1</sup> or almost  $\pi$ -finite. Then:

(2.4.1) As an object of  $\mathbf{Spc}$ , the space  $X$  admits a  $\mathbf{Set}_{\text{fin}}$ -presentation: If  $X$  is finite, then  $X$  can be written as the geometric realization of a simplicial set with finitely many nondegenerate simplices [12, Proposition 2.4].<sup>2</sup> If  $X$  is almost  $\pi$ -finite, then  $X$  can be written as the geometric realization of a Kan complex with finitely many simplices in each degree [SAG, Lemma E.1.6.5].

(2.4.2) Since the Yoneda embedding  $\mathbf{Spc} \hookrightarrow \text{Pro}(\mathbf{Spc})$  preserves colimits, when regarded as an object of  $\text{Pro}(\mathbf{Spc})$ , the space  $X$  admits a  $\mathbf{Set}_{\text{fin}}$ -presentation.

(2.4.3) Since the protruncation functor  $\tau_{<\infty} : \text{Pro}(\mathbf{Spc}) \rightarrow \text{Pro}(\mathbf{Spc}_{<\infty})$  preserves colimits, the protruncated space  $\tau_{<\infty}(X)$  admits a  $\mathbf{Set}_{\text{fin}}$ -presentation.

(2.4.4) Since the profinite completion functor  $(-)^{\wedge}_{\pi} : \text{Pro}(\mathbf{Spc}) \hookrightarrow \text{Pro}(\mathbf{Spc}_{\pi})$  preserves colimits, the profinite space  $X_{\pi}^{\wedge}$  admits a  $\mathbf{Set}_{\text{fin}}$ -presentation.

Algebraic geometry also gives rise to many examples of protruncated spaces admitting profinite presentations.

**2.5 Notation.** Given an  $\infty$ -topos  $\mathbf{X}$ , we write  $\Pi_{\infty}(\mathbf{X}) \in \text{Pro}(\mathbf{Spc})$  for the shape of  $\mathbf{X}$ . We write  $\Pi_{<\infty}(\mathbf{X})$  for the protruncation of  $\Pi_{\infty}(\mathbf{X})$ .

Recall that an  $\infty$ -topos  $\mathbf{X}$  is *spectral* in the sense of [1, Definition 9.2.1] if  $\mathbf{X}$  is bounded coherent and the  $\infty$ -category  $\text{Pt}(\mathbf{X})$  of points of  $\mathbf{X}$  has the property that every endomorphism of an object of  $\text{Pt}(\mathbf{X})$  is an equivalence. The most important example of a spectral  $\infty$ -topos is the étale  $\infty$ -topos of a qcqs scheme [1, Example 9.2.4]. **Example A.9** explains why the protruncated space  $\Pi_{<\infty}(\mathbf{X})$  of a spectral  $\infty$ -topos admits a natural  $\text{Pro}(\mathbf{Spc}_{\pi})$ -presentation.

The following is the key categorical result that we need:

**2.6 Lemma.** *Let  $\mathcal{C}$  be an  $\infty$ -category with geometric realizations and finite products, and let  $L : \mathcal{C} \rightarrow \mathcal{D}$  be a localization. Assume that geometric realizations preserve finite products in both  $\mathcal{C}$  and  $\mathcal{D}$ . If  $X, Y \in \mathcal{C}$  admit  $\mathcal{D}$ -presentations  $X_\bullet$  and  $Y_\bullet$ , then the natural map*

$$L(X \times Y) \rightarrow L(X) \times L(Y)$$

*is an equivalence.*

<sup>1</sup>I.e., in the smallest subcategory of  $\mathbf{Spc}$  containing the point and closed under finite coproducts and pushouts.

<sup>2</sup>In fact, every finite space is equivalent to the classifying space of a finite poset; see [HTT, Proposition 4.1.1.3(3) & Variant 4.2.3.16; 10, Theorem 1].

*Proof.* We compute

$$\begin{aligned}
L(X \times Y) &\simeq L(|X \cdot |_{\mathcal{C}} \times |Y \cdot |_{\mathcal{C}}) \\
&\simeq L(|X \cdot \times Y \cdot |_{\mathcal{C}}) && \text{(geometric realizations preserve finite products in } \mathcal{C} \text{)} \\
&\simeq |X \cdot \times Y \cdot |_{\mathcal{D}} \\
&\simeq |X \cdot |_{\mathcal{D}} \times |Y \cdot |_{\mathcal{D}} && \text{(geometric realizations preserve finite products in } \mathcal{D} \text{)} \\
&\simeq L |X \cdot |_{\mathcal{C}} \times L |Y \cdot |_{\mathcal{C}} \\
&\simeq L(X) \times L(Y). \quad \square
\end{aligned}$$

In the rest of this section, we record consequences of [Lemma 2.6](#).

**2.7 Corollary.** *Let  $\Sigma$  be a set of prime numbers. Then the  $\Sigma$ -completion functor*

$$(-)_{\Sigma}^{\wedge} : \text{Pro}(\mathbf{Spc}_{\pi}) \rightarrow \text{Pro}(\mathbf{Spc}_{\Sigma})$$

*preserves products.*

*Proof.* Since  $\Sigma$ -completion preserves cofiltered limits and the terminal object, it suffices to show that  $\Sigma$ -completion preserves binary products of profinite spaces. Again because  $\Sigma$ -completion preserves cofiltered limits, we are reduced to showing that if  $X$  and  $Y$  are  $\pi$ -finite spaces, then the natural map

$$(X \times Y)_{\Sigma}^{\wedge} \rightarrow X_{\Sigma}^{\wedge} \times Y_{\Sigma}^{\wedge}$$

is an equivalence. Since  $\pi$ -finite spaces admit  $\mathbf{Set}_{\text{fin}}$ -presentations ([Example 2.4](#)) and geometric realizations preserve finite products in  $\text{Pro}(\mathbf{Spc}_{\pi})$  and  $\text{Pro}(\mathbf{Spc}_{\Sigma})$  ([Example 1.15](#)), the claim follows from [Lemma 2.6](#).  $\square$

**2.8 Warning.** The functor  $(-)_{\Sigma}^{\wedge} : \text{Pro}(\mathbf{Spc}_{\pi}) \rightarrow \text{Pro}(\mathbf{Spc}_{\Sigma})$  does not generally preserve pullbacks, or even loop objects.

**2.9 Corollary.** *Let  $\Sigma$  be a set of primes and let  $X$  and  $Y$  be protruncated spaces that admit  $\text{Pro}(\mathbf{Spc}_{\pi})$ -presentations. Then the natural map*

$$(X \times Y)_{\Sigma}^{\wedge} \rightarrow X_{\Sigma}^{\wedge} \times Y_{\Sigma}^{\wedge}$$

*is an equivalence.*

*Proof.* Since geometric realizations preserve finite products in  $\text{Pro}(\mathbf{Spc}_{<\infty})$  and  $\text{Pro}(\mathbf{Spc}_{\pi})$  ([Example 1.15](#)), [Lemma 2.6](#) shows that the natural map

$$(X \times Y)_{\pi}^{\wedge} \rightarrow X_{\pi}^{\wedge} \times Y_{\pi}^{\wedge}$$

is an equivalence. The claim now follows from [Corollary 2.7](#).  $\square$

Since protruncation preserves finite products, [Corollary 2.9](#) and [Example A.9](#) show:

**2.10 Corollary.** *Let  $\Sigma$  be a set of primes and let  $\mathbf{X}$  and  $\mathbf{Y}$  be spectral  $\infty$ -topoi. Then the natural map*

$$(\Pi_{\infty}(\mathbf{X}) \times \Pi_{\infty}(\mathbf{Y}))_{\Sigma}^{\wedge} \rightarrow \Pi_{\infty}(\mathbf{X})_{\Sigma}^{\wedge} \times \Pi_{\infty}(\mathbf{Y})_{\Sigma}^{\wedge}$$

*is an equivalence.*

## A Classifying prospaces via geometric realizations

The purpose of this appendix is to explain why the protruncated shape of a spectral  $\infty$ -topos (e.g., the étale  $\infty$ -topos of a qcqs scheme) admits a presentation as a geometric realization of a simplicial profinite space (Example A.9). Using the description of the protruncated shape of a spectral  $\infty$ -topos as a protruncated classifying prospace given by Barwick–Glasman–Haine [1, Theorem 10.2.3], this is an exercise in the definitions. For the ease of the reader, we spell out the details here.

We make use of the description of  $\infty$ -categories as simplicial spaces.

**A.1 Recollection.** The nerve functor

$$\begin{aligned} N : \mathbf{Cat}_\infty &\rightarrow \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathbf{Spc}) \\ \mathcal{C} &\mapsto [I \mapsto \mathrm{Fun}(I, \mathcal{C})^\simeq] \end{aligned}$$

is fully faithful [HA, Proposition A.7.10; SAG, §A.8.2; 7; 8, §1; 11]. One can explicitly identify the essential image; objects in the image of this embedding are often called *complete Segal spaces* or *categories internal to spaces*. Under this embedding, the subcategory  $\mathbf{Spc} \subset \mathbf{Cat}_\infty$  corresponds to the constant functors  $\Delta^{\mathrm{op}} \rightarrow \mathbf{Spc}$ . Moreover, the localization  $B : \mathbf{Cat}_\infty \rightarrow \mathbf{Spc}$  is given by geometric realization.

**A.2 Notation.** Let  $\mathcal{C}$  be an  $\infty$ -category with finite limits. We write

$$\mathrm{Cat}(\mathcal{C}) \subset \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathcal{C})$$

for the full subcategory spanned by the *categories internal to  $\mathcal{C}$* . See [1, Definition 13.1.1; 9, Proposition 3.2.7] for the definition.

**A.3 Notation.** Write  $\mathbf{Cat}_{<\infty} \subset \mathbf{Cat}_\infty$  for the full subcategory spanned by those  $\infty$ -categories  $\mathcal{C}$  for which there exists an integer  $n \geq 0$  such that  $\mathcal{C}$  is an  $n$ -category. Write  $\mathbf{Cat}_{\infty, \pi} \subset \mathbf{Cat}_{<\infty}$  for the full subcategory spanned by those  $\infty$ -categories with the property that there are finitely many objects up to equivalence and all mapping spaces are  $\pi$ -finite.

**A.4 Observation.** The nerve  $N : \mathbf{Cat}_\infty \simeq \mathrm{Cat}(\mathbf{Spc})$  restricts to equivalences

$$\mathbf{Cat}_{<\infty} \simeq \mathrm{Cat}(\mathbf{Spc}_{<\infty}) \quad \text{and} \quad \mathbf{Cat}_{\infty, \pi} \simeq \mathrm{Cat}(\mathbf{Spc}_\pi).$$

In order to describe protruncated classifying spaces via geometric realizations, it is useful to describe pro- $\infty$ -categories as category objects in prospaces.

**A.5 Observation.** By [HTT, Proposition 5.3.5.11; 1, Proposition 13.1.12], the composite

$$\mathbf{Cat}_{<\infty} \xrightarrow[\mathrm{N}]{\simeq} \mathrm{Cat}(\mathbf{Spc}_{<\infty}) \hookrightarrow \mathrm{Cat}(\mathrm{Pro}(\mathbf{Spc}_{<\infty}))$$

extends along cofiltered limits to a fully faithful right adjoint

$$N : \mathrm{Pro}(\mathbf{Cat}_{<\infty}) \hookrightarrow \mathrm{Cat}(\mathrm{Pro}(\mathbf{Spc}_{<\infty})).$$

This functor restricts to a fully faithful right adjoint

$$N : \mathrm{Pro}(\mathbf{Cat}_{\infty, \pi}) \hookrightarrow \mathrm{Cat}(\mathrm{Pro}(\mathbf{Spc}_\pi)).$$



**A.6 Remark.** We do not know if the embedding  $N : \mathbf{Cat}_\infty \hookrightarrow \mathbf{Cat}(\mathrm{Pro}(\mathbf{Spc}))$  extends along cofiltered limits to a fully faithful functor  $\mathrm{Pro}(\mathbf{Cat}_\infty) \rightarrow \mathbf{Cat}(\mathrm{Pro}(\mathbf{Spc}))$ .

**A.7 Observation.** It is immediate from the definitions that the following diagram of fully faithful right adjoints commutes

$$\begin{array}{ccccccc}
\mathrm{Pro}(\mathbf{Spc}_\pi) & \xleftarrow[\perp]{B_\pi^\wedge} & \mathrm{Pro}(\mathbf{Cat}_{\infty,\pi}) & \xleftarrow[\mathrm{N}]{\perp} & \mathbf{Cat}(\mathrm{Pro}(\mathbf{Spc}_\pi)) & \xleftarrow[\perp]{\perp} & \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Pro}(\mathbf{Spc}_\pi)) \\
(-)^\wedge_\pi \updownarrow & & \updownarrow & & \updownarrow & & (-)^\wedge_\pi \circ - \updownarrow \\
\mathrm{Pro}(\mathbf{Spc}_{<\infty}) & \xleftarrow[\perp]{B_{<\infty}} & \mathrm{Pro}(\mathbf{Cat}_{<\infty}) & \xleftarrow[\mathrm{N}]{\perp} & \mathbf{Cat}(\mathrm{Pro}(\mathbf{Spc}_{<\infty})) & \xleftarrow[\perp]{\perp} & \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Pro}(\mathbf{Spc}_{<\infty})) .
\end{array}$$

The long composite left adjoints

$$\mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Pro}(\mathbf{Spc}_{<\infty})) \rightarrow \mathrm{Pro}(\mathbf{Spc}_{<\infty}) \quad \text{and} \quad \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Pro}(\mathbf{Spc}_\pi)) \rightarrow \mathrm{Pro}(\mathbf{Spc}_\pi) .$$

are simply the colimit functors. Since the diagram of left adjoints also commutes we deduce:

**A.8 Corollary.** *Let  $\mathcal{C} \in \mathrm{Pro}(\mathbf{Cat}_{\infty,\pi})$  be a profinite  $\infty$ -category. Then there are natural equivalences*

$$B_{<\infty}(\mathcal{C}) \simeq \mathrm{colim} \left( \Delta^{\mathrm{op}} \xrightarrow{N(\mathcal{C})} \mathrm{Pro}(\mathbf{Spc}_\pi) \hookrightarrow \mathrm{Pro}(\mathbf{Spc}_{<\infty}) \right)$$

and

$$B_\pi^\wedge(\mathcal{C}) \simeq \mathrm{colim} \left( \Delta^{\mathrm{op}} \xrightarrow{N(\mathcal{C})} \mathrm{Pro}(\mathbf{Spc}_\pi) \right) .$$

**A.9 Example.** Let  $\mathbf{X}$  be a spectral  $\infty$ -topos. Through [1, Theorem 9.3.1], Barwick–Glasman–Haine refined the the  $\infty$ -category  $\mathrm{Pt}(\mathbf{X})$  of points of  $\mathbf{X}$  to a profinite  $\infty$ -category

$$\widehat{\Pi}_{(\infty,1)}(\mathbf{X}) \in \mathrm{Pro}(\mathbf{Cat}_{\infty,\pi})$$

called the *stratified shape* of  $\mathbf{X}$ . In [1, Theorem 10.2.3] they show that there is a natural equivalence

$$\Pi_{<\infty}(\mathbf{X}) \simeq B_{<\infty} \left( \widehat{\Pi}_{(\infty,1)}(\mathbf{X}) \right) .$$

That is, the protruncated shape of  $\mathbf{X}$  can be recovered as the protruncated classifying space of the stratified shape  $\widehat{\Pi}_{(\infty,1)}(\mathbf{X})$ . Hence **Corollary A.8** shows that the protruncated shape  $\Pi_{<\infty}(\mathbf{X})$  admits a natural  $\mathrm{Pro}(\mathbf{Spc}_\pi)$ -presentation in the sense of **Definition 2.2**.

## References

- HTT J. Lurie, *Higher topos theory*, Annals of Mathematics Studies. Princeton, NJ: Princeton University Press, 2009, vol. 170, pp. xviii+925, ISBN: 978-0-691-14049-0; 0-691-14049-9.
- HA \_\_\_\_\_, *Higher algebra*, Preprint available at [math.ias.edu/~lurie/papers/HA.pdf](http://math.ias.edu/~lurie/papers/HA.pdf), Sep. 2017.
- SAG \_\_\_\_\_, *Spectral algebraic geometry*, Preprint available at [math.ias.edu/~lurie/papers/SAG-rootfile.pdf](http://math.ias.edu/~lurie/papers/SAG-rootfile.pdf), Feb. 2018.
1. C. Barwick, S. Glasman, and P. J. Haine, *Exodromy*, Preprint available at [arXiv:1807.03281v7](https://arxiv.org/abs/1807.03281v7), Aug. 2020.

2. P. Boavida de Brito, G. Horel, and M. Robertson, *Operads of genus zero curves and the Grothendieck–Teichmüller group*, *Geom. Topol.*, vol. 23, no. 1, pp. 299–346, 2019. DOI: 10.2140/gt.2019.23.299.
3. B. Calmès, E. Dotto, Y. Harpaz, F. Hebestreit, M. Land, K. Moi, D. Nardin, T. Nikolaus, and W. Steimle, *Hermitian K-theory for stable  $\infty$ -categories I: Foundations*, Preprint available at arXiv:2009.07223, Dec. 2021.
4. C.-Y. Chough, *Proper base change for étale sheaves of spaces*, *J. Eur. Math. Soc. (JEMS)*, vol. 24, no. 11, pp. 3679–3702, 2022. DOI: 10.4171/jems/1263.
5. P. J. Haine, T. Holzschuh, and S. Wolf, *The fundamental fiber sequence in étale homotopy theory*, Preprint available at arXiv:2209.03476, Oct. 2022.
6. L. Hesselholt and P. Pstrągowski, *Dirac geometry I: Commutative algebra*, Preprint available at arXiv:2207.09256, Jul. 2022.
7. A. Joyal and M. Tierney, *Quasi-categories vs Segal spaces*, in *Categories in algebra, geometry and mathematical physics*, *Contemp. Math. Vol. 431*, Amer. Math. Soc., Providence, RI, 2007, pp. 277–326. DOI: 10.1090/conm/431/08278.
8. J. Lurie,  *$(\infty, 2)$ -Categories and the Goodwillie Calculus I*, Preprint available at math.ias.edu/~lurie/papers/GoodwillieI.pdf, Oct. 2009.
9. L. Martini, *Yoneda’s lemma for internal higher categories*, Preprint available at arXiv:2103.17141, Apr. 2022.
10. M. C. McCord, *Singular homology groups and homotopy groups of finite topological spaces*, *Duke Math. J.*, vol. 33, pp. 465–474, 1966.
11. C. Rezk, *A model for the homotopy theory of homotopy theory*, *Trans. Amer. Math. Soc.*, vol. 353, no. 3, 973–1007 (electronic), 2001.
12. M. Volpe, *Verdier duality on conically smooth stratified spaces*, Preprint available at arXiv:2206.02728, Jun. 2022.