

# q-Operators as Geometrization of N=2 Theories

**Peter Koroteev**

2/9/2022

Talk at Aspen conference on *Geometrization of  $D \leq 6$  Theories*

# Literature

[arXiv:2108.04184]

**q-Operators, QQ-systems, and Bethe Ansatz II: Generalized Minors**

[P. Koroteev](#), [A. M. Zeitlin](#)

[arXiv:2105.00588]

**3d Mirror Symmetry for Instanton Moduli Spaces**

[P. Koroteev](#), [A. M. Zeitlin](#)

[arXiv:2007.11786] J. Inst. Math. Jussieu

**Toroidal q-Operators**

[P. Koroteev](#), [A. M. Zeitlin](#)

[arXiv:2002.07344] J. Europ. Math. Soc.

**q-Operators, QQ-Systems, and Bethe Ansatz**

[E. Frenkel](#), [P. Koroteev](#), [D. S. Sage](#), [A. M. Zeitlin](#)

[arXiv:1811.09937] Commun.Math.Phys. **381** (2021) 641


**(SL(N),q)-operators, the q-Langlands correspondence, and quantum/classical duality**

[P. Koroteev](#), [D. S. Sage](#), [A. M. Zeitlin](#)

[arXiv:1705.10419] Selecta Math. **27** (2021) 87

**Quantum K-theory of Quiver Varieties and Many-Body Systems**

[P. Koroteev](#), [P. P. Pushkar](#), [A. V. Smirnov](#), [A. M. Zeitlin](#)





# Motivation

Quantum Geometry and Integrable Systems

[Okounkov et al]

[Pushkar, Zeitlin, Smirnov]

[PK, Pushkar, Smirnov, Zeitlin]

BPS/CFT Correspondence

[Nekrasov Shatashvili]

Geometric q-Langlands Correspondence

[Frenkel]

[Aganagic, Frenkel, Okounkov]




ODE/IM Correspondence

[Bazhanov, Lukyanov, Zamolodchikov]

[Dorey, Tateo]

# I. Integrable Many-Body Systems

Calogero in 1971 introduced a new integrable system. Moser in 1975 proved its integrability using Lax pair



$$H_{CM} = \sum_{i=1}^n \frac{p_i^2}{2m} + g^2 \sum_{j \neq i} \frac{1}{(x_i - x_j)^2}$$

$$V(z) \simeq \frac{1}{z^2} \quad \wp(x_j - x_i)$$

The **Calogero-Moser (CM)** system has several generalizations

rCM  $\rightarrow$  tCM  $\rightarrow$  eCM

$$V(z) \simeq \frac{1}{\sinh z^2}$$

Another relativistic generalization called **Ruijsenaars-Schneider (RS)** family

rRS  $\rightarrow$  tRS  $\rightarrow$  eRS

Geometrically described by Hamiltonian reduction of  $T^*GL(n)$

$$H_{CM} = \lim_{c \rightarrow \infty} H_{RS} - nmc^2$$




# The ITEP Table

[Gorsky PK Koroteeva Shakirov ]

<div>p \ q</div>	rational	trigonometric	elliptic
r	<div>rational CMS</div>	<div>trigonometric CMS</div>	<div>elliptic CMS</div> <div>quantum cohomology</div>
t	<div>rational RS</div> <div>(dual trig. CMS)</div>	<div>trigonometric RS</div>	<div>elliptic RS</div> <div>quantum K-theory</div>
e	<div>dual elliptic CMS</div>	<div>dual elliptic RS</div>	<div>DELL</div> <div>Elliptic Cohomology</div>

Quantum XXZ Chain

QQ-Systems



SU(**n**) XXZ spin chain on n sites w/ **anisotropies** and **twisted periodic boundary conditions**

**twist eigenvalues**  $z_i$


**equivariant parameters** (anisotropies)  $a_i$

Bethe Ansatz Equations

$$\frac{\zeta_i}{\zeta_{i+1}} \prod_{\beta=1}^{\mathbf{v}_{i-1}} \frac{\sigma_{i,\alpha} - \hbar^{1/2} \sigma_{i-1,\beta}}{\sigma_{i-1,\beta} - \hbar^{1/2} \sigma_{i,\alpha}} \cdot \prod_{\substack{\beta=1 \\ \beta \neq \alpha}}^{\mathbf{v}_i} \frac{\hbar \sigma_{i,\alpha} - \sigma_{i,\beta}}{\hbar \sigma_{i,\beta} - \sigma_{i,\alpha}} \cdot \prod_{\beta=1}^{\mathbf{v}_{i+1}} \frac{\sigma_{i,\alpha} - \hbar^{1/2} \sigma_{i+1,\beta}}{\sigma_{i+1,\beta} - \hbar^{1/2} \sigma_{i,\alpha}} = (-1)^{\delta_i}$$

Classical tRS Model

q-Operators



**n**-particle trigonometric Ruijsenaars-Schneider model

$$\Omega = \sum_i \frac{dp_i}{p_i} \wedge \frac{dz_i}{z_i}$$
$$[T_i, T_j] = 0$$

**coordinates**  $z_i$

**energy** (eigenvalues of Hamiltonians)  $e_i(a_i)$

Energy level equations


$$T_i(\mathbf{z}, \hbar) = e_i(\mathbf{a}), \quad i = 1, \dots, n$$



# Quiver Varieties from Branes

[Nekrasov Shatashvili]  
[PK Pushkar Smirnov Zeitlin]

Quiver Variety from Hanany-Witten




moment map



$$\mu : T^*R \longrightarrow \text{Lie}(G)^* \qquad L(\mathbf{v}, \mathbf{w}) = \mu^{-1}(0)$$

$$Y = L(\mathbf{v}, \mathbf{w}) //_{\theta} G = L(\mathbf{v}, \mathbf{w})_{ss} / G$$

automorphism group  $\prod_i GL(W_i) \times \mathbb{C}_{\hbar}^{\times}$



Physically: 3d N=4 quiver gauge theory



Classical K-theory of X is formed by tensorial polynomials of tautological bundles and their duals

The equivariant K-theory of X is a module over the ring of equivariant constants  $R = K_{\mathsf{T}}(\cdot) = \mathbb{Z}[a_1^{\pm}, \dots, a_n^{\pm 1}, \hbar^{\pm 1}]$


K-theory classes  $\tau(V) = V^{\otimes 2} - \Lambda^3 V^*$

$$\tau(s_1, \dots, s_k) = (s_1 + \dots + s_k)^2 - \sum_{1 \leq i_1 < i_2 < i_3 \leq k} s_{i_1}^{-1} s_{i_2}^{-1} s_{i_3}^{-1}$$

Relations  $\prod_{j=1}^n (s_i - a_j) = 0, \quad i = 1 \dots k$

# Quantum K-theory

Quantum equivariant K-theory of Nakajima quiver varieties



$$A \circledast B = A \otimes B + \sum_{d=1}^\infty A \circledast_d B z^d$$

$$\mathbf{V}^{(\tau)}(z) = \sum_d \mathrm{ev}_{p_2,*}(\widehat{\mathcal{O}}_{\mathrm{vir}}^d \otimes \tau|_{p_1}, \mathrm{QM}_{\mathrm{nonsing}\,p_2}^d) z^d \in K_{\mathbb{T} \times \mathbb{C}_q^\times}(X)_{loc}[[z]]$$

Saddle point limit yields Bethe equations for XXZ

$$\hbar^{\frac{\Delta_i}{2}} \frac{\zeta_i}{\zeta_{i+1}} \frac{Q_{i-1}^{(1)} Q_i^{(-2)} Q_{i+1}^{(1)}}{Q_{i-1}^{(-1)} Q_i^{(2)} Q_{i+1}^{(-1)}} = -1$$

$$Q_i(u) = \prod_{\alpha=1}^{\mathbf{v}_i} (u - \sigma_{i,\alpha})$$


$$\Lambda_i(z) = \prod_{b=1}^{\mathbf{w}_i} (z - a_{i,b})$$

Can be written as QQ-system

$$\xi_i Q_i^+(\hbar z) Q_i^-(z) - \xi_{i+1} Q_i^+(z) Q_i^-(\hbar z) = \Lambda_i(z) Q_{i-1}^+(\hbar z) Q_{i+1}^+(z)$$

# Quantum/Classical Duality from Branes


[PK Gaiotto]  
[PK Zeitlin]



Quiver representation data  $\longleftrightarrow$  Linking Numbers

$$r_i^! = \#D3(R) - \#D3(L) + \#D5(L)$$

$$r_i = \#D3(L) - \#D3(R) + \#NS5(R)$$




3d N=2\* quiver theory  $\longleftrightarrow$  4d N=2\* theory on interval

$$\mathbb{R}^2 \times S^1 \times I_{L,R}^1$$

Quantum K-theory of X  $\longleftrightarrow$  Calogero-Moser Space


$$\hbar MT - TM = u \otimes v^T$$



$$QK_T(X) \cong \frac{\mathbb{C}(\{\xi_i\}, \{a_i\}, \hbar)(\{p_i\})}{(\det(u - T(\{p_i\}, \{a_i\}, \hbar)) - f(u, \{\xi_i\}, \hbar))}$$


T - tRS Lax Matrix

Cotangent bundle to complete flag variety:  
n-particle tRS



# Quantum/Classical Duality


[PK Gaiotto]  
[PK Zeitlin]



tRS momenta  $p_i^\xi = \exp \frac{\partial Y}{\partial \xi_i}, \quad p_i^a = \exp \frac{\partial Y}{\partial a_i}$

Symplectic form  $\Omega = \sum_{i=1}^N \frac{dp_i^\xi}{p_i^\xi} \wedge \frac{d\xi_i}{\xi_i} - \frac{dp_i^a}{p_i^a} \wedge \frac{da_i}{a_i}$

tRS energy relations



$\mathcal{W} = \widetilde{\mathcal{W}}$   
 $Y = Y'$   
3d mirror symmetry


$$\det(u - T) = \prod_{i=1}^N (u - a_i), \quad \det(u - M) = \prod_{i=1}^N (u - \xi_i)$$

$$\sum_{\substack{\mathcal{J} \subset \{1, \dots, L\} \\ |\mathcal{J}| = k}} \prod_{\substack{i \in \mathcal{J} \\ j \notin \mathcal{J}}} \frac{a_i - \hbar a_j}{a_i - a_j} \prod_{m \in \mathcal{J}} p_m = \ell_k(\xi_i)$$

$\mathcal{L}_\mu$  Eigenvalues of M and Slodowy form on T

$\mathcal{L}_\tau$  Eigenvalues of T and Slodowy form on M

space of vacua — intersection points



[Dimofte Gaiotto van der Veen]

XXZ/tRS duality! Can we generalize it?




# II. q-Operators — SL(2) Example

Consider vector bundle  $E$  over  $\mathbb{P}^1$

$$M_q : \mathbb{P}^1 \rightarrow \mathbb{P}^1$$

$$z \mapsto qz$$



Map of vector bundles  $A : E \longrightarrow E^q$

Upon trivialization  $A(z) \in \mathfrak{gl}(N, \mathbb{C}(z))$

q-gauge transformation  $A(z) \mapsto g(qz)A(z)g^{-1}(z)$

Difference equation  $D_q(s) = As$ .


**Definition:** A meromorphic  $(\mathrm{GL}(N), q)$ -connection over  $\mathbb{P}^1$  is a pair  $(E, A)$ , where  $E$  is a (trivializable) vector bundle of rank  $N$  over  $\mathbb{P}^1$  and  $A$  is a meromorphic section of the sheaf  $\mathrm{Hom}_{\mathcal{O}_{\mathbb{P}^1}}(E, E^q)$  for which  $A(z)$  is invertible, i.e. lies in  $\mathrm{GL}(N, \mathbb{C}(z))$ . The pair  $(E, A)$  is called an  $(\mathrm{SL}(N), q)$ -connection if there exists a trivialization for which  $A(z)$  has determinant 1.

# q-Operators

**Definition:** A  $(\mathrm{GL}(2), q)$ -oper on  $\mathbb{P}^1$  is a triple  $(E, A, \mathcal{L})$ , where  $(E, A)$  is a  $(\mathrm{GL}(2), q)$ -connection and  $\mathcal{L}$  is a line subbundle such that the induced map  $\bar{A} : \mathcal{L} \longrightarrow (E/\mathcal{L})^q$  is an isomorphism. The triple is called an  $(\mathrm{SL}(2), q)$ -oper if  $(E, A)$  is an  $(\mathrm{SL}(2), q)$ -connection.

$$\text{in a trivialization} \quad s(qz) \wedge A(z)s(z) \neq 0$$

**Definition:** A  $(\mathrm{SL}(2), q)$ -oper with regular singularities at the points  $z_1, \dots, z_L \neq 0, \infty$  with weights  $k_1, \dots, k_L$  is a meromorphic  $(\mathrm{SL}(2), q)$ -oper  $(E, A, \mathcal{L})$  for which  $\bar{A}$  is an isomorphism everywhere on  $\mathbb{P}^1 \setminus \{0, \infty\}$  except at the points  $z_m, q^{-1}z_m, q^{-2}z_m, \dots, q^{-k_m+1}z_m$  for  $m \in \{1, \dots, L\}$ , where it has simple zeros.



Finally,  $(\mathrm{SL}(2), q)$ -oper is **Z-twisted** in  $A(z)$  is gauge equivalent to a diagonal matrix  $Z$

# Miura q-Operators

**Miura (SL(2),q)-oper** is a quadruple  $(E, A, \mathcal{L}, \hat{\mathcal{L}})$  where  $(E, A, \mathcal{L})$  is an (SL(2),q)-oper and  $\hat{\mathcal{L}}$  is preserved by the q-connection A

Chose trivialization of  $\mathcal{L}$

$s(z) = \begin{pmatrix} Q_+(z) \\ Q_-(z) \end{pmatrix}$

Twist element  $Z = \text{diag}(\zeta, \zeta^{-1})$

q-Oper condition — SL(2) **QQ-system**

$$\zeta Q_-(z)Q_+(zq) - \zeta^{-1}Q_-(zq)Q_+(z) = \Lambda(z)$$

singularities

One of the polynomials  
can be made monic

$$Q_+(z) = \prod_{k=1}^m (z - w_k)$$

$$\Lambda(z) = \prod_{p=1}^L \prod_{j_p=0}^{r_p-1} (z - q^{-j_p} z_p)$$

From QQ-system to Bethe equations

$$\frac{\Lambda(w_k)}{\Lambda(q^{-1}w_k)} = -\zeta^2 \frac{Q_+(qw_k)}{Q_+(q^{-1}w_k)}, \quad k = 1, \dots, m.$$

$$q^r \prod_{p=1}^L \frac{w_k - q^{1-r_p} z_p}{w_k - q z_p} = -\zeta^2 q^m \prod_{j=1}^m \frac{q w_k - w_j}{w_k - q w_j}, \quad k = 1, \dots, m$$

# q-Miura Transformation

$$A(z) = \begin{pmatrix} g(z) & \Lambda(z) \\ 0 & g(z)^{-1} \end{pmatrix}$$

Z-twisted q-oper condition

$$A(z) = v(zq)Zv(z)^{-1}, \quad Z = \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix}$$

Gauge transformation reads

$$v(z) = \begin{pmatrix} y(z) & 0 \\ 0 & y(z)^{-1} \end{pmatrix} \begin{pmatrix} 1 & -\frac{Q_-(z)}{Q_+(z)} \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} y(z) & -y(z)\frac{Q_-(z)}{Q_+(z)} \\ 0 & y(z)^{-1} \end{pmatrix}$$

We find

$$g(z) = \zeta_i y(zq)y(z)^{-1}$$

$$\Lambda(z) = y(z)y(zq) \left( \zeta \frac{Q_-(z)}{Q_+(z)} - \zeta^{-1} \frac{Q_-(zq)}{Q_+(zq)} \right)$$

The q-oper condition becomes the **SL(2) QQ-system**

$$\zeta Q_-(z)Q_+(zq) - \zeta^{-1}Q_-(zq)Q_+(z) = \Lambda(z)$$

Difference Equation

$$D_q(s) = As.$$

$$D_q(s_1) = \Lambda(z)s_2$$

after elimination

$$\left( D_q^2 - T(qz)D_q - \frac{\Lambda(qz)}{\Lambda(z)} \right) s_1 = 0$$



# tRS Hamiltonians

Recover 2-body tRS Hamiltonian from a q-Oper

Let  $Q_- = z - p_-$  and  $Q_+ = c(z - p_+)$

$$z^2 - \frac{z}{q} \left[ \frac{\zeta - q\zeta^{-1}}{\zeta - \zeta^{-1}} p_+ + \frac{q\zeta - \zeta^{-1}}{\zeta - \zeta^{-1}} p_- \right] + \frac{p_+ p_-}{q} = (z - z_+)(z - z_-)$$

qOper condition yields  
tRS Hamiltonians!

$T_1$

$T_2$

$$\det(z - L_{tRS}) = (z - z_+)(z - z_-)$$

# III. $(G,q)$ -Connection

$G$ -simple simply-connected complex Lie group

Principal  $G$ -bundle  $\mathcal{F}_G$  over  $\mathbb{P}^1$

$$M_q : \mathbb{P}^1 \rightarrow \mathbb{P}^1 \\ z \mapsto qz$$

A meromorphic  **$(G,q)$ -connection** on  $\mathcal{F}_G$  is a section  $A$  of  $\mathrm{Hom}_{\mathcal{O}_U}(\mathcal{F}_G, \mathcal{F}_G^q)$

$U$ -Zariski open dense set

Choose  $U$  so that the restriction  $\mathcal{F}_G|_U$  of  $\mathcal{F}_G$  to  $U$  is isomorphic to a trivial  $G$ -bundle

$$A(z) \in G(\mathbb{C}(z)) \quad \text{on} \quad U \cap M_q^{-1}(U)$$

Change of trivialization  $A(z) \mapsto g(qz)A(z)g(z)^{-1}$

# (G,q)-Opers

A meromorphic (G,q)-oper on  $\mathbb{P}^1$  is a triple  $(\mathcal{F}_G, A, \mathcal{F}_{B_-})$

$A$  is a meromorphic  $(G, q)$ -connection

$\mathcal{F}_{B_-}$  is a reduction of  $\mathcal{F}_G$  to  $B_-$

**Oper condition:** Restriction of the connection on some Zariski open dense set  $U$

$$A : \mathcal{F}_G \longrightarrow \mathcal{F}_G^q \text{ to } U \cap M_q^{-1}(U)$$

takes values in the *double Bruhat cell*

$$B_-(\mathbb{C}[U \cap M_q^{-1}(U)])cB_-(\mathbb{C}[U \cap M_q^{-1}(U)])$$

Coxeter element:  $c = \prod_i s_i$

Locally

$$A(z) = n'(z) \prod_i (\phi_i(z)^{\check{\alpha}_i} s_i) n(z)$$

$\phi_i(z) \in \mathbb{C}(z)$  and  $n(z), n'(z) \in N_-(z)$

# Miura $(G,q)$ -Operators

**Definition:** A *Miura  $(G, q)$ -oper* on  $\mathbb{P}^1$  is a quadruple  $(\mathcal{F}_G, A, \mathcal{F}_{B_-}, \mathcal{F}_{B_+})$ , where  $(\mathcal{F}_G, A, \mathcal{F}_{B_-})$  is a meromorphic  $(G, q)$ -oper on  $\mathbb{P}^1$  and  $\mathcal{F}_{B_+}$  is a reduction of the  $G$ -bundle  $\mathcal{F}_G$  to  $B_+$  that is preserved by the  $q$ -connection  $A$ .

It can be shown that the two flags  $\mathcal{F}_{B_-}$  and  $\mathcal{F}_{B_+}$  are in *generic relative position* for some dense set  $V$

The fiber  $\mathcal{F}_{G,x}$  of  $\mathcal{F}_G$  at  $x$  is a  $G$ -torsor with reductions  $\mathcal{F}_{B_-,x}$  and  $\mathcal{F}_{B_+,x}$  to  $B_-$  and  $B_+$ , respectively. Choose any trivialization of  $\mathcal{F}_{G,x}$ , i.e. an isomorphism of  $G$ -torsors  $\mathcal{F}_{G,x} \simeq G$ . Under this isomorphism,  $\mathcal{F}_{B_-,x}$  gets identified with  $aB_- \subset G$  and  $\mathcal{F}_{B_+,x}$  with  $bB_+$ .

Then  $a^{-1}b$  is a well-defined element of the double quotient  $B_- \backslash G / B_+$ , which is in bijection with  $W_G$ .

We will say that  $\mathcal{F}_{B_-}$  and  $\mathcal{F}_{B_+}$  have a *generic relative position* at  $x \in X$  if the element of  $W_G$  assigned to them at  $x$  is equal to 1 (this means that the corresponding element  $a^{-1}b$  belongs to the open dense Bruhat cell  $B_- \cdot B_+ \subset G$ ).



# Structure Theorems

**Theorem 1:** *For any Miura  $(G, q)$ -oper on  $\mathbb{P}^1$ , there exists a trivialization of the underlying  $G$ -bundle  $\mathcal{F}_G$  on an open dense subset of  $\mathbb{P}^1$  for which the oper  $q$ -connection has the form*

$$A(z) \in N_-(z) \prod_i ((\phi_i(z)^{\check{\alpha}_i} s_i) N_-(z) \cap B_+(z)).$$

**Theorem 2:** *Let  $F$  be any field, and fix  $\lambda_i \in F^\times, i = 1, \dots, r$ . Then every element of the set  $N_- \prod_i \lambda_i^{\check{\alpha}_i} s_i N_- \cap B_+$  can be written in the form*

$$\prod_i g_i^{\check{\alpha}_i} e^{\frac{\lambda_i t_i}{g_i} e_i}, \quad g_i \in F^\times,$$

*where each  $t_i \in F^\times$  is determined by the lifting  $s_i$ .*

# Adding Singularities and Twists

Consider family of polynomials  $\{\Lambda_i(z)\}_{i=1,\dots,r}$

**(G,q)-oper with regular singularities** can be written as

$$A(z) = n'(z) \prod_i (\Lambda_i(z)^{\check{\alpha}_i} s_i) n(z), \quad n(z), n'(z) \in N_-(z)$$

Using structure theorem every Miura (G,q)-oper with singularities reads

$$A(z) = \prod_i g_i(z)^{\check{\alpha}_i} e^{\frac{\Lambda_i(z)}{g_i(z)} e_i}, \quad g_i(z) \in \mathbb{C}(z)^\times$$

**(G,q)-oper** is **Z-twisted** if it is equivalent to a constant element of  $G$   $Z \in H \subset H(z)$   $Z$  is regular semisimple. There are  $W_G$

$$A(z) = g(qz) Z g(z)^{-1}$$

Miura (G,q)-opers for each (G,q)-opers

**Z-twisted Miura (G,q)-oper** if gauge transform is from Borel

$$A(z) = v(qz) Z v(z)^{-1}, \quad v(z) \in B_+(z)$$

# Plucker Relations

$V_i^+$ irrep of $G$ with highest weight $\omega_i$	Line $L_i \subset V_i$ stable under $B_+$
---	---

Plucker relations: for two integral dominant weights  $L_{\lambda+\mu} \subset V_{\lambda+\mu}$  is the image of  $L_{\lambda} \otimes L_{\mu} \subset V_{\lambda} \otimes V_{\mu}$  under canonical projection  $V_{\lambda} \otimes V_{\mu} \longrightarrow V_{\lambda+\mu}$

Conversely, for a collection of lines  $L_\lambda \subset V_\lambda$  satisfying Plucker relations  $\exists B \subset G$  such that  $L_\lambda$  is stabilized by  $B$  for all  $\lambda$

A choice of  $B$  is equivalent to a choice of  $B_+$ -torsor in  $G$

Let  $\nu_{\omega_i}$  be a generator of the line  $L_i \subset V_i$ . This is a vector of weight  $\omega_i$  wrt  $H \subset B_+$

The subspace of  $V_i$  of weight  $\omega_i - \alpha_i$  is one-dimensional and spanned by  $f_i \cdot \nu_{\omega_i}$

Thus the 2d subspace spanned by  $\{\nu_{\omega_i}, f_i \cdot \nu_{\omega_i}\}$  is a  $B_+$ -invariant subspace of  $V_i$

# Miura-Plucker (G,q)-Operators

let  $(\mathcal{F}_G, A, \mathcal{F}_{B_-}, \mathcal{F}_{B_+})$  be a Miura  $(G, q)$ -oper with regular singularities  $\{\Lambda_i(z)\}_{i=1, \dots, r}$

Associated vector bundle  $\mathcal{V}_i = \mathcal{F}_{B_+} \times_{B_+} V_i = \mathcal{F}_G \times_G V_i$  contains rank-two subbundle  $\mathcal{W}_i = \mathcal{F}_{B_+} \times_{B_+} W_i$

associated to  $W_i \subset V_i$ , and  $\mathcal{W}_i$  in turn contains a line subbundle  $\mathcal{L}_i = \mathcal{F}_{B_+} \times_{B_+} L_i$

Using structure theorems we obtain  $\mathbf{r}$  Miura  $(GL(2), q)$ -operators

$$A_i(z) = \begin{pmatrix} g_i(z) & \Lambda_i(z) \prod_{j>i} g_j(z)^{-a_{ji}} \\ 0 & g_i^{-1}(z) \prod_{j \neq i} g_j(z)^{-a_{ji}} \end{pmatrix}$$

**Z-twisted Miura-Plucker (G,q)-oper** is meromorphic Miura  $(G, q)$ -oper on  $P^1$  such that for each Miura  $(GL(2), q)$ -oper

$$A_i(z) = v(zq)Zv(z)^{-1}|_{W_i} = v_i(zq)Z_i v_i(z)^{-1}$$

where  $v_i(z) = v(z)|_{W_i}$  and  $Z_i = Z|_{W_i}$

# QQ-System

**Theorem:**

*There is a one-to-one correspondence between the set of nondegenerate  $Z$ -twisted Miura-Plücker  $(G, q)$ -opers and the set of nondegenerate polynomial solutions of the QQ-system*

$$\begin{aligned} \tilde{\xi}_i Q_-^i(z) Q_+^i(qz) - \xi_i Q_-^i(qz) Q_+^i(z) = \\ \Lambda_i(z) \prod_{j>i} \left[ Q_+^j(qz) \right]^{-a_{ji}} \prod_{j<i} \left[ Q_+^j(z) \right]^{-a_{ji}}, \quad i = 1, \dots, r, \end{aligned}$$

$$\tilde{\xi}_i = \zeta_i \prod_{j>i} \zeta_j^{a_{ji}}, \quad \xi_i = \zeta_i^{-1} \prod_{j<i} \zeta_j^{-a_{ji}}$$

Proof uses

$$v(z) = \prod_{i=1}^r y_i(z)^{\check{\alpha}_i} \prod_{i=1}^r e^{-\frac{Q_-^i(z)}{Q_+^i(z)} e_i} \dots, \qquad g_i(z) = \zeta_i \frac{Q_+^i(qz)}{Q_+^i(z)}.$$



# XXZ Bethe Ansatz Equations for G

roots of Q+

$$\frac{Q_+^i(qw_i^k)}{Q_+^i(q^{-1}w_i^k)} \prod_j \zeta_j^{a_{ji}} = - \frac{\Lambda_i(w_k^i) \prod_{j>i} [Q_+^j(qw_k^i)]^{-a_{ji}} \prod_{j<i} [Q_+^j(w_k^i)]^{-a_{ji}}}{\Lambda_i(q^{-1}w_k^i) \prod_{j>i} [Q_+^j(w_k^i)]^{-a_{ji}} \prod_{j<i} [Q_+^j(q^{-1}w_k^i)]^{-a_{ji}}}$$

Space of nondegenerate solutions of  
QQ-system for G

Nondegenerate **Z-twisted Miura-Plucker** (G,q)-opers  
with regular singularities



?

Space of nondegenerate solutions of  
XXZ for G

?

Nondegenerate **Z-twisted Miura** (G,q)-opers  
with regular singularities



# Quantum Backlund Transformation

**Theorem:** Consider the following q-gauge transformation

$$A \mapsto A^{(i)} = e^{\mu_i(qz)f_i} A(z) e^{-\mu_i(z)f_i}, \quad \text{where} \quad \mu_i(z) = \frac{\prod_{j \neq i} [Q_+^j(z)]^{-a_{ji}}}{Q_+^i(z) Q_-^i(z)}$$

changes the set  
of Q-functions

$$\begin{aligned} Q_+^j(z) &\mapsto Q_+^j(z), & j \neq i, \\ Q_+^i(z) &\mapsto Q_-^i(z), & Z \mapsto s_i(Z) \end{aligned}$$

$$\begin{aligned} \{\tilde{Q}_+^j\}_{j=1,\dots,r} &= \{Q_+^1, \dots, Q_+^{i-1}, Q_-^i, Q_+^{i+1}, \dots, Q_+^r\} \\ \{\tilde{z}_j\}_{j=1,\dots,r} &= \{z_1, \dots, z_{i-1}, z_i^{-1} \prod_{j=i+1}^r z_j^{-a_{ji}}, \dots, z_r\} \end{aligned}$$

Now the strategy is to successively apply Backlund transformations according to the reduced decomposition of the element of the Weyl group

Consider longest element  $w_0 = s_{i_1} \dots s_{i_\ell}$

**Theorem:** Every Z-twisted Miura-Plucker (G,q)-oper is Z-twisted Miura (G,q)-oper

The proof based on properties of double Bruhat cells addresses existence of the diagonalizing element v(z) (to be constructed later)

# (SL(N),q)-Operators

The QQ-system

$$\xi_i \phi_i(z) - \xi_{i+1} \phi_i(qz) = \rho_i(z)$$

$$\phi_i(z) = \frac{Q_i^-(z)}{Q_i^+(z)}, \qquad \rho_i(z) = \Lambda_i(z) \frac{Q_{i-1}^+(qz) Q_{i+1}^+(z)}{Q_i^+(z) Q_i^+(qz)}$$

q-Oper condition

$$v(qz)^{-1} A(z) = Z v(z)^{-1}$$

Diagonalizing element

Polynomials  $Q_{i,\dots,j}^-(z)$

form extended QQ-system

$$v(z)^{-1} = \left( \begin{array}{cccccc} \frac{1}{Q_1^+(z)} & \frac{Q_1^-(z)}{Q_2^+(z)} & \frac{Q_{12}^-(z)}{Q_3^+(z)} & \cdots & \frac{Q_{1,\dots,r-1}^-(z)}{Q_r^+(z)} & Q_{1,\dots,r}^-(z) \\ 0 & \frac{Q_1^+(z)}{Q_2^+(z)} & \frac{Q_2^-(z)}{Q_3^+(z)} & \cdots & \frac{Q_{2,\dots,r-1}^-(z)}{Q_r^+(z)} & Q_{2,\dots,r}^-(z) \\ 0 & 0 & \frac{Q_2^+(z)}{Q_3^+(z)} & \cdots & \frac{Q_{3,\dots,r-1}^-(z)}{Q_r^+(z)} & Q_{3,\dots,r}^-(z) \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & \cdots & \cdots & \frac{Q_{r-1}^+(z)}{Q_r^+(z)} & Q_r^-(z) \\ 0 & \cdots & \cdots & \cdots & 0 & Q_r^+(z) \end{array} \right)$$