

Moment Bounds for the Smoluchowski Equation and their Consequences

Alan Hammond*
New York University
and
Fraydoun Rezakhanlou†
UC Berkeley

November 24, 2009

Abstract

We prove $L^\infty(\mathbb{R}^d \times [0, \infty))$ bounds on moments $X_a := \sum_{m \in \mathbb{N}} m^a f_m(x, t)$ of the Smoluchowski coagulation equations with diffusion, in any dimension $d \geq 1$. If the collision propensities $\alpha(n, m)$ of mass n and mass m particles grow more slowly than $(n+m)(d(n)+d(m))$, and the diffusion rate $d(\cdot)$ is non-increasing and satisfies $m^{-b_1} \leq d(m) \leq m^{-b_2}$ for some b_1 and b_2 satisfying $0 \leq b_2 < b_1 < \infty$, then any weak solution satisfies $X_a \in L^\infty(\mathbb{R}^d \times [0, T]) \cap L^1(\mathbb{R}^d \times [0, T])$ for every $a \in \mathbb{N}$ and $T \in (0, \infty)$, (provided that certain moments of the initial data are finite). As a consequence, we infer that these conditions are sufficient to ensure uniqueness of a weak solution and its conservation of mass.

1 Introduction

The Smoluchowski coagulation equation is a coupled system of partial differential equations that describes the evolving densities of a system of diffusing particles that are prone to coagulate in pairs. A sequence of functions $f_n : \mathbb{R}^d \times [0, \infty) \rightarrow [0, \infty)$, $n \in \mathbb{N}$, is a solution of the Smoluchowski coagulation equation if it satisfies

$$(1.1) \quad \frac{\partial}{\partial t} f_n(x, t) = d(n) \Delta f_n(x, t) + Q_n(f)(x, t),$$

*This work was performed while A.H. held a postdoctoral fellowship in the Department of Mathematics at U.B.C.

†This work is supported in part by NSF grant DMS0307021

with $Q_n(f) = Q_n^+(f) - Q_n^-(f)$, where

$$Q_n^+(f)(x, t) = \sum_{m=1}^{n-1} \alpha(m, n-m) f_m(x, t) f_{n-m}(x, t)$$

and

$$Q_n^-(f)(x, t) = 2f_n(x, t) \sum_{m=1}^{\infty} \alpha(n, m) f_m(x, t).$$

We will interpret this solution in weak sense. Namely, we will assume that Q_n^+ and Q_n^- belong to $L^1(\mathbb{R}^d \times [0, T])$ for each $T \in [0, \infty)$ and $n \in \mathbb{N}$, and that

$$f_n(x, t) = S_t^{d(n)} f_n^0(x) + \int_0^t S_{t-s}^{d(n)} Q_n(x, s) ds,$$

where $\{f_n^0 : n \in \mathbb{N}\}$ denotes the initial data, S_t^D the semigroup associated with the equation $u_t = D\Delta u$, and where $Q_n(x, s)$ means $Q_n(f)(x, s)$.

The system (1.1) has two sets of parameter values, the sequence $d : \mathbb{N} \rightarrow [0, \infty)$, where $d(n)$ denotes the diffusion rate of the Brownian particle of mass n , and the collection $\alpha : \mathbb{N}^2 \rightarrow [0, \infty)$, where $\alpha(m, n)$ models the average propensity of particles of masses m and n to coagulate. The terms $Q_n^+(f)$ and $Q_n^-(f)$ are gain and loss terms for the presence of particles of mass n that arise from the binary coagulation of particles.

The system (1.1) may be augmented by considering the fragmentation of particles into two or more sub-particles. A continuous version of the system, in which particles have real (rather than integer) mass, has also been considered. The data are defined $d : [0, \infty) \rightarrow [0, \infty)$ and $\alpha : [0, \infty)^2 \rightarrow [0, \infty)$, and the sums are replaced integrals in the definitions of $Q_n^+(f)$ and $Q_n^-(f)$. The spatially homogeneous version of the equations, in which each f_n is a function of time alone, are better understood (the work [1] resolved many of the central questions for the discrete case without fragmentation).

Lang and Nguyen [9] considered a system of mass-bearing diffusing particles, whose diffusion rate was chosen to be independent of the mass, that are prone to coagulate in pairs at close range, and demonstrated that, in a kinetic limit, the density of particles evolves macroscopically as a solution of (1.1). A kinetic limit means that, with an initial number N of particles, the order of the interaction range $\epsilon = \epsilon(N)$ of any given particle is chosen so that a typical particle experiences a rate of collision that is bounded away from zero and infinity for all high N . In [7], [8], this kinetic limit derivation was extended to permit more general diffusion rates, and to include a stochastic mechanism of interaction. Suppose that particles of mass $n \in \mathbb{N}$ have a range of interaction given by $r(n)\epsilon$, for some increasing function $r : \mathbb{N} \rightarrow (0, \infty)$. That is, $\epsilon = \epsilon(N)$ determines the order of the range as a function of the initial particle number, whereas the function r specifies the relative interaction range of particles of differing masses. The variations to the derivation required

by introducing the radial dependence r have been discussed in [12]. It is shown there that, if the dimension $d \geq 3$, the macroscopic collision propensity $\alpha : \mathbb{N}^2 \rightarrow (0, \infty)$ that appears in the Smoluchowski coagulation equation satisfied by the macroscopic density profile of the particle system satisfies

$$(1.2) \quad \alpha(n, m) \leq c(d(n) + d(m)) \left(r(n) + r(m) \right)^{d-2},$$

where c is a constant without dependence on the details of the stochastic interaction. (Note that β rather than α was the notation used in [7],[8]. The change in notation in this paper ensures consistency with the PDE literature.) In fact, the left and right-hand-sides of (1.2) are of the same order provided that the interaction mechanism is strong enough to ensure that a uniformly positive fraction of pairs of particles that come within the interaction range coagulate.

The growth rate of $r : \mathbb{N} \rightarrow (0, \infty)$ presumably depends on the internal structure of the particles. If they are simply balls, each of the same density, then $r(n) = c_0 n^{1/d}$. On the other hand, an internal structure that is fractal might give rise to a relation of the form $r(n) = n^\chi$, for some $\chi \geq 1/d$. The physically reasonable range of values of χ would seem to be contained in $[1/d, 1]$. This is because, whatever fractal structure a particle of mass $n \in \mathbb{N}$ may have, if contact with another particle is required for the pair to coagulate, then the interaction range $r(n)$ is at most of order n , (with the extremal case being that in which the particle takes the form of a line segment). Indeed, non-trivial fractal structure would suggest that χ is strictly less than one.

Monotonically decreasing choices of the diffusion rates $d : \mathbb{N} \rightarrow [0, \infty)$ seem to be physically realistic, since the diffusive motion is presumably stimulated by the bombardment of much smaller elements of an ambient gas. The choice $d(n) = \frac{1}{n}$ in three dimensional space is justified if the particles are modelled as balls. (For a mathematical treatment, see [4], which derives Brownian motion as the long-term behaviour of a ball being struck by elements in a Poisson cloud of point particles of Gaussian velocity.)

In this paper, we examine the behaviour of solutions of (1.1). We have directed our attention to parameter values α and d that seem to be justified by the existing kinetic limit derivations of (1.1) (though, in principle, other choices may arise from a derivation of the equations from a quite different model).

We mention firstly that, under the assumption

$$(1.3) \quad \lim_{m \rightarrow \infty} \frac{\alpha(n, m)}{m} = 0 \quad \text{for each } n \in \mathbb{N},$$

the global existence of a weak solution of (1.1) has been established in a bounded domain in [11] and for \mathbb{R}^d in [15], each of these works including fragmentation in the equations. From the physically reasonable assumptions that $d(\cdot)$ is uniformly bounded and $r(n) = o(n)$, we see that (1.3) is satisfied in dimension $d = 3$ by choices of α satisfying (1.2). (We mention

also that, using an assumption analogous to (1.3) as well as either a monotonicity condition on the coagulation rates or a detailed balance condition between the rates of coagulation and fragmentation, [10] proves the global

existence of a weak solution in the case of continuous mass.)

An important formal property of solutions of (1.1) is mass-conservation.

Definition 1.1 *Let f solve (1.1) weakly. We say that f conserves mass on the time interval $[0, T]$ provided that $I(t) = I(0)$ for each $t \in [0, T]$, where*

$$I(t) = \sum_{m \in \mathbb{N}} m \int_{\mathbb{R}^d} f_m(x, t) dx.$$

While mass conservation holds formally, the only estimate that is readily available is $I(t) \leq I(0)$ for $t \geq 0$. Indeed, the inequality may be strict, in which case, *gelation* is said to occur (at the infimum of times at which the inequality is strict). If this happens for a solution of (1.1) which is obtained as a kinetic limit of a particle system, then, after the gelation time, a positive fraction of the mass of particles is contained in particles whose mass is greater than some function that grows to infinity as the initial particle number tends to infinity.

In the spatially homogeneous setting, much progress has been made in establishing when gelation occurs. In [6] and [5], continuous versions of the spatially homogeneous equations with fragmentation are considered. If the continuous analogue $\alpha(x, y)$ of the coagulation rates α is supposed to satisfy

$$(1.4) \quad \alpha(x, y) = x^a y^b + x^b y^a, \quad x, y \in \mathbb{R}$$

with $a, b \in (0, 1)$ and $a + b > 1$, then gelation occurs, unless the inhibiting effect of fragmentation is strong enough. It is natural to postulate from these results that if the microscopic interaction range $r(n)$ we have discussed behaves like $r(n) = n^\chi$ with $\chi > \frac{1}{d-2}$, with a diffusion rate d uniformly bounded below and a bounded domain Ω in place of \mathbb{R}^d , then gelation will occur. This is because the formula (1.2), (which, as already noted, may be written as an equality in the case of a reaction mechanism that is not particularly weak), is bounded below by the discrete analogue of the coagulation propensity given in (1.4). As we have commented, however, we do not anticipate such behaviour for α in equations arising from a three-dimensional particle system of the type considered in [12]. Moreover, as discussed in [12], the derivation of (1.1) from a random model performed in [7] is valid only if $\chi < \frac{1}{d-2}$: it is thus merely a conjecture that gelation occurs if $\chi > \frac{1}{d-2}$.

Rigorous sufficient conditions for mass-conservation, or for gelation, have been available in the spatially inhomogeneous setting only under stringent assumptions on parameters. See [3] for the case of constant diffusion rates, and [14] for a criterion that requires uniform boundedness of α and further information about the behaviour of solutions of the system

(however, each of these papers includes fragmentation in the equations). In Theorem 1.3, we present a more applicable sufficient condition.

The result largely depends on new moment bounds on solutions of (1.1). Theorem 1.1 presents bounds on the L^1 norm of moments of a solution, and Theorem 1.2 provides L^∞ bounds on such moments. Previously, L^∞ estimates on solutions of (1.1), with the effect of fragmentation included, have been obtained (see [13],[14],[2],[3]), under fairly restrictive assumptions on coefficients for coagulation and fragmentation propensity. The dependence on $i \in \mathbb{N}$ of the bounds obtained on L^∞ norms of f_i does not generally permit deductions about the L^∞ -norm of moments $\sum_{m=1}^\infty m^a f_m(x, t)$ for any $a \geq 0$ (note however that such inferences are made in [3] if the diffusion rate is identically constant, or in [13] if the coagulation rates $\alpha(n, m)$ decay quickly enough).

Our final result, Theorem 1.4, provides a criterion for uniqueness of solutions of (1.1). It also relies principally on the moment bounds and is a straightforward adaptation of the uniqueness proof of [1] which applies to the homogeneous case. To our knowledge, the uniqueness question has been largely unsettled: [14] treats the one-dimensional case including fragmentation under quite restrictive hypotheses, while [2] treats the case where $d(n)$ is eventually constant, with a specific choice of $\alpha(n, m)$ whose growth is of the order of nm .

We provide a corollary after the theorems that summarises the principal conditions on the coagulation and diffusion coefficients, and on the initial data, under which we may infer uniqueness and mass conservation of the solution.

Our results are valid for each dimension $d \geq 1$. Each deduction, moment bound, mass conservation, or uniqueness, depends on some regularity in the initial data, and some assumption on the parameters of the system. We now state the various assumptions that we require.

Hypothesis 1.1

$$\lim_{n+m \rightarrow \infty} \frac{\alpha(n, m)}{(n+m)(d(n) + d(m))} = 0.$$

More precisely, for every $\delta > 0$, there exists $k_0 = k_0(\delta) > 0$ such that if $n + m > k_0$, then

$$\alpha(n, m) \leq \delta(n+m)(d(n) + d(m)).$$

In addition, the function d is uniformly bounded.

Hypothesis 1.2 The function $d(n)$ is a non-increasing function of n and that $\alpha(n, m) \leq C_0(n+m)$ for a constant C_0 . Moreover, there exist positive constants r_1 and r_2 and nonnegative constants $b_2 \leq b_1$ such that,

$$r_1 n^{-b_1} \leq d(n) \leq r_2 n^{-b_2}.$$

Hypothesis 1.3 The function d is uniformly positive and non-increasing, and there exists a constant C_0 such that

$$\alpha(n, m) \leq C_0(n+m).$$

Our notation for the various moments of f will be

$$(1.5) \quad X_a = X_a(x, t) = \sum_n n^a f_n(x, t), \quad \hat{X}_a = \hat{X}_a(x, t) = \sum_n n^a d(n)^{d/2} f_n(x, t).$$

and

$$(1.6) \quad \begin{aligned} Y_a(x, t) &= \sum_{n,m} nm(n^a + m^a)(d(n) + d(m))f_n(x, t)f_m(x, t), \\ \hat{Y}_a(x, t) &= \sum_{n,m} (n^a m + m^a n)\alpha(n, m)f_n f_m. \end{aligned}$$

We also set

$$(1.7) \quad \phi_0(x) = \begin{cases} |x|^{2-d} & \text{if } d \geq 3, \\ -\frac{1}{2\pi} \log |x| \mathbb{1}(|x| \leq 1) & \text{if } d = 2, \\ \frac{1}{2}(1 - |x|) \mathbb{1}(2|x| \leq 1) & \text{if } d = 1. \end{cases}$$

We now state the four theorems.

Theorem 1.1 (Moment bound I) *Assume Hypothesis 1.1. Then for every $a \geq 2$ and positive A and T , there exists a constant $C = C(a, A, T)$ such that, if*

$$(1.8) \quad \iint \sum_{n,m} X_a(x, 0)X_1(y, 0)\phi_0(x - y)dxdy \leq A,$$

and

$$(1.9) \quad \operatorname{ess\,sup}_x \int X_a(y, 0)\phi_0(x - y)dy \leq A, \quad \int X_a(x, 0)dx \leq A,$$

then

$$(1.10) \quad \int_0^T \int Y_{a-1}dxdt \leq C, \quad \int_0^T \int \hat{Y}_{a-1}dxdt \leq C,$$

and

$$(1.11) \quad \sup_{t \in [0, T]} \int X_a(x, t)dx \leq C.$$

Moreover, the constant C can be chosen to be independent of T when $d > 2$.

Theorem 1.2 (Moment bound II) *Assume Hypothesis 1.2. Then there exists a function $\gamma(a, b_1, b_2)$ with $\lim_{a \rightarrow \infty} \gamma(a, b_1, b_2) = \infty$ such that if*

$$(1.12) \quad \sum_n n^e \|f_n^0\|_{L^\infty(\mathbb{R}^d)} < \infty, \quad X_a \in L^1(\mathbb{R}^d \times [0, T]),$$

then

$$(1.13) \quad \sum_n n^e \|f_n\|_{L^\infty(\mathbb{R}^d \times [0, T])} < \infty,$$

for every $e \leq \gamma(a, b_1, b_2)$.

Remark 1.1 In particular, if Hypotheses 1.1–1.2 hold, $X_a(\cdot, 0) \in L^1(\mathbb{R}^d)$ and

$$\sum_n n^a \|f_n^0\|_{L^\infty(\mathbb{R}^d)} < \infty,$$

for every $a \in \mathbb{N}$, then $X_a \in L^\infty(\mathbb{R}^d \times [0, T]) \cap L^1(\mathbb{R}^d \times [0, T])$ for every $a \in \mathbb{N}$ and $T \in (0, \infty)$.

We refer to (4.15) for the explicit form of the function γ . Also note that Theorem 1.1 offers sufficient conditions to ensure $X_a \in L^1(\mathbb{R}^d \times [0, T])$ (so that this theorem has been invoked in Remark 1.1).

Theorem 1.3 (Conservation of Mass) *Let f be a weak solution of (1.1). Assume that (1.10) holds for $a = 2$. Then f conserves mass on the time interval $[0, T]$. Assume instead that $X_1(\cdot, 0) \in L^\infty(\mathbb{R}^d)$, $X_2(\cdot, 0) \in L^1(\mathbb{R}^d)$ and Hypothesis 1.3 holds. Then f conserves mass on the time interval $[0, \infty)$.*

Theorem 1.4 (Uniqueness) *Suppose that, for some $c_0 > 0$, we have that $\alpha(n, m) \leq c_0 nm$ for each $n, m \in \mathbb{N}$. Then there is a unique weak solution of (1.1) on the interval $[0, T]$ among those satisfying $X_2 \in L^\infty(\mathbb{R}^d \times [0, T])$.*

The following corollary states an important consequence of our results.

Corollary 1.1 *Assume that there exist positive constants a, b, c_1 and c_2 , with $a + b < 1$, such that $\alpha(n, m) \leq c_1(n^a + m^a)$ and $d(n) \geq c_2 n^{-b}$ for all $n, m \in \mathbb{N}$. If $\sum_n n^e \|f_n^0\|_{L^\infty(\mathbb{R}^d)} < \infty$ and $\|\sum_n n^e f_n^0\|_{L^1(\mathbb{R}^d)} < \infty$ for sufficiently large e , then (1.1) has a unique solution, and this solution is mass conserving.*

Remark 1.2 Note that Hypothesis 1.1 implies the existence of a weak solution, because (1.3) holds in this case, whereas Hypotheses 1.2 and 1.3 do not. The solution f referred to in Theorems 1.2 and 1.3 is thus not known to exist under the hypotheses of these results, and it should be considered that these results make no assertion if the choice of parameters

α and d is such that no solution exists.

Remark 1.3 Theorem 1.1 is also true for the continuous version of the system (1.1) with a verbatim proof. In the continuous version, all the summations over n and m are replaced with integrations with respect to dn and dm . On the other hand, we can derive the continuous version of Theorems 1.2 and 1.3 only for a particular solution. The proof of Theorem 1.4 does not readily adapt to the continuous setting. See Remark 4.1 for further comments.

If the parameters α and d derived from the kinetic limit of a particle system are such that uniqueness among solutions of (1.1) is unknown, then, in principle at least, the particle system may not approximate a single solution of (1.1). It might, for example, approximate several different solutions, each with a positive probability. This would be very peculiar, and so, it is pleasing to be able to rule out the possibility by establishing uniqueness.

If we adopt the relation $r(n) = n^\chi$ for particle interaction range, then, recalling (1.2), we have that, in \mathbb{R}^d with $d \geq 3$, the macroscopic coagulation propensity arising from the microscopic random model satisfies

$$\alpha(n, m) \leq c(d(n) + d(m)) \max\{n, m\}^{\chi(d-2)}.$$

In view of Theorems 1.3 and 1.4, we have a unique solution and mass conservation throughout time provided that $\chi \in [0, \frac{1}{d-2})$ and suitable moments of the initial densities are finite. As such, the discussion following (1.2) rules out the occurrence of gelation in three dimensions in particle systems of the type considered in [7], [8] and [12].

In a similar vein to the comment that follows (1.4), we mention that, in the case where d is uniformly positive, Theorem 1.3 gives a sufficient condition for mass conservation of a solution of (1.1) that is close to being sharp. Indeed, choices of $\alpha(n, m)$ that grow much more quickly than Hypothesis 1.2 permits are bounded below by the expression in (1.4), for some choice of $a, b \in (0, 1)$ with $a + b > 1$. Such a choice of α would thus be expected to show gelation in the spatially homogeneous case. Corollary 8.2 of [6] adapts the argument of the homogeneous case to assert that gelation occurs for any weak solution of the continuous version of (1.1) in a bounded subset of \mathbb{R}^d , provided that (1.4) is satisfied for such a and b as above. The diffusive motion of particles in \mathbb{R}^d may act to inhibit gelation of a solution of (1.1), though a dense initial condition is likely to ensure it.

Acknowledgment. We thank Gábor Pete for comments on a draft version.

2 The tracer particle approach

Each of the L^∞ and moment bounds on solutions of (1.1) that we present in this paper will be proved by PDE methods. However, for each of our results, we earlier derived a similar assertion by a quite different approach. A given solution of (1.1) is understood in terms of the random trajectory of a tracer particle, whose behaviour is typical of the many particles that form the density profile of the solution. We have certainly found this random method

to be intuitively appealing, and it may find application to other PDE for which kinetic limit derivations have been made. We have thus devoted this section to explaining the tracer particle approach.

It will in fact be helpful to recall in more detail the model analysed in [7]. A sequence of microscopic random models indexed by their initial number $N \in \mathbb{N}$ of particles is given. In the N -th model, each of the N particles has an initial location $x(0)$ and integer mass $m(0)$ set independently according to

$$(2.1) \quad \mathbb{P}(m(0) = k) = \frac{\int_{\mathbb{R}^d} f_k^0(x) dx}{\sum_{n=1}^{\infty} \int_{\mathbb{R}^d} f_n^0(x) dx},$$

with $x(0)$ having law

$$(2.2) \quad \frac{f_k^0(\cdot)}{\int_{\mathbb{R}^d} f_k^0(x) dx},$$

conditional on $m(0) = k$. At any given moment of time $t \in [0, \infty)$, particles of mass $k \in \mathbb{N}$ evolve as Brownian motions with diffusion rate $d(k)$, with $d : \mathbb{N} \rightarrow (0, \infty)$ a given collection of constants. Particles are liable to coagulate in pairs when their displacement is of order $\epsilon = \epsilon(N)$ (we refer the reader to the introduction of [7] for the details of the interaction mechanism). We set $\epsilon \approx N^{-\frac{1}{d-2}}$ (for $d \geq 3$), to ensure that a typical particle experiences a rate of collision that is bounded away from zero and ∞ uniformly in N . At the collision event, the two incoming particles disappear, to be replaced by a third, that assumes the sum of the masses of the ingoing two, and is located in an ϵ -vicinity of either of the colliding particles. While the macroscopic behaviour of the system is likely to be independent of the choice of placement of the new particle within this microscopic vicinity of the colliding pair, it was convenient for the derivation performed in [7] to assume that, if the masses of the colliding pair are n and m , then the location of the new particle is taken to be that of one or other of the pair with probabilities $\frac{n}{n+m}$ and $\frac{m}{n+m}$. For what follows, it is convenient to regard a particle that has a collision as surviving it, and becoming the outgoing particle, with a probability proportional to its mass, and disappearing from the model in the other event.

Theorem 1 of [7] specifies the macroscopic behaviour of this particle system, when the initial particle number N is taken to be high. To summarise the result without recourse to equations, at typical points $(x, t) \in \mathbb{R}^d \times [0, \infty)$ of space-time, the number of particles of given mass $m \in \mathbb{N}$ located in a vicinity of x at time t , normalized appropriately, approximates the *density* $f_m(x, t)$, for some solution $\{f_m : m \in \mathbb{N}\}$ of (1.1). From the result, we may infer the law of the trajectory of a typical particle, in the limit of high initial particle number, as follows.

Consider a particle picked uniformly at random from the N particles that are initially present. We call this the tracer particle in the N -th model. In accordance with the preceding

description, the initial mass $m(0)$ and location $x(0)$ of the particle are given by (2.1) and (2.2). We know from [7] that $\{f_n : \mathbb{R}^d \times [0, \infty) \rightarrow [0, \infty), n \in \mathbb{N}\}$ gives the asymptotic particle densities of the microscopic models. Hence, asymptotically in high N , the trajectory of the particle is such that, if at time $t \in [0, \infty)$, the particle has location $x \in \mathbb{R}^d$ and mass m , it evolves as a Brownian motion at rate $d(m)$, and experiences collision with a particle of mass n at rate $\alpha(m, n)f_n(x, t)$. The details of the collision event that we specified imply that the tracer particle $(x(t), m(t))$, on colliding with a mass n particle, survives the collision and assumes mass $m(t^+) = m(t) + n$ with probability $\frac{m(t)}{m(t)+n}$, and disappears from the model otherwise.

The limit in high N of the law of the tracer particle in the N -th model is in fact dependent only on the given solution $\{f_n : n \in \mathbb{N}\}$ that the microscopic particle densities approximate, and not on other details of the random models, such as the location of other particles. We now formalise this definition: a tracer particle specified in terms of a given solution $\{f_n : n \in \mathbb{N}\}$ of (1.1) and not in terms of the data of any microscopic model.

Definition 2.1 *Let a solution $\{f_n : n \in \mathbb{N}\}$ of (1.1) be given. The tracer particle governed by f the random process $z = (x, m) : [0, \infty) \rightarrow \mathbb{R}^d \times \mathbb{N} \cup \{c\}$ whose initial law is given by*

$$\mathbb{P}(m(0) = m) = \frac{\int_{\mathbb{R}^d} f_m^0(x) dx}{\sum_{n=1}^{\infty} \int_{\mathbb{R}^d} f_n^0(x) dx},$$

with the initial location $x(0)$ having density $f_m^0(\cdot) / \int_{\mathbb{R}^d} f_m^0(x) dx$, conditional on $m(0) = m$. At time t , the particle's location $x(t)$ evolves as a Brownian motion at rate $d(m)$, provided that $m(t) = m$. Moreover, at time t , and for any $n \in \mathbb{N}$, the particle is said to undertake a mass transition $m \rightarrow n + m$ at rate $\alpha(n, m)f_n(x(t), t)$. Such a transition succeeds with probability $m/(n+m)$, in which case, $m(t)$ is set equal to $n+m$, and fails in the other event, in which case, the particle is relegated to the cemetery state c : $z(s)$ is set equal to c for all $s \geq t$.

Remark 2.1 Certain smoothness (or measurability) assumptions are in fact required to ensure that the process $z = (x, m) : [0, \infty) \rightarrow \mathbb{R}^d \times \mathbb{N} \cup \{c\}$ exists, even locally in time.

Let $z : [0, \infty) \rightarrow \mathbb{R}^d \times \mathbb{N} \cup \{c\}$ denote the tracer particle governed by a given solution $\{f_n : n \in \mathbb{N}\}$ of (1.1). Let $g_n(x, t)$ denote the density of its location at time t :

$$\mathbb{P}(x(t) \in A, m(t) = n) = \int_A g_n(x, t) dx.$$

The evolution equation for the system $\{g_n : n \in \mathbb{N}\}$ is given by

$$(2.3) \quad \begin{aligned} \frac{\partial}{\partial t} g_n(x, t) &= d(n) \Delta g_n(x, t) \\ &+ \sum_{m=1}^{n-1} \alpha(m, n-m) f_m(x, t) g_{n-m}(x, t) - 2g_n(x, t) \sum_{m=1}^{\infty} \alpha(n, m) f_m(x, t). \end{aligned}$$

Note that the choice $g_n \equiv f_n$ solves this equation. Assuming that there is a unique non-zero solution of (2.3), we find that

$$(2.4) \quad g_n \equiv f_n.$$

Before commenting further on (2.4), we want to emphasise how we have changed our point of view of the tracer particle. At first, we regarded it as a typical particle in a microscopic random model, whose behaviour in the large is described by a solution of (1.1), and then, secondly, as a random trajectory defined purely in terms of such a solution. The former point of view motivates the study of the system (1.1). We will now discuss how the latter is valuable in studying a given solution of (1.1). Indeed, that the tracer particle defined by a given solution of (1.1) may be a useful tool for studying that solution is apparent from (2.4): if we understand the likely behaviour of the tracer particle, we infer bounds on the density $\{g_n : n \in \mathbb{N}\}$ of its location, and, by (2.4), on the solution $\{f_n : n \in \mathbb{N}\}$ itself.

We have stated the relation (2.4) because doing so permits a more succinct summary of how the tracer particle approach works. However, in making the approach rigorous, we take a different route, which we now summarise. We construct a sequence $\{f_n^N : \mathbb{R}^d \times [0, \infty) \rightarrow [0, \infty), n \in \mathbb{N}\}$, indexed by N , which will approximate the density of the tracer particle of some solution of (1.1) when N is high. For each $N \in \mathbb{N}$, the functions f_n^N , for $n \in \mathbb{N}$, are constructed inductively, on the domains $\mathbb{R}^d \times [0, \frac{i}{N})$, for each $i \in \mathbb{N}$. They are extended to $\mathbb{R}^d \times [0, \frac{i+1}{N})$ by an inductive step in which $f_n^N(t)$ for $n \in \mathbb{N}$ and $t \in [\frac{i}{N}, \frac{i+1}{N})$ is defined as the density of the location of a tracer particle governed by the constant data $f_n^N(i/N)$ during the time interval $[\frac{i}{N}, t)$. For each $i \in \mathbb{N}$ and $T \in [0, \infty)$, the function f_n^N is shown to converge in $L^1(\mathbb{R}^d \times [0, T])$ as $N \rightarrow \infty$ to a limit $f_n : \mathbb{R}^d \times [0, \infty) \rightarrow [0, \infty)$. By mimicking the proof of the weak stability result of [11], we infer that the sequence $\{f_n : n \in \mathbb{N}\}$ is a solution of (1.1). Tracer particle arguments, of which an example will shortly be given, are then applied directly to the tracer particle densities $\{f_n^N : n \in \mathbb{N}\}$, and the resulting bounds, which hold uniformly in N , are inherited in the high N limit by the solution $\{f_n : n \in \mathbb{N}\}$. This method for making the tracer particle approach rigorous does have the drawback of applying to only one solution of (1.1).

How may the tracer particle approach be used to prove an L^∞ -bound on the solution $\{f_n : n \in \mathbb{N}\}$ of (1.1) whose construction we have just discussed? To simplify the exposition, suppose that the initial condition takes the form $\int_{\mathbb{R}^d} f_1^0(x) dx = 1$, $f_m^0(x) = 0$ for $x \in \mathbb{R}^d$ and $m > 1$. We will now sketch a proof of the following reformulation of Lemma 4.1. Suppose that $d : \mathbb{N} \rightarrow (0, \infty)$ is decreasing, and that $d(m) > cm^{-\frac{2}{d}(1-\alpha)}$ for some $\alpha \in [0, 1]$ and $c > 0$. Then

$$(2.5) \quad \sum_{m=1}^{\infty} m^\alpha f_m(x, t) \leq Cu(x, t),$$

where $u : \mathbb{R}^d \times [0, \infty) \rightarrow [0, \infty)$ solves,

$$\frac{\partial u}{\partial t} = d(1)\Delta u,$$

$$u(x, 0) = f_1(x, 0).$$

The solution $\{f_n : n \in \mathbb{N}\}$ of (1.1), being given by the density of the tracer particle (2.4), has the following interpretation: for $A \subseteq \mathbb{R}^d$ a Borel set, the quantity

$$\sum_{m=1}^{\infty} m^\alpha \int_A f_m(x, t) dx$$

is equal to the expected value of the random variable $R = R(A)$ equal to m^α if the tracer particle at time t has mass m and lies inside the set A .

In comparing the left- and right-hand-sides of (2.5), we are thus assessing the degree to which the dynamics of the tracer particle may increase the expected values of the random variables $R(A)$ over those obtained by using a simple Brownian particle (for numerous choices of the set A). For example, if the set A is a small ball about $x \in \mathbb{R}^d$, and the tracer particle is close to x at some time s satisfying $0 < s \ll t$, then a mass transition undertaken by the tracer particle at times shortly after s will serve to increase the expected value of R , because the slower diffusion rate produced by the transition is more likely to leave the particle nearby to x at the later time t . However, if a transition occurs that sharply increases the mass of the particle, then it is likely to fail, in which case, it contributes zero to the expected value of R . This latter effect limits the capacity of the tracer particle to focus towards x .

Phrasing the question quantitatively, we ask: what is the expected value of $R(A)$ at time $s < t$ if a mass transition $m_1 \rightarrow m_2$ occurs at time s ? Assuming that there is no other mass transition, and supposing that the tracer particle is at $y \in \mathbb{R}^d$ at time s , the expected value of R is

$$m_1^\alpha \frac{1}{(2\pi(t-s)d(m_1))^{d/2}} \int_A \exp\left\{-\frac{(y-x)^2}{2(t-s)d(m_1)}\right\} dx$$

if no mass transition occurs, whereas, it is

$$(2.6) \quad \frac{m_1}{m_2} m_2^\alpha \frac{1}{(2\pi(t-s)d(m_2))^{d/2}} \int_A \exp\left\{-\frac{(y-x)^2}{2(t-s)d(m_2)}\right\} dx$$

if the transition does occur. (The first factor in (2.6) is the survival probability for the transition). Given that $d(m_2) \leq d(m_1)$, the latter exponential term may be bounded pointwise by the former, and we find that

$$\frac{\mathbb{E}(R \text{ if mass transition occurs})}{\mathbb{E}(R \text{ if it does not})} \leq \frac{m_1}{m_2} \frac{d(m_1)^{d/2}}{d(m_2)^{d/2}} \frac{m_2^\alpha}{m_1^\alpha} = \frac{m_1^{1-\alpha} d(m_1)^{d/2}}{m_2^{1-\alpha} d(m_2)^{d/2}}.$$

If a whole sequence of mass transition occurs, $m_i \rightarrow m_{i+1}$ at time t_i for $i \in \{1, \dots, n\}$, with $t_i \in [0, t]$ an increasing sequence, we similarly find that the ratio of the expected values of R in the case where the sequence of mass transitions occurs and in that where no transition takes place, is bounded above by

$$\frac{1}{m^{1-\alpha}} \frac{d(1)^{d/2}}{d(m)^{d/2}}.$$

(We have omitted some details of the argument: to make it rigorous, we might use an induction on the total number of mass transitions undertaken prior to time t , and invoke the strong Markov property.)

By comparison, the expected value of R if no transition occurs is $u(x, t)$, where

$$\frac{\partial u}{\partial t} = d(1)\Delta u,$$

with $u(x, 0) = f_1(x, 0)$. By choosing $A = B(x, r)$ for each $r > 0$, we learn that

$$\sum_{m=1}^{\infty} m^\alpha f_m(x, t) \leq \frac{d(1)^{d/2}}{\inf_{m \in \mathbb{N}} m^{1-\alpha} d(m)^{d/2}} u(x, t).$$

From the hypothesis $d(m) > cm^{-2/d(1-\alpha)}$, we find that

$$\sum_{m=1}^{\infty} m^\alpha f_m(x, t) \leq \frac{d(1)^{d/2}}{c} u(x, t),$$

as we sought.

In fact, a more careful tracer particle argument improves this result: the same conclusion (2.5) may be reached under the weaker assumption that d is decreasing and satisfies $d(m) > m^{-(1-\alpha)+\epsilon}$ for some $\alpha \in (0, 1)$ and $\epsilon \in (0, \alpha)$. This is a reformulation of Corollary 1.1 which is a consequence of Theorem 1.2.

We have mentioned that each of our results has an analogue with a derivation that considers the tracer particle governed by a given solution of (1.1). We will not outline the method of proof of the result on mass conservation (see Theorem 1.3), stating only the characterization of gelation in terms of a tracer particle. We alter the definition of the tracer particle governed by $\{f_n : n \in \mathbb{N}\}$, so that the particle survives every mass transition. Given (2.4), mass conservation of the solution f until a given time $T \in [0, \infty)$ occurs if and only if the tracer particle experiences only finitely many collisions on the time interval $[0, T]$.

3 Moment bounds under Hypothesis 1.1

Let us first construct an auxiliary function H that will be needed for the proof of Theorem 1.1. Before doing so, let us make an observation. Define

$$(3.1) \quad H_0(x) = \begin{cases} c(d)|x|^{2-d} & \text{if } d \geq 3, \\ -\frac{1}{2\pi} \log |x| & \text{if } d = 2, \\ -\frac{1}{2}|x| & \text{if } d = 1, \end{cases}$$

where $c(d) = (d-2)^{-1}\omega_d^{-1}$ with ω_d denoting the $(d-1)$ -dimensional measure of the unit $S^{d-1} = \{x \in \mathbb{R}^d : |x| = 1\}$. We then have that $\Delta H_0 = -\delta_0$, where δ_0 denotes the Dirac's measure at 0. More precisely, for a test function g , the function

$$u(x) = \int H_0(x-y)g(y)dy,$$

satisfies $\Delta u = -g$. Note that $H_0 \geq 0$ when $d \geq 3$ and this property is lacking when $d \leq 2$. Because of this we can only hope for the existence of a suitable function H such that $H \geq 0$ but now $\Delta H = -\delta_0 + \text{Error}$ for an *Error* that can be controlled. This is the content of our first lemma.

Lemma 3.1 *Assume $d \leq 2$. There exist functions H and K such that $H \geq 0$, K is bounded, K is of compact support,*

$$(3.2) \quad -\Delta H(x) = \delta_0 - K(x),$$

and the function $H - \phi_0$ is bounded. (The function ϕ_0 was defined in (1.7).)

Proof. The construction of H for $d = 1$ is straightforward; we can readily find a nonnegative function H such that $H = \phi_0$ in $[-1/2, 1/2]$, $H = 0$ outside $[-1, 1]$, and H is smooth off the origin.

For the construction of the function H when $d = 2$, let us start from the function ϕ_0 and make an important observation. Note that if

$$(3.3) \quad R(x) = J * \phi_0(x) = \frac{-1}{2\pi} \int_{|x-y| \leq 1} \log |x-y| J(y) dy,$$

with J continuous and nonnegative, then $R \geq 0$ and

$$(3.4) \quad -\Delta R = J - \tilde{J},$$

where

$$(3.5) \quad \tilde{J}(x) = \frac{1}{2\pi} \int_{|z|=1} J(x-z) dS(z) = J * \tilde{\delta}_0,$$

where dS denotes the Lebesgue measure on the unit circle and $\tilde{\delta}_0$ denotes the normalized Lebesgue measure on the unit circle. In (3.3), we may replace $J(y)dy$ with a measure $J(dy)$. Then (3.4) is still valid weakly for an obvious interpretation for (3.5). In particular if we choose $J(dy) = \delta_0(dy)$, then

$$-\Delta\phi_0 = \delta_0 - \tilde{\delta}_0.$$

Our goal is to replace $\tilde{\delta}_0$ with a bounded function of compact support. For this we set $\phi_1 = \tilde{\delta}_0 * \phi_0$ to obtain $\phi_1 \geq 0$ and

$$-\Delta(\phi_0 + \phi_1) = \delta_0 - \hat{\delta}_0,$$

where $\hat{\delta}_0 = \tilde{\delta}_0 * \tilde{\delta}_0$ is now a “function” and is weakly given by

$$\begin{aligned} \int h(z) \hat{\delta}_0(dz) &= \frac{1}{(2\pi)^2} \int_{|z|=1} \int_{|a|=1} h(z+a) dS(z) dS(a) \\ &= \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} h(e^{i\theta_1} + e^{i\theta_2}) d\theta_1 d\theta_2. \end{aligned}$$

Note that the Jacobian of the transformation $\psi : (\theta_1, \theta_2) \mapsto (e^{i\theta_1} + e^{i\theta_2})$ is given by $|\sin(\theta_1 - \theta_2)|$. Since $\cos(\theta_1 - \theta_2) = \frac{1}{2}|e^{i\theta_1} + e^{i\theta_2}|^2 - 1$, we learn that if $\hat{\delta}_0(dz) = \hat{\delta}_0(z)dz$, then

$$\begin{aligned} \hat{\delta}_0(z) &= 2 \frac{1}{(2\pi)^2} \frac{1}{\sqrt{1 - (\frac{1}{2}|z|^2 - 1)^2}} \mathbb{1}(|z| \leq 2) \\ &= 2 \frac{1}{(2\pi)^2} \frac{2}{|z|\sqrt{4 - |z|^2}} \mathbb{1}(|z| \leq 2). \end{aligned}$$

Here the factor 2 comes from the fact that ψ maps (exactly) two points to one point because $\psi(\theta_1, \theta_2) = \psi(\theta_2, \theta_1)$. The function $\hat{\delta}$ is not bounded. Let us apply the above procedure one more time to define $\phi_2 = \phi_0 * \hat{\delta}_0$ so that $\phi_2 \geq 0$ and

$$-\Delta(\phi_0 + \phi_1 + \phi_2) = \delta_0 - \bar{\delta}_0,$$

where

$$(3.6) \quad \bar{\delta}_0(z) = \frac{4}{(2\pi)^3} \int_{|a|=1} \frac{1}{|z-a|\sqrt{4-|z-a|^2}} \mathbb{1}(|z-a| \leq 2) dS(a).$$

As we will see, the function $\bar{\delta}_0$ is not bounded either. However, the function $\bar{\delta}_0$ is less “singular” than $\hat{\delta}_0$. For this we show that in fact $\bar{\delta}_0$ has a logarithmic singularity. To do this, first observe that $\bar{\delta}_0(z)$ is radially symmetric because both dS and $\hat{\delta}_0$ are rotationally invariant. Hence we may assume that z lies on the x_1 -axis and $z > 0$. Observe that the integrand in (3.6) is singular when either $|z - a| = 0$ or $|z - a| = 2$. It is not hard to show that there exists a positive constant c_1 such that if $a = e^{i\theta}$ with $\theta \in (-\pi, \pi]$, then

$$(3.7) \quad c_1(|\theta| + |1 - |z||) \leq |z - a| \leq |\theta| + |1 - |z||.$$

The first inequality will be used to treat $1/|z - a|$ singularity in (3.6). We now turn to the singularity which comes from the factor $(2 - |z - a|)^{-1/2}$. For this observe that if $a = e^{i\theta}$, then we have

$$\gamma(\theta) := 4 - |z - a|^2 = 3 - z^2 + 2z \cos \theta.$$

Let us choose $\theta_0 \in [0, \pi]$ so that

$$3 - z^2 + 2z \cos \theta_0 = 0 \quad \text{or} \quad \cos \theta_0 = \frac{z^2 - 3}{2z}.$$

This means that the integrand in (3.6) is singular at θ_0 and $-\theta_0$. Note that when $z \geq 0$, the singular point θ_0 exists only if z belongs to the interval $[1, 3]$. Also note that if z is neither close to 1 nor 3, then θ_0 is neither close to 0 nor π . From the elementary inequality

$$\gamma(\theta) = \gamma(\theta) - \gamma(\theta_0) = 2z(\cos \theta - \cos \theta_0) = -4z \sin \frac{\theta + \theta_0}{2} \sin \frac{\theta - \theta_0}{2},$$

we learn that if z is neither close to 1 nor 3, then we can find a positive constant c_2 such that

$$\sqrt{4 - |z - a|^2} \geq c_2 \sqrt{|\theta - \theta_0|},$$

for θ close to θ_0 . (Recall that $z \in [1, 3]$.) Since this is an integrable singularity with respect to $d\theta$ -integration, we deduce that $\bar{\delta}_0$ is bounded if z stays away from the circles $|z| = 1$ and $|z| = 3$. We now assume that $|z|$ is close to 1. In this case θ_0 is close to π and $|\theta_0 - \pi|$ is comparable to $\sqrt{|1 - |z||}$. Also, because of $|z - a| \leq 2$ and $\gamma(\theta) - \gamma(\theta_0) > 0$ we learn that $\theta < \theta_0$. We have

$$(3.8) \quad 4 - |z - a|^2 \geq c_3 [(\pi - \theta) + (\pi - \theta_0)] [(\pi - \theta) - (\pi - \theta_0)],$$

for a positive constant c_3 . When $|z|$ is close to 1, the integrand in (3.6) is singular at $\pm\theta_0$ and “almost” singular (see (3.7)) at 0. From (3.7) and (3.8) we deduce

$$\bar{\delta}_0(z) \leq c_4 |\log |1 - |z|||,$$

whenever $|z|$ is close to 1. (Here we used the fact that $\int_0^{\theta_0} [(\pi - \theta)^2 - (\pi - \theta_0)^2]^{-1/2} d\theta$ is of order $|\log(\pi - \theta_0)|$.)

When $|z|$ is close to 3, θ_0 is small (in fact of order $O(\sqrt{3 - |z|})$), and the condition $|z - a| \leq 2$ forces $\theta \in [-\theta_0, \theta_0]$. On the other hand,

$$4 - |z - a|^2 \geq c_5 |\theta + \theta_0| |\theta - \theta_0|,$$

for a positive constant c_5 , implies that $\bar{\delta}_0(z)$ is bounded for $|z|$ close to 3. Here we are using

$$\int_{-\theta_0}^{\theta_0} (\theta_0^2 - \theta^2)^{-\frac{1}{2}} d\theta < \infty.$$

Putting all the pieces together, we deduce

$$(3.9) \quad \bar{\delta}_0(z) \leq c [|\log |1 - |z||] \mathbf{1}(|z| \leq 3).$$

We set $\phi_3 = \bar{\delta}_0 * \phi_0$ to obtain $\phi_3 \geq 0$ and

$$(3.10) \quad -\Delta(\phi_0 + \phi_1 + \phi_2 + \phi_3) = \delta_0 - K$$

where

$$K(z) = \frac{1}{2\pi} \int_{|a|=1} \bar{\delta}_0(z + a) dS(a).$$

It is straightforward to use (3.9) to show that K is uniformly bounded. Now (3.2) follows from (3.10) by choosing $H = \phi_0 + \phi_1 + \phi_2 + \phi_3$. It is also straightforward to check that the functions ϕ_1 , ϕ_2 , and ϕ_3 are bounded. \square

Let ζ be a nonnegative smooth function of compact support with $\int \zeta = 1$ and set $\zeta^\delta(x) = \delta^{-d} \zeta(x/\delta)$. We also define $f_n^\delta = f_n * \zeta^\delta$ and $Q_n^\delta = Q_n * \zeta^\delta$. We certainly have

$$(3.11) \quad f_n^\delta(x, t) = f_n^\delta(x, 0) + \int_0^t d(n) \Delta f_n^\delta(x, s) ds + \int_0^t Q_n^\delta(x, s) ds.$$

Also, as it is well-known,

$$(3.12) \quad \sum_n \phi(n) Q_n = \sum_{n,m} \alpha(n, m) (\phi(n+m) - \phi(n) - \phi(m)) f_n f_m.$$

The same identity is valid if we replace $f_n f_m$ with $(f_n f_m) * \zeta^\delta$ and Q with Q^δ . Using the fact that for $\phi(n) = n \mathbf{1}(n \leq \ell)$, we have $\sum_n \phi(n) Q_n \leq 0$ we can readily deduce that

$$(3.13) \quad \sup_\delta \sup_\ell \sup_t \int \sum_{n=1}^\ell n f_n^\delta(x, t) dx < \infty.$$

Lemma 3.2 *Let H be as in Lemma 2.1. Then there exists a constant c_0 such that*

$$\sup_x \sum_{n=1}^{\ell} n(f_n^\delta *_x H)(x, t) \leq \sup_x \sum_{n=1}^{\ell} n(f_n^\delta *_x H)(x, 0) + c_0 t$$

for every positive δ . We may choose $c_0 = 0$ when $d \geq 3$.

Proof. We have

$$\begin{aligned} \sum_{n=1}^{\ell} n(f_n^\delta *_x H)(x, t) &= \sum_{n=1}^{\ell} n(f_n^\delta *_x H)(x, 0) - \int_0^t \sum_{n=1}^{\ell} nd(n) f_n^\delta(x, s) ds \\ &\quad + \int_0^t \sum_{n=1}^{\ell} nd(n) f_n^\delta *_x K(x, s) ds + \int_0^t \sum_{n=1}^{\ell} nQ_n^\delta *_x H(x, s) ds. \end{aligned}$$

From the boundedness of K , $H \geq 0$ and $\sum_{n=1}^{\ell} nQ_n \leq 0$ we deduce

$$\sum_{n=1}^{\ell} n(f_n^\delta *_x H)(x, t) \leq \sum_{n=1}^{\ell} n(f_n^\delta *_x H)(x, 0) + c_1 \int_0^t \int \sum_{n=1}^{\ell} nd(n) f_n^\delta(x, s) dx ds.$$

We now use (3.13) to bound the last term to complete the proof. □

Proof of Theorem 1.1. Set

$$\begin{aligned} Z^\delta(t) &= \int \left(\sum_{n=1}^{\ell} n^a f_n^\delta(x, t) \right) \left(\sum_{n=1}^{\ell} n f_n^\delta *_x H(x, t) \right) dx, \\ Z(t) &= \int \left(\sum_{n=1}^{\ell} n^a f_n(x, t) \right) \left(\sum_{n=1}^{\ell} n f_n *_x H(x, t) \right) dx. \end{aligned}$$

We have that weakly,

$$\begin{aligned}
\frac{d}{dt}Z^\delta(t) &= - \int \left(\sum_{n=1}^{\ell} n^a f_n^\delta(x, t) \right) \left(\sum_{n=1}^{\ell} nd(n) f_n^\delta(x, t) \right) dx \\
&\quad - \int \left(\sum_{n=1}^{\ell} n^a d(n) f_n^\delta(x, t) \right) \left(\sum_{n=1}^{\ell} n f_n^\delta(x, t) \right) dx \\
&\quad + \int \left(\sum_{n=1}^{\ell} n^a f_n^\delta(x, t) \right) \left(\sum_{n=1}^{\ell} n Q_n^\delta *_x H(x, t) \right) dx \\
&\quad + \int \left(\sum_{n=1}^{\ell} n^a Q_n^\delta(x, t) \right) \left(\sum_{n=1}^{\ell} n f_n^\delta *_x H(x, t) \right) dx \\
&\quad + \int \left(\sum_{n=1}^{\ell} n^a f_n^\delta(x, t) \right) \left(\sum_{n=1}^{\ell} nd(n) f_n^\delta *_x K(x, t) \right) dx \\
&\quad + \int \left(\sum_{n=1}^{\ell} n^a d(n) f_n^\delta(x, t) \right) \left(\sum_{n=1}^{\ell} n f_n^\delta *_x K(x, t) \right) dx \\
&=: \Omega_1 + \Omega_2 + \Omega_3 + \Omega_4 + \Omega_5 + \Omega_6.
\end{aligned}$$

We now study the various terms which appear on the right-hand side. We certainly have

$$\begin{aligned}
\Omega_1 + \Omega_2 &= -\frac{1}{2} \int \sum_{n,m} \mathbb{1}(n, m \leq \ell) [n^a m d(m) + n^a d(n) m + m^a n d(n) + m^a d(m) n] f_n^\delta f_m^\delta dx \\
&= -\frac{1}{2} \int \sum_{n,m} \mathbb{1}(n, m \leq \ell) n m (d(n) + d(m)) (n^{a-1} + m^{a-1}) f_n^\delta f_m^\delta dx.
\end{aligned}$$

From $\sum_{n=1}^{\ell} n Q_n \leq 0$, we learn that $\Omega_3 \leq 0$. By boundedness of K , (3.13) and the boundedness of $d(\cdot)$ we deduce that

$$|\Omega_5 + \Omega_6| \leq c_1 \int X_a^\delta dx,$$

where X_a is defined in (1.5) and $X_a^\delta = X_a * \zeta^\delta$. It remains to bound Ω_4 . Note that

$$\begin{aligned}
\sum_{n=1}^{\ell} n^a Q_n(x, t) &= \sum_{n,m} [(n+m)^a \mathbb{1}(n+m \leq \ell) - n^a \mathbb{1}(n \leq \ell) - m^a \mathbb{1}(m \leq \ell)] \alpha(n, m) f_n f_m \\
(3.14) \quad &\leq c_2 \sum_{n,m} (n^{a-1} m + m^{a-1} n) \alpha(n, m) \mathbb{1}(n+m \leq \ell) f_n f_m =: c_2 Z_4.
\end{aligned}$$

From this, Lemma 3.2, the boundedness of $H - \phi_0$ and (1.9),

$$\Omega_4 \leq c_2 \int Z_4^\delta \left(\sum_{n=1}^{\ell} n f_n^\delta *_x H(x, t) \right) dx \leq c_3 \int Z_4^\delta dx,$$

where $Z_4^\delta = Z_4 * \zeta^\delta$. On the other hand, for $\delta_0 > 0$, we can find $k_0 = k_0(\delta_0)$ such that if $k_0 < \ell$, then

$$\begin{aligned} (3.15) \quad Z_4 &= \sum_{n,m} \mathbb{1}(k_0 \leq n+m \leq \ell) (n^{a-1}m + m^{a-1}n) \alpha(n, m) f_n f_m \\ &+ \sum_{n,m} \mathbb{1}(n+m < k_0) (n^{a-1}m + m^{a-1}n) \alpha(n, m) f_n f_m \\ &\leq \delta_0 \sum_{n,m} \mathbb{1}(n+m \leq \ell) (n^{a-1}m + m^{a-1}n) (n+m) (d(n) + d(m)) f_n f_m \\ &+ 2k_0^a \sum_{n,m} \mathbb{1}(n+m \leq \ell) \alpha(n, m) f_n f_m. \end{aligned}$$

As a result,

$$\begin{aligned} \frac{d}{dt} Z^\delta(t) &\leq -\frac{1}{2} \int \sum_{n,m} \mathbb{1}(n, m \leq \ell) nm (n^{a-1} + m^{a-1}) (d(n) + d(m)) f_n^\delta f_m^\delta dx \\ &+ 2c_3 \delta_0 \int \sum_{n,m} \mathbb{1}(n, m \leq \ell) nm (n^{a-1} + m^{a-1}) (d(n) + d(m)) (f_n f_m) * \zeta^\delta dx \\ &+ 2c_3 k_0^a \int \sum_{n,m} \mathbb{1}(n+m \leq \ell) \alpha(n, m) (f_n f_m) * \zeta^\delta dx + c_1 \int X_a^\delta(x, t) dx. \end{aligned}$$

Here, we are using the identity,

$$(n+m)(n^{a-2} + m^{a-2}) \leq 2(n^{a-1} + m^{a-1}),$$

which is valid provided that $a \geq 2$.

We now send δ to 0 to yield

$$\begin{aligned} (3.16) \quad \frac{d}{dt} Z(t) &\leq \left(2c_3 \delta_0 - \frac{1}{2} \right) \int \sum_{n,m} \mathbb{1}(n, m \leq \ell) nm (n^{a-1} + m^{a-1}) (d(n) + d(m)) f_n f_m dx \\ &+ 2c_3 k_0^a \int \sum_{n,m} \mathbb{1}(n+m \leq \ell) \alpha(n, m) f_n f_m dx + c_1 \int X_a(x, t) dx. \end{aligned}$$

Note that the time integral of the second integral is bounded because

$$\frac{d}{dt} \int \sum_{n=1}^{\ell} f_n dx \leq - \int \sum_{n,m} \mathbb{1}(n+m \leq \ell) \alpha(n,m) f_n f_m dx,$$

which implies,

$$(3.17) \quad \int_0^T \int \sum_{n,m} \mathbb{1}(n+m \leq \ell) \alpha(n,m) f_n f_m dx dt \leq \int \sum_{n=1}^{\ell} n f_n^0 dx.$$

Furthermore, the equality

$$(3.18) \quad \frac{d}{dt} \int \sum_{n=1}^{\ell} n^a f_n(x,t) dx = \int \sum_{n=1}^{\ell} n^a Q_n(x,t) dx,$$

and (3.14) imply that

$$\begin{aligned} \int_0^t \int \sum_{n=1}^{\ell} n^a f_n(x,s) dx ds &\leq t \int \sum_{n=1}^{\ell} n^a f_n(x,0) dx + c_2 \int_0^t \int_0^s \int Z_4(x,\theta) dx d\theta \\ &\leq t \int \sum_{n=1}^{\ell} n^a f_n(x,0) dx + t c_2 \int_0^t \int Z_4(x,\theta) dx d\theta. \end{aligned}$$

From this, (3.17) and (3.16) we deduce that $Z(t) - Z(0)$ is bounded above by

$$\left((2c_3 + c_1 c_2 t) \delta_0 - \frac{1}{2} \right) \int_0^t \int \sum_{n,m \leq \ell} nm(n^{a-1} + m^{a-1})(d(n) + d(m)) f_n f_m dx ds + c_4(1+t).$$

We now choose $\delta_0 = \delta_0(t)$ so that $1/2 > (2c_3 + c_1 c_2 t) \delta_0$. With this choice, the bounds in (1.10) follow. From (1.10), (3.14) and (3.18) we conclude (1.11). \square

We end this section with a variant of Theorem 1.1 that holds under Hypothesis 1.3.

Lemma 3.3 *Under Hypothesis 1.3, there exists a constant C such that*

$$(3.19) \quad \int X_2(x,t) dx \leq \left(\int X_2(x,0) dx \right) \exp \left(CT \left\| \sum_n n f_n^0 \right\|_{L^\infty} \right).$$

Proof. We start from

$$\begin{aligned}
\frac{d}{dt} \int \sum_{n=1}^{\ell} n^2 f_n(x, t) dx &= \int \sum_{n=1}^{\ell} n^2 Q_n(x, t) dx \\
&\leq \int 2 \sum_{n,m} nm \alpha(n, m) \mathbb{1}(n+m \leq \ell) f_n f_m dx \\
&\leq \int 2C_0 \sum_{n,m} nm(n+m) \mathbb{1}(n+m \leq \ell) f_n f_m dx \\
&\leq \int 4C_0 \left(\sum_n n \mathbb{1}(n \leq \ell) f_n \right) \left(\sum_m m^2 \mathbb{1}(m \leq \ell) f_m \right) dx.
\end{aligned}$$

This and Gronwall's inequality imply (3.19) because we can use the uniform positivity of $d(\cdot)$ and Lemma 3.1 of Section 3 to assert that $X_1 \in L^\infty$. \square

4 Moment bounds when $d(\cdot)$ is non-increasing

This section is devoted to the proof of Theorem 1.2. We start with a lemma.

Lemma 4.1 *Assume $d(\cdot)$ is non-increasing. Then*

$$(4.1) \quad \hat{X}_1(x, t) = \sum_{n=1}^{\infty} nd(n)^{d/2} f_n(x, t) \leq d(1)^{d/2} u(x, t).$$

where u is the unique solution to $u_t = d(1)\Delta u$ subject to the initial condition $u(x, 0) = \sum_{n=1}^{\infty} n f_n(x, 0)$.

Proof. We first establish

$$(4.2) \quad \sum_1^{\ell} nd(n)^{d/2} f_n(t) \leq d(1)^{d/2} S_t^{d(1)} \left(\sum_1^{\ell} n f_n^0 \right) + d(\ell)^{d/2} \int_0^t S_{t-s}^{d(\ell)} \left(\sum_1^{\ell} n Q_n(s) \right) ds.$$

Here for simplicity, we do not display the dependence on the x -variable. Note that (4.2) implies

$$(4.3) \quad \sum_1^{\ell} nd(n)^{d/2} f_n \leq d(1)^{d/2} S_t^{d(1)} \left(\sum_1^{\ell} n f_n^0 \right),$$

because $\sum_1^\ell nQ_n \leq 0$. Evidently (4.3) implies (4.1).

We establish (4.2) by induction. (4.2) is obvious when $\ell = 1$ by definition; in fact we have equality. Suppose (4.2) is valid. We would like to deduce (4.2) with ℓ replaced with $\ell + 1$. To do so, first observe that if $D_1 \geq D_2$ and $g \geq 0$, then

$$(4.4) \quad D_1^{d/2} S_t^{D_1} g \geq D_2^{d/2} S_t^{D_2} g.$$

From this and (4.2) we learn

$$(4.5) \quad \sum_1^\ell nd(n)^{d/2} f_n \leq d(1)^{d/2} S_t^{d(1)} \left(\sum_1^\ell n f_n^0 \right) + d(\ell + 1)^{d/2} \int_0^t S_{t-s}^{d(\ell+1)} \left(\sum_1^\ell n Q_n(s) \right) ds$$

because $d(\ell) \geq d(\ell + 1)$ and $\sum_1^\ell nQ_n \leq 0$. Applying (4.4) to

$$f_{\ell+1}(t) = S_t^{d(\ell+1)} f_{\ell+1}^0 + \int_0^t S_{t-s}^{d(\ell+1)} Q_{\ell+1}(s) ds,$$

yields

$$(4.6) \quad f_{\ell+1}(t) \leq \left(\frac{d(1)}{d(\ell + 1)} \right)^{d/2} S_t^{d(1)} f_{\ell+1}^0 + \int_0^t S_{t-s}^{d(\ell+1)} Q_{\ell+1}(s) ds.$$

We multiply both sides of (4.6) by $(\ell + 1)d(\ell + 1)^{d/2}$ and add the result to (4.5). The outcome is

$$\sum_1^{\ell+1} nd(n)^{d/2} f_n \leq d(1)^{d/2} S_t^{d(1)} \left(\sum_1^{\ell+1} n f_n^0 \right) + d(\ell + 1)^{d/2} \int_0^t S_{t-s}^{d(\ell+1)} \left(\sum_1^{\ell+1} n Q_n(s) \right) ds.$$

This completes the proof. \square

Remark 4.1. For the continuous version of (1.1), a similar proof leads to an L^∞ bound on $\int_0^\infty nd(n)^{d/2} f_n dn$ provided that the function $d(\cdot)$ is piecewise constant, uniformly positive and nonincreasing. If we assume only the second and third conditions on $d(\cdot)$, we can establish the same bound on a solution provided that this solution can be approximated by solutions corresponding to piecewise constant $d(\cdot)$. Of course, if we already know the uniqueness of solutions to (1.1), then our L^∞ bound applies to all solutions. But we do not: when uniqueness is proved in Section 4, we will use Lemma 3.1 and its consequence Theorem 1.2. However, if we only postulate the uniqueness for those solutions with $\int_0^\infty nd(n)^{d/2} f_n dn \in L^\infty$, then Theorem 1.4 is also valid in the continuous case.

Proof of Theorem 1.2.

Step 1. Let us simply write L^ℓ for $L^\ell(\mathbb{R}^d \times [0, T])$. We first show that if $\ell \geq 1$, then

$$(4.7) \quad \hat{X}_{a\ell+1} \in L^1 \Rightarrow \hat{X}_{a+1} \in L^\ell.$$

Indeed,

$$\begin{aligned}\hat{X}_{a+1} &= \sum_n n^a nd(n)^{d/2} f_n = \hat{X}_1 \sum_n n^a \frac{nd(n)^{d/2} f_n}{\hat{X}_1}, \\ \hat{X}_{a+1}^\ell &\leq \hat{X}_1^\ell \sum_n n^{a\ell} \frac{nd(n)^{d/2} f_n}{\hat{X}_1} = \hat{X}_1^{\ell-1} \hat{X}_{a\ell+1},\end{aligned}$$

by Hölder's inequality. This implies (4.7) because by Lemma 3.1, $\hat{X}_1 \in L^\infty$.

Recall that we are assuming

$$r_1 n^{-b_1} \leq d(n) \leq r_2 n^{-b_2}.$$

From this and (4.7) we deduce that

$$X_{(a\ell+1-b_2\frac{d}{2})} \in L^1 \Rightarrow X_{(a+1-b_1\frac{d}{2})} \in L^\ell.$$

This means that

$$(4.8) \quad X_a \in L^1 \Rightarrow X_b \in L^{\frac{2a+b_2d-2}{2b+b_1d-2}},$$

provided that $2b + b_1d - 2 > 0$ and $(2a + b_2d - 2)/(2b + b_1d - 2) \geq 1$.

Step 2. We now try to bound Q_n^+ with the aid of (4.8). Using $\alpha(n, m) \leq C_0(n + m)$, we certainly have

$$\begin{aligned}Q_n^+ &= \sum_{n_1+n_2=n} \alpha(n_1, n_2) f_{n_1} f_{n_2} \\ &\leq \sum_{n_1, n_2} \mathbb{1}(n_1 \geq n/2 \text{ or } n_2 \geq n/2) \alpha(n_1, n_2) f_{n_1} f_{n_2} \\ &\leq 2C_0 [X_1 X_0(n/2) + X_1(n/2) X(0)]\end{aligned}$$

where $X_a(N) = \sum_{m \geq N} m^a f_m$. Hence

$$\begin{aligned}Q_n^+ &\leq c_2 n^{-\ell} [X_1 X_\ell + X_{1+\ell} X_0], \\ \|Q_n^+\|_{L^p} &\leq c_2 n^{-\ell} [\|X_1\|_{L^{\ell_1}} \|X_\ell\|_{L^{\ell_2}} + \|X_{1+\ell}\|_{L^{\ell_3}} \|X_0\|_{L^{\ell_4}}]\end{aligned}$$

provided that $\frac{1}{p} = \frac{1}{\ell_1} + \frac{1}{\ell_2} = \frac{1}{\ell_3} + \frac{1}{\ell_4}$. To use (4.8), let us first assume that $b_1d > 2$ and that $2a + b_2d - 2 \geq b_1d + 2\ell$. Assume $X_a \in L^1$. Then we use (4.8) to assert that

$$\begin{aligned}X_1 &\in L^{\frac{2a+b_2d-2}{b_1d}}, & X_0 &\in L^{\frac{2a+b_2d-2}{b_1d-2}} \\ X_\ell &\in L^{\frac{2a+b_2d-2}{2\ell+b_1d-2}}, & X_{\ell+1} &\in L^{\frac{2a+b_2d-2}{2\ell+b_1d}}.\end{aligned}$$

Hence if we set

$$(4.9) \quad p = \frac{2a + b_2d - 2}{2b_1d + 2\ell - 2},$$

and assume that $p \geq 1$, $b_1d > 2$, then we have that

$$(4.10) \quad X_a \in L^1 \Rightarrow \|Q_n\|_{L^p} \leq c n^{-\ell}.$$

However, if $b_1d \leq 2$, then we have that $X_0 \in L^\infty$ because $\hat{X}_1 \in L^\infty$. From this, we learn that in this case (4.9) is true but now for

$$(4.11) \quad p = \frac{2a + b_2d - 2}{2\ell + b_1d}.$$

Step 3. Note that if

$$p_D(x, t) = \begin{cases} (4\pi Dt)^{-d/2} \exp\left(-\frac{|x|^2}{4Dt}\right) & \text{if } t > 0, \\ 0 & \text{if } t < 0, \end{cases}$$

and g is a function with $g(x, t) = 0$ for $t < 0$, then $\int_0^t S_{t-s}^D g(x, s) ds = (p_D * g)(x, t)$ where the convolution is in both x and t variables. Also note that

$$\begin{aligned} \int_0^T \int (p_D(x, t))^r dx dt &= \int_0^T \int \left(\frac{1}{\sqrt{Dt}}\right)^{dr} p_1\left(\frac{x}{\sqrt{Dt}}, 1\right)^r dx dt \\ &= \int_0^T (Dt)^{\frac{d}{2}(1-r)} dt = c(T, r) D^{\frac{d}{2}(1-r)} \end{aligned}$$

with $c(T, r) < \infty$ if and only if $r < \frac{2}{d} + 1$. We certainly have

$$f_n(x, t) = (S_t^{d(n)} f_n^0)(x) + (p_{d(n)} * Q_n)(x, t).$$

So,

$$(4.12) \quad \|f_n\|_{L^\infty} \leq \|(S_t^{d(n)} f_n^0)(x)\|_{L^\infty} + \|p_{d(n)}\|_{L^r} \|Q_n\|_{L^p}$$

provided that $\frac{1}{r} + \frac{1}{p} = 1$. Since $p_D \in L_r$ with $r < \frac{2}{d} + 1$, it suffices to have

$$\frac{1}{1 + 2/d} + \frac{1}{p} < 1.$$

Choosing p as in (4.9) or (4.11) requires

$$\frac{2b_1d + 2\ell - 2}{2a + b_2d - 2} \text{ or } \frac{b_1d + 2\ell}{2a + b_2d - 2} < \frac{2}{d + 2}.$$

More precisely,

$$(4.13) \quad \ell < \begin{cases} \frac{1}{d+2}(2a + b_2d - 2) - b_1d + 1 & \text{if } b_1d > 2, \\ \frac{1}{d+2}(2a + b_2d - 2) - \frac{1}{2}b_1d & \text{if } b_1d \leq 2. \end{cases}$$

In summary, we need ℓ to satisfy (4.13) and r to satisfy $\frac{d}{2}(1-r) > -1$. As a result, if $X_a \in L^1$ and ℓ satisfies (4.13), then

$$\|f_n\|_{L^\infty} \leq A_n + c_3 n^{-\ell} d(n)^{d(1-r)/2} \leq A_n + c_4 n^{-\ell} d(n)^{-1}$$

where

$$A_n = \|(S_t^{d(n)})f_n^0(x)\|_{L^\infty} \leq \|f_n^0\|_{L^\infty(\mathbb{R}^d)}.$$

Final Step. We have that $\sum_n n^e \|f_n\|_{L^\infty} \leq \infty$ if $X_a \in L^1$,

$$(4.14) \quad \sum_n n^{e-\ell} d(n)^{-1} < \infty,$$

and,

$$\sum_n n^e \|f_n^0\|_{L^\infty(\mathbb{R}^d)} < \infty.$$

For (4.14) it suffices to have

$$e - \ell + b_1 < -1.$$

In other words,

$$(4.15) \quad e \leq \gamma(a, b_1, b_2) := \begin{cases} \frac{1}{d+2}(2a + b_2d - 2) - b_1(d+1) & \text{if } b_1d > 2, \\ \frac{1}{d+2}(2a + b_2d - 2) - \frac{1}{2}b_1d - b_1 - 1 & \text{if } b_1d \leq 2. \end{cases}$$

□

5 Uniqueness

The main result of this section is Theorem 5.1. Theorem 1.4 is an immediate consequence of Theorem 5.1.

Theorem 5.1 *Assume that $\alpha(n, m) \leq c_0 nm$ and let f and g be two solutions with*

$$\left\| \sum_n n^2 f_n \right\|_{L^\infty}, \left\| \sum_n n^2 g_n \right\|_{L^\infty} \leq A,$$

where L^p abbreviates $L^p(\mathbb{R}^d \times [0, T])$. Then

$$X(t) := \int \sum_n n |f_n - g_n|(x, t) dx$$

satisfies

$$(5.1) \quad X(t) \leq e^{4c_0 A t} X(0),$$

for $t \leq T$. In particular, if $f_n(\cdot, 0) = g_n(\cdot, 0)$ for all n , then $f_n(\cdot, t) = g_n(\cdot, t)$ for all n and $t \in [0, T]$.

We first state a straightforward lemma:

Lemma 5.1 *Let u be a weak solution of*

$$u_t = D\Delta u + h$$

with u and $h \in L^1$. Assume that ψ is a continuously differentiable convex function with $|\psi'(a)| \leq c_1$ for a constant c_1 and all $a \in \mathbb{R}$. Then

$$(5.2) \quad \int \psi(u(x, t)) dx \leq \int \psi(u(x, s)) dx + \int_s^t \int \psi'(u(x, \theta)) h(x, \theta) dx d\theta$$

whenever $0 < s < t$.

Lemma 5.1 is established by choosing a smooth mollifier ρ_ϵ , and showing the inequality (5.2) for $u_\epsilon = u * \rho_\epsilon$ and $h_\epsilon = h * \rho_\epsilon$. We then pass to the limit $\epsilon \rightarrow 0$. We omit the details.

Proof of Theorem 5.1. Choose a continuously differentiable convex function ψ_δ so that $\psi'_\delta(r) = \text{sgn}(r)$ for $r \notin (-\delta, \delta)$, $\psi_\delta \in C^2$, $|\psi'_\delta(r)| \leq 1$, $\psi_\delta(0) = 0$ and $\psi_\delta \geq 0$. We then apply Lemma 5.1 to assert that the expression

$$(5.3) \quad \int \sum_{n=1}^N n \psi_\delta(f_n(x, t) - g_n(x, t)) dx,$$

is bounded above by

$$(5.4) \quad \begin{aligned} & \int \sum_{n=1}^N n \psi_\delta(f_n(x, s) - g_n(x, s)) ds \\ & + \int_s^t \int \sum_{n=1}^N n \psi'_\delta(f_n(x, \theta) - g_n(x, \theta)) (Q_n(f)(x, \theta) - Q_n(g)(x, \theta)) dx d\theta \\ & = \int \sum_{n=1}^N n \psi_\delta(f_n(x, s) - g_n(x, s)) dx \\ & + \int_s^t \int \sum_{n,m} \alpha(n, m) (\Gamma_{n+m} - \Gamma_n - \Gamma_m) (f_n f_m - g_n g_m) dx d\theta. \end{aligned}$$

where $\Gamma_n = n \psi'_\delta(f_n - g_n)\mathbb{1}(n \leq N)$. Observe that if $n, m \leq N$, then

$$(\Gamma_{n+m} - \Gamma_n - \Gamma_m)(f_n f_m - g_n g_m),$$

equals

$$\begin{aligned} & (\Gamma_{n+m} - \Gamma_n - \Gamma_m)(f_n - g_n)f_m + (\Gamma_{n+m} - \Gamma_n - \Gamma_m)(f_m - g_m)g_n \\ & \leq (n+m)|f_n - g_n|f_m - n\psi'_\delta(f_n - g_n)(f_n - g_n)f_m + m|f_n - g_n|f_m \\ & \quad + (n+m)|f_m - g_m|g_n + n|f_m - g_m|g_n - m\psi'_\delta(f_m - g_m)(f_m - g_m)g_n \\ & \leq 2m|f_n - g_n|f_m + 2n\mathbb{1}(|f_n - g_n| < \delta)|f_n - g_n|f_m \\ & \quad + 2n|f_m - g_m|g_n + 2m\mathbb{1}(|f_m - g_m| < \delta)|f_m - g_m|g_n. \end{aligned}$$

From this, (5.4) and $\alpha(n, m) \leq c_0 nm$, we learn that the expression (5.3) is bounded above by

$$\begin{aligned} & \int \sum_{n=1}^N n\psi_\delta(f_n(x, s) - g_n(x, s))dx \\ & + 2c_0 \int_s^t \int \left[\sum_{n=1}^N n|f_n - g_n| \right] \left[\sum_{m=1}^N m^2(f_m + g_m) \right] dx d\theta \\ & + 2c_0 \delta \int_s^t \int \left[\sum_{n=1}^N n^2\mathbb{1}(|f_n - g_n| < \delta) \right] \left[\sum_{m=1}^N m(f_m + g_m) \right] dx d\theta. \end{aligned}$$

We then send $\delta \rightarrow 0$ and $N \rightarrow \infty$, in this order to obtain

$$\begin{aligned} \int \sum_{n=1}^{\infty} n|f_n(x, t) - g_n(x, t)|dx & \leq \int \sum_{n=1}^{\infty} n|f_n(x, s) - g_n(x, s)|dx \\ & + 2c_0 \int_s^t \int \left[\sum_{n=1}^{\infty} n|f_n - g_n| \right] \left[\sum_{m=1}^{\infty} m^2(f_m + g_m) \right] dx d\theta. \end{aligned}$$

The theorem now follows from this and Gronwall's inequality. \square

6 Mass conservation

Proof of Theorem 1.3. We first assume that $\hat{Y}_1 \in L^1$. Evidently,

$$\begin{aligned} \frac{d}{dt} \int \left(\sum_{n=1}^N n f_n \right) dx &= - \sum_{n,m} \{ \mathbb{1}(n \leq N < n+m)n + \mathbb{1}(m \leq N < n+m)m \} \alpha(n,m) f_n f_m \\ &= -2 \sum_{n,m} \mathbb{1}(n \leq N < n+m) n \alpha(n,m) f_n f_m \\ &\geq -2 \sum_{n,m} \mathbb{1}(n \geq N/2 \text{ or } m > N/2) n m \alpha(n,m) f_n f_m. \end{aligned}$$

The limit $N \rightarrow \infty$ of the time average of the right-hand side is 0 because $\hat{Y}_1 \in L^1$. From this we can readily deduce that

$$\lim_{N \rightarrow \infty} \left[\int \left(\sum_{n=1}^N n f_n(x,t) \right) dx - \int \left(\sum_{n=1}^N n f_n(x,0) \right) dx \right] = 0.$$

This completes the proof when $\hat{Y}_1 \in L^1$.

We now assume that Hypothesis 1.3 holds. We have,

$$\begin{aligned} \frac{d}{dt} \int \left(\sum_{n=1}^N n f_n \right) dx &= - \sum_{n,m} \{ \mathbb{1}(n \leq N < n+m)n + \mathbb{1}(m \leq N < n+m)m \} \alpha(n,m) f_n f_m \\ &\geq -2C_0 \sum_{n,m} \mathbb{1}(n \leq N < n+m) n(n+m) f_n f_m \\ &\geq -2C_0 \sum_{n,m} \mathbb{1}(n \leq N/2, m > N/2) n(n+m) f_n f_m \\ &\quad -2C_0 \sum_{n,m} \mathbb{1}(n > N/2) n(n+m) f_n f_m \\ &= -\Omega_1 - \Omega_2. \end{aligned}$$

We certainly have,

$$\begin{aligned}
\|\Omega_1\|_{L^1} &\leq c \left\| \sum_n n^2 f_n \right\|_{L^1} \left\| \sum_{m>N/2} f_m \right\|_{L^\infty} \\
&+ c \left\| \sum_n n f_n \right\|_{L^2} \left\| \sum_{m>N/2} m f_m \right\|_{L^2}, \\
\|\Omega_2\|_{L^1} &\leq c \left\| \sum_{n>N/2} n f_n \right\|_{L^2} \left\| \sum_m m f_m \right\|_{L^2} \\
&+ c \left\| \sum_{n>N/2} n^2 f_n \right\|_{L^1} \left\| \sum_m f_m \right\|_{L^\infty},
\end{aligned}$$

where L^p abbreviates $L^p(\mathbb{R}^d \times [0, T])$. Since $\sum_m m f_m \in L^2$, $\sum_m m^2 f_m \in L^1$, and $\sum_n n f_n \in L^\infty$, by Lemmas 4.1 and 2.3, we are done. \square

References

- [1] J. M. Ball and J. Carr. The discrete coagulation-fragmentation equations: existence, uniqueness, and density conservation. *J. Statist. Phys.*, 61(1-2):203–234, 1990.
- [2] P. Bénilan and D. Wrzosek. On an infinite system of reaction-diffusion equations. *Adv. Math. Sci. Appl.*, 7(1):351–366, 1997.
- [3] J. F. Collet and F. Poupaud. Asymptotic behaviour of solutions to the diffusive fragmentation-coagulation system. *Phys. D*, 114(1-2):123–146, 1998.
- [4] D. Dürr, S. Goldstein, and J. L. Lebowitz. A mechanical model of Brownian motion. *Comm. Math. Phys.*, 78(4):507–530, 1980/81.
- [5] M. Escobedo, Ph. Laurençot, S. Mischler, and B. Perthame. Gelation and mass conservation in coagulation-fragmentation models. *J. Differential Equations*, 195(1):143–174, 2003.
- [6] M. Escobedo, S. Mischler, and B. Perthame. Gelation in coagulation and fragmentation models. *Comm. Math. Phys.*, 231(1):157–188, 2002.
- [7] A. M. Hammond and F. Rezakhanlou. The kinetic limit of a system of coagulating Brownian particles. *To appear in Arch. Rational Mech. Anal.* Available at www.arxiv.org/math.PR/0408395.

- [8] Alan Hammond and Fraydoun Rezakhanlou. Kinetic limit for a system of coagulating planar Brownian particles. *J. Stat. Phys.*, 124(2-4):997–1040, 2006.
- [9] R. Lang and X.-X. Nyugen. Smoluchowski’s theory of coagulation in colloids holds rigorously in the Boltzmann-Grad limit. *Z. Wahrsch. Verw. Gebiete*, 54:227–280, 1980.
- [10] Ph. Laurençot and S. Mischler. The continuous coagulation-fragmentation equations with diffusion. *Arch. Ration. Mech. Anal.*, 162(1):45–99, 2002.
- [11] Ph. Laurençot and S. Mischler. Global existence for the discrete diffusive coagulation-fragmentation equations in L^1 . *Rev. Mat. Iberoamericana*, 18(3):731–745, 2002.
- [12] F. Rezakhanlou. The coagulating Brownian particles and Smoluchowski’s equation. *Markov Process. Related Fields*, 12:425–445, 2006.
- [13] D. Wrzosek. Existence of solutions for the discrete coagulation-fragmentation model with diffusion. *Topol. Methods Nonlinear Anal.*, 9(2):279–296, 1997.
- [14] D. Wrzosek. Mass-conserving solutions to the discrete coagulation-fragmentation model with diffusion. *Nonlinear Anal.*, 49(3, Ser. A: Theory Methods):297–314, 2002.
- [15] D. Wrzosek. Weak solutions to the Cauchy problem for the diffusive discrete coagulation-fragmentation system. *J. Math. Anal. Appl.*, 289(2):405–418, 2004.