GSI name: Roy Zhao

Circle True or False or leave blank. (1 point for correct answer, -1 for incorrect answer, 0 if left blank)

1. **TRUE** False It is possible to determine the equation for the PDF, given the equation for the CDF.

Solution: Taking the derivative of the CDF gives the PDF.

2. True **FALSE** Suppose that f(x) = x for $-0.5 \le x \le 1.5$ and 0 everywhere else. Since $\int_{-0.5}^{1.5} x dx = 1$ (you can assume the integral is correct), then f is a PDF.

Solution: This is false since f(-0.5) = -0.5 which is negative and PDFs cannot be negative.

Show your work and justify your answers. Please circle or box your final answer.

3. (10 points) (a) (5 points) Suppose that $f(x) = Cxe^{-x^2}$ for $x \ge 0$ and f(x) = 0 for x < 0 for some constant C. If f is a PDF, then find C.

Solution: Since f is a PDF, we require that

$$\int_{-\infty}^{\infty} f(x)dx = \int_{0}^{\infty} Cxe^{-x^{2}}dx = 1.$$

We u sub with $u = x^2$ and du = 2xdx to get

$$= \lim_{t \to \infty} \int_0^{t^2} Ce^{-u}/2du = \lim_{t \to \infty} -Ce^{-u}/2|_0^{t^2} = C/2.$$

Therefore C=2.

(b) (2 points) Find the CDF of f from above. (Hint: the CDF will be piecewise)

Solution: For $x \leq 0$, then the CDF is 0 because the PDF is 0 there. Then for $x \geq 0$, we have that the CDF is

$$F(x) = \int_{-\infty}^{x} f(t)dt = \int_{0}^{x} f(t)dt = -e^{-u}|_{0}^{x^{2}} = 1 - e^{-x^{2}}.$$

(c) (3 points) Find the probability that a randomly picked value from the PDF f from above is between 0 and 1.

Solution: The probability is

$$F(1) - F(0) = (1 - e^{-1^2}) - (1 - e^{-0^2}) = 1 - e^{-1}.$$