Name: \_\_\_\_\_

Circle True or False or leave blank. (1 point for correct answer, -1 for incorrect answer, 0 if left blank)

1. **TRUE** False If  $A \subset B$ , then P(B|A) = 1 (assuming all quantities are well defined).

**Solution:** Writing the formula, we get  $P(A \cap B) = P(A)$ .

2. True **FALSE** If  $P(A), P(B) \neq 0$ , then P(A|B) = P(B|A).

**Solution:** See the quiz problem for a counter-example.

Show your work and justify your answers. Please circle or box your final answer.

- 3. (10 points) Suppose a new cancer test has a 90% chance of correctly identifying that a sick patient has cancer and a 10% chance of incorrectly identifying that a healthy patient has cancer. Assume that 20% of the population has this form of cancer.
  - (a) (2 points) Let A be the event that a random person has cancer and B being the event that a person tests positive for cancer. Write the probabilities you are given in terms of A and B.

**Solution:** Then we are told that  $P(B|A) = \frac{9}{10}$ ,  $P(B|\overline{A}) = \frac{1}{10}$ ,  $P(A) = \frac{1}{5}$ .

(b) (3 points) What is the probability that the test says a random person has cancer?

Solution: We are asked to calculate P(B) and we get  $P(B) = P(B|A)P(A) + P(B|\bar{A})P(\bar{A})$   $= \frac{9}{10}\frac{1}{5} + \frac{1}{10}\frac{4}{5} = \frac{13}{50}.$ 

(c) (5 points) What is the probability that a person who tests positive does not have cancer?

Solution: We are asking for  $P(\bar{A}|B)$  and we use Bayes theorem to get

$$P(\bar{A}|B) = \frac{P(B|\bar{A})P(\bar{A})}{P(B)}$$
$$= \frac{\frac{1}{10}\frac{4}{5}}{\frac{13}{50}} = \frac{4}{13}.$$