

ALGEBRAIC DIFFERENTIAL EQUATIONS FROM COVERING MAPS

THOMAS SCANLON

ABSTRACT. Let Y be a complex algebraic variety, $G \curvearrowright Y$ an action of an algebraic group on Y , $U \subseteq Y(\mathbb{C})$ a complex submanifold, $\Gamma < G(\mathbb{C})$ a discrete, Zariski dense subgroup of $G(\mathbb{C})$ which preserves U , and $\pi : U \rightarrow X(\mathbb{C})$ an analytic covering map of the complex algebraic variety X expressing $X(\mathbb{C})$ as $\Gamma \backslash U$. We note that the theory of elimination of imaginaries in differentially closed fields produces a generalized Schwarzian derivative $\tilde{\chi} : Y \rightarrow Z$ (where Z is some algebraic variety) expressing the quotient of Y by the action of the constant points of G . Under the additional hypothesis that the restriction of π to some set containing a fundamental domain is definable in an o-minimal expansion of the real field, we show as a consequence of the Peterzil-Starchenko o-minimal GAGA theorem that the *prima facie* differentially analytic relation $\chi := \tilde{\chi} \circ \pi^{-1}$ is a well-defined, differential constructible function. The function χ nearly inverts π in the sense that for any differential field K of meromorphic functions, if $a, b \in X(K)$ then $\chi(a) = \chi(b)$ if and only if after suitable restriction there is some $\gamma \in G(\mathbb{C})$ with $\pi(\gamma \cdot \pi^{-1}(a)) = b$.

1. INTRODUCTION

As is well-known, the complex exponential function $\exp : \mathbb{C} \rightarrow \mathbb{C}^\times$ admits a local analytic inverse, the logarithm function, but the logarithm cannot be made into a globally defined analytic function. The ambiguity in the choice of a branch of the logarithm comes from addition by an element of the discrete group $2\pi i\mathbb{Z}$. Hence, if we regard the logarithm as acting on meromorphic function via $f \mapsto \log \circ f$, then while the operator $f \mapsto \log(f)$ is not well-defined, the logarithmic derivative, $f \mapsto \frac{d}{dz}(\log(f))$ is. Of course, more is true in that the logarithmic derivative is given by the simple differential algebraic formula $\frac{d}{dz}(\log(f)) = \frac{f'}{f}$.

There are many ways to see that the logarithmic derivative is a differential rational function. This fact follows from a direct computation, or from Kolchin's general theory of logarithmic differentiation on algebraic groups [12], or from the techniques we employ in this paper. In Section 5 we discuss the algebraic construction of the logarithmic derivative in detail.

The purpose of this paper is to show that under very general hypotheses, differential analytic operators constructed by inverting analytic covering maps and then

The author was partially supported by NSF grants DMS-1001550 and DMS-1363372. This paper was completed while the author was in residence at MSRI during the Spring 2014 semester for the Model Theory and Number Theory program which was partially supported by NSF grant 0932078 000. The author thanks D. Bertrand, A. Buium, J. Freitag, B. Mazur, Y. Peterzil, M. Singer, and S. Starchenko for their very helpful discussions about this work. The author thanks the anonymous referee for a close reading of an earlier version of this paper and for numerous useful suggestions for improvements of this text.

applying differential operators to kill the action of the constant points of some algebraic group are actually differential algebraic. Let us describe more precisely what we have in mind. We are given an algebraic group G over \mathbb{C} , a complex algebraic variety Y , a regular action of G on Y , a complex submanifold $U \subseteq Y(\mathbb{C})$ of $Y(\mathbb{C})$, a Zariski dense subgroup $\Gamma < G(\mathbb{C})$ for which Γ preserves U and an analytic covering map $\pi : U \rightarrow X(\mathbb{C})$ expressing the complex algebraic variety X as the quotient $\Gamma \backslash U$. Because π is a covering map, the inverse $\pi^{-1} : X(\mathbb{C}) \rightarrow U$ is a many-valued analytic function, well-defined up to the action of Γ . Using the theory of elimination of imaginaries in differential fields, we produce a differential constructible function, which we call a generalized Schwarzian associated to the action of G on Y , $\tilde{\chi} : Y \rightarrow Z$ (where Z is an algebraic variety) so that for any differential field M having field of constants \mathbb{C} and points $a, b \in Y(M)$ one has $\tilde{\chi}(a) = \tilde{\chi}(b)$ if and only if there is some $\gamma \in G(\mathbb{C})$ with $\gamma \cdot a = b$. Hence, the differential analytic operator $\chi := \tilde{\chi} \circ \pi^{-1}$ gives a well defined function $X(M) \rightarrow Z(M)$ for M any field of meromorphic functions.

Under a mild hypothesis on π , namely that there is an o-minimal expansion of \mathbb{R} in which there is a definable subset $F \subseteq Y(\mathbb{C})$ for which the restriction of π is definable and surjective onto $X(\mathbb{C})$, we then deduce from a remarkable theorem of Peterzil-Starchenko that the *a priori* differential *analytically* constructible function χ is in fact differential *algebraically* constructible. That is, the map χ may be expressed piecewise as a rational function of its argument and some its derivatives and pieces in the decomposition of its domain are defined by finitely many algebraic differential equations and inequations. The definability hypothesis holds in many cases of interest, such as for the covering maps associated to moduli spaces of abelian varieties and of the universal families of abelian varieties over these moduli spaces.

Our work on this problem was motivated by our attempt to understand Buium's construction of differential rational functions on moduli spaces of abelian varieties whose fibres are finite dimensional differential varieties containing the isogeny classes encoded by such moduli points [5]. His construction is algebraic in the style of Kolchin's construction of logarithmic derivations, though much more sophisticated. Buium's maps are differential rational, meaning that they have a non-trivial indeterminacy locus; ours are defined everywhere, though they are intrinsically differential constructible, meaning that they are only piecewise given by differential regular functions. We seek an algebraic interpretation, for example in terms of a variant of the notion of δ -Hodge structure, for our maps on Buium's indeterminacy locus. More generally, knowing that the map χ is differential algebraic, one expects a direct algebraic construction. We speculate about this in Section 6.

This paper is organized as follows. We begin in Section 2 by recalling some of the basics of differential algebra, complex analysis and especially the Peterzil-Starchenko theory of o-minimal complex analysis. In Section 3 we state precisely and prove our main theorem. In Section 4 we develop some of the basic theory of generalized logarithmic derivatives. In Section 5 we discuss specific examples of covering maps to which our main theorem applies. In particular, we note the existence of differential constructible functions whose fibres are the Kolchin closures of isogeny classes in moduli spaces of abelian varieties. We also discuss the problem of extending our main theorem to the context of Picard-Fuchs equations associated to families of varieties and to the analytic construction of Manin homomorphisms

for nonconstant abelian varieties coming from families of covering maps. We close in Section 6 with some natural questions.

2. PRELIMINARIES

As our main theorem involves a comparison of three different kinds of structures, namely, complex manifolds, differential algebraic functions, and o-minimally definable sets, it should come as no surprise that sometimes the same object may be considered differently in each of these domains. In this section we establish our notation and explain how these theories interact with each other.

2.1. Jet spaces. We shall use a construction of higher order tangent spaces which goes variously under the names of spaces of jets, arc spaces, and prolongation spaces. We begin by recalling the differential geometric jet spaces modified slightly for the complex analytic category. The reader can find details of the space of jets construction in Chapter 12 of [4], though the discussion there covers only jets of maps from one manifold to another while we shall consider jets of germs. The extension to our case is routine.

We begin with our notation for balls and polydiscs.

Definition 2.1. We denote the unit disc in the complex plane by $\mathbb{B} := \{z \in \mathbb{C} : |z| < 1\}$. For a natural number n , the polydisc $\mathbb{B}^n := \{(z_1, \dots, z_n) \in \mathbb{C}^n : |z_i| < 1 \text{ for } i \leq n\}$ is the n^{th} Cartesian power of \mathbb{B} . More generally, for $0 < r < 1$ we denote the disc of radius r centered at the origin by \mathbb{B}_r and its n^{th} Cartesian power by \mathbb{B}_r^n .

Let us recall the construction of the space of jets.

Definition 2.2. If M is a complex manifold and $f : \mathbb{B}_r^n \rightarrow M$ and $g : \mathbb{B}_s^n \rightarrow M$ are two analytic maps into M from polydiscs of some radii r and s and m is a natural number, then we say that f and g have the same jets up to order m if there is a coordinate neighborhood $U \ni f(0) = g(0)$ with $\eta : U \cong \mathbb{B}^{\dim(M)}$ so that for each $j \leq \dim(M)$ and each multi-index $\alpha = (\alpha_1, \dots, \alpha_n)$ with $|\alpha| = \sum \alpha_i \leq m$ one has $\frac{\partial^{|\alpha|}}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}}(\eta_j \circ f)(0) = \frac{\partial^{|\alpha|}}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}}(\eta_j \circ g)(0)$. We write $[f]_m$ for the equivalence class of f with respect to the equivalence relation of having the same jets up to order m . Note that $[f]_m$ depends only on the germ of f at the origin.

The set of equivalence classes of maps from n -dimensional polydiscs into M may be given the structure of a complex manifold, $J_{m,n}(M)$, and comes equipped with an analytic map $J_{m,n}(M) \rightarrow M$ making $J_{m,n}(M)$ into an affine bundle over M . The construction is functorial in that if $\phi : M \rightarrow N$ is a map of complex manifolds, then there is an induced morphism $J_{m,n}(\phi) : J_{m,n}(M) \rightarrow J_{m,n}(N)$ given by $J_{m,n}([f]_m) := [\phi \circ f]_{m,n}$.

The space of jets construction has an algebraic geometric counterpart with the notion of arc spaces. For an introduction to arc spaces see [7] in which what we call *arc spaces* are called *jet spaces*. For details of these constructions at the level of generality we use in this paper, see [17].

Definition 2.3. If X is a scheme over \mathbb{C} , then for each n and m the functor from \mathbb{C} -algebras to sets given by $R \mapsto X(R[\epsilon_1, \dots, \epsilon_n]/(\epsilon_1, \dots, \epsilon_n)^{m+1})$ is represented by a scheme $\mathcal{A}_{m,n}(X)$. In the literature, this construction is usually limited to the case of $n = 1$ for which $\mathcal{A}_{m,1}(X)$ is the m^{th} arc bundle of X .

Remark 2.4. We shall use the more general n to discuss partial differential equations. To ease readability, when no confusion would arise we shall suppress the subscript n writing J_m for $J_{m,n}$ and \mathcal{A}_m for $\mathcal{A}_{m,n}$.

In general, even if X is an algebraic variety, it may happen that $\mathcal{A}_m(X)$ is non-reduced. However, when X is a smooth variety, one has $\mathcal{A}_m(X)(\mathbb{C}) = J_m(X(\mathbb{C}))$ and we will need the arc bundle construction only in the case of smooth varieties.

2.2. Differential algebra. Let us recall some of the basics of differential algebra for which the book [12] is the standard reference. For an introduction to the model theory of differential fields, see [14].

Definition 2.5. A *differential ring* (K, Δ) is a (commutative) ring K given together with a finite sequence $\Delta = \langle \partial_1, \dots, \partial_n \rangle$ of commuting derivations. If K is a field, then we call (K, Δ) a *differential field*. Generally, we write K for the tuple (K, Δ) . When $|\Delta| = 1$, we call K an *ordinary differential field* and otherwise K is a *partial differential field*. In what follows we shall work only with differential fields of characteristic zero and the phrase *differential field* shall mean *differential field of characteristic zero*. A map of differential rings $f : (A, \partial_1, \dots, \partial_n) \rightarrow (B, \partial_1, \dots, \partial_n)$ is given by a map of rings $f : A \rightarrow B$ which respects the derivations in the sense that $f \circ \partial_i = \partial_i \circ f$ for all $i \leq n$. By a K - Δ -*algebra* we mean a differential ring (A, Δ) given together with a K -algebra structure for which the map $K \rightarrow A$ is a map of differential rings.

A standard example of a differential field is given by taking a connected open domain $U \subseteq \mathbb{C}^n$ and setting $(K, \partial_1, \dots, \partial_n) = (\mathcal{M}(U), \frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n})$ where $\mathcal{M}(U)$ the field of meromorphic functions on U where z_1, \dots, z_n are the standard coordinates on U . The Seidenberg embedding theorem asserts that every finitely generated differential field of characteristic zero may be realized as a subdifferential field of $\mathcal{M}(U)$ for some connected domain $U \subseteq \mathbb{C}^n$. In fact, a little more is true.

Theorem 2.6 (Seidenberg [31, 32]). *If $K \subseteq \mathcal{M}(U)$ is a finitely generated differential subfield of the differential field of meromorphic functions on some connected domain $U \subseteq \mathbb{C}^n$ and $L = K\langle u \rangle$ is a simple differential extension of K (meaning that L is a differential field extending K which is generated as a differential field by K and the single element $u \in L$), then there is a connected domain $V \subseteq U$ and an embedding of differential fields $\iota : L \hookrightarrow \mathcal{M}(V)$ compatible with the embedding $\mathcal{M}(U) \rightarrow \mathcal{M}(V)$.*

To speak of differential equations on an algebraic variety, we need to generalize the arc space construction.

Consider $(K, \partial_1, \dots, \partial_n)$ a differential field. For each $m \in \mathbb{N}$ there are two natural K -algebra structures on $R_m = K[\epsilon_1, \dots, \epsilon_n]/(\epsilon_1, \dots, \epsilon_n)^{m+1}$. First, there is the standard structure $\iota : K \rightarrow R_m$ given by $a \mapsto a + \sum_{\mathbf{0} \neq \alpha} 0 \cdot \epsilon^\alpha$. Secondly, using the derivations, we have an exponential map $E_m : K \rightarrow R_m$ given by

$$a \mapsto \sum_{|\alpha| \leq m} \frac{1}{\alpha_1! \cdots \alpha_n!} \partial_1^{\alpha_1} \cdots \partial_n^{\alpha_n}(a) \epsilon_1^{\alpha_1} \cdots \epsilon_n^{\alpha_n} .$$

Note that the standard map ι is itself an exponential map with respect to the trivial derivations.

If X is an algebraic variety over K , then the functor from K -algebras to Set defined by

$$A \mapsto (X \times_{E_m^*} \text{Spec}(R_m))(A \otimes_\iota R_m)$$

is representable by a scheme $\tau_m X$ called the m^{th} *prolongation space* of X . Note that when the derivations are trivial, then $E_m = \iota$ and $\tau_m X = \mathcal{A}_{m,n}(X)$.

From the description of $\tau_m X$ as a functor, one sees that there are natural projection maps $\pi_{m,m'} : \tau_m X \rightarrow \tau_{m'} X$ for $m \geq m'$ corresponding to the reduction maps $R_m \rightarrow R_{m'}$ and that $\tau_0 X = X$ canonically. On the other hand, corresponding to the exponential map $E_m : K \rightarrow R_m$ one has a differentially defined map $\nabla_m : X \rightarrow \tau_m X$. That is, for any K - Δ -algebra A we have a map of sets $\nabla_m : X(A) \rightarrow \tau_m X(A)$ corresponding to the map

$$X(A) \rightarrow (X \times_{E_m^*} \text{Spec}(R_m))(A \otimes_\iota R_m)$$

coming from $1_A \otimes_K E_m$.

Using the prolongation spaces, one can make sense of the notions of differential regular (respectively, rational or constructible) functions on the algebraic variety X . That is, a differential regular (respectively, rational or constructible) function $f : X \rightarrow Y$ (where Y is another algebraic variety over K) is a map from X to Y , considered as functors from the category of K - Δ -algebras to Set which is given by a regular (respectively, rational or constructible) function $f_m : \tau_m X \rightarrow Y$ for some $m \in \mathbb{N}$ in the sense that for any K - Δ -algebra A and point $a \in X(A)$ one has $f(a) = f_m(\nabla_m(a))$.

A differential subvariety Y of X is given by a subvariety $Y_m \subseteq \tau_m X$ for some $m \in \mathbb{N}$. For any K - Δ -algebra A , the set of A -points on Y is

$$Y(A) := \{a \in X(A) : \nabla_m(a) \in Y_m(A)\}.$$

Since here are only interested in differential subvarieties rather than more general differential subschemes, two differential subvarieties Y and Z of X are to be regarded as the same if for every differential field L extending K we have $Y(L) = Z(L)$.

Sets of the form $Y(K) \subseteq X(K)$ are called *Kolchin closed* (or, sometimes, Δ -closed) and as the name suggests are the closed sets of a noetherian topology on $X(K)$. If Y is an irreducible differential subvariety of the algebraic variety X , then the set of differential rational functions on Y forms a differential field denoted by $K\langle Y \rangle$. If $\text{tr. deg}_K(K\langle Y \rangle)$ is finite, then we say that Y is finite dimensional and define $\dim Y := \text{tr. deg}_K(K\langle Y \rangle)$.

An important class of Kolchin closed sets comes from the constants.

Definition 2.7. If (R, Δ) is a differential ring, then

$$R^\Delta := \{a \in R : \partial(a) = 0 \text{ for all } \partial \in \Delta\}$$

is the *ring of constants*. We sometimes write $C = C_R$ for R^Δ .

If $(K, \partial_1, \dots, \partial_n)$ is a differential field and X is an algebraic variety over the constants K^Δ , then as we noted above $\tau_m(X_K) = \mathcal{A}_{m,n}(X_K)$, canonically, and there is a natural regular section $z_m : X_K \rightarrow \tau_m(X_K)$ of the projection map $\tau_m(X_K) \rightarrow \tau_0(X_K) = X_K$ corresponding to the standard map $\iota : K \rightarrow R_m$. We define X^Δ to be the differential subvariety of X given by $z(X_K) \subseteq \tau_1(X_K)$. At the level of points, if A is K - Δ -algebra, then $X^\Delta(A) = X(A^\Delta) \subseteq X(A)$. Note that

if X is an algebraic variety over the constants, then its dimension as an algebraic variety is equal to the dimension of X^Δ as a differential variety.

Just as algebraically closed fields play the role of universal domains for ordinary algebraic varieties, differentially closed fields serve as universal domains for differential algebraic geometry. Here a *differentially closed field* $(\mathbb{U}, \partial_1, \dots, \partial_n)$ is an existentially closed differential field in the sense that if X is an algebraic variety over \mathbb{U} and $Y \subseteq X$ is a differential subvariety for which there is some differential field extension $(L, \partial_1, \dots, \partial_n) \supseteq (\mathbb{U}, \partial_1, \dots, \partial_n)$ with $Y(L) \neq \emptyset$, then $Y(\mathbb{U}) \neq \emptyset$. The theory of differentially closed fields (of characteristic zero in n commuting derivations) is axiomatized by a first-order theory $\text{DCF}_{0,n}$ in the language of rings augmented by n unary function symbols to be interpreted as the distinguished derivations. For us, the crucial facts about $\text{DCF}_{0,n}$ are:

- $\text{DCF}_{0,n}$ eliminates quantifiers. Geometrically, this means that if \mathbb{U} is a differentially closed field, $Y \subseteq X$ is a differential subvariety of an algebraic variety over X and $f : Y \rightarrow Z$ is a differential rational function, then $f(Y(\mathbb{U}))$ is a differentially constructible subset of $Z(\mathbb{U})$. That is, it is a finite Boolean combination of Kolchin closed sets.
- It follows from Theorem 2.6 that if $K \subseteq \mathcal{M}(U)$ is a finitely generated differential field of meromorphic functions on some domain U , then there is a differentially closed field \mathbb{U} containing K which may be realized as a subfield of a differential field of germs of meromorphic functions at some point in U .
- Finally, $\text{DCF}_{0,n}$ *eliminates imaginaries*.

The precise content of elimination of imaginaries is given by the following theorem of Poizat in the case of $n = 1$ and of McGrail in general.

Theorem 2.8 (Poizat [28], McGrail [15]). *The theory $\text{DCF}_{0,n}$ of differentially closed fields of characteristic zero with n commuting derivations eliminates imaginaries. More concretely, if \mathbb{U} is a differentially closed field, $X \subseteq \mathbb{U}^m$ is a definable set and $E \subseteq X \times X$ is a definable equivalence relation, then there is a differential constructible function $\eta : X \rightarrow \mathbb{U}^\ell$ so that for $(a, b) \in X \times X$ one has $\eta(a) = \eta(b) \iff aEb$. Moreover, η may be defined over the same parameters required for the definitions of X and E .*

Prima facie, Theorem 2.8 applies only to differentially closed fields, but the map η may be taken to be defined over the parameters required to define X and E and then from the constructibility of η it is easy to see that for any differential field K over which everything is defined, if $a, b \in X(K)$, then $aEb \iff \eta(a) = \eta(b)$. The proof of Theorem 2.8 passes through the corresponding elimination of imaginaries theorem for algebraically closed fields (which is also proven in detail in [28]) and for our applications, we shall need to unwind this connection between elimination of imaginaries for differentially closed fields and for ordinary algebraically closed fields.

2.3. O-minimality. Finally, we shall make use of the theory of o-minimality for which the book [35] is a good introduction.

Definition 2.9. An o-minimal structure on the real numbers $\mathbb{R}_{\mathcal{F}}$ is given by the choice of a distinguished set \mathcal{F} of functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$, where n may depend on

f , having the property that in the first-order structure

$$\mathbb{R}_{\mathcal{F}} := (\mathbb{R}, +, \cdot, 0, 1, <, \{f\}_{f \in \mathcal{F}})$$

every definable subset of \mathbb{R} is a finite union of points and intervals.

Remark 2.10. With the usual definition of an o-minimal structure, any expansion of the language of ordered sets is allowed. However, it follows from the existence of definable choice functions that every o-minimal expansion of an ordered field is bi-definable with one obtained by expanding the language of ordered rings by function symbols.

That interesting o-minimal structures exist at all is highly non-trivial. For us, the most important o-minimal structure is $\mathbb{R}_{\text{an,exp}}$ in which the set of distinguished functions consists of the real exponential function $\exp : \mathbb{R} \rightarrow \mathbb{R}$ and local analytic functions. That is, for each real analytic function f defined on some open neighborhood of the unit box $[-1, 1]^n$ we are given the function $\bar{f} : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by $\bar{f}(x_1, \dots, x_n) = f(x_1, \dots, x_n)$ if $(x_1, \dots, x_n) \in [-1, 1]^n$ and $\bar{f}(x_1, \dots, x_n) = 0$ otherwise. O-minimality of $\mathbb{R}_{\text{an,exp}}$ is established in [36].

O-minimality implies many strong regularity properties of the definable sets in any number of variables. One result which we shall use is the existence of definable choice functions. If $f : X \rightarrow Y$ is a definable (in the o-minimal structure $\mathbb{R}_{\mathcal{F}}$) surjective function, then there is a definable right inverse $g : Y \rightarrow X$ (see Proposition 1.2 in [35]).

Since o-minimality is fundamentally a theory of ordered structures, it does not directly apply to complex analysis. However, by realizing \mathbb{C} as \mathbb{R}^2 via the real and imaginary parts, one may interpret complex analysis within an o-minimal structure and in the series of papers [19, 20, 22, 23, 24, 25] Peterzil and Starchenko do just this. For us, the most important result from their work is the following strengthening of Chow's Theorem.

Theorem 2.11 (Peterzil-Starchenko, Corollary 4.5 of [23]). *Let $\mathbb{R}_{\mathcal{F}}$ be some o-minimal structure on the real numbers. Let X be a quasiprojective algebraic variety over \mathbb{C} . If $Y \subseteq X(\mathbb{C})$ is an $\mathbb{R}_{\mathcal{F}}$ -definable, closed complex analytic set in X , then Y is algebraic.*

Remark 2.12. The statement of Corollary 4.5 in [23] takes $X(\mathbb{C}) = \mathbb{C}^n$, but the proof uses only the fact that X embeds into a projective space.

Remark 2.13. Note that in Theorem 2.11 there is no hypothesis that X be projective nor that f satisfy any kind of growth condition towards the boundary of X in some compactification. On the other hand, one cannot completely avoid such considerations in that to establish that the relevant analytic sets are definable it may be necessary to study their boundary behavior.

The proof of Theorem 2.11 requires much of the Peterzil-Starchenko theory of o-minimal complex analysis, but the basic idea is clear. Many standard theorems in the theory of complex analysis assert that if some closed subset of a complex manifold is generically analytic in some precise sense (for example, with respect to some reasonable dimension) and the ambient space is sufficiently nice, then that set must be analytic (see the theorems of Bishop [3], Remmert and Stein [29], and Shiffman [34]). Sets definable in o-minimal structure enjoy a very smooth dimension

theory as do all of the sets naturally associated to them through standard geometric and elementary analytic constructions because these are also definable.

As a consequence of the flexibility of o-minimal definability, we have the following useful natural generalization of Theorem 2.11.

Corollary 2.14. *Let $\mathbb{R}_{\mathcal{F}}$ be an o-minimal expansion of the real field, X a complex algebraic variety and $Z \subseteq X(\mathbb{C})$ a definable subset which is also analytically constructible, that is, a finite Boolean combination of complex analytic subsets of $X(\mathbb{C})$. Then Z is algebraically constructible.*

Proof. We work by induction on the dimension of the analytic-Zariski closure of Z , $\overline{Z}^{\text{an-Zar}}$, that is, the smallest complex analytic set containing Z . Using the Boolean disjunctive normal form and the local Noetherianity of the analytic-Zariski topology, we may express Z as a finite union of sets of the form $A \setminus B$ where A and B are complex analytic and $B \subsetneq A$. Replacing A with the union of its components which are not contained in B , we may assume that no component of A is contained in B . Then B is nowhere dense in A with respect to the Euclidean topology so that the Euclidean closure of $A \setminus B$, $\overline{A \setminus B}^{\text{Euc}}$, is A , which is also equal to $\overline{A \setminus B}^{\text{an-Zar}}$ as the Euclidean topology is finer than the analytic-Zariski topology. For any set definable in an o-minimal expansion of the real field, the Euclidean closure of that set is also definable using the usual first-year analysis definition of the closure. Thus, we see that $\overline{Z}^{\text{an-Zar}}$ is also definable, and, hence, an algebraic variety by Theorem 2.11. The set $\overline{Z}^{\text{an-Zar}} \setminus Z$ is also analytically constructible and definable and its analytic-Zariski closure has strictly smaller dimension. Hence, it is algebraically constructible and so is $Z = \overline{Z}^{\text{an-Zar}} \setminus (\overline{Z}^{\text{an-Zar}} \setminus Z)$. \square

3. MAIN THEOREM

With our preliminaries in place, we flesh out the sketch of our main theorem from the introduction. Throughout this section, we fix a natural number n and when we speak of a differential field we mean one with n distinguished derivations. Likewise, when speaking of arc and jet spaces and differential fields we suppress the index n . That is, we write J_m for $J_{m,n}$ and \mathcal{A}_m for $\mathcal{A}_{m,n}$.

In what follows we shall make the following hypotheses.

- $\mathbb{R}_{\mathcal{F}}$ is a fixed o-minimal structure on the reals. Throughout the rest of this section by “definable” we mean “definable in $\mathbb{R}_{\mathcal{F}}$ ”.
- G is an algebraic group over \mathbb{C} .
- Y is a complex algebraic variety and $G \times Y \rightarrow Y$ is a regular function expressing a faithful action $G \curvearrowright Y$ of G on Y .
- $U \subseteq Y(\mathbb{C})$ is a complex submanifold of the \mathbb{C} -points of Y .
- $\Gamma < G(\mathbb{C})$ is a Zariski dense, discrete subgroup of $G(\mathbb{C})$.
- Via the restriction of the action of $G(\mathbb{C})$ on $Y(\mathbb{C})$ to U , Γ acts as a group of automorphisms of U .
- $\pi : U \rightarrow X(\mathbb{C})$ is a complex analytic covering map of the algebraic variety X expressing $X(\mathbb{C})$ as the quotient $\Gamma \backslash U$.
- $F \subseteq Y(\mathbb{C})$ is an open $\mathbb{R}_{\mathcal{F}}$ -definable subset of $Y(\mathbb{C})$ for which the restriction $\pi \upharpoonright F : F \rightarrow X(\mathbb{C})$ is $\mathbb{R}_{\mathcal{F}}$ -definable and surjective onto $X(\mathbb{C})$.

Remark 3.1. Let us note that for each $\gamma \in \Gamma$, the set $\gamma \cdot F$ is definable since the action of $G(\mathbb{C})$ on $Y(\mathbb{C})$ is algebraic and hence *a fortiori* definable. Thus, we may cover U by a set of definable sets indexed by Γ .

Remark 3.2. It follows from the existence of definable choice functions in o-minimal expansions of ordered fields that there is a definable $F' \subseteq F$ so that the restriction of π to F' is definable and a bijection between F' and $X(\mathbb{C})$. It is convenient for us to take F to be open in which case we cannot assume that the restriction of π to F is one-to-one.

From these data we construct a differential analytic map χ on X which nearly inverts π . We shall call this resulting map the *generalized logarithmic derivative associated to π* .

We begin with two lemmata on jet spaces, the first showing that jets of covering maps are themselves covering maps and the second showing that jets of definable functions are also definable.

Lemma 3.3. *For each natural number m , there is a natural action of Γ on $J_m(U)$ and with respect to this action, $J_m(\pi)$ expresses $\mathcal{A}_m(X)(\mathbb{C})$ as $\Gamma \backslash J_m(U)$.*

Proof. Define the action of Γ on $J_m(U)$ by $\gamma \cdot x := J_m(\gamma \cdot)(x)$. By the functoriality of the space of jets construction, we see that for each $\gamma \in \Gamma$ we have $J_m(\pi) \circ \gamma \cdot = J_m(\pi) \circ J_m(\gamma \cdot) = J_m(\pi \circ \gamma \cdot) = J_m(\pi)$. That is, $J_m(\pi)$ is invariant under precomposition with the action of Γ . Let us check now that if x and y are two points of $J_m(U)$ having the same image under $J_m(\pi)$, then there is some $\gamma \in \Gamma$ with $\gamma \cdot x = y$. Taking the images in U , we see that their images \bar{x} and \bar{y} in U have the same image under π . Hence, there is some $\gamma \in \Gamma$ with $\gamma \cdot \bar{x} = \bar{y}$. As π is a covering map, we can find a neighborhood $V \ni \bar{y}$ on which $\pi \upharpoonright V : V \rightarrow \pi(V) \subseteq X(\mathbb{C})$ is biholomorphic. By functoriality of J_m , $J_m(\pi) \upharpoonright J_m(V) : J_m(V) \rightarrow J_m(\pi(V)) \subseteq \mathcal{A}_m(X)(\mathbb{C})$ is also biholomorphic. In particular, because $J_m(\pi) \upharpoonright J_m(V)(y) = J_m(\pi)(y) = J_m(\pi)(x) = J_m(\pi)(\gamma \cdot x) = J_m(\pi) \upharpoonright J_m(V)(\gamma \cdot x)$, we must have $y = \gamma \cdot x$. \square

With our second result on jets we note that the jet of a definable function is itself definable.

Lemma 3.4. *Fix an o-minimal expansion $\mathbb{R}_{\mathcal{F}}$ of the real field. Let Z be a complex algebraic variety, $V \subseteq Z(\mathbb{C})$ an open subset of $Z(\mathbb{C})$, W a complex algebraic variety, m a natural number and $f : V \rightarrow W(\mathbb{C})$ a definable analytic function. Then $J_m(f) : J_m(V) \rightarrow \mathcal{A}_m(W)(\mathbb{C})$ is a definable analytic function.*

Proof. For the sake of legibility, we suppress the subscripts from J and \mathcal{A} . Let us write $\nu : \mathcal{A}(Z) \rightarrow Z$ for the natural projection map.

Note that because the function f is definable, so is its domain V . Using the identification $\mathcal{A}(Z)$ with $J(Z)$, we see $J(V)$ as the definable set $\nu^{-1}(V)$.

Analyticity of $J(f)$ is a general feature of the jet of an analytic function (see [4]). Definability of $J(f)$ follows from the facts that for any definable complex analytic function its complex derivatives are definable and the jet space $J(V) = \nu^{-1}V$ is definable as a subset of $\mathcal{A}(Z)(\mathbb{C})$. Indeed, the usual limit definition of a complex derivative may be naturally expressed using the formula defining f and the field operations. As f is analytic, it follows that all of its derivatives of all orders are definable. Read in coordinates, the map $J(f)$ is given by the usual Faà di

Bruni formula which is polynomial in the derivatives of f up to order m and the coordinates on the jet space. \square

Remark 3.5. One could generalize Lemma 3.4 to the case that $V \subseteq Z(\mathbb{C})$ is merely a complex submanifold by noting that the jet space of V at a point as a submanifold of $\mathcal{A}(Z)(\mathbb{C})$ may be identified by iterating the standard definitions of tangent spaces from calculus.

The relation on Y defined by $x \sim y$ if and only if $(\exists g \in G(\mathbb{C}))g \cdot x = y$ is an equivalence relation which on \mathbb{C} -points identifies everything, but when interpreted in a differential field M with field of constants \mathbb{C} is a nontrivial definable equivalence relation. In particular, when \mathbb{U} is a differentially closed field with field of constants \mathbb{C} , Theorem 2.8 says that there is a differential constructible function $\tilde{\chi}$ defined on Y having the property that for $a, b \in Y(\mathbb{U})$ one has $\tilde{\chi}(a) = \tilde{\chi}(b)$ if and only if $(\exists g \in G(\mathbb{C}))g \cdot a = b$. Since $\tilde{\chi}$ is differential constructible, if $\mathbb{C} \subseteq M \subseteq \mathbb{U}$ is some intermediate differential field, it is still the case that for $a, b \in Y(M)$ the equality $\tilde{\chi}(a) = \tilde{\chi}(b)$ holds if and only if $(\exists g \in G(\mathbb{C}))g \cdot a = b$.

Remark 3.6. In the case that $Y = \mathbb{P}^1$ and $G = \mathrm{PGL}_2$ acting by fractional linear transformations, then the function $\tilde{\chi}$ may be identified with the classical Schwarzian derivative. Similar maps, sometimes with explicit formulae, appear in the literature for other quotients of algebraic varieties by the constant points of an algebraic group (see, for example, the generalized Schwarzians for $Y = \mathbb{P}^n$ and $G = \mathrm{PGL}_{n+1}$ in [5] or [30]). In analogy with the classical Schwarzian, we shall call the map $\tilde{\chi}$ a *generalized Schwarzian derivative*.

Our generalized Schwarzian $\tilde{\chi}$ admits a more algebraic description. Being a differential constructible function, there is some $m \in \mathbb{Z}_+$ and a constructible function $\tilde{\chi}_m$ defined on $\tau_m Y$ for which $\tilde{\chi} = \tilde{\chi}_m \circ \nabla_m$. The algebraic group G acts algebraically on $\tau_m Y = \mathcal{A}_m(Y)$ for precisely the same reason that Γ acts on $J_m U$. While quotients in the category of algebraic varieties with regular maps as morphisms do not always exist, they do always exist constructibly. That is, there is some constructible function ξ_m on $\mathcal{A}_m(Y)$ expressing the quotient $G \backslash \mathcal{A}_m(Y)$. We aim to show that for $m \gg 0$ one may take $\tilde{\chi}_m = \xi_m$ so that $\tilde{\chi} = \xi_m \circ \nabla_m$.

To this end we first prove two basic lemmata on differential algebraic geometry, one result about definable equivalence relations and a second about the connection between the Kolchin topology on a variety and the Zariski topology on its prolongations.

Lemma 3.7. *Let \mathbb{U} be a differentially closed field. If X is a differential variety over \mathbb{U} and \sim is a definable equivalence relation on X , then for a and b elements of $X(\mathbb{U})$, one has $a \sim b$ if and only if $\overline{[a]_\sim}^{\mathrm{Kolchin}} = \overline{[b]_\sim}^{\mathrm{Kolchin}}$ where we write $\overline{A}^{\mathrm{Kolchin}}$ for the Kolchin closure of the set A .*

Proof. Certainly, if $a \sim b$, then $[a]_\sim = [b]_\sim$ so that $\overline{[a]_\sim}^{\mathrm{Kolchin}} = \overline{[b]_\sim}^{\mathrm{Kolchin}}$. Conversely, by quantifier elimination in differentially closed fields we know that the sets $[a]_\sim$ and $[b]_\sim$ are differentially constructible. Hence, there is a (Kolchin) dense open in $\overline{[a]_\sim}^{\mathrm{Kolchin}}$ set $V \subseteq [a]_\sim$ and a dense open in $\overline{[b]_\sim}^{\mathrm{Kolchin}}$ set $W \subseteq [b]_\sim$. If $\overline{[a]_\sim}^{\mathrm{Kolchin}} = \overline{[b]_\sim}^{\mathrm{Kolchin}}$, then $V \cap W$ is a dense open subset of $\overline{[a]_\sim}^{\mathrm{Kolchin}}$. In

particular, it is nonempty so that $[a]_{\sim} \cap [b]_{\sim} \supseteq V \cap W$ is non-empty implying that $[a]_{\sim} = [b]_{\sim}$. \square

The Kolchin topology and the Zariski topology are related through the prolongation spaces. With the next lemma we show how to compute a Kolchin closure via Zariski closures in prolongation spaces.

Lemma 3.8. *Let \mathbb{U} be a differentially closed field and X be an algebraic variety over \mathbb{U} and let $A \subseteq X(\mathbb{U})$ be a set of \mathbb{U} -points on X . Then*

$$\overline{A}^{\text{Kolchin}} = \bigcap_{m \geq 0} \nabla_m^{-1}(\overline{\nabla_m(A)}^{\text{Zariski}}).$$

Moreover, this set may be realized as $\nabla_M^{-1}(\overline{\nabla_M(A)}^{\text{Zariski}})$ for M sufficiently large.

Proof. Each of the sets $\nabla_m^{-1}(\overline{\nabla_m(A)}^{\text{Zariski}})$ is a Kolchin closed set containing A . Hence, $\overline{A}^{\text{Kolchin}}$ is contained in $\bigcap_{m \geq 0} \nabla_m^{-1}(\overline{\nabla_m(A)}^{\text{Zariski}})$.

Before embarking on the proof of the other inclusion let us note that the intersection defining the righthand side of our purported equation is actually a descending intersection. That is, for each m we have $\nabla_{m+1}^{-1}(\overline{\nabla_{m+1}(A)}^{\text{Zariski}}) \subseteq \nabla_m^{-1}(\overline{\nabla_m(A)}^{\text{Zariski}})$. Indeed, the prolongation spaces form a projective system and the ∇ maps respect this system in the sense that if $\nu_{m+1,m} : \tau_{m+1}X \rightarrow \tau_m X$ is the projection map from level $m+1$ to level m , then $\nabla_m = \pi_{m+1,m} \circ \nabla_{m+1}$. Thus, if $Y \subseteq \tau_m X$ is a Zariski closed set containing $\nabla_m(A)$, then $\nu_{m+1,m}^{-1}(Y)$ is a Zariski closed subset of $\tau_{m+1}X$ containing $\nabla_{m+1}(A)$ so that

$$\nabla_{m+1}^{-1}(\overline{\nabla_{m+1}(A)}^{\text{Zariski}}) \subseteq \nabla_{m+1}^{-1}(\nu_{m+1,m}^{-1}(\overline{\nabla_m(A)}^{\text{Zariski}})) = \nabla_m^{-1}(\overline{\nabla_m(A)}^{\text{Zariski}}).$$

For the inclusion of the righthand side in the left, consider a Kolchin closed set $Y \subseteq X$ which contains A . By the definition of the Kolchin topology, there is some m and a Zariski closed $Y_m \subseteq \tau_m X$ with $Y = \nabla_m^{-1}Y_m$. Since $A \subseteq Y(\mathbb{U})$, we have $\nabla_m(A) \subseteq Y_m$. Thus, $\overline{\nabla_m(A)}^{\text{Zariski}} \subseteq Y_m$ so that $\nabla_m^{-1}(\overline{\nabla_m(A)}^{\text{Zariski}}) \subseteq Y$. Clearly, for any $\ell \geq m$ if we set $Y_\ell := \nu_{\ell,m}^{-1}Y_m$, then $Y = \nabla_\ell^{-1}Y_\ell$. Thus, for all $\ell \geq m$ we have $\nabla_\ell^{-1}(\overline{\nabla_\ell(A)}^{\text{Zariski}}) \subseteq Y$. On the other hand, by the observation above, we have $\bigcap_{M \geq 0} \nabla_M^{-1}(\overline{\nabla_M(A)}^{\text{Zariski}}) = \bigcap_{M \geq m} \nabla_M^{-1}(\overline{\nabla_M(A)}^{\text{Zariski}}) \subseteq Y$.

The moreover clause is an immediate consequence of the noetherianity of the Kolchin topology. \square

We employ the above lemmata to relate the quotient of an algebraic variety by the constant points of an algebraic group to algebraic quotients.

Proposition 3.9. *Let \mathbb{U} be a differentially closed field, G an algebraic group over the constant field $C = \mathbb{U}^\Delta$, X an algebraic variety over C and $G \curvearrowright X$ an action of G on X over C . For \mathbb{U} -points $a, b \in X(\mathbb{U})$ the following are equivalent.*

- There is some $g \in G(C)$ with $g \cdot a = b$.
- For all $m \in \mathbb{N}$ there is some $g_m \in G(\mathbb{U})$ with $g_m \cdot \nabla_m(a) = \nabla_m(b)$.

Moreover, there is some natural number M so that these conditions are equivalent to

- There is some $g_M \in G(\mathbb{U})$ with $g_M \cdot \nabla_M(a) = \nabla_M(b)$.

Proof. For this proof, we write \overline{A} for the Zariski closure of a set A and $\overline{A}^{\text{Kolchin}}$ for its Kolchin closure.

Define the equivalence relation \sim on $X(\mathbb{U})$ by

$$a \sim b : \iff (\exists g \in G(C)) g \cdot a = b$$

and let \approx be given by

$$a \approx b : \iff (\forall m \in \mathbb{N})(\exists g_m \in G(\mathbb{U})) g_m \cdot \nabla_m(a) = \nabla_m(b).$$

Note that \sim is a definable equivalence relation while *prima facie* \approx is merely type-definable. Clearly, $a \sim b \Rightarrow a \approx b$ for if $g \in G(C)$ satisfies $g \cdot a = b$, then for each m we may take $g_m := \nabla_m(g)$ (which belongs to the image of the zero section as the constant points are precisely those whose image under ∇_m agree with the zero section) to witness that there is some $g_m \in G(\mathbb{U})$ with $g_m \cdot \nabla_m(a) = \nabla_m(g) \cdot \nabla_m(a) = \nabla_m(g \cdot a) = \nabla_m(b)$.

On the other hand, we observe that for each m the Zariski closure of $\nabla_m([a]_{\sim})$ is the Zariski closure of the $G(\mathbb{U})$ -orbit of $\nabla_m(a)$. Indeed, consider the algebraic group S defined as the stabilizer of $\overline{\nabla_m([a]_{\sim})}$ in G :

$$S(\mathbb{U}) := \{g \in G(\mathbb{U}) : g \cdot \overline{\nabla_m([a]_{\sim})} \subseteq \overline{\nabla_m([a]_{\sim})}\}$$

As $\nabla_m([a]_{\sim})$ is a homogeneous space for $G(C)$ under the natural action of $G(\mathbb{U})$ on $\tau_m X$, we see that $G(C) \subseteq S(\mathbb{U}) \subseteq G(\mathbb{U})$. As $G(C)$ is Zariski dense in $G(\mathbb{U})$ and S is closed, we conclude that $S = G$. Thus, $\overline{\nabla_m([a]_{\sim})} \supseteq \overline{G \cdot \nabla_m(a)}$. On the other hand, since $\nabla_m([a]_{\sim}) = G(C) \cdot \nabla_m(a) \subseteq G(\mathbb{U}) \cdot \nabla_m(a)$, we must have equality.

Therefore, if $a \approx b$, then for each $m \in \mathbb{N}$ by definition over \approx we have that $G(\mathbb{U}) \cdot \nabla_m(a) = G(\mathbb{U}) \cdot \nabla_m(b)$ so that $\overline{\nabla_m([a]_{\sim})} = \overline{G(\mathbb{U}) \cdot \nabla_m(a)} = \overline{G(\mathbb{U}) \cdot \nabla_m(b)} = \overline{\nabla_m([b]_{\sim})}$. As this equality is true for every m , $\overline{[a]_{\sim}}^{\text{Kolchin}} = \overline{[b]_{\sim}}^{\text{Kolchin}}$. By Lemma 3.7, $a \sim b$.

The moreover clause follows by the compactness theorem. \square

It follows from elimination of imaginaries in algebraically closed fields [28] that if the algebraic group H acts on an algebraic variety V , then the quotient $V \rightarrow H \backslash V$ may be realized constructibly. In fact, while the map expressing the quotient cannot be taken to be regular, it may be assumed to have some regularity properties.

Lemma 3.10. *Let K be an algebraically closed field, H an algebraic group over K , V an algebraic variety over K , and $H \curvearrowright V$ an action of H on V , also defined over K . Then there are*

- a chain of closed algebraic varieties $\emptyset = V_0 \subsetneq V_1 \subsetneq \cdots \subsetneq V_m = V$,
- an algebraic variety W , and
- regular maps $\xi_i : V_i \setminus V_{i-1} \rightarrow W$ for each positive $i \leq m$

so that

- each V_i is H -invariant and
- for $a, b \in (V_i \setminus V_{i-1})(K)$ one has $\xi_i(a) = \xi_i(b) \iff (\exists g \in H(K)) g \cdot a = b$.

Proof. We work by noetherian induction on V with the case that $\dim(V) = 0$ being immediate. By elimination of imaginaries we find a constructible function $\psi : V \rightarrow W$ to some algebraic variety W expressing $H \backslash V$. (Note: we are not claiming that $W = H \backslash V$.) As ψ is a constructible function, we can find a dense open $U \subseteq V$ for which $\psi \upharpoonright U$ is regular. Set $U' := H \cdot U$ which is again a dense

open subset of V . We claim that $\psi \upharpoonright U'$ is regular. Indeed, we may cover U' with charts of the form gS where $S \subseteq U$ is an open affine in U and $g \in G(K)$. By the H -invariance of ψ , we see that on gS , ψ satisfies $\psi(gx) = \psi(x)$. That is, $\psi \upharpoonright gS$ agrees with $\psi \upharpoonright S$ via the isomorphism $g \cdot : S \rightarrow gS$. Thus, $\psi \upharpoonright gS$ is regular and therefore $\psi \upharpoonright U'$ is regular. Set $V' := V \setminus U'$. By induction, V' admits the requisite chain, of length m , say. Let $V_{m+1} := X$. \square

We are now in a position to prove an algebraic counterpart of our main theorem.

Proposition 3.11. *With the notation and hypotheses as introduced at the beginning of this section, if $\eta : Y \rightarrow Z$ is a constructible function from Y to some algebraic variety Z expressing the quotient $Y \rightarrow G \backslash Y$ as in Lemma 3.10, then $\chi : X \rightarrow Z$ defined by $\chi := \eta \circ \pi^{-1}$ (for any choice of a branch of π^{-1}) is a constructible function.*

Proof. Consider the following set.

$$\Xi := \{(x, z) \in X(\mathbb{C}) \times Z(\mathbb{C}) : (\exists y \in Y(\mathbb{C})) \pi(y) = x \ \& \ \eta(y) = z\}$$

Let us observe that the set Ξ is definable. Indeed, because π is Γ -invariant and η is G -invariant (hence, also Γ -invariant), in the definition of Ξ we may restrict y to F . That is, we have

$$\Xi = \{(x, z) \in X(\mathbb{C}) \times Z(\mathbb{C}) : (\exists y \in F) \pi(y) = x \ \& \ \eta(y) = z\}.$$

As $\pi \upharpoonright F$ is definable, this expression presents Ξ as a definable set.

Not only is the set Ξ definable, but it is the graph of a function. Indeed, using definable choice in $\mathbb{R}_{\mathcal{F}}$, we see that there is a definable function $\zeta : X(\mathbb{C}) \rightarrow F \subseteq Y(\mathbb{C})$ which is a right inverse to π . Again using the Γ -invariance of η we see that Ξ is the graph of $\eta \circ \zeta$. Let us write $\chi : X \rightarrow Z$ for the function whose graph is Ξ . Let us note that while this definition of χ expresses its definability, there are other ways it could be presented. Indeed, from the Γ -invariance of η for any $x \in X(\mathbb{C})$ we have $\chi(x) = \eta(\hat{\zeta}(x))$ where $\hat{\zeta}$ is *any* branch of π^{-1} near x .

We will now show that Ξ may be expressed as a finite union of definable complex analytic subvarieties. By Lemma 3.10, there is a sequence of a closed subvarieties $\emptyset = V_0 \subsetneq V_1 \subsetneq \dots \subsetneq V_\ell = Y$ so that each V_i is G -invariant and the restriction of η to $V_i \setminus V_{i-1}$ is regular. For each $i \leq \ell$, define $\xi_i := (\eta \upharpoonright V_i) \circ \zeta : X \rightarrow Z$. Note that $\xi_\ell = \chi$. We show by induction on i that ξ_i is constructible. The case of $i = 0$ is trivial. Let us consider now the case that $i > 0$. Let us define $W_i := \eta(V_i) \subseteq Z$. Since η is a constructible function, W_i is a constructible subset of Z . Consider $\Xi_i := \Xi \cap (X \times (W_i \setminus W_{i-1}))$. The set Ξ_i being the intersection of two definable sets is definable. Moreover, our second presentation of χ shows that Ξ_i is the graph of an analytic function. Indeed, if $(x, y) \in \Xi_i$, then fix a branch $\hat{\zeta}$ of π^{-1} near x . We have $y = \eta(\hat{\zeta}(x))$ and, in fact, near (x, y) , Ξ_i is the graph of $\eta \circ \hat{\zeta}$. As $y \in W_i \setminus W_{i-1}$ and V_i and V_{i-1} are G -invariant, necessarily $\hat{\zeta}(x) \in (V_i \setminus V_{i-1})(\mathbb{C})$. Thus, near (x, y) , Ξ_i is the graph of the analytic function $(\eta \upharpoonright (V_i \setminus V_{i-1})) \circ \hat{\zeta}$. By the Peterzil-Starchenko o-minimal GAGA Theorem 2.11, Ξ_i is algebraic. By induction, ξ_{i-1} is algebraically constructible and the graph of ξ_i is simply the union of Ξ_i and the graph of ξ_{i-1} . Hence, ξ_i is itself algebraically constructible.

Taking $i = \ell$, we see that χ is algebraically constructible. \square

We deduce our main theorem from Proposition 3.11.

Theorem 3.12. *Work with the notation and hypotheses as introduced at the beginning of this section. Let $\tilde{\chi} : Y \rightarrow Z$ be a generalized Schwarzian for the action of G^Δ on Y , that is, it is a differential constructible function from Y to some algebraic variety Z expressing the quotient $Y \rightarrow G^\Delta \backslash Y$ given by Theorem 2.8. Then $\chi : X \rightarrow Z$ defined by $\chi := \tilde{\chi} \circ \pi^{-1}$ (for any choice of a branch of π^{-1}) is a differential constructible function, which we shall call the generalized logarithmic derivative associated to π .*

Proof. For this proof, by a differential field M we mean one with field of constants $M^\Delta = \mathbb{C}$.

By Proposition 3.9 for $N \gg 0$ there is a constructible function $\tilde{\chi}_N : \mathcal{A}_N(Y) \rightarrow Z$ so that $\tilde{\chi} : Y \rightarrow Z$ takes the form $\tilde{\chi} := \tilde{\chi}_N \circ \nabla_N$ and for any differential field M with field of constants \mathbb{C} one has that

- for points $a, b \in Y(M)$

$$\tilde{\chi}(a) = \tilde{\chi}(b) \iff (\exists g \in G(\mathbb{C})) g \cdot a = b$$

- for points $a, b \in \mathcal{A}_N(Y)(M)$

$$\tilde{\chi}_N(a) = \tilde{\chi}_N(b) \iff (\exists g \in G(M^{\text{alg}})) g \cdot a = b.$$

With this choice of N , we may realize our generalized logarithmic derivative $\chi : X \rightarrow Z$ as $\chi = \tilde{\chi} \circ \pi^{-1} = \tilde{\chi}_N \circ J_N(\pi)^{-1} \circ \nabla_N$ as in the following diagram.

$$\begin{array}{ccccc}
 Y & \xleftrightarrow{\nabla_N} & \mathcal{A}_N(Y) & \xrightarrow{\tilde{\chi}_N} & Z \\
 \uparrow & & \uparrow & & \uparrow \\
 U & \xleftarrow{\quad} & J_N(U) & \xrightarrow{\chi_N} & Z \\
 \uparrow \pi^{-1} & & \uparrow J_N(\pi)^{-1} & & \uparrow \\
 X & \xleftrightarrow{\nabla_N} & \mathcal{A}_N(X) & \xrightarrow{\chi} & Z \\
 & & \downarrow J_N(\pi) & & \downarrow \\
 & & & &
 \end{array}$$

Indeed, the hypotheses of Proposition 3.11 apply with $\mathcal{A}_N(Y)$ in place of Y , $J_N(U)$ in place of U , $J_N(F)$ in place of F , $J_N(\pi)$ in place of π , $\mathcal{A}_N(X)$ in place of X , and $\tilde{\chi}_N$ in place of η . Thus, by Proposition 3.11, $\chi_N := \tilde{\chi}_N \circ J_N(\pi)^{-1} : \mathcal{A}_N(X) \rightarrow Z$ is constructible. Hence, $\chi = \chi_N \circ \nabla_N : X \rightarrow Z$ is differential constructible. \square

Remark 3.13. The construction of χ as a differential meromorphic function is classical. The new content of Theorem 3.12 is that χ is differential *constructible* and that it is defined everywhere on X .

Remark 3.14. The function χ depends on the choice of $\tilde{\chi}$, which is itself well-defined only up to differential constructible isomorphism. However, the equivalence relation defined by $a \sim b : \iff \chi(a) = \chi(b)$ is intrinsic.

4. TOWARDS A THEORY OF GENERALIZED LOGARITHMIC DERIVATIVES

Using our construction of the generalized logarithmic derivatives it is possible to deduce differential algebraic properties of these maps from their analytic interpretation. In this section we draw some of these conclusions though we leave a fine analysis for a future work.

Throughout this section we shall work inside a differentially closed field (\mathbb{U}, Δ) which we shall realize as a differential subfield of some differential field M of germs of meromorphic functions.

We shall work with the conventions and notation of Section 3. By way of notation, for $a \in X(\mathbb{U})$ we denote by F_a the differential constructible subset of X defined by $\chi(x) = \chi(a)$.

Let us begin with a simple observation about the meaning of two points having the same image under χ .

Proposition 4.1. *For $a, b \in X(\mathbb{U})$, one has $\chi(a) = \chi(b)$ if and only if for any choice of a branch of π^{-1} there is some $g \in G(\mathbb{C})$ so that $\pi(g \cdot \pi^{-1}(a)) = b$.*

Proof. By construction, $\chi = \tilde{\chi} \circ \pi^{-1}$ where $\tilde{\chi}(x) = \tilde{\chi}(y)$ if and only if there is some $g \in G(M^\Delta) = G(\mathbb{C})$ with $g \cdot x = y$. Hence, $\chi(a) = \chi(b)$ if and only if there is some $g \in G(\mathbb{C})$ with $g \cdot \pi^{-1}(a) = \pi^{-1}(b)$, or equivalently, $\pi(g \cdot \pi^{-1}(a)) = b$. \square

Using again the analytic interpretation of χ we compute the dimension of a fibre of χ .

Proposition 4.2. *Let $a \in X(\mathbb{U})$. Then $\dim(F_a) \leq \dim(G^\Delta)$ (which is the dimension of G as an algebraic group). In fact, if π^{-1} is any branch of the inverse of π , then recalling that \mathbb{U} is a field of germs of meromorphic functions, $\dim(F_a) = \dim(G^\Delta) - \dim(G_{\pi^{-1}(a)}^\Delta)$, where $G_{\pi^{-1}(a)}^\Delta$ is the stabilizer of $\pi^{-1}(a)$ in G^Δ .*

Proof. As in the proof of Theorem 3.12, we take $N \gg 0$ large enough so that $\chi = \tilde{\chi}_N \circ J_N(\pi)^{-1} \circ \nabla_N$ where $\tilde{\chi}_N : \mathcal{A}_N(Y) \rightarrow Z$ is an algebraically constructible map expressing the quotient $\mathcal{A}_N(Y) \rightarrow G \backslash \mathcal{A}_N(Y)$. Then the differential variety F_a is the pullback by ∇_N of the algebraic subvariety $(F_a)_N$ of $\mathcal{A}_N(X)$ defined by $\chi_N(x) = \chi_N(\nabla_N(a))$. The preimage \tilde{F}_a of $(F_a)_N$ in $J_N(U)$ is a complex analytic variety which contains $G(\mathbb{C}) \cdot J_N(\pi)^{-1}(\nabla_N(a))$. As the map $J_N(\pi)$ is a covering map, the analytic variety \tilde{F}_a has the same dimension as that of $(F_a)_N$. Since the fibre of $\tilde{\chi}_N$ over $\chi(a)$ is precisely the $G(\mathbb{C})$ orbit of $\nabla_N \pi^{-1}(a) = J_N(\pi)^{-1}(\nabla_N(a))$, we see that $\tilde{F}_a = \tilde{\chi}_N^{-1}(\chi(a))$. Hence, the dimension of F_a is equal to the dimension of the orbit of $\nabla_N(\pi^{-1}(a)) = \dim G^\Delta - \dim G_{\pi^{-1}(a)}^\Delta$. \square

Remark 4.3. A similar calculation occurs in [2] in which Bertrand and Zudilin show that the partial differential fields generated by Siegel modular forms have transcendence degrees governed by the groups acting on the covering spaces.

Our calculation of the dimension of the fibres suggests an alternate method to describe the fibres of the generalized logarithmic derivative. Let us introduce the notion of a generalized Hecke correspondence.

Definition 4.4. The commensurability group of Γ is the subgroup of $G(\mathbb{C})$ defined by

$$\Gamma^{\text{comm}} := \{ \gamma \in G(\mathbb{C}) : \gamma(U) = U \text{ \& } [\Gamma : \Gamma^\gamma \cap \Gamma] < \infty \text{ \& } [\Gamma^\gamma : \Gamma^\gamma \cap \Gamma] < \infty \} .$$

For $\gamma \in \Gamma^{\text{comm}}$, the image of the graph of $\gamma \cdot : U \rightarrow U$ under (π, π) is an algebraic correspondence $T_\gamma \subseteq X \times X$ which we shall call the generalized Hecke correspondence associated to γ .

Remark 4.5. That the generalized Hecke correspondences are algebraic seems to be well-known, but for lack of a reference, we explain how to prove this. Let $\gamma \in G(\mathbb{C})$ and write T_γ for the image of $\{(x, \gamma \cdot x) : x \in U\}$ under $\pi \times \pi$. Let D be a set of coset representatives of $\Gamma \cap \Gamma^\gamma$ in Γ^γ . Then one sees that $(\pi \times \pi)^{-1}(T_\gamma)$ is the union over $\delta \in D$ and $\epsilon \in \Gamma$ of the graphs of $\epsilon\delta\gamma$. From this description, it is clear that if $\gamma \notin \Gamma^{\text{comm}}$, then T_γ is not algebraic since for any $x \in X(\mathbb{C})$ there are countably infinitely many y with $(x, y) \in T_\gamma$. On the other hand, for $\gamma \in \Gamma^{\text{comm}}$ we see that T_γ is a definable analytic set since it may be realized as $\bigcup \delta \in D(\pi \times \pi)(\{(x, \delta\gamma x) : x \in F\})$ so that algebraicity of T_γ follows from Theorem 2.11.

Let us note that the fibres of χ are closed under the generalized Hecke operators.

Proposition 4.6. *If $a, b \in X(\mathbb{U})$ and there is some $\gamma \in \Gamma^{\text{comm}}$ with $(a, b) \in T_\gamma(\mathbb{U})$, then $\chi(a) = \chi(b)$.*

Proof. By the construction of T_γ , for some choice of a branch of π^{-1} and for some $\delta \in \Gamma$, we have $\gamma \cdot \pi^{-1}(a) = \delta \cdot \pi^{-1}(b)$. Hence, $\pi^{-1}(a)$ and $\pi^{-1}(b)$ are in the same $G(\mathbb{C})$ orbit so that $\tilde{\chi}(\pi^{-1}(a)) = \tilde{\chi}(\pi^{-1}(b))$ implying that $\chi(a) = \chi(b)$. \square

If the commensurability group is large, then the fibres of χ are precisely the Kolchin closures of the generalized Hecke orbits.

Proposition 4.7. *Assume that Γ^{comm} is large in the sense that every set of coset representatives of Γ in Γ^{comm} is Zariski dense in $G(\mathbb{C})$. Then for each $a \in X(\mathbb{U})$, the generalized Hecke orbit of a is Kolchin dense in F_a .*

Proof. Let $a \in X(\mathbb{U})$ and let $V \subseteq X$ be the Kolchin closure of the generalized Hecke orbit of a . Let $N \gg 0$ be large enough so that $\chi = \chi_N \circ \nabla_N$ and there is an algebraic variety $V_N \subseteq \mathcal{A}_N(X)$ with $V(\mathbb{U}) = \nabla_N^{-1}(V_N(\mathbb{U}))$ and $\nabla_N(V(\mathbb{U}))$ is Zariski dense in V_N . Let \tilde{V}_N be a component of $J_N^{-1}V_N$. Let $H := \{\gamma \in G(\mathbb{C}) : \gamma \cdot \tilde{V}_N = \tilde{V}_N\}$. Since $J_N(\pi)^{-1}V_N$ is closed under the action of Γ^{comm} , we see that H contains a set of coset representatives for Γ in Γ^{comm} . By our hypothesis on Γ^{comm} , H is dense in $G(\mathbb{C})$, and, hence, is equal to $G(\mathbb{C})$. Thus, the \tilde{V}_N is the Zariski closure of $\tilde{\chi}_N^{-1}(\chi(a))$ so that V is the Kolchin closure of the constructible set defined by $\chi(x) = \chi(a)$, namely, F_a . \square

One might ask for a description of the set of points in F_a which are algebraic over a . We solve this problem only for the case that a is the generic point (in the sense of the Zariski topology) of X and that $\mathbb{C}(a) = \mathbb{C}(X)$ is given a differential structure for which $\mathbb{C}(X)^\Delta = \mathbb{C}$.

Proposition 4.8. *Let $K = \mathbb{C}(X)$, the field of rational functions on X , given with a basis of \mathbb{C} -derivations $\Delta = \{\partial_1, \dots, \partial_n\}$. Fix an embedding of K into \mathbb{U} over \mathbb{C} . Let $a \in X(K)$ be the generic point, that is, the K -rational point corresponding to the identity map $\text{id} : X \rightarrow X$. Then*

$$F_a(K^{\text{alg}}) = \{b \in X(\mathbb{U}) : (\exists \gamma \in \Gamma^{\text{comm}})(a, b) \in T_\gamma(\mathbb{U})\} .$$

Proof. We have already noted $F_a(\mathbb{U})$ contains all points which are in generalized Hecke correspondence with a and because each generalized Hecke correspondence is a finite-to-finite correspondence, all such points are algebraic over K . Hence, we need only verify the left to right inclusion.

Suppose that $b \in F_a(K^{\text{alg}})$. Geometrically, we may represent b by a dominant, generically finite, rational map $b : X' \dashrightarrow X$ where X' is an irreducible complex algebraic variety. Consider some nonempty connected open set $V' \subseteq X'(\mathbb{C})$ for which the covering $\pi : U \rightarrow X(\mathbb{C})$ trivializes over $V := b(V')$ and $V \cong \mathbb{B}^{\dim(X)}$ is a coordinate chart. Let $\tilde{V} \subseteq U$ be one of the components of $\pi^{-1}V$. Let us work with the branch of π^{-1} which on V takes values in \tilde{V} . By Proposition 4.1, there is some $g \in G(\mathbb{C})$ with $\pi^{-1}(b) = g \cdot \pi^{-1}(a)$. Hence, (a, b) (regarded now a point in $\text{Mor}(V', V \times V) \subseteq (X \times X)(\mathcal{M}(V')) \subseteq (X \times X)(\mathbb{U})$) lies on the analytic subset of $V \times V$, which we shall call $\Upsilon := (\pi, \pi)(\{(x, y) \in \tilde{V} \times \tilde{V} : g \cdot x = y\})$. Let $D \subseteq X \times X$ be the algebraic locus of (a, b) over \mathbb{C} , that is, D is the smallest \mathbb{C} -variety which contains the point (a, b) . The analytic variety Υ being the graph of a function is irreducible. Hence, either $\dim(\Upsilon \cap D) < \dim(X)$ or $\Upsilon \subseteq D$. The former condition is impossible by our hypothesis that $\mathbb{C}(a)^\Delta = \mathbb{C}$: writing $a = (a_1, \dots, a_{\dim(X)})$ and $b = (b_1, \dots, b_{\dim(X)})$, with respect to coordinates on V , then the hypothesis on the constants is equivalent to the assertion that the matrix $(\partial_i a_j)$ has rank $\dim(X)$. A further analytic relation would force the rank to be $\leq \dim(\Upsilon \cap D) < \dim(X)$. Thus, $\Upsilon \subseteq D$ implying that $\{(x, y) \in U \times U : g \cdot x = y\} \subseteq (\pi \times \pi)^{-1}D$ so that the image of the graph of the action of g is algebraic which is only possible for $g \in \Gamma^{\text{comm}}$. \square

5. SOME COVERING MAPS WITH ALGEBRAIC GENERALIZED LOGARITHMIC DERIVATIVES

In this section we specialize Theorem 3.12 to some concrete cases of complex algebraic varieties admitting suitable covering maps. In each case, we are required to check that the covering map is definable in some o-minimal expansion of the real field on some fundamental domain.

5.1. Logarithmic derivatives. Let us begin with the example with which we introduced this paper. As we have already noted, the usual logarithmic derivative $\frac{d}{dz} \log : \mathbb{G}_m \rightarrow \mathbb{G}_a$ defined by $f \mapsto \frac{f'}{f}$ is clearly differential algebraic and may be obtained from Theorem 3.12 using the fact that the restriction of $\exp : \mathbb{C} \rightarrow \mathbb{C}^\times$ to the set

$$F := \{z \in \mathbb{C} : -2\pi < \text{Im}(z) < 2\pi\}$$

is surjective and definable in $\mathbb{R}_{\text{an}, \text{exp}}$ via the formula

$$\exp(z) = e^{\text{Re}(z)} \cos(\text{Im}(z)) + ie^{\text{Re}(z)} \sin(\text{Im}(z)) .$$

Of course, in $\mathbb{R}_{\text{an}, \text{exp}}$ the real exponential function is explicitly allowed as a definable function while the functions $\cos(x)$ and $\sin(x)$ restricted to $[-2\pi, 2\pi]$ are expressible as restricted analytic terms.

Let us recall the theory of logarithmic derivatives for algebraic groups over the constants; for further details see [12]. We shall work with differential fields (K, Δ) with $\Delta = \{\partial_1, \dots, \partial_n\}$. If G is an algebraic group defined over the constants $C = K^\Delta$, then the tangent bundle of G splits as a semi-direct product $TG = G \rtimes T_e G$ where $T_e G$ is the tangent space to G at the identity element $e \in G$. The second part of the splitting $\eta : TG \rightarrow T_e(G)$ is given by $(g, v) \mapsto d(g^{-1} \cdot)v$ where we write the points on TG as pairs (g, v) consisting of a point g of G and a vector v in the tangent space to G at g . Because G is defined over C , we may identify $\tau_1 G$ with the n^{th} fibre power of TG over G so that $\nabla : G(K) \rightarrow \tau_1 G(K)$ gives a map of

groups $\nabla : G(K) \rightarrow (TG \times_G \cdots \times_G TG)(K)$. The logarithmic derivative $\partial \log_G : G(M) \rightarrow (T_e G)^n(M)$ is given by $x \mapsto \eta^{\times n}(\nabla(x))$. The differential algebraicity of this logarithmic derivative is clear from this construction, but as with the usual logarithmic derivative this map may be interpreted analytically.

If G is a connected complex algebraic group, then Lie theory supplies a complex analytic exponential map $\exp_G : T_e G(\mathbb{C}) \rightarrow G(\mathbb{C})$. In fortuitous cases, for example when G is commutative, the exponential map is a covering map, but this is not always so. With the next lemma, we note that in the case that G is commutative, there is a definable (in $\mathbb{R}_{\text{an,exp}}$) subset $F \subseteq T_e G(\mathbb{C})$ for which $\exp_G \upharpoonright F$ is definable and surjective.

Lemma 5.1. *If G is a connected commutative complex algebraic group of dimension g , then there is a semialgebraic subset $F \subseteq \mathbb{C}^g \cong T_e G(\mathbb{C})$ for which $\exp_G \upharpoonright F : F \rightarrow G(\mathbb{C})$ is surjective and definable in $\mathbb{R}_{\text{an,exp}}$.*

Proof. By the structure theory of algebraic groups, G fits into an exact sequence

$$0 \longrightarrow L \longrightarrow G \longrightarrow A \longrightarrow 0$$

where L is a commutative, connected linear algebraic group and A is an abelian variety. As \mathbb{C} has characteristic zero, L is isomorphic to a product of additive and multiplicative groups (see [33]). The tangent space at the origin of the additive group may be identified with the additive group itself and relative to this identification, the exponential map may be taken to be the identity map. As noted above, the usual complex exponential function admits a semialgebraic fundamental domain on which the exponential function restricts to an $\mathbb{R}_{\text{an,exp}}$ -definable function. Hence, we may find a semialgebraic set $F_L \subseteq T_e L(\mathbb{C}) \cong \mathbb{C}^{\dim(L)}$ for which \exp_L restricted to F_L is surjective onto $L(\mathbb{C})$ and definable in $\mathbb{R}_{\text{an,exp}}$.

Choose a subspace $V \leq T_e G(\mathbb{C})$ for which $V \cap T_e L(\mathbb{C}) = 0$ and $V + T_e L(\mathbb{C}) = T_e G(\mathbb{C})$. Since the fundamental domain for \exp_A is bounded, if $F_A \subseteq V$ is any closed box for which the image of F_A under the natural map $T_e G(\mathbb{C}) \rightarrow T_e A(\mathbb{C})$ contains a fundamental domain, then the restriction of \exp_G to $\{0_{T_e L(\mathbb{C})}\} \times F_A$ is explicitly \mathbb{R}_{an} -definable and its image under $G \rightarrow A$ maps onto A . Let us set $F := F_L \times F_A$. Then on F , \exp_G may be expressed as the sum (in G) of $\exp_L \upharpoonright F_L$ and $\exp_G \upharpoonright \{0_{T_e L(\mathbb{C})}\} \times F_A$ and is therefore definable. A simple diagram chase shows that the restriction of \exp_G to F is surjective. \square

It follows from Lemma 5.1 that for a connected commutative complex algebraic group G Kolchin's logarithmic derivative for G may be realized as a generalized logarithmic derivative given by Theorem 3.12.

5.2. Moduli spaces of abelian varieties and general arithmetic varieties.

In this section we consider the application of Theorem 3.12 to various classical functions and then also discuss the relation between this theorem and classical Picard-Fuchs theory.

5.2.1. *Moduli of elliptic curves.* Let us recall some of the basic theory of modular functions and moduli spaces of elliptic curves. The reader may wish to consult [16] for more details. We write the upper half plane as $\mathfrak{h} := \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ and regard \mathfrak{h} as an open subset of $\mathbb{P}^1(\mathbb{C})$ on which the algebraic group GL_2 acts by linear fractional transformations. Via this action, \mathfrak{h} is preserved by $\text{GL}_2^+(\mathbb{R})$,

the subgroup of $\mathrm{GL}_2(\mathbb{R})$ on which the determinant is positive. In particular, \mathfrak{h} is preserved by $\mathrm{SL}_2(\mathbb{Z})$.

For each $N \in \mathbb{Z}_+$, let $\Gamma(N)$ be the kernel of the reduction modulo N map $\mathrm{SL}_2(\mathbb{Z}) \rightarrow \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$. The quotient $\Gamma(N)\backslash\mathfrak{h}$ has the structure of an affine algebraic variety which is usually called $Y(N)(\mathbb{C})$. When $N > 2$, the quotient map $j_N : \mathfrak{h} \rightarrow Y(N)(\mathbb{C})$ is a covering map, but for $N = 1, j_1$, which is usually denoted by j , the usual analytic j -function, fails to be a covering map because it is ramified over two points. The curve $Y(1)$ may be identified with \mathbb{A}^1 and with the usual normalizations, the two points over which j is ramified are 0 and 1728 (with preimages $\exp(\frac{\pi i}{3})$ and i , respectively).

If one takes $F := \{z \in \mathfrak{h} : |\mathrm{Re}(z)| \leq \frac{1}{2} \text{ and } \mathrm{Im}(z) \geq \frac{\sqrt{3}}{2}\}$, then F contains a fundamental domain for j and the restriction of j to F is definable in $\mathbb{R}_{\mathrm{an}, \mathrm{exp}}$. Indeed, the function $\tau \mapsto q_\tau := \exp(2\pi i\tau)$ maps F to the disc of radius $\exp(-\pi\sqrt{3})$. It is known that the q -expansion of the j -function is meromorphic on the unit disc with a simple pole at the origin. Hence, j may be expressed as the composition of the restriction of a meromorphic function to the closed disc of radius $\exp(-\pi\sqrt{3})$ with $\exp(2\pi i\tau)$, which as we have noted above, is definable in $\mathbb{R}_{\mathrm{an}, \mathrm{exp}}$ on vertical strips of bounded width. As each of the modular functions j_N factor through j , they, too, are definable when restricted to F .

Consequently, for $N > 1$, Theorem 3.12 applies directly showing that the generalized logarithmic derivative $\chi : Y_0(N) \rightarrow Z$ defined as the usual Schwarzian derivative applied to j_N^{-1} is differential constructible. For the j -function itself, we may use the flexibility provided by the hypotheses of Theorem 3.12: take U to be the open subset of \mathfrak{h} on which j is a covering map and X to be $\mathbb{A}^1 \setminus \{0, 1728\}$.

5.2.2. Universal abelian schemes. The definability of j on a fundamental domain is part of a theorem of Peterzil and Starchenko to the effect that the two-variable Weierstraß \wp -function is definable (in $\mathbb{R}_{\mathrm{an}, \mathrm{exp}}$) on a fundamental domain [21]. That is, the covering maps associated to universal families of elliptic curves may be treated by Theorem 3.12. In [26], Peterzil and Starchenko extend this result showing that the covering maps associated to the universal families of abelian varieties are definable in $\mathbb{R}_{\mathrm{an}, \mathrm{exp}}$.

Let $g \in \mathbb{Z}_+$ be a positive integer and write \mathfrak{h}_g for the Siegel space of symmetric $g \times g$ complex matrices whose imaginary parts are positive definite. For a fixed sequence $D = (d_1, \dots, d_g)$ of positive integers with $d_1|d_2|\dots|d_g$, we obtain the notion of a polarization of type D . For $\tau \in \mathfrak{h}_g$, we associate the complex torus $X_\tau^D(\mathbb{C}) := \mathbb{C}^g / (\tau\mathbb{Z}^g + \mathrm{diag}(d_1, \dots, d_g)\mathbb{Z}^g)$ which, because of the choice of polarization, is actually an abelian variety. There is a discrete subgroup G_D of the symplectic group $\mathrm{Sp}_{2g}(\mathbb{Q})$ for which the quotient $G_D\backslash\mathfrak{h}_g$ has the structure of an algebraic variety $\mathcal{A}_g^D(\mathbb{C})$ (in this case, “ \mathcal{A} ” does not refer to an arc space). As in the case of $g = 1$, the quotient map is ramified, but in passing to congruence subgroups one obtains covering maps. Theorem 1.1 of [26] asserts that is a fundamental domain on which the covering map $\mathfrak{h}_g \rightarrow \mathcal{A}_{g,D}(\mathbb{C})$ is definable.

The covering map for the abelian variety associated to $\tau \in \mathfrak{h}_g$ is itself uniformly defined. That is, if we were to take $U := \mathfrak{h}_g \times \mathbb{C}^g$, then to a pair (τ, z) we may associate the abelian variety X_τ^D and the point $z \bmod (\tau\mathbb{Z}^g + \mathrm{diag}(d_1, \dots, d_g)\mathbb{Z}^g) \in X_\tau^D(\mathbb{C})$. The quotient $(G_D \times \mathbb{Z}^{2g})\backslash(\mathfrak{h}_g \times \mathbb{C}^g)$ has the structure of an algebraic variety \mathcal{X}_g^D and fits into the following commutative diagram

$$\begin{array}{ccc}
\mathfrak{h}_g \times \mathbb{C}^g & \longrightarrow & \mathfrak{h}_g \\
\downarrow & & \downarrow \\
\mathcal{X}_g^D(\mathbb{C}) & \longrightarrow & \mathcal{A}_g^D(\mathbb{C})
\end{array}$$

Theorem 1.3 of [26] asserts that the covering map of \mathcal{X}_g^D is definable on a fundamental domain. Passing to a finite cover of \mathcal{A}_g^D corresponding to a suitable congruence subgroup (or, equivalently, to the choice of some level structure when interpreted in terms of the moduli problem), the resulting map is a covering map to which Theorem 3.12 applies directly.

5.2.3. Arithmetic varieties. Klingler, Ullmo and Yafaev have shown [11] that for *any* arithmetic variety, the associated covering map is definable in $\mathbb{R}_{\text{an,exp}}$ on some fundamental domain. Here, an arithmetic variety is a complex algebraic variety expressible as $\Gamma \backslash D$ where D is a symmetric Hermitian domain and Γ is an arithmetic group. This definability theorem supersedes the Peterzil-Starchenko definability result for the covering maps for the moduli spaces of abelian varieties and includes the covering maps for all Shimura varieties. However, the requirement that D be a symmetric Hermitian domain precludes its application to the case of the universal abelian variety or more generally to mixed Shimura varieties. Gao has shown [9] that the covering maps associated to all mixed Shimura varieties are $\mathbb{R}_{\text{an,exp}}$ -definable when restricted to some fundamental domain. Thus, from Theorem 3.12 and the Klingler-Ullmo-Yafaev and Gao definability theorems, we deduce that the generalized logarithmic derivatives associated to arithmetic varieties and to mixed Shimura varieties are differential constructible.

5.3. Picard-Fuchs equations, periods and Manin homomorphisms. Families of complex algebraic varieties often come equipped with period mappings. As we cannot say as much as we would like in full generality, we shall confine ourselves to a naïve discussion referring the reader to [6] for more details. Our classical approach to period mappings is explained clearly in the first section of [18].

If X is a complex algebraic variety and $\omega_1, \dots, \omega_g$ is a basis of global one-forms on X and $\gamma_1, \dots, \gamma_\ell$ is a basis of the free part of the integral homology group $H_1(X, \mathbb{Z})$, then the matrix $(\int_{\gamma_i} \omega_j) \in M_{\ell \times g}(\mathbb{C})$ is called a period matrix of X . Of course, changing bases would result in a different matrix so that the period matrix of X is well defined only up to the action of $\text{GL}_g(\mathbb{C})$ on one side and $\text{GL}_\ell(\mathbb{Z})$ on the other. If X varies in an algebraic family, $X \rightarrow S$ for which this map is smooth, and the forms $\omega_1, \dots, \omega_g$ are also taken to vary algebraically, then locally in the Euclidean topology this family is trivial so that the cycles γ_i on some reference fibre X_0 may be transferred to nearby fibers. It is a theorem of Griffiths [10] that the components of the period matrix are analytic functions on S . The theory of Picard-Fuchs equations shows that period matrices satisfies linear differential equations with coefficients from the function field of S .

We explain now how to deduce the existence of the Picard-Fuchs equations with coefficients in $\mathbb{C}(S)$ from the global $\mathbb{R}_{\text{an,exp}}$ -definability of some branch of functions extending $s \mapsto \int_{\gamma_i} (\omega_j)_s$. In fact, we shall show a little more: not only do the period matrices satisfy linear differential equations over $\mathbb{C}(S)$, but one may differentially constructibly find linear differential operators so that for every differential field M

of meromorphic functions extending $\mathbb{C}(S)$ and every point $a \in S(M)$ the kernel of the associated linear differential operator is exactly the \mathbb{C} -vector space generated by the periods of X_s .

Let us note that it follows from the Peterzil-Starchenko theorem on the definability of the covering maps for universal families of abelian varieties that for every algebraic family of polarized abelian varieties [26, Theorem 1.3], global branches of the period functions are definable. However, we do not know whether such an hypothesis on definability holds for all families of varieties.

With the following definition we make precise what we mean by a local system and what it means for such a local system to definably trivialize.

Definition 5.2. Let S be a complex algebraic variety and N and g be two natural numbers. A \mathbb{Z}^g -local system over S is a subset $\Gamma \subseteq \mathbb{G}_{a,S}^N(\mathbb{C}) = S(\mathbb{C}) \times \mathbb{C}^N$ such that

- for each $s \in S(\mathbb{C})$, $\Gamma_s \subseteq \mathbb{C}^N$ is a subgroup isomorphic to \mathbb{Z}^g and
- $S(\mathbb{C})$ is covered by open sets $U \subseteq S(\mathbb{C})$ on which there are analytic functions $\gamma_{1,U}, \dots, \gamma_{g,U} : U \rightarrow \mathbb{C}^N$ so that for all $s \in S(\mathbb{C})$ one has $\Gamma_s = \sum_{i=1}^g \mathbb{Z}\gamma_{i,U}(s)$.

Fix an o-minimal structure $\mathbb{R}_{\mathcal{F}}$ expanding the real field. In practice, one may take $\mathbb{R}_{\text{exp,an}}$. We say that Γ *definably trivializes* if there are definable (though not necessarily continuous) functions $\nu_1, \dots, \nu_g : S(\mathbb{C}) \rightarrow \mathbb{C}^N$ so that at each point $s \in S(\mathbb{C})$ one has $\Gamma_s = \sum_{i=1}^g \mathbb{Z}\nu_i(s)$.

We begin with an algebraic version of our theorem on Picard-Fuchs equations similar to Proposition 3.11.

Proposition 5.3. *Let S be a complex algebraic variety and $\Gamma \subseteq \mathbb{G}_{a,S}^N(\mathbb{C}) = S(\mathbb{C}) \times \mathbb{C}^N$ a \mathbb{Z}^g -local system over S which definably trivializes via definable functions $\nu_1, \dots, \nu_g : S(\mathbb{C}) \rightarrow \mathbb{C}^N$. Then there is a constructible function $\Phi : S \rightarrow \text{Mat}_{N \times N}$ so that for each $s \in S(\mathbb{C})$ the \mathbb{C} -vector subspace of \mathbb{C}^N generated by Γ_s is the kernel of $\Phi(s)$.*

Proof. For each $j \leq N$, we denote by $G(j, N)$ the Grassmannian of j -dimensional linear subspaces of \mathbb{G}_a^N . We shall prove this proposition by first showing that the association $\sigma : S \rightarrow \bigcup_{j=1}^g G(j, N)$ given by sending $s \in S(\mathbb{C})$ to the \mathbb{C} -vector space generated by Γ_s is constructible. Then, we shall observe that there is a constructible function $\Psi : G(j, N) \rightarrow \text{Mat}_{N \times N}$ so that for each $[V] \in G(j, N)(\mathbb{C})$ the kernel of $\Phi([V])$ is precisely V . Our map Φ is then $\Psi \circ \sigma$.

For $1 \leq j \leq g$, let $S_j(\mathbb{C}) := \{s \in S(\mathbb{C}) : \dim_{\mathbb{C}} \mathbb{C} \cdot \Gamma_s \leq j\}$. Working in an open set $U \subseteq S(\mathbb{C})$ on which Γ trivializes, we see that $S_j(\mathbb{C}) \cap U$ is an analytic set as it is defined by the vanishing of a collection of determinants of certain minors of the matrix $(\gamma_{1,U}(s), \dots, \gamma_{g,U}(s))$. Hence, S_j is analytic. On the other hand, using the natural definition of dimension and the functions ν_1, \dots, ν_g , one sees that S_j is definable as well. Hence, by Theorem 2.11, S_j is an algebraic subvariety of S .

Likewise, for each j , the function $\sigma_j : (S_j \setminus S_{j-1}) \rightarrow G(j, N)$ given by sending $s \in (S_j \setminus S_{j-1})(\mathbb{C})$ to the code for the vector space generated by Γ_s is both analytic and definable, and, hence, a regular map. The function $\sigma : S(\mathbb{C}) \rightarrow \bigcup_{j=1}^g G(j, N)(\mathbb{C})$ is simply the union of the various σ_j .

Let us prove by noetherian induction that if $Z \subseteq G(j, N)$ is an irreducible subvariety, then there is a constructible function $\Psi_Z : Z \rightarrow \text{Mat}_{N \times N}$ so that if $[V] \in Z$ is the code for the vector space $V \leq \mathbb{G}_a^N$, then $V = \ker \Psi_Z([V])$. Taking

$Z = G(j, N)$, we obtain our desired Ψ . Let $[V] \in Z(\mathbb{C}(Z))$ be the generic point. By basic linear algebra, there is some linear map $L : \mathbb{C}(Z)^N \rightarrow \mathbb{C}(Z)^N$ for which $\ker L = V(\mathbb{C}(Z))$. Localizing, we find $Z' \subsetneq Z$ a proper subvariety of Z for which the coefficients of L are regular functions on $Z \setminus Z'$. By induction, there is a constructible function $\Psi_{Z'} : Z' \rightarrow \text{Mat}_{N \times N}$ with the desired properties. Let Ψ_Z be given by $\Psi_{Z'}$ on Z' and by associating $z \in Z \setminus Z'$ the specialization of L to z on $Z \setminus Z'$. \square

As with our proof of Theorem 3.12, we find our tighter Picard-Fuchs operators from the ostensibly weaker algebraic Proposition 5.3.

Theorem 5.4. *Let S be a complex algebraic variety. Give $\mathbb{C}(S)$ the structure of a differential field by fixing a basis of \mathbb{C} -derivations. Let N and g be two natural numbers. Suppose that $\Gamma \subseteq \mathbb{G}_{a,S}^N(\mathbb{C}) = S(\mathbb{C}) \times \mathbb{C}^N$ is a \mathbb{Z}^g -local system over S with local trivializations as in Proposition 5.3. Also as in Proposition 5.3, we suppose moreover that Γ definably trivializes via functions $\nu_1, \dots, \nu_g : S(\mathbb{C}) \rightarrow \mathbb{C}^N$. Then there is a differential constructible function Θ on S taking values in a space of linear differential operators so that for any differential field M of meromorphic functions extending $\mathbb{C}(S)$ and point $s \in S(M)$ the \mathbb{C} -vector subspace of \mathbb{C}^N generated by Γ_s is the kernel of $\Theta(s)$.*

Proof. Working by noetherian induction on the Kolchin topology on S , one sees that there is some number $\ell \gg 0$ so that for any differential field M of meromorphic function extending $\mathbb{C}(S)$ and point $s \in S(M)$ the \mathbb{C} -vector space generated by Γ_s is the pre-image under ∇_ℓ of the M -vector space generated by $\nabla_\ell(\Gamma_s)$ in $\mathcal{A}_\ell(\mathbb{G}_a^N)$.

Let us define $\Gamma^{(\ell)}$. For $U \subseteq S(\mathbb{C})$ an open set over which Γ trivializes and $t \in J_\ell(U) \subseteq \mathcal{A}_\ell(S)(\mathbb{C})$, let $\Gamma_t^{(\ell)} := \sum_{i=1}^g \mathbb{Z} J_\ell(\gamma_{i,U})(t)$. From the local patching for Γ , it is easy to see that $\Gamma^{(\ell)}$ is a \mathbb{Z}^g -local system over $\mathcal{A}_\ell(S)$.

Replacing S by $\mathcal{A}_\ell(S)$, Γ by $\Gamma^{(\ell)}$, and N by $N' := \dim \mathcal{A}_\ell(\mathbb{G}_a^N)$, Proposition 5.3 applies giving a constructible function $\Phi : \mathcal{A}_\ell(\mathbb{G}_a^N) \rightarrow \mathbb{G}_a^{N'}$ so that for any $t \in \mathcal{A}_\ell(S)(M)$, the kernel of $\Phi(t)$ is the M -vector space generated by $\Gamma_t^{(\ell)}$. Define Θ on S by $\Theta(s) := \Phi(\nabla(s)) \circ \nabla$. Then for $s \in S(M)$, $\ker \Theta(s)$ is the \mathbb{C} -vector space generated by Γ_s . \square

Remark 5.5. If one drops either the hypothesis that Γ is a \mathbb{Z}^g -local system (in the sense that it locally analytically trivializes) or the hypothesis that Γ definably discontinuously trivializes, then the conclusion of Theorem 5.4 fails. For example, if $E \rightarrow S$ is an elliptic scheme over an algebraic curve S , then there is a definable function $\rho : S(\mathbb{C}) \rightarrow \mathbb{C}$ so that for each $s \in S(\mathbb{C})$, $\rho(s)$ is a nonzero element of the kernel of the exponential map $\exp_{E_s} : \mathbb{C} \rightarrow E_s(\mathbb{C})$. It is well-known that if $E \rightarrow S$ is not isotrivial, then ρ cannot satisfy an order one algebraic differential equation (even where it is analytic). Theorem 5.4 does not apply as $\Gamma := \mathbb{Z} \cdot \rho \subseteq S(\mathbb{C}) \times \mathbb{C}$ is not a local system since it does not locally analytically trivialize near the discontinuities of ρ . On the other hand, the Bessel function does give rise to a \mathbb{Z} -local system, but it too satisfies an order two algebraic differential equation, but not one of order one. The issue is that the local system does not definably trivialize.

I thank D. Bertrand for suggesting these examples. In the same communication he suggests that in line with the classical Picard-Fuchs theory the hypotheses of Theorem 5.4 should correspond to the assertions that the associated differential equations are invariant under monodromy and have at worst regular singularities.

Let us apply Theorem 5.4 to the case of periods of abelian varieties.

Corollary 5.6. *Let S be an irreducible complex algebraic variety and $X \rightarrow S$ a polarized abelian scheme over S of relative dimension g . We regard $\mathbb{C}(S)$ as a differential field by a fixing a basis $\partial_1, \dots, \partial_n$ of commuting derivations on $\mathbb{C}(S)$. Let $\Gamma := \{(s, z) \in S(\mathbb{C}) \times \mathbb{C}^g : z \in \ker \exp_{X_s}\}$ be the \mathbb{Z}^{2g} -local system over S given by the lattices of the abelian varieties X_s regarded as complex tori. Then there is a differential constructible map Θ on S taking values in a space of linear differential operators so that for any field M of meromorphic functions extending $\mathbb{C}(S)$ and any point $s \in S(M)$, the kernel of $\Theta(s)$ is the \mathbb{C} -vector space generated by Γ_s .*

Proof. As a consequence of [26, Theorem 1.2], when $X \rightarrow S$ is a universal family of polarized abelian varieties, the function which associates to $s \in S(\mathbb{C})$ the coordinates of generators for a fundamental parallelootope of the covering $\exp_{X_s} : \mathbb{C}^g \rightarrow X_s(\mathbb{C})$ is definable in $\mathbb{R}_{an, \exp}$. That is, there is a definable (discontinuous) splitting of Γ . The Peterzil-Starchenko theorem does not apply directly to a general $X \rightarrow S$, but after passing to a finite base extension, we may assume that $X \rightarrow S$ is a family of polarized abelian varieties with sufficient level structure. Then this family comes from the universal such family by pullback, so that the definability of a global (discontinuous) splitting of Γ for $X \rightarrow S$ follows from the definability for the universal family. This corollary is now a special case of Theorem 5.4. \square

Corollary 5.6 justifies Manin's analytic construction of differential additive characters on abelian varieties [13].

Corollary 5.7. *Let S be an irreducible complex algebraic variety and $X \rightarrow S$ a polarized abelian scheme over S of relative dimension g . Then there is a map of differential algebraic groups over S , $\mu : X \rightarrow (\mathbb{G}_a^N)_S$ (for some N) so that for any differentially closed field $(\mathbb{U}, \partial_1, \dots, \partial_n) \supseteq (\mathbb{C}(S), \partial_1, \dots, \partial_n)$ and any point $s \in S(\mathbb{U})$, the differential function field of the kernel of $\mu_s : X_s(\mathbb{U}) \rightarrow \mathbb{G}_a^N(\mathbb{U})$ has transcendence degree at most $2g$.*

Proof. The exponential map $\exp_X : S(\mathbb{C}) \times \mathbb{C}^g \rightarrow X(\mathbb{C})$ is an analytic covering map over S . As such, it has a local analytic inverse $\log_X : X(\mathbb{C}) \rightarrow S(\mathbb{C}) \times \mathbb{C}^g$. The many valued function \log_X is well defined up to addition by $\Gamma := \ker \exp_X$, which is a \mathbb{Z}^{2g} -local system over S . Let Θ be given by Corollary 5.6. Fibrewise, Θ is a linear differential map. Thus, $\mu := \Theta \circ \log_X$ is a well-defined, fiberwise additive differentially analytically constructible mapping. As such, μ may be represented as $\mu = \tilde{\mu} \circ \nabla$ where $\tilde{\mu}$ is a definable analytically constructible function on some prolongation space of X . By the Peterzil-Starchenko GAGA Theorem 2.11, $\tilde{\mu}$ is algebraically constructible, so that μ is differentially constructible. From its construction because on each fibre the kernel of Θ is a vector spaces over the constants of dimension at most $2g$, it follows that the fibres of μ have dimension at most $2g$. \square

Remark 5.8. Rather than working with local systems, we could have proven Corollary 5.7 from our main Theorem 3.12 by viewing the Manin maps as components of the generalized logarithmic derivative associated to the analytic covering of the universal families of polarized abelian varieties.

6. CONCLUDING REMARKS AND QUESTIONS

Let us conclude with some natural questions raised by our construction of generalized logarithmic derivations.

6.1. Differential equations for covering maps. Let $\pi : U \rightarrow X(\mathbb{C})$ be a covering map as in the statement of Theorem 3.12. Let $M = \mathcal{M}(U)$ be the field of meromorphic functions on U treated as differential field with respect to some choice of a basis of derivations $\partial_1, \dots, \partial_n$. Let $t \in U(M)$ be the M -rational point corresponding to the identity map $U \rightarrow U$. Then $\pi(t) \in X(M)$ “is” π regarded as an M -rational point of X . From Theorem 3.12 we know that $\pi(t)$ satisfies the algebraic differential equation $\chi(\pi(t)) = \tilde{\chi}(t)$. Noting that t is the restriction of the generic point of Y to U and that the derivations on M may be chosen to be the restriction of a basis of derivations of $\mathbb{C}(Y)$, we see that the differential field generated by $\pi(t)$ over $\mathbb{C}(Y)$ has transcendence degree at most $\dim G$ over $\mathbb{C}(Y)$. Since $\mathbb{C}(Y)$ is a differential field of finite transcendence degree over \mathbb{C} , it follows that the differential field generated by $\pi(t)$ over \mathbb{C} has finite transcendence degree.

However, in some cases, for example for the analytic j -function $j : \mathfrak{h} \rightarrow \mathbb{A}^1(\mathbb{C})$ and for exponential functions $\exp_G : \mathbb{C}^g \rightarrow G(G)$ of connected, commutative algebraic groups, with the usual differential structure, the differential field generated by $\pi(t)$ over \mathbb{C} has transcendence degree exactly $\dim(G)$ over \mathbb{C} by Proposition 4.2 and the observation that $\tilde{\chi}(t)$ is a constant point.

Question 6.1. *Is it always the case that there is some choice of a basis of $\text{Der}(\mathbb{C}(Y)) = \text{Hom}_{\mathbb{C}(Y)}(\Omega_{\mathbb{C}(Y)/\mathbb{C}}, \mathbb{C}(Y))$ so that $\tilde{\chi}(t)$ is a constant point? If not, under what conditions is this the case?*

6.2. Classification theory and the fibres of χ . In [8], Freitag and the present author show that the differential equation satisfied by the j -function is strongly minimal and has trivial forking geometry. It is not unreasonable to expect that other generalized logarithmic derivatives will produce examples of types with trivial forking geometry but complicated binary structure coming from generalized Hecke correspondences. We do not propose a precise statement of a conjecture on the classification theoretic structure of fibres of generalized logarithmic derivatives as that would involve delineating the ways in which sets nonorthogonal to the constants and to Manin kernels may arise.

Problem 6.2. *Describe the classification theoretic structure (i.e., an analysis in minimal types placing these types into their positions relative to the Zilber trichotomy) of fibres of generalized logarithmic derivatives in terms of group theoretic data of the covering spaces.*

6.3. Ax-Schanuel problems. In [1], Ax proved a function field version of Schanuel’s conjecture on algebraic relations amongst exponentials as a purely differential algebraic theorem. In [27], Pila and Tsimerman prove a variant of Ax’s theorem for the j -function. They observe that via the Seidenberg embedding theorem, their Schanuel-like theorem for the j -function admits a differential algebraic formulation, though their proof involves other techniques. Our differential equations permit the differential algebraic formulation of general transcendence conjectures for covering maps. As we have not resolved the question of which “obvious” relations might hold, we do not suggest a precise general conjecture here.

REFERENCES

- [1] J. AX, *On Schanuel's conjectures*, Ann. of Math. (2), 93 (1971), pp. 252–268.
- [2] D. BERTRAND AND W. ZUDILIN, *On the transcendence degree of the differential field generated by Siegel modular forms*, J. Reine Angew. Math., 554 (2003), pp. 47–68.
- [3] E. BISHOP, *Conditions for the analyticity of certain sets*, Michigan Math. J., 11 (1964), pp. 289–304.
- [4] N. BOURBAKI, *Éléments de mathématique. Fasc. XXXVI. Variétés différentielles et analytiques. Fascicule de résultats (Paragraphes 8 à 15)*, Actualités Scientifiques et Industrielles, No. 1347, Hermann, Paris, 1971.
- [5] A. BUIUM, *Geometry of differential polynomial functions. III. Moduli spaces*, Amer. J. Math., 117 (1995), pp. 1–73.
- [6] J. CARLSON, S. MÜLLER-STACH, AND C. PETERS, *Period mappings and period domains*, vol. 85 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2003.
- [7] J. DENEFF AND F. LOESER, *Geometry on arc spaces of algebraic varieties*, in European Congress of Mathematics, Vol. I (Barcelona, 2000), vol. 201 of Progr. Math., Birkhäuser, Basel, 2001, pp. 327–348.
- [8] J. FREITAG AND T. SCANLON, *Strong minimality and the j -function*, J. Eur. Math. Soc., ((to appear)).
- [9] Z. GAO, *Towards the André-Oort conjecture for mixed Shimura varieties: The Ax-Lindemann theorem and lower bounds for Galois orbits of special points*. [arXiv:1310.1302v4](https://arxiv.org/abs/1310.1302v4), 2014.
- [10] P. A. GRIFFITHS, *Periods of integrals on algebraic manifolds. I. Construction and properties of the modular varieties*, Amer. J. Math., 90 (1968), pp. 568–626.
- [11] B. KLINGLER, E. ULLMO, AND A. YAFAEV, *The hyperbolic Ax-Lindemann-Weierstrass conjecture*. 2016.
- [12] E. R. KOLCHIN, *Differential algebra and algebraic groups*, Academic Press, New York-London, 1973. Pure and Applied Mathematics, Vol. 54.
- [13] J. I. MANIN, *Proof of an analogue of Mordell's conjecture for algebraic curves over function fields*, Dokl. Akad. Nauk SSSR, 152 (1963), pp. 1061–1063.
- [14] D. MARKER, M. MESSMER, AND A. PILLAY, *Model theory of fields*, vol. 5 of Lecture Notes in Logic, Association for Symbolic Logic, La Jolla, CA; A K Peters, Ltd., Wellesley, MA, second ed., 2006.
- [15] T. MCGRAIL, *The model theory of differential fields with finitely many commuting derivations*, J. Symbolic Logic, 65 (2000), pp. 885–913.
- [16] J. S. MILNE, *Modular functions and modular forms (v1.30)*, 2012. Available at www.jmilne.org/math/.
- [17] R. MOOSA AND T. SCANLON, *Jet and prolongation spaces*, J. Inst. Math. Jussieu, 9 (2010), pp. 391–430.
- [18] D. R. MORRISON, *Picard-Fuchs equations and mirror maps for hypersurfaces*, in Essays on mirror manifolds, Int. Press, Hong Kong, 1992, pp. 241–264.
- [19] Y. PETERZIL AND S. STARCHENKO, *Expansions of algebraically closed fields in o -minimal structures*, Selecta Math. (N.S.), 7 (2001), pp. 409–445.
- [20] ———, *Expansions of algebraically closed fields. II. Functions of several variables*, J. Math. Log., 3 (2003), pp. 1–35.
- [21] ———, *Uniform definability of the Weierstrass \wp functions and generalized tori of dimension one*, Selecta Math. (N.S.), 10 (2004), pp. 525–550.
- [22] ———, *Complex analytic geometry in a nonstandard setting*, in Model theory with applications to algebra and analysis. Vol. 1, vol. 349 of London Math. Soc. Lecture Note Ser., Cambridge Univ. Press, Cambridge, 2008, pp. 117–165.
- [23] ———, *Complex analytic geometry and analytic-geometric categories*, J. Reine Angew. Math., 626 (2009), pp. 39–74.
- [24] ———, *Mild manifolds and a non-standard Riemann existence theorem*, Selecta Math. (N.S.), 14 (2009), pp. 275–298.
- [25] ———, *Tame complex analysis and o -minimality*, in Proceedings of the International Congress of Mathematicians. Volume II, Hindustan Book Agency, New Delhi, 2010, pp. 58–81.

- [26] ———, *Definability of restricted theta functions and families of abelian varieties*, Duke Math. J., 162 (2013), pp. 731–765.
- [27] J. PILA AND J. TSIMERMAN, *Ax-Schanuel for the j -function*. 2016.
- [28] B. POIZAT, *Une théorie de Galois imaginaire*, J. Symbolic Logic, 48 (1983), pp. 1151–1170 (1984).
- [29] R. REMMERT AND K. STEIN, *Über die wesentlichen Singularitäten analytischer Mengen*, Math. Ann., 126 (1953), pp. 263–306.
- [30] T. SASAKI AND M. YOSHIDA, *The uniformizing differential equation of the complex hyperbolic structure on the moduli space of marked cubic surfaces*, Proc. Japan Acad. Ser. A Math. Sci., 75 (1999), pp. 129–133.
- [31] A. SEIDENBERG, *Abstract differential algebra and the analytic case*, Proc. Amer. Math. Soc., 9 (1958), pp. 159–164.
- [32] ———, *Abstract differential algebra and the analytic case. II*, Proc. Amer. Math. Soc., 23 (1969), pp. 689–691.
- [33] J.-P. SERRE, *Groupes algébriques et corps de classes*, Publications de l’Institut Mathématique de l’Université de Nancago [Publications of the Mathematical Institute of the University of Nancago], 7, Hermann, Paris, second ed., 1984. Actuelles Scientifiques et Industrielles [Current Scientific and Industrial Topics], 1264.
- [34] B. SHIFFMAN, *On the removal of singularities of analytic sets*, Michigan Math. J., 15 (1968), pp. 111–120.
- [35] L. VAN DEN DRIES, *Tame topology and o -minimal structures*, vol. 248 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 1998.
- [36] L. VAN DEN DRIES AND C. MILLER, *On the real exponential field with restricted analytic functions*, Israel J. Math., 85 (1994), pp. 19–56.

E-mail address: scanlon@math.berkeley.edu

UNIVERSITY OF CALIFORNIA, BERKELEY, DEPARTMENT OF MATHEMATICS, EVANS HALL, BERKELEY, CA 94720-3840, USA