

ALGEBRAIC EQUATIONS ON THE ADÈLIC CLOSURE OF A DRINFELD MODULE

DRAGOS GHIOCA AND THOMAS SCANLON

ABSTRACT. Let k be a field of positive characteristic and $K = k(V)$ a function field of a variety V over k and let \mathbf{A}_K be the ring of adèles of K with respect to the places on K corresponding to the divisors on V . Given a Drinfeld module $\Phi : \mathbb{F}[t] \rightarrow \text{End}_K(\mathbb{G}_a)$ over K and a positive integer g we regard both K^g and \mathbf{A}_K^g as $\Phi(\mathbb{F}_p[t])$ -modules under the diagonal action induced by Φ . For $\Gamma \subseteq K^g$ a finitely generated $\Phi(\mathbb{F}_p[t])$ -submodule and an affine subvariety $X \subseteq \mathbb{G}_a^g$ defined over K , we study the intersection of $X(\mathbf{A}_K)$, the adèlic points of X , with $\bar{\Gamma}$, the closure of Γ with respect to the adèlic topology, showing under various hypotheses that this intersection is no more than $X(K) \cap \bar{\Gamma}$.

1. INTRODUCTION

Fix a field K with a set of places Ω so that each $x \in K$ is v -integral at all but finitely many $v \in \Omega$. We say that the Hasse principle holds for a given class \mathcal{C} of algebraic varieties over K if for each $X \in \mathcal{C}$ the set $X(K)$ of K -rational points on X is nonempty just in case for each $v \in \Omega$ the set $X(K_v)$ of K_v -rational points is nonempty where K_v is the completion of K with respect to v .

In some ways, the Hasse principle is both too strong and too weak. It is too strong in the sense that the classes of varieties for which it is known to hold are quite restrictive, for example, Brauer-Severi varieties when K is a number field and Ω is the set of all places, while there are many classes of varieties for which it is known to fail. It is too weak in the sense that it merely says that one may test for existence of a K -rational point by checking local conditions. It does not say that the set $X(K)$ may be computed from purely local data.

Let us suppose now that \mathcal{C} is a class of pairs (X, Y) where each of Y is a variety over K and X is a subvariety of Y also defined over K . Suppose moreover that for each such pair $(X, Y) \in \mathcal{C}$ that we know $Y(K)$. We wish to describe $X(K)$ as a subset of $Y(K)$ via strictly local data. Consider the ring of adèles of K :

$$\mathbf{A}_K := \left\{ (x_v)_{v \in \Omega} \in \prod_{v \in \Omega} K_v \mid x_v \in \mathcal{O}_{K_v} \text{ for all but finitely many } v \in \Omega \right\}.$$

Date: November 8, 2010.

From the diagonal inclusion $K \hookrightarrow \mathbf{A}_K$ we obtain inclusions $X(K) \subseteq Y(K) \subseteq Y(\mathbf{A}_K)$ and $X(K) \subseteq X(\mathbf{A}_K)$. Indeed, we may express $X(K) = Y(K) \cap X(\mathbf{A}_K)$. The set $Y(K)$ itself is not described by local conditions, but we may think of its closure with respect to the adèlic topology as a locally defined set. Replacing $Y(K)$ by its closure $\overline{Y(K)}$, in general we have only the inclusion $X(K) \subseteq \overline{Y(K)} \cap X(\mathbf{A}_K)$. When equality holds for all pairs $(X, Y) \in \mathcal{C}$ then we would say that the set of K -rational points on X may be computed locally. In [9], Poonen and Voloch show that when K is a function field of a curve, then the local conditions determine $X(K)$ whenever Y is an abelian variety and X is a subvariety. Moreover, in the case that the field of constants of K is finite, they relate the possible inequality of $X(K)$ with $\overline{Y(K)} \cap X(\mathbf{A}_K)$ to the Brauer-Manin obstruction.

With the above problem, we took the full set $Y(K)$ of K -rational points on Y as given. There is no good reason to do so, especially as in practice, even when Y is an abelian variety it might be possible to determine some points on Y and possibly thereby compute the group they generate, but the full computation of $Y(K)$ may be difficult. In other cases, some proper subset of $Y(K)$ might have more arithmetic significance. Let us say that a class \mathcal{C} of triples (X, Y, Γ) consisting of an algebraic variety Y over K , a subvariety $X \subseteq Y$ also over K and a set $\Gamma \subseteq Y(K)$ is locally determined if for each $(X, Y, \Gamma) \in \mathcal{C}$ we have $X(K) \cap \Gamma = X(\mathbf{A}_K) \cap \overline{\Gamma}$ in $Y(\mathbf{A}_K)$.

In this paper we take K to be a function field in positive characteristic and we address the question of when it happens that (X, Y, Γ) is locally determined for $Y = \mathbb{A}_K^n$ affine space over K , $X \subseteq Y$ an arbitrary subvariety and $\Gamma \subseteq Y(K)$ a finitely generated submodule with respect to the action of a Drinfeld module.

Let p be a prime number, $R := \mathbb{F}_p[t]$ and $\mathbb{F} := \mathbb{F}_p(t)$. For the remainder of this introduction we shall assume that K is a field extension of \mathbb{F} . We denote by $\text{End}_K(\mathbb{G}_a)$ the ring of endomorphisms of the additive group scheme over K . Concretely, $\text{End}_K(\mathbb{G}_a)$ may be identified with the ring of additive polynomials over K under addition and composition. A Drinfeld module of generic characteristic defined over K is a ring homomorphism $\Phi : R \rightarrow \text{End}_K(\mathbb{G}_a)$ for which $\Phi_t := \Phi(t)$ has differential t , but is not scalar multiplication. That is,

$$\Phi_t(x) = tx + c_1x^p + \cdots + c_Dx^{p^D} \in K[x] \text{ where } c_D \neq 0 \text{ (and } D \geq 1).$$

Let g be any positive integer, then we let Φ act diagonally on \mathbb{G}_a^g . This induces a natural structure of a $\Phi(R)$ -module on K^g where each element $a \in R$ acts via $\Phi_a := \Phi(a)$; when we want to emphasize the $\Phi(R)$ -module structure of K^g we will denote it by K_Φ^g . We say that $\Gamma \subseteq K^g$ is a $\Phi(R)$ -module if it is a submodule of K_Φ^g with respect to this action. Our main theorems (Theorem 2.1 and Theorem 2.3) assert that under appropriate hypotheses, for a subvariety $X \subseteq \mathbb{G}_a^g$ of a power of the additive group and a finitely generated $\Phi(R)$ -submodule $\Gamma \subseteq K^g$, the local conditions determine the intersection of Γ with X . That is, $X(K) \cap \Gamma = X(\mathbf{A}_K) \cap \overline{\Gamma}$.

We had embarked on this project hoping to thereby prove Denis's Mordell-Lang conjecture for Drinfeld modules [2].

Conjecture 1.1. *Let Φ be a Drinfeld module of generic characteristic. If $X \subseteq \mathbb{G}_a^g$ is any subvariety of a Cartesian power of the additive group and $\Gamma \subseteq K^g$ is a finitely generated $\Phi(R)$ -module, then $X(K) \cap \Gamma$ is a finite union of translates of $\Phi(R)$ -submodules of Γ .*

Instead, we have used the known partial results towards Conjecture 1.1 to reduce the problem of computing $X(\mathbf{A}_K) \cap \bar{\Gamma}$ to the case that X is zero dimensional. This is very much in the spirit of the proofs of the analogous results about semiabelian varieties [9, 12].

While the overall structure of our proofs is similar to those for semiabelian varieties, there are some essential differences; the most important of which concerns the problem of showing that the topology induced on Γ from the adèlic topology is at least as strong as the topology given by the subgroups $\Phi_a(\Gamma)$ as a ranges through the nonzero elements of R . In the case of semiabelian varieties, the corresponding result is almost immediate, but here, because the ambient geometry of the additive group does not determine the Drinfeld module structure, our proof is substantially more complicated.

This paper is organized as follows. In Section 2 we establish our notation and state our main results. In Section 3 we prove a simple case of our theorem in which the adèlic topology on Γ is discrete. In Section 4 we reduce the overall problem to that of zero dimensional X . We complete the proof in Section 5 by proving a uniform version of a version of the Drinfeld module Mordell-Lang conjecture.

Acknowledgments. The first author is grateful to Thomas Tucker for many stimulating conversations and for his constant encouragement for writing this paper.

2. STATEMENT OF OUR RESULTS

Let K be a finitely generated extension of \mathbb{F} such that $\text{trdeg}_{\mathbb{F}} K \geq 1$. We study the intersection of the adèlic points of V with the adèlic closure of Γ for both cases: generic characteristic and special characteristic Drinfeld modules $\Phi : R \rightarrow \text{End}_K(\mathbb{G}_a)$. First we need to set up our notation.

We let \bar{K} be a fixed algebraic closure of K , and we denote by K^{sep} the separable closure of K inside \bar{K} . We also let $\bar{\mathbb{F}}$ be the algebraic closure of \mathbb{F} inside \bar{K} . At the expense of replacing K by a finite extension and then replacing \mathbb{F} by a finite extension, we may assume (see [1, Remark 4.2]) there exists a projective, smooth, geometrically irreducible variety \mathcal{V} defined over \mathbb{F} such that its function field equals K (note that \mathbb{F} is the algebraic closure of $\mathbb{F}_p(t)$ inside K). We let $\Omega := \Omega_{\mathcal{V}}$ be the set of *all* valuations of K associated with the irreducible divisors of \mathcal{V} . By abuse of language, we will also call each $v \in \Omega$ a place of K ; note that for each nonzero $x \in \mathbb{F}$, we have $|x|_v = 1$, where $|\cdot|_v$ is the corresponding norm for the place v . Also note that \mathbb{F}

embeds naturally into the residue field \mathbb{F}_v of each place $v \in \Omega$; furthermore, each \mathbb{F}_v is a finite extension of \mathbb{F} and the reduction modulo v may be seen as a *specialization* of K to $\overline{\mathbb{F}}$. Finally, we note that K admits a product formula with respect to the set of places Ω (see [11]); i.e., there exists a set of positive integers $\{N_v\}_{v \in \Omega}$ such that for all $x \in K^*$ we have

$$(2.1) \quad \prod_{v \in \Omega} |x|_v^{N_v} = 1.$$

Now, for each $v \in \Omega$, we let K_v be the completion of K at v . We let \mathcal{O}_v be the ring of v -adic integers contained in K_v , and we equip K_v with the v -adic topology. We define the topological ring of adèles \mathbf{A}_K as the restricted direct product $\prod_{v \in \Omega_1} (K_v, \mathcal{O}_v)$ where Ω_1 is a cofinite subset of Ω . More precisely, each element of \mathbf{A}_K is of the form $\{x_v\}_{v \in \Omega_1}$, where $x_v \in K_v$ for all v , and moreover, for all but finitely many $v \in \Omega_1$ we actually have that $x_v \in \mathcal{O}_v$. In particular, $\prod_{v \in \Omega_1} \mathcal{O}_v$ is an open subset of \mathbf{A}_K whose induced topology is the usual product topology. We note that one could take instead the adèles as the restricted product of (K_v, \mathcal{O}_v) for *all* places $v \in \Omega$; i.e.,

$$\mathbf{A}_{K, \text{all}} := \prod_{v \in \Omega} (K_v, \mathcal{O}_v).$$

The restriction on K of the topology from $\mathbf{A}_{K, \text{all}}$ yields a stronger topology than the topology induced from \mathbf{A}_K . Therefore, the results we are proving regarding the closures in the adèlic topology of subsets of K are stronger with our definition for the adèles; see also the next result which is a purely topological statement.

Proposition 2.2. *Let $\Gamma \subset K$ be a set of points which are all integral at all places in Ω_1 . Let $y = (y_v)_{v \in \Omega} \in \mathbf{A}_{K, \text{all}}$ be a point in the closure of Γ inside $\mathbf{A}_{K, \text{all}}$. Assume we know that there exists $y_0 \in K$ such that $y_v = y_0$ for all $v \in \Omega_1$. Then $y_v = y_0$ for all $v \in \Omega$; i.e., $y = y_0$.*

Proof. Clearly, y_0 is integral at each place $v \in \Omega_1$. At the expense of replacing y by $y - y_0$ we may assume $y_0 = 0$. Therefore, we know that there exists an infinite sequence $\{x_n\}_{n \in \mathbb{N}} \subset K$ satisfying the following conditions:

- (a) $|x_n|_v \leq 1$ for all $v \in \Omega_1$;
- (b) $|x_n|_v \rightarrow 0$ as $n \rightarrow \infty$, for all $v \in \Omega_1$; and
- (c) $|x_n - y_v|_v \rightarrow 0$ as $n \rightarrow \infty$, for all $v \in \Omega \setminus \Omega_1$.

Using condition (c) above, we conclude that there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ we have $|x_n|_v = |y_v|_v$ for all $v \in \Omega \setminus \Omega_1$. In particular, there exists a positive real number C_0 such that for all $n \geq n_0$, we have

$$(2.3) \quad \prod_{v \in \Omega \setminus \Omega_1} |x_n|_v^{N_v} \leq C_0.$$

On the other hand, using conditions (a)-(b) above we conclude that

$$(2.4) \quad \lim_{n \rightarrow \infty} \prod_{v \in \Omega_1} |x_n|_v^{N_v} = 0.$$

Now, if the sequence $\{x_n\}_{n \in \mathbb{N}}$ contains infinitely many nonzero elements, then equations (2.3) and (2.4) provide a contradiction to the product formula (2.1). Therefore, we obtain that there exists $n_1 \in \mathbb{N}$ such that $x_n = 0$ for all $n \geq n_1$. Hence for each $v \in \Omega \setminus \Omega_1$ we have $y_v = 0$, which concludes our proof. \square

Naturally, $\mathbb{G}_a^g(\mathbf{A}_K) = \mathbf{A}_K^g$, and we define the product topology on \mathbf{A}_K^g . Then for any subset $S \subseteq \mathbf{A}_K^g$, we let \overline{S} be the topological closure of S in \mathbf{A}_K^g . Clearly, \mathbf{A}_K (and \mathbf{A}_K^g) have natural structures of $\Phi(R)$ -modules denoted $\mathbf{A}_{K,\Phi}$ (and respectively $\mathbf{A}_{K,\Phi}^g$).

We will prove the following result.

Theorem 2.5. *Let K be a function field over \mathbb{F} of arbitrary finite transcendence degree, and let Φ be a Drinfeld module of generic characteristic defined over K . Then for each affine subvariety $V \subseteq \mathbb{G}_a^g$ defined over K , and for each finitely generated $\Phi(R)$ -submodule $\Gamma \subseteq K^g$, we have $V(K) \cap \Gamma = V(\mathbf{A}_K) \cap \overline{\Gamma}$.*

So, Theorem 2.5 shows that the intersection $V(K) \cap \Gamma$ can be recovered from the local information (as given by the adèles). Theorem 2.5 is an immediate consequence of the fact that for all but finitely many places $v \in \Omega$, the induced v -adic topology on Γ is the discrete topology (see Proposition 3.1).

Our Theorem 2.5 is the Drinfeld modules analogue of [9, Theorem A]. Similarly, we can prove an analogue of [9, Theorem B] for Drinfeld modules of special characteristic.

A Drinfeld module of special characteristic is a ring homomorphism $\Phi : R \rightarrow \text{End}_K(\mathbb{G}_a)$ such that $\Phi_t = \Phi(t)$ is an inseparable endomorphism of the additive group scheme; more precisely,

$$\Phi_t(x) = \sum_{i=d}^D c_i x^{p^i} \text{ for some } c_i \in K, \text{ where } 1 \leq d \leq D.$$

In order to state our next result we also need the following definition.

Definition 2.6. *For each $\Phi(R)$ -submodule $\Gamma \subseteq K^g$, we call its full divisible hull inside K , the set of all $x \in \mathbb{G}_a^g(K)$ such that for some nonzero $a \in R$ we have $\Phi_a(x) \in \Gamma$.*

We say that the subgroup $\Gamma \subseteq K_\Phi^g$ is full in K , if it equals its full divisible hull inside K_Φ^g . When the field K is understood from the context, we will simply say that Γ is full.

As proved in [13] (see also [8] and [3, Theorem 5.12] for similar results), the $\Phi(R)$ -module structure of K_Φ^g is that of a direct sum of a finite torsion module with a free $\Phi(R)$ -module of rank \aleph_0 . So, replacing Γ with the $\Phi(R)$ -submodule containing all $x \in K^g$ such that $\Phi_a(x) \in \Gamma$ for some nonzero $a \in R$, we may assume Γ is both full in K and finitely generated.

Let $\Phi : R \rightarrow \text{End}_K(\mathbb{G}_a)$ be any Drinfeld module, and let L be a field containing K . For each nonzero $a \in R$ we let $\Phi[a](L)$ be the set of all $x \in L$

such that $\Phi_a(x) = 0$. We also let

$$\Phi[t^\infty](L) := \bigcup_{n \in \mathbb{N}} \Phi[t^n](L),$$

and

$$\Phi_{\text{tor}}(L) := \bigcup_{a \in R \setminus \{0\}} \Phi[a](L).$$

When $L = \overline{K}$, we simply use $\Phi[a]$ and Φ_{tor} instead of $\Phi[a](\overline{K})$ and respectively $\Phi_{\text{tor}}(\overline{K})$. In general, for any $\Phi(R)$ -submodule Γ we denote by Γ_{tor} the torsion submodule of Γ .

We also recall the notion of modular transcendence degree, first introduced by the second author in [10]. The *modular transcendence degree* of Φ is the smallest nonnegative integer d such that there exists a subfield $L \subseteq \overline{K}$ of transcendence degree d over \mathbb{F}_p , and there exists $\gamma \in \overline{K}$ such that

$$(2.7) \quad \gamma^{-1} \Phi_t(\gamma \cdot x) \in L[x].$$

Theorem 2.8. *Let K be a function field over \mathbb{F} of arbitrary finite transcendence degree, and let Φ be a Drinfeld module of special characteristic defined over K . Assume $\Phi[t^\infty](K^{\text{sep}})$ is finite, and also assume that Φ has positive modular transcendence degree. Let V be an affine subvariety of \mathbb{G}_a^g defined over K which contains no translate of a positive dimensional algebraic subgroup of \mathbb{G}_a^g . Then for each finitely generated full $\Phi(R)$ -submodule $\Gamma \subseteq K^g$, we have $V(K) \cap \Gamma = V(\mathbf{A}_K) \cap \overline{\Gamma}$.*

Since we assumed that V contains no translate of a positive dimensional algebraic subgroup of \mathbb{G}_a^g , the main result of [10] (see also our Theorem 5.3) yields that $V(K) \cap \Gamma$ is finite. However, the novelty of our result is that we can show that each of these finitely many points from the intersection $V(K) \cap \Gamma$ can be recovered purely from the local information given by the adèlic topology. The strategy for proving Theorem 2.8 is as follows:

- (a) in Section 4 we show that Theorem 2.8 holds under the additional hypothesis that V is zero-dimensional; and
- (b) then in Section 5 we show that for each subvariety V , there exists a zero-dimensional subvariety V_0 of V also defined over K such that $V(\mathbf{A}_K) \cap \overline{\Gamma} = V_0(\mathbf{A}_K) \cap \overline{\Gamma}$. This reduces Theorem 2.8 to part (a) above.

3. PROOF OF THEOREM 2.5

We are working with the notation and the hypotheses of Theorem 2.5 (see Section 2).

For each finitely generated $\Phi(R)$ -submodule Γ of K_Φ^g , we have $\Gamma \subseteq \Gamma_0^g$, where $\Gamma_0 \subseteq K_\Phi$ is the $\Phi(R)$ -submodule generated by all projections of Γ on each coordinate of \mathbb{G}_a^g . Clearly, Γ_0 is also a finitely generated $\Phi(R)$ -submodule. We will show that each finitely generated $\Phi(R)$ -submodule of

K_Φ is discrete with respect to all but finitely many places $v \in \Omega$. In particular this yields that Γ is discrete v -adically for almost all places v . Thus $\bar{\Gamma} = \Gamma$, and hence Theorem 2.5 follows immediately.

Proposition 3.1. *Let $\Gamma \subseteq K_\Phi$ be a finitely generated $\Phi(R)$ -submodule, and let v be a place in Ω at which all coefficients of Φ_a (for each $a \in R$) and also all elements of Γ are integral. Then Γ is discrete as a subset of K_v .*

Proof. We have to show that there is no infinite sequence of points in Γ converging to 0 with respect to the v -adic topology. We prove Proposition 3.1 by induction on the number r of generators for the $\Phi(R)$ -module Γ .

Assume $r = 1$, i.e. Γ is the cyclic $\Phi(R)$ -module generated by x_1 , and also assume Γ is not discrete with respect to the v -adic topology. In particular, this yields that x_1 is not torsion since otherwise Γ would be finite (and thus discrete v -adically).

Therefore, there exists a sequence of distinct nonzero elements $a_n \in R$ (for each $n \in \mathbb{N}$) such that $|\Phi_{a_n}(x_1)|_v \rightarrow 0$ as $n \rightarrow \infty$. Now, since each coefficient of Φ_a , for each $a \in R$ is integral at v , we obtain that the set

$$I_1 := \{a \in R : |\Phi_a(x_1)|_v < 1\}$$

is an ideal (see also [4, Lemma 3.11]). Since R is a PID, we conclude that I_1 is generated by some polynomial c_1 . Furthermore, since we assumed that $|\Phi_{a_n}(x_1)|_v \rightarrow 0$, we obtain that I_1 is nontrivial, and thus $c_1 \neq 0$. So, for each $n \in \mathbb{N}$ such that $|\Phi_{a_n}(x_1)|_v < 1$ we have that $c_1 \mid a_n$, and thus there exists a nonzero $b_n \in R$ such that $a_n = b_n \cdot c_1$. On the other hand, $|b_n|_v = 1$ since each element of $\mathbb{F}_p(t)^* \subseteq \mathbb{F}^*$ is a v -adic unit. According to [4, Lemma 3.10], we obtain then that

$$|\Phi_{a_n}(x_1)|_v = |\Phi_{b_n}(\Phi_{c_1}(x_1))|_v = |b_n \cdot \Phi_{c_1}(x_1)|_v = |\Phi_{c_1}(x_1)|_v,$$

which contradicts our assumption that $\Phi_{a_n}(x_1)$ converges v -adically to 0.

Now, assume we proved that Proposition 3.1 holds for each finitely generated $\Phi(R)$ -module spanned by less than r elements; next we prove that it holds when Γ is generated by r elements. Since Γ is a finitely generated $\Phi(R)$ -submodule, it is a finite union of cosets of free finitely generated $\Phi(R)$ -submodules. Therefore, it suffices to prove Proposition 3.1 for free submodules. Let x_1, \dots, x_r be a basis for the free $\Phi(R)$ -module Γ .

Assume there exists an infinite sequence $\{(a_{n,1}, \dots, a_{n,r})\}_{n \in \mathbb{N}}$ of nonzero tuples of elements of R (i.e. for each $n \in \mathbb{N}$ not all $a_{n,i} = 0$) such that

$$\left| \sum_{i=1}^r \Phi_{a_{n,i}}(x_i) \right|_v \rightarrow 0 \text{ as } n \rightarrow \infty.$$

The set I_r of tuples $(c_1, \dots, c_r) \in R^r$ such that $|\sum_{i=1}^r \Phi_{c_i}(x_i)|_v < 1$ is an R -submodule of R^r (again using [4, Lemma 3.11] since all coefficients of each Φ_a are v -adic integers). Since R is a PID and R^r is a free R -module, then I_r is a free R -module of rank $s \leq r$. Let $\{(c_{j,1}, \dots, c_{j,r})\}_{1 \leq j \leq s}$ be a basis for

the free R -module I_r . For each $j = 1, \dots, s$, we let

$$y_j := \sum_{i=1}^r \Phi_{c_{j,i}}(x_i).$$

Note that each $|y_j|_v < 1$ since $(c_{j,1}, \dots, c_{j,r}) \in I_r$. Also, each y_j is nonzero since the elements x_1, \dots, x_r form a basis for the free R -module Γ , and not all $c_{j,i}$ are 0 (for $i = 1, \dots, r$).

We know that for n sufficiently large, $(a_{n,1}, \dots, a_{n,r}) \in I_r$, and thus each $\sum_{i=1}^r \Phi_{a_{n,i}}(x_i)$ is in the $\Phi(R)$ -module generated by y_1, \dots, y_s . So, if $s < r$, we are done by the induction hypothesis. Therefore, assume from now on that $s = r$. So, for each $n \in \mathbb{N}$ such that $|\sum_{i=1}^r \Phi_{a_{n,i}}(x_i)|_v < 1$, there exist $b_{n,1}, \dots, b_{n,r} \in R$ such that

$$a_{n,i} = \sum_{j=1}^r b_{n,j} c_{j,i} \text{ for all } i = 1, \dots, r.$$

Therefore $\sum_{i=1}^r \Phi_{a_{n,i}}(x_i) = \sum_{j=1}^r \Phi_{b_{n,j}}(y_j)$. Since each $|y_j|_v < 1$, we get that for each $b \in R$ we have

$$(3.2) \quad \Phi_b(y_j) = by_j + L(b, y_j) \text{ where } |L(b, y_j)|_v < |y_j|_v,$$

since each coefficient of Φ_b is a v -adic integer. Without loss of generality assume that

$$|y_1|_v = |y_2|_v = \dots = |y_\ell|_v = |u|_v^m > |y_{\ell+1}|_v \geq \dots |y_r|_v \text{ for some } 1 \leq \ell \leq r,$$

where u is some fixed uniformizer of the maximal ideal of \mathcal{O}_v , and $m \in \mathbb{N}$. Using (3.2) we conclude that

$$\sum_{j=1}^r \Phi_{b_{n,j}}(y_j) = \sum_{j=1}^{\ell} b_{n,j} y_j + L(n, \bar{y}),$$

where $|L(n, \bar{y})|_v < |u|_v^m$. For each $j = 1, \dots, \ell$, we let d_j be the first nonzero coefficient when expanding y_j with respect to u inside K_v . So,

$$y_j = d_j \cdot u^m + \text{larger powers of } u,$$

with $|d_j|_v = 1$ for each $j = 1, \dots, \ell$. Thus

$$\sum_{j=1}^r \Phi_{b_{n,j}}(y_j) = \left(\sum_{j=1}^{\ell} b_{n,j} d_j \right) \cdot u^m + L_1(n, \bar{y}),$$

where $|L_1(n, \bar{y})|_v < |u|_v^m$. Now, since

$$\sum_{i=1}^r \Phi_{a_{n,i}}(x_i) = \sum_{j=1}^r \Phi_{b_{n,j}}(y_j) \rightarrow 0 \text{ as } n \rightarrow \infty,$$

we conclude that for all $n \in \mathbb{N}$ sufficiently large we have that $|\sum_{j=1}^{\ell} b_{n,j} d_j|_v < 1$. For each $j = 1, \dots, \ell$, we let e_j be the reduction of d_j modulo v , and since

$\mathbb{F}_p(t) \subseteq \mathbb{F}$ embeds naturally into the residue field at the place v , we conclude that

$$(3.3) \quad \sum_{j=1}^{\ell} e_j b_{n,j} = 0,$$

for n sufficiently large. Since each d_j is a v -adic unit, we conclude that each e_j is nonzero. The set S of tuples (b_1, \dots, b_r) for which $\sum_{j=1}^{\ell} e_j b_j = 0$ is a submodule of R^r of rank $r - 1$. Let $\{(f_{j,1}, \dots, f_{j,r})\}_{1 \leq j \leq r-1}$ be generators of S , and let $\Gamma_S \subseteq K_{\Phi}$ be the $\Phi(R)$ -module spanned by

$$\sum_{i=1}^r \Phi_{f_{j,i}}(y_i) \text{ for each } j = 1, \dots, r - 1.$$

By the induction hypothesis on the rank of the R -module Γ_S (which is less than r), we know that Γ_S is discrete v -adically. Since for n sufficiently large we have that all $\sum_{i=1}^r \Phi_{a_{n,i}}(x_i) = \sum_{j=1}^r \Phi_{b_{n,j}}(y_j)$ are in Γ_S , we are done. \square

Proof of Theorem 2.5. Let Γ_0 be the $\Phi(R)$ -submodule of K spanned by the projections of Γ on each coordinate of K_{Φ}^g . Since Γ is finitely generated, then also Γ_0 is finitely generated. We let v be a place in Ω at which all coefficients of Φ_t and also each of the finitely many generators of Γ_0 are integral. In particular, this means that *all* elements of Γ are integral at the place v , and also that each coefficient of each Φ_a is a v -adic integer. Note that all but finitely many places $v \in \Omega$ satisfy these conditions. Proposition 3.1 yields that Γ_0 (and thus Γ_0^g) is discrete v -adically. Therefore $\bar{\Gamma} = \Gamma$ and so, $V(K) \cap \Gamma = V(\mathbf{A}) \cap \bar{\Gamma}$, as desired. \square

4. PROOF OF THEOREM 2.8 WHEN $\dim(V) = 0$

In this Section, we work under the hypotheses from Theorem 2.8. So, Φ is a Drinfeld module of special characteristic such that $\Phi[t^{\infty}](K^{\text{sep}})$ is finite. In addition, we assume Φ has positive modular transcendence degree. First we will prove several preliminary results which describe the topology induced on each finitely generated $\Phi(R)$ -module Γ by the adélic topology.

We let Ω_0 be the set of all places $v \in \Omega$ at which all coefficients of Φ_t are integral, while the first and the last nonzero coefficients of Φ_t are v -adic units. Clearly, $\Omega \setminus \Omega_0$ is finite. Note that by our assumption, we obtain that for *all* nonzero $a \in R$, each coefficient of Φ_a is a v -adic integer, while the first and the last coefficients of Φ_a are v -adic units.

Proposition 4.1. *For each $v \in \Omega_0$ and for each nonzero $a \in R$, the quotient $K_v^g / \Phi_a(K_v^g)$ is Hausdorff.*

Proof. Let $v \in \Omega_0$, and let $a \in R$ be nonzero. It suffices to show that $K_v / \Phi_a(K_v)$ is Hausdorff, which is equivalent with showing that $\Phi_a(K_v)$ is closed in K_v . For this, we need to show that if $\Phi_a(x_n) \rightarrow y$ as $n \rightarrow \infty$ for some $x_n \in K_v$ and $y \in K_v$, then actually $y \in \Phi_a(K_v)$.

Let $z_n := x_{n+1} - x_n$. We have that $\Phi_a(z_n) \rightarrow 0$ as $n \rightarrow \infty$. We claim that the distance between z_n and the nearest torsion point from $\Phi[a]$ tends to 0 with respect to the v -adic topology.

If $t \nmid a$, this is an immediate consequence of Hensel's Lemma. Indeed, reducing modulo v , we obtain that the reduction \bar{z}_n modulo v of z_n is a root of the reduced polynomial $\bar{\Phi}_a$ modulo v . Now, since $\Phi'_a(x) = a_0 \neq 0$ (where $a_0 \in \mathbb{F}_p$ such that $t \mid a - a_0$), and $|a_0|_v = 1$, we conclude using Hensel's Lemma that there exists a torsion point $u_n \in \Phi[a](K_v)$ such that $|z_n - u_n|_v < 1$. Note also that each torsion point is integral at each place v in Ω_1 due to our assumptions regarding the v -adic absolute values of the coefficients of Φ_t . Then

$$|z_n - u_n|_v = |a_0(z_n - u_n)|_v = |\Phi_a(z_n - u_n)|_v = |\Phi_a(z_n)|_v \rightarrow 0, \text{ as desired.}$$

Now, assume $t \mid a$. Let $a = c_m t^m + \dots + c_n t^n$ (with $1 \leq m \leq n$), where each $c_i \in \mathbb{F}_p$, and $c_m, c_n \neq 0$. Letting $\Phi_t(x) = \sum_{j=\ell}^D \gamma_j x^{p^j}$, where $\gamma_\ell, \gamma_D \neq 0$, then

$$\Phi_a(x) = c_m \gamma_\ell^{(p^{m\ell}-1)/(p^\ell-1)} x^{p^{m\ell}} + \sum_{j=m\ell+1}^{nD} \delta_j x^{p^j}.$$

We let $f := \Phi_a^{1/p^{m\ell}} \in K^{1/p^{m\ell}}[x]$. It is immediate to see that $f(z_n) \rightarrow 0$ in the v -adic topology (by abuse of language, we call also v the induced valuation on $K_v^{1/p^{m\ell}}$ extending v). Also, $f(x)$ is a separable additive polynomial in $K^{1/p^{m\ell}}[x]$ whose derivative equals $\delta_0 := c_m \gamma_\ell^{(p^{m\ell}-1)/(p^{(m+1)\ell}-p^{m\ell})}$. By our definition of the set Ω_0 , we know that δ_0 is a v -adic unit. Therefore, by Hensel's Lemma, just as before, we conclude that the distance from z_n to the nearest root $u_n \in K_v^{1/p^{m\ell}}$ of $f(x) = 0$ (which is also a torsion point in $\Phi[a]$) converges to 0 in the v -adic topology.

So, for each $n \in \mathbb{N}$, there exists a suitable torsion point $z'_n \in \Phi[a](K_v^{1/p^{m\ell}})$ such that if $y_n = x_n + z'_n$, then the sequence $\{y_n\}_{n \in \mathbb{N}}$ is convergent in $K_v^{1/p^{m\ell}}$. Because $\Phi[a]$ is a finite set, it means that we can extract an infinite sequence $\{n_i\}_{i \in \mathbb{N}}$ of positive integers such that $z'_{n_i} = z'_{n_{i+1}}$ for all $i \in \mathbb{N}$. Hence the subsequence $\{x_{n_i}\}_{i \in \mathbb{N}}$ is convergent in K_v ; since K_v is a complete space, then $\{x_{n_i}\}_{i \in \mathbb{N}}$ converges to a point x such that $\Phi_a(x) = y$, as desired. \square

Let $\Gamma \subseteq \mathbb{G}_a^g(K)$ be a finitely generated $\Phi(R)$ -submodule; we also assume that Γ is full in K . Let $\Omega_1 \subseteq \Omega_0$ be the set of all places v at which the finitely many generators of $\Gamma \subseteq \mathbb{G}_a^g(K)$ are integral. In particular, this means that all points of Γ are v -adic integral for $v \in \Omega_1$; also note that $\Omega \setminus \Omega_1$ is finite. Without loss of generality (see also Proposition 2.2) we may assume \mathbf{A}_K is the restricted product $\prod_{v \in \Omega_1} (K_v, \mathcal{O}_v)$.

Proposition 4.2. *Let Γ be a full, finitely generated $\Phi(R)$ -submodule of K^g , and let $v \in \Omega_1$. Then the v -adic topology on Γ is at least as strong as the topology induced by all subgroups $\Phi_{t^n}(\Gamma)$ for $n \in \mathbb{N}$.*

Proof. All we need to show is that for each $n \in \mathbb{N}$, there exists an open subset U_n of K_v^g such that $U_n \cap \Gamma \subseteq \Phi_{t^n}(\Gamma)$.

Let $n \in \mathbb{N}$. We know that $\Phi[t^\infty](K^{\text{sep}})$ is finite, which means that there exists $m \in \mathbb{N}$ such that $\Phi[t^\infty](K^{\text{sep}}) \subseteq \Phi[t^m]$. Let

$$\pi : \Gamma \longrightarrow K_v^g / \Phi_{t^{n+m}}(K_v^g)$$

be the composition of the natural injection of Γ into K_v^g followed by the canonical projection map on the quotient $K_v^g / \Phi_{t^{n+m}}(K_v^g)$; clearly, π is a continuous map. According to Proposition 4.1, the quotient $K_v^g / \Phi_{t^{n+m}}(K_v^g)$ has an induced Hausdorff topology, which means that there exists an open subset U_n of K_v^g such that $U_n \cap \Gamma = \ker(\pi)$ (we also use the fact that $\Gamma / \Phi_{t^{n+m}}(\Gamma)$ is finite, and hence the induced topology on $\Gamma / \Phi_{t^{n+m}}(\Gamma)$ as a subset of $K_v^g / \Phi_{t^{n+m}}(K_v^g)$ is the discrete topology).

Now, for any $y \in U_n \cap \Gamma$, there exists $x \in K_v^g \cap \overline{K}^g$ such that $y = \Phi_{t^{n+m}}(x)$. However, $\overline{K} \cap K_v \subseteq K^{\text{sep}}$ (see [9, Lemma 3.1]); so, $x \in (K^{\text{sep}})^g$. Now, for any automorphism $\sigma \in \text{Gal}(K^{\text{sep}}/K)$, we have that $\Phi_{t^{n+m}}(x^\sigma - x) = 0$ (since $y = \Phi_{t^{n+m}}(x) \in K^g$), which means that $x^\sigma - x \in (\Phi[t^{n+m}](K^{\text{sep}}))^g$. So, in particular $x^\sigma - x \in (\Phi[t^m](K^{\text{sep}}))^g$, which means that $\Phi_{t^m}(x) \in K^g$. However, since $\Phi_{t^m}(x) \in K^g$ and $\Phi_{t^n}(\Phi_{t^m}(x)) = y \in \Gamma$, while Γ is a full $\Phi(R)$ -submodule in K , we conclude that also $\Phi_{t^m}(x) \in \Gamma$, and thus $y = \Phi_{t^n}(\Phi_{t^m}(x)) \in \Phi_{t^n}(\Gamma)$. Therefore $U_n \cap \Gamma \subseteq \Phi_{t^n}(\Gamma)$, as desired. \square

Proposition 4.3. *Let $\Gamma \subseteq K^g$ be a finitely generated $\Phi(R)$ -submodule which is full in K . For each $v \in \Omega_1$ we let Γ_v be the closure of Γ in K_v^g . Then $\Gamma_v \cap K^g = \Gamma$.*

Proof. One inclusion is obvious. Now, let $y \in \Gamma_v \cap K^g$. We let Γ_1 be the full divisible hull in K of the finitely generated $\Phi(R)$ -module Γ_0 spanned by Γ and y . There exists an infinite sequence of distinct elements $x_n \in \Gamma$ such that $x_n \rightarrow y$ in K_v^g . So, by Proposition 4.2, we have that for all $j \in \mathbb{N}$, there exists $n_j \in \mathbb{N}$ such that

$$(4.4) \quad x_{n_j} - y = \Phi_{t^j}(y_j) \text{ for some } y_j \in \Gamma_1.$$

Since Γ_1 is finitely generated as a $\Phi(R)$ -module, and Γ_1 is the full divisible hull in K of Γ_0 , we conclude that there exists a nonzero $b \in R$ such that $\Phi_b(\Gamma_1) \subseteq \Gamma_0$. We assume $b = t^{j_0} \cdot b_0$, for some $j_0 \in \mathbb{N}$, and some $b_0 \in R$ which is not divisible by t . Now, we use (4.4) for $j = j_0 + 1$, and obtain

$$(4.5) \quad \Phi_{b_0}(x_{n_j}) - \Phi_{b_0}(y) = \Phi_{b_0}(\Phi_{t^{j_0+1}}(y_j)).$$

Because $\Phi_{b_0 t^{j_0}}(y_j) \in \Gamma_0$, we conclude that there exist $z_j \in \Gamma$ and $c_j \in R$ such that $\Phi_{b_0 t^{j_0}}(y_j) = z_j + \Phi_{c_j}(y)$. Therefore, (4.5) reads

$$\Phi_{b_0}(x_{n_j}) - \Phi_{b_0}(y) = \Phi_t(z_j) + \Phi_{t c_j}(y).$$

Hence, $\Phi_{t c_j + b_0}(y) = \Phi_{b_0}(x_{n_j}) - \Phi_t(z_j) \in \Gamma$. Since $t \nmid b_0$, we get that $t c_j + b_0 \neq 0$, which yields that $y \in \Gamma$ because Γ is full in K . \square

Proposition 4.6. *The topology induced on Γ by the adèlic topology is at least as strong as the one given by all subgroups $\Phi_a(\Gamma)$ for $a \in R$ relative prime to t .*

Proof. Let $a \in R$ relative prime to t . We have to show that there exists an open subset U_a in the adèlic topology such that $U_a \cap \Gamma \subseteq \Phi_a(\Gamma)$. Since Φ_a is separable (more precisely, Φ'_a is a constant v -adic unit for all $v \in \Omega$), using Hensel's Lemma, we obtain that $\Phi_a(\mathcal{O}_v^g)$ is an open subgroup of K_v^g for each $v \in \Omega_1$. Therefore it suffices to show that there exist finitely many places $v_i \in \Omega_1$ (with $1 \leq i \leq n_a$ for some $n_a \in \mathbb{N}$) such that for all $y \in \Gamma$ satisfying $y \in \Phi_a(\mathcal{O}_{v_i}^g)$ for $1 \leq i \leq n_a$, we have $y \in \Phi_a(\Gamma)$. Since Γ is full in K , we only have to show that $y \in \Phi_a(K^g)$.

Now, assume there is no such finite set of places v_i as above. This means that for any finite subset S of places contained in Ω_1 , we may find $y = y_S \in \Gamma \setminus \Phi_a(\Gamma)$ such that $y \in \Phi_a(\mathcal{O}_v^g)$ for each $v \in S$. Note that the quotient $\Gamma/\Phi_a(\Gamma)$ is finite (since Γ is a finitely generated $\Phi(R)$ -module and $a \neq 0$); let $y_1, \dots, y_s \in \Gamma$ be a set of representatives for the nonzero cosets of $\Gamma/\Phi_a(\Gamma)$.

For each coset $y_i + \Phi_a(\Gamma)$ for $i = 1, \dots, s$, we let S_{y_i} be the set of places $v \in \Omega_1$ such that $y_i \notin \Phi_a(\mathcal{O}_v^g)$; note that our definition does not depend on the particular choice of the class representative for the particular coset of $\Phi_a(\Gamma)$ in Γ . If each S_{y_i} is nonempty, then for each $i = 1, \dots, s$, choose $v_i \in S_{y_i}$ and let $S := \{v_1, \dots, v_s\}$. By our assumption, if $y \in \Phi_a(\mathcal{O}_{v_i}^g)$ for each $i = 1, \dots, s$, then $y \in \Phi_a(\Gamma)$, as desired.

So, from now on assume there exists a coset $y + \Phi_a(\Gamma)$ of Γ (as above) such that S_y is empty, i.e. for all places $v \in \Omega_1$, we have $y \in \Phi_a(\mathcal{O}_v^g)$. So, even though $y \notin \Phi_a(K^g)$ (note that Γ is full in K), for all but finitely many places $v \in \Omega$ we still have that $y \in \Phi_a(\mathcal{O}_v^g)$. We will show next that this is impossible. At the expense of passing to the coordinates in \mathbb{G}_a^g of each point, we may assume $g = 1$. In particular this means that the polynomial $\Phi_a(X) - y \in K[X]$ has no linear irreducible factor over K . Since \mathbb{F} is a finite extension of $\mathbb{F}_p(t)$, then \mathbb{F} is a Hilbertian field (see [6, Ch. 9]). This yields that there exist infinitely many places $v \in \Omega_1$ such that the reduction modulo v of the polynomial $\Phi_a(X) - y$ has no irreducible linear factor over \mathbb{F}_v . This contradicts our assumption that for all but finitely many places $v \in \Omega_1$ we have $y \in \Phi_a(\mathcal{O}_v)$ since that would mean that for all but finitely many places $v \in \Omega_1$, the reduction modulo v of y belongs to $\Phi_a(\mathbb{F}_v)$ (since \mathbb{F}_v is the residue field of K and also of \mathcal{O}_v modulo v). This concludes the proof of our Proposition 4.6. \square

Combining Propositions 4.2 and 4.6 we obtain the following result.

Proposition 4.7. *The induced adèlic topology on Γ is at least as strong as the topology given by all subgroups $\Phi_a(\Gamma)$ for all $a \in R$.*

Proposition 4.8. *Let $\Gamma \subseteq K_{\mathbb{F}}^g$ be any finitely generated $\Phi(R)$ -submodule which is full in K . Then $\overline{\Gamma}_{\text{tor}} = \Gamma_{\text{tor}}$, i.e. all torsion points contained in $\overline{\Gamma}$ are actually contained in Γ .*

Proof. The proof is similar to the proof of [9, Lemma 3.7]. Let Γ_0 be the subgroup of all $y \in \Gamma$ such that for all $v \in \Omega_1$ the reduction of y modulo v is in $(\overline{\Phi}_{\text{tor}}(\mathbb{F}_v))^g$, where $\overline{\Phi}$ is the reduction of Φ modulo v . It is immediate to see that Γ_0 is full in K , and thus it contains Γ_{tor} .

Now, since Γ is a finitely generated $\Phi(R)$ -submodule, there exist finitely many places $v_i \in \Omega_1$ for $i = 1, \dots, \ell$ such that Γ_0 is the kernel of the natural reduction map

$$\Gamma \longrightarrow \prod_{i=1}^{\ell} (\mathbb{F}_{v_i} / \overline{\Phi}_{\text{tor}}(\mathbb{F}_{v_i}))^g.$$

Therefore Γ/Γ_0 embeds into the free $\Phi(R)$ -submodule

$$\prod_{i=1}^{\ell} (\mathbb{F}_{v_i} / \overline{\Phi}_{\text{tor}}(\mathbb{F}_{v_i}))^g.$$

We conclude that there exists a free submodule $\Gamma_1 \subseteq \Gamma$ such that $\Gamma = \Gamma_0 \oplus \Gamma_1$. We claim that

$$(4.9) \quad \overline{\Gamma} = \overline{\Gamma_0} \oplus \Gamma_1$$

Indeed, for each sequence $\{x_n + y_n\}_{n \in \mathbb{N}}$ where $x_n \in \Gamma_0$ and $y_n \in \Gamma_1$ which converges in \mathbf{A}_K^g , we obtain that for m and n sufficiently large we have that $y_n - y_m$ and $x_n - x_m$ have the same reduction modulo v_i for each $i = 1, \dots, \ell$. Therefore, $y_n - y_m \in \Gamma_0$, and thus $y_n = y_m$ since $\Gamma = \Gamma_0 \oplus \Gamma_1$.

So $\overline{\Gamma} = \overline{\Gamma_0} \oplus \Gamma_1$ and since Γ_1 is a free $\Phi(R)$ -module, we obtain that $\overline{\Gamma}_{\text{tor}} = (\overline{\Gamma_0})_{\text{tor}}$. We claim that the induced adèlic topology on Γ_0 is the one given by all subgroups $\Phi_a(\Gamma_0)$ for all $a \in R$. By Proposition 4.7 we know that the adèlic topology induces on Γ_0 a topology at least as strong as the one given by all subgroups $\Phi_a(\Gamma_0)$. Now we check the converse statement. Indeed, for each $y \in \Gamma_0$ and for each $v \in \Omega_1$, there exists a nonzero $a_v \in R$ such that $\Phi_{a_v}(y)$ reduces to 0 modulo v . Then $\Phi_{t^n a_v}(y)$ converges v -adically to 0 as $n \rightarrow \infty$, because v is a place of integrality for all coefficients of $\Phi_{t^n a_v}$ and moreover Φ_t has no linear or constant term. Moreover, let $\{y_i\}_{1 \leq i \leq s}$ be a finite set of generators for Γ_0 as a $\Phi(R)$ -module, and let $a_{i,v} \in R$ be as above for each corresponding y_i and each place $v \in \Omega_1$. Letting $b_v := \prod_{i=1}^s a_{i,v}$, we obtain that

$$\Phi_{t^n b_v}(y) \text{ covers } v\text{-adically to } 0, \text{ as } n \rightarrow \infty, \text{ for all } y \in \Gamma_0.$$

This proves our claim about the induced adèlic topology on Γ_0 . Thus

$$(4.10) \quad \overline{\Gamma_0} \xrightarrow{\sim} \Gamma_0 \otimes_R \hat{R},$$

where \hat{R} is the restricted direct product of all completions of R at its maximal ideals. Since $R/aR \xrightarrow{\sim} \hat{R}/a\hat{R}$ for each nonzero $a \in R$, we conclude that

$$\overline{\Gamma}_{\text{tor}} = (\overline{\Gamma_0})_{\text{tor}} \xrightarrow{\sim} (\Gamma_0)_{\text{tor}} = \Gamma_{\text{tor}},$$

which concludes the proof of Proposition 4.8. \square

The following result is an immediate consequence of our proof of Proposition 4.8.

Corollary 4.11. *For each nonzero $a \in R$, we have $\Gamma/\Phi_a(\Gamma) \xrightarrow{\sim} \bar{\Gamma}/\Phi_a(\bar{\Gamma})$.*

Proof. The claimed result follows immediately from (4.9) and (4.10) using that $\hat{R}/a\hat{R} \xrightarrow{\sim} R/aR$. \square

Now we are ready to prove Theorem 2.8 when $\dim(V) = 0$.

Proposition 4.12. *Theorem 2.8 holds if V is a zero-dimensional subvariety of \mathbb{G}_a^g .*

Proof. Clearly, it suffices to show that $V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq V(K) \cap \Gamma$ since the reverse inclusion is obvious.

First we show that it suffices to prove our result when we replace K by a finite extension L , and replace Γ by its full divisible hull Γ_L inside L_{Φ}^g . Indeed, we denote by \mathbf{A}_L the adèles for the function field L (with respect to the places which extend the places contained in Ω_1), and we assume that

$$V(\mathbf{A}_L) \cap \bar{\Gamma}_L \subseteq V(L) \cap \Gamma_L.$$

Then

$$V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq V(\mathbf{A}_L) \cap \bar{\Gamma}_L \subseteq V(L) \cap \Gamma_L \subseteq V(L).$$

Therefore $V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq V(L) \cap V(\mathbf{A}_K) = V(K)$ because $L \cap \mathbf{A}_K = K$ by [9, Lemma 3.2]. So, $V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq K^g \cap \bar{\Gamma} = \Gamma$ by Proposition 4.3. Therefore

$$V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq V(K) \cap \Gamma, \text{ as desired.}$$

Secondly, we show that for any finite extension L of K , the hypothesis that $\Phi[t^\infty](L^{\text{sep}})$ is finite holds. Indeed, this claim is obvious when L is a separable extension of K ; so it suffices to prove it when L is a purely inseparable extension of K . Hence, assume $L = K^{1/p^r}$ for some positive integer r . Since Φ_t is inseparable, we obtain that $\Phi_{t^r}(L^{\text{sep}}) \subseteq K^{\text{sep}}$. Because $\Phi[t^\infty](K^{\text{sep}})$ is finite, we know that there exists a positive integer m such that $\Phi[t^\infty](K^{\text{sep}}) \subseteq \Phi[t^m]$. Therefore, $\Phi[t^\infty](L^{\text{sep}}) \subseteq \Phi[t^{r+m}]$, as desired.

Now, since we proved that we may replace K by a finite extension (and replace Γ by its full divisible hull inside this extension), we may assume that $V(\bar{K}) = V(K)$ (recall our hypothesis is that $\dim(V) = 0$). Let $\{x_n\}_{n \in \mathbb{N}} \subseteq \Gamma$ be a sequence which converges adèlically to a point $(y_v)_{v \in \Omega_1} \in V(\mathbf{A}_K)$, where $y_v \in V(K_v)$ for all places v . Actually, $y_v \in V(K)$ since V is a zero dimensional variety whose algebraic points are all defined over K . So, we know that for all places $v \in \Omega_1$, we have

$$x_n \rightarrow y_v \text{ in the } v\text{-adic topology.}$$

We claim that $y_v - y_w \in \Gamma_{\text{tor}}$ for any given two places v and w from Ω_1 . Since Γ is full in K , and since $y_v, y_w \in K^g$, all we need to show is that $y_v - y_w$ is a torsion point for Φ . By Proposition 4.2, we conclude that for each $j \in \mathbb{N}$, there exists $n_j \in \mathbb{N}$ and there exist $z_{v,j}, z_{w,j} \in K^g$ such that

$$x_{n_j} - y_v = \Phi_{t^j}(z_{v,j}) \text{ and } x_{n_j} - y_w = \Phi_{t^j}(z_{w,j}).$$

So, $y_v - y_w = \Phi_{t^j}(z_{w,j} - z_{v,j}) \in \Phi_{t^j}(K^g)$. Since we can do this for *all* $j \in \mathbb{N}$, and K_{Φ}^g is a direct product of a finite torsion submodule with a free submodule of rank \aleph_0 , we conclude that $y_v - y_w \in \Phi_{\text{tor}}(K)^g = \Gamma_{\text{tor}}$, as claimed.

Because Γ_{tor} is a finite set, and because $y_v - y_w \in \Gamma_{\text{tor}}$ for each pair $(v, w) \in \Omega_1 \times \Omega_1$, we conclude that there exists $y \in V(K)$ and there exist infinitely many places $v \in \Omega_1$ such that $y = y_v$. Then, as shown above, the sequence $\{x_n - y\}_{n \in \mathbb{N}}$ converges to an adèlic torsion point for Φ . By Proposition 4.8 applied to the full divisible hull in K of the $\Phi(R)$ -submodule generated by Γ and y , we conclude that $\{x_n - y\}_{n \in \mathbb{N}}$ converges adèlically to a torsion point $z_0 \in \Phi_{\text{tor}}(K)^g = \Gamma_{\text{tor}}$. However, since the reduction of z_0 modulo v equals 0 for infinitely many places v , we conclude that $z_0 = 0$. Therefore, the sequence $\{x_n\}_{n \in \mathbb{N}}$ converges adèlically to the point $y \in V(K)$. Using Proposition 4.3, we obtain that $y \in \Gamma$, which yields the desired conclusion: $V(\mathbf{A}_K) \cap \bar{\Gamma} = V(K) \cap \Gamma$. \square

5. A UNIFORM MORDELL-LANG VERSION FOR DRINFELD MODULES

In this Section we continue with the same notation as in Section 4. Our main goal is to prove the following result, which follows from the unpublished notes of the second author.

Proposition 5.1. *Let K be a function field over \mathbb{F} of arbitrary finite transcendence degree, and let Φ be a Drinfeld module of special characteristic defined over K . Assume Φ has positive modular transcendence degree, and that $\Phi[t^\infty](K^{\text{sep}})$ is finite. Let Γ be a full, finitely generated $\Phi(R)$ -submodule of K_{Φ}^g . Let $V \subseteq \mathbb{G}_a^g$ be an affine subvariety defined over K , and assume that V contains no translate of a positive dimensional algebraic subgroup of \mathbb{G}_a^g .*

Then there exists a zero-dimensional subvariety $W \subseteq V$ defined over K such that $V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq W(\mathbf{A}_K)$.

Proposition 5.1 allows us to reduce Theorem 2.8 to the case $\dim(V) = 0$, which was already proven in Proposition 4.12.

In order to prove Proposition 5.1 we shall require a uniformity result used by Hrushovski in his proof [5] of the positive characteristic function field Mordell-Lang conjecture but adapted here for Drinfeld modules. Our results are valid for any inseparable endomorphism ψ of \mathbb{G}_a defined over K ; for our application, we have $\psi = \Phi_t$. For each $n \in \mathbb{N}$ we denote by ψ^n the n -th iterate of ψ when composed with itself, and we extend the action of ψ on \mathbb{G}_a^g for any $g \in \mathbb{N}$.

First we let \mathcal{B} be a p -basis for K , i.e. a basis for K as K^p -vector space. We say that an extension of fields L/K is *fully separable* if \mathcal{B} continues to be a p -basis for L . With this definition, a subextension of a fully separable extension need not be fully separable itself.

For L/K a fully separable, separably closed extension field of K , define $\psi^\infty(L) := \bigcap_{n \geq 0} \psi^n(\mathbb{G}_a(L)) \subseteq \mathbb{G}_a(L)$.

Proposition 5.2. *Given an algebraic subvariety $X \subseteq \mathbb{G}_a^g$ having the property that X does not contain the translate of any positive dimensional algebraic subgroup of \mathbb{G}_a^g , there are numbers ℓ and m such that for any fully separable field extension L/K and point $a \in \mathbb{G}_a^g(L)$ the cardinality of $(a + X(L)) \cap \psi^m(\mathbb{G}_a^g(L))$ is at most ℓ .*

The proof of this proposition relies on the main dichotomy theorem on minimal groups type definable in separably closed fields, though in this particular case the Pillay-Ziegler [7] geometric argument using jet spaces may be employed rather than the Zariski geometries argument needed in general. To apply the Pillay-Ziegler argument following their proof of the positive characteristic Mordell-Lang conjecture for ordinary abelian varieties, we need to know that the generic type of ψ^∞ is *very thin* in their language. That is, if L/K is a fully separable extension of K and $a \in \psi^\infty(L)$ then the smallest fully separable subfield of L containing a and K is finitely generated as a field over K . The argument of [7, Lemma 6.4] in which it is shown that if A is ordinary then $p^\infty A$ is very thin applies to ψ^∞ . That is, we may write $\psi = \tilde{\psi} \circ F$ where F is a Frobenius map corresponding to some power of p and $\tilde{\psi}$ is separable. Following the proof of [7, Lemma 6.4] replacing the Verschiebung by $\tilde{\psi}$, one sees that $K(a)$ is already fully separable.

With this preamble, let us state precisely the required result and then explain how to deduce Proposition 5.1 from the main theorem.

Theorem 5.3. *With our hypotheses on ψ , the type definable group ψ^∞ is modular. In particular, for any separably closed, fully separable extension field L/K , and any algebraic variety $Y \subseteq \mathbb{G}_a^g$ defined over L , the set $Y(L) \cap \psi^\infty(L)^g$ is a finite union of translates of subgroups (possibly trivial) of $\psi^\infty(L)^g$.*

The proof of Theorem 5.3 follows identically as the proof of the main theorem of [5] and is sketched by the second author in the Arizona Winter School notes [10]. As noted above, in this case one could also give a geometric proof using a variant of the Gauss map.

Given Theorem 5.3, let us show how to deduce Proposition 5.2. This argument also appears in [5].

Proof of Proposition 5.2. If no such bounds ℓ and m existed, then the following set of sentences would be consistent:

- L is a separably closed, fully separable extension of K . (This is expressible by a countable set of sentences in the language of rings augmented by constant symbols for the elements of K .)
- $a \in \mathbb{G}_a^g(L)$
- $c_i \in \mathbb{G}_a^g(L)$ (for $i \in \mathbb{N}$)
- $c_i \neq c_j$ for $i \neq j$
- $(\exists y \in \mathbb{G}_a^g(L)) \psi^m(y) = c_i$ for each $i \in \mathbb{N}$ and $m \in \mathbb{N}$
- $c_i \in a + X(L)$

By the compactness theorem, we obtain a fully separable, separably closed field extension L/K and a point $a \in \mathbb{G}_a^g(L)$ so that $(a + X(L)) \cap \psi^\infty(L)^g$ is infinite. By the modularity of $\psi^\infty(L)$ it follows that there is an infinite group $H \leq \psi^\infty(L)^g$ and a point $b \in \psi^\infty(L)^g$ such that $b + H \subseteq a + X(L)$. But then X contains the translate of the Zariski closure of H by $(b - a)$ contrary to our hypothesis on X . \square

Now we are ready to prove Proposition 5.1.

Proof of Proposition 5.1. Let m and ℓ be the numbers given by Proposition 5.2 for $X = V$ and $\psi = \Phi_t$. By Corollary 4.11, the group $\bar{\Gamma}/\Phi_{t^m}\bar{\Gamma}$ is finite and each coset of $\Phi_{t^m}\bar{\Gamma}$ contains an element of Γ . Let $\gamma_1, \dots, \gamma_s \in \Gamma$ so that $\bar{\Gamma} = \bigcup_{i=1}^s \gamma_i + \Phi_{t^m}\bar{\Gamma}$. By Proposition 5.2, for each i there is a zero dimensional variety W_i defined over K for which $V(L) \cap (\gamma_i + \Phi_{t^m}\mathbb{G}_a^g(L)) \subseteq W_i(L)$ for any fully separable extension field L/K ; clearly we may assume $W_i \subset V$. Let

$$W := \bigcup_{i=1}^s W_i.$$

Clearly each K_v is a fully separable extension of K . So, if $b = (b_v)_{v \in \Omega_1} \in V(\mathbf{A}_K) \cap \bar{\Gamma}$, we have $b_v \in V(K_v) \cap (\bigcup_{i=1}^s \gamma_i + (\Phi_{t^m}\mathbb{G}_a^g(K_v))) \subseteq W(K_v)$. Hence, $b \in W(\mathbf{A}_K)$ as desired. \square

Now we are ready to prove Theorem 2.8 for any subvariety V which contains no translate of a positive dimensional algebraic subgroup of \mathbb{G}_a^g .

Proof of Theorem 2.8. By Proposition 5.1, there exists a zero-dimensional affine subvariety W of V also defined over K such that $V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq W(\mathbf{A}_K)$. On the other hand, by Proposition 4.12, we have that $W(\mathbf{A}_K) \cap \bar{\Gamma} = W(K) \cap \Gamma$. So, $V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq W(\mathbf{A}_K) \cap \bar{\Gamma} = W(K) \cap \Gamma$. On the other hand, since $W \subseteq V$, we conclude that $V(\mathbf{A}_K) \cap \bar{\Gamma} \subseteq V(K) \cap \Gamma$. But always $V(K) \cap \Gamma \subseteq V(\mathbf{A}_K) \cap \bar{\Gamma}$, which concludes the proof of Theorem 2.8. \square

REFERENCES

- [1] A. J. de Jong, *Smoothness, semi-stability and alterations*, Inst. Hautes Études Sci. Publ. Math **83** (1996), 51–93.
- [2] L. Denis, *Géométrie diophantienne sur les modules de Drinfeld*, in: The arithmetic of function fields (Columbus, OH, 1991) 285–302, de Gruyter, Berlin, 1992
- [3] D. Ghioca, *The Lehmer inequality and the Mordell-Weil theorem for Drinfeld modules*, J. Number Theory **122** (2007), 37–68.
- [4] D. Ghioca and T. J. Tucker, *Equidistribution and integrality for Drinfeld modules*, Tran. Amer. Math. Soc. **350** (2008), 4863–4887.
- [5] E. Hrushovski, *The Mordell-Lang conjecture for function fields*, J. Amer. Math. Soc. **9** (1996), 667–690.
- [6] S. Lang, *Fundamentals of Diophantine Geometry*, Springer-Verlag, New York, 1983.
- [7] A. Pillay and M. Ziegler, *Jet spaces of varieties over differential and difference fields*, Selecta Math. (N.S.) **9** (2003), 579–599.
- [8] B. Poonen, *Local height functions and the Mordell-Weil theorem for Drinfeld modules*, Compositio Math. **97** (1995), 349–368.

- [9] B. Poonen and J. F. Voloch, *The Brauer-Manin obstruction for subvarieties of abelian varieties over function fields*, Ann. Math. **171** (2010), 511–532.
- [10] T. Scanlon, *Model theory and diophantine geometry*, Arizona Winter School, 2003, <http://math.arizona.edu/~swc/oldaws/03Notes.html>
- [11] J. P. Serre, *Lectures on the Mordell-Weil theorem*, Translated from the French and edited by Martin Brown from notes by Michel Waldschmidt. Aspects of Mathematics, E15. friedr. Vieweg & Sohn, Braunschweig, 1989.
- [12] C.-L. Sun, *Adèlic points of subvarieties of isotrivial semi-abelian varieties over a global field of positive characteristic*, preprint, [arXiv:1005.4998](https://arxiv.org/abs/1005.4998).
- [13] J. T.-Y. Wang, *The Mordell-Weil theorems for Drinfeld modules over finitely generated function fields*, Manuscripta Math. **106** (2001), 305–314.

DRAGOS GHIOCA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA,
1984 MATHEMATICS ROAD, VANCOUVER, BC V6T 1Z2, CANADA
E-mail address: dghioca@math.ubc.ca

THOMAS SCANLON, DEPARTMENT OF MATHEMATICS, EVANS HALL, UNIVERSITY OF
CALIFORNIA, BERKELEY, CA 94720-3840, USA
E-mail address: scanlon@math.berkeley.edu