

KRONECKER WEBS, BIHAMILTONIAN STRUCTURES, AND THE METHOD OF ARGUMENT TRANSLATION

ILYA ZAKHAREVICH

ABSTRACT. We show that manifolds which parameterize values of first integrals of integrable finite-dimensional bihamiltonian systems carry a geometric structure which we call a *Kronecker web*. We describe two functors between Kronecker webs and integrable bihamiltonian structures, one is left inverse to another one. Conjecturally, these two functors are mutually inverse (for “small” open subsets).

The above conjecture is proven provided the bihamiltonian structure allows an antiinvolution of a particular form. This implies the conjecture of [10] that on a dense open subset the bihamiltonian structure on \mathfrak{g}^* for a semisimple \mathfrak{g} is flat.

CONTENTS

0. Basic notions	2
1. Introduction	5
2. Linear relations and pencils	9
3. Kronecker relations	10
4. Kronecker webs	12
5. Pairs of skewsymmetric forms	18
6. Micro-Kronecker bihamiltonian structures	19
7. Lattice of kernels	22
8. Lagrangian foliations	24
9. Two functors	26
10. Conjecture on classification	28
11. Antiinvolutions	29
12. Real and complex bihamiltonian structures	32
13. Endomorphisms of bihamiltonian structures and the principal theorem	33
14. Method of argument translation	35
15. Appendix on Kronecker decompositions	41
References	46

0. BASIC NOTIONS

We postpone discussion of what is done in this paper until Section 1, starting with introduction of notations and conventions used throughout this text.

Many results of this paper may be stated in more generality, but for simplicity we assume that all the vector spaces we consider here are finite-dimensional¹ vector spaces over a field \mathbb{K} which is either \mathbb{R} or \mathbb{C} . A *manifold* is a C^∞ -manifold or real-analytic manifold in the case $\mathbb{K} = \mathbb{R}$, and an analytic manifold in the case $\mathbb{K} = \mathbb{C}$. Use the word *smooth* to mean C^∞ -smooth, real-analytic, or complex-analytic correspondingly.

For a vector space V over \mathbb{K} denote by V^* the space of \mathbb{K} -linear functionals on V . (Note that throughout this paper we do *not* consider semilinear functional or Hermitian forms on complex vector spaces.)

Recall basic notions and notations of Poisson geometry (see [2, 15, 3]). In what follows if f is a function or a tensor field on M , $f|_m$ denotes the value of f at $m \in M$.

Remark 0.1. Throughout the paper the phrase “*at generic points*” means “at points of an appropriate open dense subset”. Similarly, a “*small open subset*” is used instead of “an appropriate neighborhood of any given point”. A *local isomorphism* between two geometric structures on M and M' is an isomorphism of a neighborhood of any point on M with a neighborhood of a given point on M' .

Definition 0.2. A *bracket* on a manifold M is a \mathbb{K} -bilinear skewsymmetric mapping $f, g \mapsto \{f, g\}$ from pairs of smooth functions² on M to smooth functions on M . This mapping should satisfy the Leibniz identity $\{f, gh\} = g\{f, h\} + h\{f, g\}$. A bracket is *Poisson* if it satisfies Jacobi identity too (thus defines a structure of a Lie algebra on functions on M).

A *Poisson structure* is a manifold M equipped with a Poisson bracket.

Remark 0.3. Leibniz identity implies $\{f, g\}|_m = 0$ if f has a zero of second order at $m \in M$. Thus a bracket is uniquely determined by describing functions $\{f_i, f_j\}$, here $\{f_i\}_{i \in I}$ is an arbitrary collection of smooth functions on M such that for any $m \in M$ the collection $\{df_i|_m\}_{i \in I}$ of vectors in \mathcal{T}_m^*M spans \mathcal{T}_m^*M as a vector space.

Definition 0.4. Consider a bracket $\{, \}$ on a manifold M . The *associated bivector*³ *field* η is the section of $\Lambda^2\mathcal{T}M$ given by $\{f, g\}|_m = \langle \eta|_m, df \wedge dg|_m \rangle$, $m \in M$, here \langle, \rangle denotes the canonical pairing between $\Lambda^2\mathcal{T}_mM$ and Ω_m^2M .

Given $m_0 \in M$, the *associated pairing* $(,)$ in $\mathcal{T}_{m_0}^*M$ is defined as $(\alpha, \beta) = \{f, g\}|_{m_0}$ if $\alpha = df|_{m_0}$, $\beta = dg|_{m_0}$.

Obviously, the associated bivector field uniquely determines the bracket and visa versa. The associated pairing is a skewsymmetric bilinear pairing.

¹With obvious exceptions of vector spaces of functions on manifolds.

²In the complex analytic case one should consider functions on open subsets $U \subset M$ and require that the brackets on these subsets are compatible on intersections.

³A *bivector field* is a skewsymmetric contravariant tensor of valence 2.

Definition 0.5. Given a skewsymmetric bilinear form $(,)$ in a vector space V , let $\text{Ker}(,) \stackrel{\text{def}}{=} \{v \in V \mid (v, v') = 0 \forall v' \in V\}$, call $\dim \text{Ker}(,)$ the *corank* of $(,)$. The *rank* of $(,)$ is $\dim V - \dim \text{Ker}(,)$.

Say that the *rank* of the bracket $\{, \}$ at $m \in M$ is r if the associated skewsymmetric bilinear pairing on \mathcal{T}_m^*M has rank r . In this case the *corank* of the bracket is $\dim M - r$.

A bracket has a *constant (co)rank* if its rank does not depend on the point $m \in M$. A Poisson bracket is *symplectic* if the corank is constant and equal to 0.

The associated tensor field η of a bracket on M can be considered as a mapping $H: \mathcal{T}^*M \rightarrow \mathcal{T}M$ (the *Hamiltonian mapping* of a bracket). If the bracket is symplectic, this mapping is invertible, and the inverse mapping $H^{-1}: \mathcal{T}M \rightarrow \mathcal{T}^*M$ can be considered as bilinear pairing on $\mathcal{T}M$, or a tensor field ω which is a section of Ω^2M of corank 0. Call ω the *symplectic 2-form* of the symplectic bracket, in local coordinates tensors η and ω are given by mutually inverse matrices.

Inversely, given a section ω of Ω^2M of corank 0, putting $\eta = \omega^{-1}$ gives a bracket on M . It is easy to check that η is Poisson iff⁴ $d\omega = 0$.

Example 0.6. Given a manifold N , let $M = \mathcal{T}^*N$, and $\pi: M \rightarrow N$ be the natural projection. Let $m \in M$, $m = (n, \nu)$, $n \in N$, $\nu \in \mathcal{T}_n^*N$. Consider $\pi^*: \mathcal{T}_n^*N \rightarrow \mathcal{T}_m^*M$, then $\pi^*\nu$ is an element of \mathcal{T}_m^*M which depends on m only. Denote this element $\alpha(m)$, α is a canonically defined section of Ω^1M .

Local coordinates (n_1, \dots, n_d) on N define local coordinates $(n_1, \dots, n_d, \nu_1, \dots, \nu_d)$ on \mathcal{T}^*N , and $\alpha = \sum_i \nu_i dn_i$. Take $\omega = d\alpha$. In local coordinates $\omega = \sum_i d\nu_i \wedge dn_i$, hence ω is of corank 0, thus defines a (symplectic) Poisson structure η on $M = \mathcal{T}^*N$.

Recall that any symplectic Poisson structure is locally isomorphic to the structure of Example 0.6.

Definition 0.7. Call two Poisson brackets $\{, \}_1$ and $\{, \}_2$ on M *compatible* if the bracket $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ is Poisson for any λ_1, λ_2 .

A *bihamiltonian structure* is a manifold M with a pair of compatible Poisson brackets.

In fact it is possible to show that if *one* linear combination $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ of two Poisson brackets is Poisson, and $\lambda_1 \neq 0$, $\lambda_2 \neq 0$, then *any* linear combination $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ is Poisson. Even if M is a C^∞ -manifold, the coefficients λ_1, λ_2 may be taken to be complex numbers. Indeed, if M is a C^∞ -manifold with a bracket, one may consider the extension of the bracket to the \mathbb{C} -vector space of complex-valued functions on M . In this case $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ is well-defined even for complex values of λ_1, λ_2 . By the above remarks, complex linear combinations of brackets of

⁴This condition is linear in ω , as opposed to the quadratic condition (of Jacobi identity) on η . This linearity makes it much easier to study symplectic Poisson brackets.

a bihamiltonian structure are also Poisson. In what follows we can always consider brackets as acting on the spaces of complex-valued functions.

Given a pair of brackets $\{, \}_1$ and $\{, \}_2$, one obtains two bivector fields η_1, η_2 . Analogously, one obtains two skewsymmetric bilinear pairings $(,)_1, (,)_2$ on \mathcal{T}_m^*M , so that $(\alpha, \beta)_a = \{f, g\}_a|_m$ if $\alpha = df|_m, \beta = dg|_m, a = 1, 2$.

Definition 0.8. Say that the Poisson structure $\{, \}$ on a vector space V is *translation-invariant* if for any parallel translation $\mathfrak{T}: V \rightarrow V$ and any two functions f, g on V one has $\mathfrak{T}^* \{f, g\} = \{\mathfrak{T}^*f, \mathfrak{T}^*g\}$. Bihamiltonian structure on V is *translation-invariant* if both Poisson brackets are translation-invariant.

Say that a bihamiltonian structure on a manifold M is *flat* if it is *locally isomorphic* to a translation-invariant bihamiltonian structure. In other words, for any $m \in M$ there is a neighborhood $U \ni m$ such that the restriction of the bihamiltonian structure to U is isomorphic to a restriction of an appropriate translation-invariant bihamiltonian structure to an appropriate open subset.

The tensor field η of a translation-invariant Poisson bracket on V has constant coefficients in any vector-space coordinate system on V . Consider an interesting example of a translation-invariant bihamiltonian structure.

Example 0.9. Consider a vector space V with coordinates x_0, \dots, x_{2k-2} and the Poisson brackets of coordinates

$$(0.1) \quad \{x_{2l}, x_{2l+1}\}_1 = 1, \quad \{x_{2l+1}, x_{2l+2}\}_2 = 1, \quad 0 \leq l \leq k-2,$$

any other brackets of coordinate functions x_0, \dots, x_{2k-2} vanishing. This pair of brackets is in fact a translation-invariant bihamiltonian structure.

The following example is the simplest of classical examples of bihamiltonian structures arising in theory of integrable systems.

Example 0.10. Given a Lie algebra \mathfrak{g} and an element $c_1 \in \mathfrak{g}^*$, define a bihamiltonian structure on \mathfrak{g}^* as in [23, 4]. An element $X \in \mathfrak{g}$ defines a linear function f_X on \mathfrak{g}^* . Due to Remark 0.3, to define a bihamiltonian structure on \mathfrak{g}^* it is enough to describe brackets $\{f_X, f_Y\}_a, a = 1, 2, X, Y \in \mathfrak{g}$.

Let $\{f_X, f_Y\}_1$ be a constant function on \mathfrak{g}^* and $\{f_X, f_Y\}_2$ be a linear function on \mathfrak{g}^* given by the formulae

$$\{f_X, f_Y\}_1 = f_{[X, Y]}(c_1), \quad \{f_X, f_Y\}_2 = f_{[X, Y]}.$$

The bracket $\{, \}_2$ is the natural Lie–Kirillov–Kostant–Souriau Poisson bracket on \mathfrak{g}^* . The bracket $\{, \}_1$ is translation-invariant. The bracket $\{, \}_2$ is translation-invariant only if \mathfrak{g} is abelian (and then $\{, \}_{1,2}$ vanish).

In fact instead of taking $c_1 \in \mathfrak{g}^*$ one can consider any 2-cocycle $c_2 \in \Lambda^2 \mathfrak{g}^*$, and define $\{f_X, f_Y\}_1 = c_2(X, Y)$. The above definition is reconstructed if one puts $c_2 = \partial c_1$.

Definition 0.11. Say that a smooth function F on a manifold M with a Poisson bracket $\{, \}$ is a *Casimir* function, if $\{F, f\} = 0$ for any smooth function f on M .

Obviously, any function $\varphi(F_1, F_2, \dots, F_k)$ of several Casimir functions is again Casimir.

Definition 0.12. A collection of smooth functions F_1, \dots, F_r on M is *dependent* if $\varphi(F_1, \dots, F_r) \equiv 0$ for an appropriate smooth function $\varphi \neq 0$.

Remark 0.13. Consider a manifold M with a Poisson bracket $\{, \}$. The local classification of Poisson structures of *constant rank* [16, 33] shows that for an arbitrary Poisson bracket there is an open (and in interesting cases dense) subset $U \subset M$ and $k \in \mathbb{Z}_{\geq 0}$ such that on U there are k independent Casimir functions F_1, \dots, F_k , and any Casimir function on U may be written as a function of F_1, \dots, F_k (we do not exclude the case $k = 0$). The common level sets $F_1 = C_1, \dots, F_k = C_k$ form an invariantly defined foliation on U , which is called the *symplectic foliation*. Moreover, one can construct additional functions $n_i, \nu_i, i = 1, \dots, d$, on U such that $\{n_i, n_j\} = \{\nu_i, \nu_j\} = 0$, $\{n_i, \nu_j\} = \delta_{ij}$, and the functions $F_\bullet, n_\bullet, \nu_\bullet$ form a coordinate system on U .

This shows that any Poisson structure of constant rank is *flat*, i.e., locally isomorphic to a translation-invariant Poisson structure. Say, this implies that any analytic Poisson structure is flat on an open dense subset. Moreover, the leaves of the symplectic foliation of such Poisson structures can be simultaneously equipped with coordinates as in Example 0.6.

Definition 0.14. Consider a foliation \mathcal{F} on B . Define the *tangent bundle* \mathcal{TF} to \mathcal{F} as a vector subbundle E of \mathcal{TB} such that E_b coincides with $\mathcal{T}_b L_b$ for any $b \in B$, here L_b is the leaf of \mathcal{F} which passes through b . Let the *normal bundle* $\mathcal{NF} \subset \mathcal{T}^*B$ to \mathcal{F} be the orthogonal complement to \mathcal{TF} , and the *cotangent bundle* $\mathcal{T}^*\mathcal{F}$ to \mathcal{F} be the dual bundle to \mathcal{TF} .

The total space of the bundle $\mathcal{T}^*\mathcal{F}$ is a union of total spaces of cotangent bundles \mathcal{T}^*L of the leaves L of the foliation, and each \mathcal{T}^*L carries a natural symplectic Poisson structure (Example 0.6). Hence

Proposition 0.15. *The total space of the cotangent bundle to a foliation carries a natural Poisson structure of constant rank.*

Moreover, the above foliation on $\mathcal{T}^*\mathcal{F}$ is the symplectic foliation of this Poisson structure. Due to Remark 0.13, any Poisson structure of constant rank is locally isomorphic to the Poisson structure on $\mathcal{T}^*\mathcal{F}$ for an appropriate foliation \mathcal{F} .

1. INTRODUCTION

Among other approaches to integrable system the so-called *bihamiltonian approach* is especially interesting from the geometric point of view. In this approach all the

properties of an integrable bihamiltonian system are deduced basing on the bihamiltonian structure on the phase manifold⁵. (The principal tool for this deduction is Lennard scheme, see [18, 19, 9, 7, 17, 10].) Since the structure of a bihamiltonian system is nothing more than a pair of tensor fields satisfying some invariantly-defined conditions, this approach puts the integrable system into the standard framework of differential geometry.

One of the most powerful approaches of differential geometry is classification of objects up to isomorphism, and description of automorphisms of a given object. Say, in the case of symplectic structures or Poisson structures of constant rank the local classification is “trivial”: locally there are some discrete parameters only, thus any structure is locally isomorphic to one from a finite list (for a given dimension). This reduces all the question on geometry of symplectic manifolds to questions of global nature, and all the question on geometry of Poisson manifolds to questions of global nature, and questions related to subsets where the rank drops.

There are many results on classifications of bihamiltonian structures. The case of general position on even-dimensional manifolds was done in [30, 20, 21, 22, 13] (in different assumptions), while the case of general position in odd-dimensional case was analyzed in [12, 13, 27, 26, 28]. However, until recently these results had little direct impact in the theory of integrable systems, since it was not known which integrable systems are subject to these conditions of general position.

On the other hand, the analysis of [11, 14] had shown that the periodic KdV system should be considered as an infinite-dimensional analogue of an odd-dimensional bihamiltonian structure in general position. Up to some extend Toda lattice is a finite-dimensional analogue of the KdV system, so one should have expected that the periodic Toda lattice might have similar properties. Unfortunately, the periodic Toda lattice is an even-dimensional bihamiltonian system, thus the direct analogy breaks down.

However, an open Toda lattice *is* an odd-dimensional bihamiltonian system, and in [10] it was shown that the open Toda lattice is an odd-dimensional bihamiltonian structure in general position. Moreover, in [10] it was also shown that the even-dimensional manifold of the periodic Toda lattice has a foliation of codimension 1 such that the bihamiltonian structure is trivial across the foliation, while generic leaves of the foliation carry an odd-dimensional bihamiltonian structure in general position.

These results show that Toda lattices are subject to the local classification of bihamiltonian systems of [12, 13]. Let us stress out that there are infinitely many non-isomorphic bihamiltonian systems of these types (with parameters being several functions of two variables), thus it is meaningful to ask *which* system of the above classification is the open Toda lattice (or the leaves of the periodic Toda lattice).

⁵There is a widespread belief that most (or all) integrable systems which arise in problems of mathematical physics allow a natural bihamiltonian structure.

Recall that the simplest possible odd-dimensional bihamiltonian structure (in general position) is the structure of Example 0.9. One of the principal results of [10] is that the bihamiltonian structure of the open Toda lattice *is locally isomorphic* (on an open dense subset) to the structure of Example 0.9 (here k is the number of “atoms” in the Toda lattice). Moreover, the bihamiltonian structure of the periodic Toda lattice is locally isomorphic (on an open dense subset) to a direct product of two copies of the structure of Example 0.9. (If periodic Toda lattice has n atoms, then one copy has $k = n$, another $k = 1$.)

Extending this observation, one of conjectures of [10] says that other integrable systems of mathematical physics are also locally isomorphic to a direct product of several copies of Example 0.9. In other words, these systems are flat on an open dense subsets. Note similarities and differences of this (meta)conjecture with the result on flatness on an open dense subset of analytic Poisson manifolds. As in the Poisson case, this (meta)conjecture reduces questions on geometry of these integrable systems to two questions: the description of the behaviour in the points outside the above dense opens subsets (points where the bihamiltonian structure *degenerates*), and the description of the global structure. The question of local geometry in generic points mostly disappear: the system is locally isomorphic to one from a finite list⁶ (for a fixed dimension).

As we stressed it out before, there is *no general result* on local triviality of bihamiltonian geometry. The above metaconjecture is a *selection principle*⁷: out of a huge variety of different integrable bihamiltonian systems the systems studied in mathematical physics fall into a very thin subclass of *flat bihamiltonian systems*.

The paper [10] also contains more concrete particular cases of the above metaconjecture: it is conjectured that some particular bihamiltonian systems of mathematical physics are also flat on an open dense subset. In this list are the complete Toda lattice, the multi-dimensional Euler top, and semisimple case of Example 0.10.

In addition to the results of [10] on the geometry of Toda lattices, in this paper we prove the above conjecture in the semisimple case of Example 0.10 (see Corollary 14.24). This describes several examples of “classical” integrable bihamiltonian systems which locally look like a product of structures of Example 0.9. On the other hand, [4] formalized the notion of *integrability* of a bihamiltonian system⁸, and all the above examples (as most of other bihamiltonian systems of mathematical physics) happen to be integrable in this strict sense. (Below we always use the word *integrable* meaning the definition of [4].)

⁶Say, due to the above result they are locally isomorphic to direct product of several open Toda lattices.

⁷Say, any proof of this metaconjecture (if possible) would need to concentrate on the question *why* mathematical physics study some systems and do not study some other systems.

⁸This notion is very close to the property of being *micro-Kronecker* we introduce below: integrable (or *complete*) bihamiltonian system is one which is micro-Kronecker on an open dense subset.

This makes it very important to investigate the local geometry of arbitrary integrable bihamiltonian systems. This investigation is one of the principal target of this paper.

The odd-dimensional bihamiltonian structures in general position are a particular case of integrable bihamiltonian structures. One of principal results of [12] says that such structures are locally classified by a structure of a so-called *Veronese web* on a manifold of (approximately) half the dimension of the initial manifold.

In [25] (and independently—and more general, but in less details—in [10]) the construction of the Veronese web was generalized to the case of more general structures integrable in the sense of [4]. This paper starts with introduction of a geometric structure of a *Kronecker web*, which is simultaneously a generalization of the construction of *Veronese web of higher codimension* of [25] and a more structured variant of the construction of a *web* of [10]. Each Kronecker web has a *rank*, and Veronese webs of [12] coincide with Kronecker webs of rank 1.

Similar to what was done in [12] in the case of rank 1, we show how to associate to any integrable bihamiltonian structures its Kronecker web, and show how to construct an integrable bihamiltonian structure basing on an arbitrary Kronecker web. Conjecturally (Conjecture 10.1), these functors are mutually inverse (as in the case of rank 1, see [12, 31]), but here we prove only that one is left inverse to another.

Provided these functors are mutually inverse, this reduces the question of local classification of integrable bihamiltonian structures to the question of local classification of Kronecker webs. As in Definition 0.8, one can define what is a *translation-invariant* Kronecker web, and a *flat* Kronecker web. In particular, to show that a given bihamiltonian structure is flat, it would be enough to show that its Kronecker web is flat⁹.

Say, in [25] it was shown that in semisimple case of Example 0.10 the corresponding Kronecker web is flat on an open dense subset. Together with our construction of two functors the above conjecture would show that the bihamiltonian structure of Example 0.9 is flat on an open dense subset. For another example, the (known) case of rank 1 of this conjecture is used in [10] to demonstrate flatness of Toda lattices.

Moreover, note that proofs of flatness of webs of particular bihamiltonian systems are very simple (see [25, 10], and Theorem 14.22). Thus the theory of Kronecker webs allows one to condense all the problems of geometric classification of bihamiltonian structures into a proof of Conjecture 10.1 (or a particular case of this conjecture).

Section 11 introduces a special subclass of bihamiltonian structures, structures which allow a special antiinvolutions. In Section 13 we show that in the case of such structures the Conjecture 10.1 holds (on a large open subset).

Section 14 uses this approach to show that the semisimple case of Example 0.10 is flat on an open dense subset. As already explained, the flatness of the Kronecker web is already proved in [25]. We use standard tools of the theory of semisimple Lie

⁹As [12, 13, 10] show, there are plenty of examples of non-flat Kronecker webs, even in the case of rank 1.

algebras to construct appropriate antiinvolutions, which provides an ad hoc way to prove this particular case of Conjecture 10.1.

We believe that the same trick will also work with complete Toda lattice and the multi-dimensional Euler top. However, we do not know whether this approach would work with the open and periodic Toda lattice. The proof of [10] used the established case of rank 1 of Conjecture 10.1.

Section 12 shows that the complex variant of Conjecture 10.1 implies the real-analytic case as well. The appendix describes which invariant structures one can associate to a Kronecker web.

With pleasure we thank V. Serganova for fruitful discussions and an important contribution: the proof of Proposition 14.19. This paper would not appear without long-term joint research with I. M. Gelfand, which molded the mindset of this paper. Special thanks go to A. Panasyuk for providing preprints of [24] and [25], and to B. Okun for patiently discussing results of [10], which lead to a cristallization of crucial ideas of the current paper.

2. LINEAR RELATIONS AND PENCILS

Definition 2.1. A *linear relation* W between vector spaces V_1 and V_2 is a vector subspace $W \subset V_1 \oplus V_2$. Say that W is *bisurjective* if projections of W to V_1 and V_2 are both surjective. The *left kernel* $\text{Ker}_L W$ of W is the kernel of projection $W \rightarrow V_2$ considered as a vector subspace of V_1 , define similarly the *right kernel* $\text{Ker}_R W \subset V_2$.

A *linear relation in* V is a linear relation between V and V .

Obviously, if W is a linear relation, then $W \supset \text{Ker}_L W \oplus \text{Ker}_R W$, thus W induces a vector subspace \widetilde{W} of $V_1/\text{Ker}_L W \oplus V_2/\text{Ker}_R W$. If W is bisurjective, then \widetilde{W} is a graph of a bijective linear mapping $V_1/\text{Ker}_L W \xrightarrow{\sim} V_2/\text{Ker}_R W$. In particular, $\dim V_1 - \dim V_2 = \dim \text{Ker}_L W - \dim \text{Ker}_R W$.

Definition 2.2. Fix once forever a two-dimensional vector space \mathcal{S} with a basis $\mathbf{s}_1, \mathbf{s}_2$. A *pencil* of linear operators between vector spaces V and W is a linear mapping $V \otimes \mathcal{S} \xrightarrow{\mathcal{P}} W$. This induces two linear mappings $\mathcal{P}_{1,2}: V \rightarrow W$ defined as $\mathcal{P}_i(v) \stackrel{\text{def}}{=} \mathcal{P}(v \otimes \mathbf{s}_i)$.

Given a pencil \mathcal{P} , one obtains a linear 2-parametric family of linear mappings $\lambda_1 \mathcal{P}_1 + \lambda_2 \mathcal{P}_2$ between V and W . Inversely, any two linear mappings $\mathcal{P}_{1,2}: V \rightarrow W$ correspond to a pencil and to a linear 2-parametric family of linear mappings.

Analyze the connections between pencils and linear relations in vector spaces. Given a linear relation W in V , two projections $\pi_{1,2}$ of $W \subset V \oplus V$ to V define a pencil π of operators $W \rightarrow V$. In the other direction, given any pencil $\mathcal{P}: W \rightarrow V$, one can construct a linear relation $\widetilde{W} \subset V \oplus V$ as the image of $\mathcal{P}_1 \oplus \mathcal{P}_2: W \rightarrow V \oplus V$. The pencils \mathcal{P} which may be obtained from linear relations are those for which $\text{Ker } \mathcal{P}_1 \cap \text{Ker } \mathcal{P}_2 = 0$. For such pencils two above constructions are mutually inverse.

Introduce *another*¹⁰ connection between bisurjective linear relations in V and pencils of linear operators $V \rightarrow V'$. Given such linear relation W in V , one obtains an identification $\alpha: V/\text{Ker}_R W \rightarrow V/\text{Ker}_L W$. Denote by $\pi_{1,2}$ the natural projections of V to $V/\text{Ker}_L W$ and to $V/\text{Ker}_R W$. Then $\mathcal{P}_1 \stackrel{\text{def}}{=} \pi_1$, $\mathcal{P}_2 = \alpha \circ \pi_2$ give a pencil of operators $V \rightarrow V/\text{Ker}_L W$. In the other direction, given a pencil \mathcal{P} of linear operators $V \rightarrow V'$, one can obtain a linear relation $W = \mathcal{P}_2^{-1}\mathcal{P}_1$ (in other words, $(v_1, v_2) \in W$ iff $\mathcal{P}_1 v_1 = \mathcal{P}_2 v_2$).

Obviously, a pencil \mathcal{P} of operators $V \rightarrow V'$ can be obtained from a bisurjective linear relation in V iff both linear operators $\mathcal{P}_1, \mathcal{P}_2$ of the pencil are surjective. Call such pencils *bisurjective*. In fact, one can obtain a much stronger result:

Definition 2.3. Given a linear relation W_1 in V_1 and a linear relation W_2 in V_2 , say that a linear mapping $\varphi: V_1 \rightarrow V_2$ is an *isomorphism* between W_1 and W_2 if φ is an isomorphism, and $(\varphi \oplus \varphi)(W_1) = W_2$.

Given a pencil \mathcal{P} of operators $V \rightarrow W$ and a pencil \mathcal{P}' of operators $V' \rightarrow W'$ say that linear mappings $\varphi: V \rightarrow V'$ and $\psi: W \rightarrow W'$ form a *morphism* between \mathcal{P} and \mathcal{P}' if $\mathcal{P}'_1 \varphi = \psi \mathcal{P}_1$, $\mathcal{P}'_2 \varphi = \psi \mathcal{P}_2$. Say that (φ, ψ) is an *isomorphism* if φ and ψ form a morphism and φ and ψ are isomorphisms of vector spaces.

Proposition 2.4. Consider a category $bs\mathfrak{R}$ of bisurjective linear relations in vector spaces with isomorphisms of relations as $\text{Mor } bs\mathfrak{R}$, and the category $bs\mathfrak{P}$ of bisurjective pencils of linear mappings with isomorphisms of pencils as $\text{Mor } bs\mathfrak{P}$. The defined above mappings $bs\mathfrak{R} \rightarrow bs\mathfrak{P}$ and $bs\mathfrak{P} \rightarrow bs\mathfrak{R}$ give an equivalence of these categories. Moreover, let Vect be the category of vector spaces, consider the functor $bs\mathfrak{R} \rightarrow \text{Vect}$ which sends a relation W in V to the vector space V , and the functor $bs\mathfrak{P} \rightarrow \text{Vect}$ which sends a pencil \mathcal{P} of operators $V \rightarrow W$ to V . The defined above equivalence of categories commutes with the mappings to Vect .

In plain words, it is “the same” to consider bisurjective linear relations in V and bisurjective pencils of operators $V \rightarrow V'$ up to isomorphisms of V' .

3. KRONECKER RELATIONS

Recall that to any bisurjective linear relation W in V we associated a pencil \mathcal{P} of operators $V \rightarrow V'$ (for an appropriate vector space V'). Thus one can consider the corresponding linear family $\lambda_1 \mathcal{P}_1 + \lambda_2 \mathcal{P}_2$ of mappings $V \rightarrow V'$. In particular, for any λ_1 and λ_2 one obtains a linear subspace $\text{Ker}(\lambda_1 \mathcal{P}_1 + \lambda_2 \mathcal{P}_2) \subset V$. Denote this subspace by $\text{Ker}_{\lambda_1, \lambda_2} W$. Obviously, for $(\lambda_1, \lambda_2) \neq (0, 0)$ this subspace depends only on the ratio $(\lambda_1 : \lambda_2) \in \mathbb{P}^1$, and $\text{Ker}_L = \text{Ker}_{1:0}$, $\text{Ker}_R = \text{Ker}_{0:1}$. Moreover, if the vector spaces we consider are defined over a field \mathbb{K} , in fact one can consider λ_1 and λ_2 to be in any extension \mathbb{E} of \mathbb{K} , then $\text{Ker}_{\lambda_1, \lambda_2} W \subset V \otimes_{\mathbb{K}} \mathbb{E}$.

In particular, if \mathbb{K} is the algebraic closure of \mathbb{K} , then for any $\lambda = (\lambda_1 : \lambda_2) \in \overline{\mathbb{K}}\mathbb{P}^1$ one can consider a correctly defined number $\dim \text{Ker}_\lambda W$.

¹⁰In fact, we will not use the previous connection between linear relations and pencils.

Definition 3.1. Say that a bisurjective linear relation W in V is *Kronecker* if $\dim \text{Ker}_\lambda W$ does not depend on $\lambda \in \overline{\mathbb{K}}\mathbb{P}^1$. Call this common dimension the *rank* of the relation.

Example 3.2. Assume that W is a graph of a linear mapping $V \rightarrow V$. Then W is Kronecker iff $\dim V = 0$.

Example 3.3. Let $V = \mathbb{K}^n$. Define $W \subset V \oplus V$ by $(v, v') \in W$ iff $v_k = v'_{k+1}$, $k = 1, \dots, n-1$. This is a Kronecker linear relation of rank 1.

Definition 3.4. A *Kronecker block* is a linear relation isomorphic to the relation of Example 3.3.

Definition 3.5. Given a linear relation W in V and a linear relation W' in V' , $W \oplus W'$ can be considered as a linear relation in $V \oplus V'$. Call this linear relation a *direct sum* of W and W' .

Definition 3.6. For $\lambda \in \mathbb{K}$ say that a relation W in V is a Jordan block with eigenvalue λ if W is a graph of a mapping $V \rightarrow V$ which is a Jordan block with eigenvalue λ . Say that W is a Jordan block with eigenvalue ∞ if W^{-1} is a Jordan block with eigenvalue 0, here W^{-1} is the image of W under transposition of summands in $V \oplus V$.

Theorem 3.7. *Suppose that \mathbb{K} is algebraically closed and $\dim V < \infty$.*

1. *any Kronecker linear relation in V of rank 1 is a Kronecker block.*
2. *Any Kronecker linear relation in V is isomorphic to a direct sum of Kronecker blocks.*
3. *Any bisurjective linear relation is isomorphic to a direct sum of Kronecker and Jordan blocks.*

The collection of dimensions (and—for Jordan blocks—eigenvalues) of these blocks is uniquely determined by V .

Proof. See classification of pencils of finite-dimensional linear operators, say in [8] or [32]. □

Given a Kronecker relation W in V of rank r , one obtains a natural mapping $\text{Ker}_\bullet: \mathbb{P}^1 \rightarrow \text{Gr}_r V$. Call this parameterized curve in $\text{Gr}_r V$ the *spectral curve* of W .

Remark 3.8. Given a bisurjective relation W in a finite-dimensional vector space V , $\dim \text{Ker}_\lambda W$ is constant on a Zariski open subset of $\overline{\mathbb{K}}\mathbb{P}^1$, i.e., outside of a finite subset $\Lambda \subset \overline{\mathbb{K}}\mathbb{P}^1$. In particular, there is a natural mapping $\text{Ker}_\bullet: \mathbb{P}^1 \setminus \Lambda \rightarrow \text{Gr}_r V$. Since $\text{Gr}_r V$ is complete, one can extend this mapping to a mapping $\mathbb{P}^1 \rightarrow \text{Gr}_r V$, which one can call a *spectral curve* as well.

Proposition 3.9. *Consider two matrices $A_1, A_2 \in \text{Mat}(m, n)$. They define a pencil of mappings $\mathbb{K}^m \rightarrow \mathbb{K}^n$. Suppose that $m \geq n$ and $\text{rk}(\lambda_1 A_1 + \lambda_2 A_2) = n$ for any $(\lambda_1, \lambda_2) \in \overline{\mathbb{K}}^2 \setminus (0, 0)$. Then the same is true for any pair (A'_1, A'_2) which is close to (A_1, A_2) .*

Proof. Consider the projectivization $\mathbb{P}\text{Mat}(m, n)$ of the vector space of matrices. Let $Z \subset \mathbb{P}\text{Mat}(m, n)$ be the projection of the set of matrices of rank $< n$ to $\mathbb{P}\text{Mat}(m, n)$, denote projections of A_1, A_2 on $\mathbb{P}\text{Mat}(m, n)$ by α_1, α_2 . Then Z is a closed subset, and the line through α_1, α_2 does not intersect Z . Due to compactness of $\mathbb{P}\text{Mat}(m, n)$, close lines do not intersect Z as well. \square

Remark 3.10. Due to the correspondence of Proposition 2.4, one can restate Proposition 3.9 as the fact that Kronecker relations in a vector space V form an open subset of all relations in V . Here we identify the set of relations with the union of Grassmannians of subspaces of different dimensions in $V \oplus V$.

4. KRONECKER WEBS

Definition 4.1. A *preweb* on a manifold B is a bisurjective linear relation in \mathcal{T}^*B , in other words, it is a linear bundle \mathcal{W} on B with an inclusion $\mathcal{W} \hookrightarrow \mathcal{T}^*B \oplus \mathcal{T}^*B$ which makes a bisurjective linear relation in each fiber of \mathcal{T}^*B .

Given a preweb \mathcal{W} and $\lambda \in \mathbb{K}\mathbb{P}^1$, one can consider $\text{Ker}_\lambda \mathcal{W}$, which is a collection of linear subspaces $\text{Ker}_\lambda \mathcal{W}_b \subset \mathcal{T}_b^*B$, $b \in B$. Recall that for a foliation \mathcal{F} Section 0 defined \mathcal{NF} , \mathcal{TF} , and $\mathcal{T}^*\mathcal{F}$.

Definition 4.2. Consider a vector subbundle E of \mathcal{T}^*B . Say that E is *integrable* if there is a foliation \mathcal{F} on B such that $E = \mathcal{NF}$.

Definition 4.3. Say that a preweb \mathcal{W} is a *web* if for any given $\lambda \in \mathbb{K}\mathbb{P}^1$ the number $\dim \text{Ker}_\lambda \mathcal{W}_b$ does not depend on $b \in B$, and this collection of subspaces is integrable. In other words, there is a foliation \mathcal{F}_λ on B such that $\text{Ker}_\lambda \mathcal{W}_b$ coincides with the normal spaces to \mathcal{F}_λ at $b \in B$. Call this foliation the λ -*integrating foliation* (or just *integrating foliation*, if λ is clear from the context) of \mathcal{W} .

Say that a preweb \mathcal{W} is *Kronecker* of rank r if for any $b \in B$ the linear relation \mathcal{W}_b in \mathcal{T}_b^*B is Kronecker of rank¹¹ r .

Remark 4.4. Note that Theorem 7.1 implies that the notion of Kronecker is a generalization of the notion of *Veronese webs (of higher codimension)* introduced in [25]. Since the definition of [25] cannot not stated in a coordinate-independent form, Definition 4.3 is much more convenient to work with.

Due to Proposition 2.4, to describe a preweb \mathcal{W} on B is “the same” as to define a vector bundle $\Phi(\mathcal{W})$ on B and a pencil \mathcal{P} of bisurjective mappings of vector bundles $\mathcal{T}^*B \rightarrow \Phi(\mathcal{W})$. If the preweb \mathcal{W} is clear from the context, we may denote $\Phi(\mathcal{W})$ as Φ as well.¹² Recall that given \mathcal{W} , $\Phi(\mathcal{W}) = \mathcal{T}^*B / \text{Ker}_\lambda \mathcal{W}$, given Φ and \mathcal{P} , the

¹¹Since rank of a Kronecker relation W in V equals $\dim W - \dim V$, rank of \mathcal{W}_m does not depend on m .

¹²The choice of notation is related to the fact that in applications the coordinates on fibers of Φ are angle-coordinates of an integrable system. (The *action* variables are coordinates on Φ coming from coordinates on M .)

vector bundle $\mathcal{W} \subset \mathcal{T}^*B \oplus \mathcal{T}^*B$ can be described by the condition $\mathcal{P}_{1,b}v_1 = \mathcal{P}_{2,b}v_2$, $(v_1, v_2) \in \mathcal{T}_b^*B$.

This preweb is Kronecker if $\text{Ker}(\lambda_1\mathcal{P}_{1,b} + \lambda_2\mathcal{P}_{2,b})$, $b \in B$, is an family of subspaces of \mathcal{T}^*B (or $\mathcal{T}^*B \otimes \mathbb{C}$) of the same dimension for any $b \in B$ and $(\lambda_1 : \lambda_2) \in \mathbb{C}\mathbb{P}^1$. If \mathcal{W} is a Kronecker preweb, \mathcal{W} is a Kronecker web if $\text{Ker}(\lambda_1\mathcal{P}_{1,b} + \lambda_2\mathcal{P}_{2,b})$ is an integrable family of subspaces of \mathcal{T}^*B of the same dimension for any $b \in B$ and $(\lambda_1 : \lambda_2) \in \mathbb{K}\mathbb{P}^1$. Due to Remark 3.10, the condition of being Kronecker is a condition of being in general position.

For a given $b \in B$ one can define $\Lambda_b \subset \mathbb{C}\mathbb{P}^1$ as in Remark 3.8. Obviously, if $\lambda_0 \notin \Lambda_{b_0}$, then there is a neighborhood $U \ni b_0$ and $\mathbf{U} \ni \lambda_0$ such that the vector subspaces $\text{Ker}_\lambda \mathcal{W}_b$ depend smoothly on $b \in U$ and $\lambda \in \mathbf{U}$. For any given $(\lambda_1 : \lambda_2) \in \mathbf{U} \subset \mathbb{C}\mathbb{P}^1$ the bundle $\mathcal{T}^*B / \text{Ker}_{(\lambda_1:\lambda_2)} \mathcal{W}|_U$ is canonically isomorphic to $\Phi|_u$. On the other hand, if $\lambda \in \mathbb{K}\mathbb{P}^1$, and \mathcal{W} is a web, then $\mathcal{T}^*B / \text{Ker}_{(\lambda_1:\lambda_2)} \mathcal{W} = \mathcal{T}^*\mathcal{F}_{(\lambda_1:\lambda_2)}$.

We see that given $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$ such that $\lambda = (\lambda_1 : \lambda_2) \in \mathbf{U}$ the vector bundle $\Phi|_U$ is canonically identified with $\mathcal{T}^*\mathcal{F}_\lambda|_U$. Since $\mathcal{T}^*\mathcal{F}$ has a natural Poisson structure (see Section 0), we see that

Proposition 4.5. *The total space $\Phi|_U$ carries a natural Poisson structure for any $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$ such that $\lambda = (\lambda_1 : \lambda_2) \in \mathbf{U}$. This Poisson structure depends smoothly on (λ_1, λ_2) .*

Call this Poisson structure $\eta_{\lambda_1, \lambda_2}$.

Proposition 4.6. *The Poisson structure $\eta_{\lambda_1, \lambda_2}$ is homogeneous in $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$ of degree 1.*

Proof. Indeed, multiplication of (λ_1, λ_2) by a constant changes the identification of $\Phi|_U$ with $\mathcal{T}^*\mathcal{F}_\lambda|_U$ by the same constant. For a diffeomorphism $\alpha: M_1 \rightarrow M_2$ and a Poisson structure η on M_1 denote by $\alpha_*\eta$ the Poisson structure η transferred to M_2 via α . Thus the only thing to prove is that if N is a manifold, η is the Poisson structure on \mathcal{T}^*N , and μ_c is multiplication by c in \mathcal{T}^*N , then $(\mu_c)_*\eta = c\eta$. One can check it momentarily in local coordinates on N (as in Example 0.6). \square

Corollary 4.7. *If $\mathbb{K} = \mathbb{C}$, and the web \mathcal{W} is Kronecker, then the Poisson structure $\eta_{\lambda_1, \lambda_2}$ on the total space of Φ depends linearly on λ_1, λ_2 .*

Proof. Since for a Kronecker web $\Lambda_b = \emptyset$, one can take $\mathbf{U} = \mathbb{C}\mathbb{P}^1$, and $U = B$, thus the Poisson structure $\eta_{\lambda_1, \lambda_2}$ is defined on the whole total space of Φ for any $(\lambda_1, \lambda_2) \neq (0, 0)$. Fix $x \in \Phi$. Since a Poisson structure is a bivector field, one can associate to $(\lambda_1, \lambda_2) \neq (0, 0)$ an element $\eta_{\lambda_1, \lambda_2}|_x$ of $\Lambda^2\mathcal{T}_x\Phi$. We know that this element depends analytically on $(\lambda_1, \lambda_2) \neq (0, 0)$ and is homogeneous of degree 1 in (λ_1, λ_2) . However, any mapping of $\mathbb{C}^2 \setminus (0, 0)$ to a vector space which is of homogeneity degree 1 is linear, which finishes the proof. \square

Theorem 4.8. *If a web \mathcal{W} is Kronecker, then the Poisson structure $\eta_{\lambda_1, \lambda_2}$ on the total space of Φ depends linearly on λ_1, λ_2 .*

Proof. The only case which remains to be proven is $\mathbb{K} = \mathbb{R}$. If a Kronecker web \mathcal{W} is real-analytic, then one can consider the complex-analytic continuation, and one returns to the case $\mathbb{K} = \mathbb{C}$. Thus in real-analytic case Corollary 4.7 implies the theorem.

Consider now C^∞ -case. First, note that the theorem follows from some “abstract nonsense” remarks. Recall that one of the contributions of algebraic geometry to differential geometry is the understanding of the importance of considering “formal” objects as tools for investigation of “geometric” objects. In particular, given a C^∞ -manifold, one can consider “an ∞ -jet of the complex-analytic neighborhood” (or a “formal neighborhood”) of this manifold. This formal neighborhood is canonically defined, and carries many properties of complex-analytic manifolds.

Readers familiar with the above language may immediately recognize that all the objects needed for the proof of Corollary 4.7 (foliations, tensors, cotangent bundles, Poisson structures) make sense in settings of “formal geometry”, thus one can finish the proof in C^∞ -case in the same way we did it in real-analytic case (when we had a no-nonsense complex neighborhood instead of a formal one). For other readers we provide a stripped-down version of the proof below. This version will not use “hard” notions of formal geometry (such as jets of manifolds etc.), but will use only notions of (finite-order) jets of sections of “real” bundles on “real” manifolds.¹³ This simplified version of the proof goes until the end of this section.

Consider a vector bundle E over B , fix a point $b \in B$. To describe a vector subbundle F of E of rank d is the same as to describe a section of the bundle $\text{Gr}_d(E)$ (the fibers of this bundle are Grassmannians of fibers of E). Recall that a k -jet near b of a section f of a bundle is the collection of Taylor coefficients of f of the order k or less.¹⁴ If a geometric object can be described locally by a section of some bundle, one can define a k -jet of this object near b as being a k -jet of a section of this bundle¹⁵. Thus one can define a k -jet of a vector subbundle F of E near b .

Remark 4.9. In what follows we will also need to consider E/F , here F is a (k -jet of) a subbundle of E . To avoid introducing new “abstract nonsense”, equip E with a norm, and identify E/F with the orthogonal complement to F . Obviously, given a k -jet of F , k -jet of E/F is well-defined as a k -jet of a vector (sub)bundle.

In particular, one can define a k -jet \mathcal{W} of a preweb. Consider two prewebs \mathcal{W} and \mathcal{W}' which have the same k -jet near b . Then the subbundles $\text{Ker}_\lambda \mathcal{W} \subset \mathcal{T}^*B$ and $\text{Ker}_\lambda \mathcal{W}' \subset \mathcal{T}^*B$ have the same k -jet near b . Thus there is a canonically defined k -jet of an isomorphism ι between linear bundles $\mathcal{T}^*B/\text{Ker}_\lambda \mathcal{W}$ and $\mathcal{T}^*B/\text{Ker}_\lambda \mathcal{W}'$. This

¹³As opposed to “formal” manifolds.

¹⁴In a coordinate-less form, k -jets are equivalence classes of sections, here equivalence is having the same Taylor coefficients of order k or less in any coordinate system.

¹⁵However, it may happen that there are different descriptions of the same object which lead to a shifted enumeration of jets. Say, locally one can describe a closed 1-form α by a function f such that $df = \alpha$, or by components of α considered as a tensor field. The k -jet in the first description is a $k - 1$ -jet in the second description.

leads to a k -jet¹⁶ of an isomorphism between the total spaces of these bundles. On the other hand, given a k -jet of an isomorphism ι between two manifolds F and F' , and a tensor field τ on F , one can define a $k - 1$ -jet of the tensor field $\iota_*\tau$ on F' .

Suppose that both prewebs \mathcal{W} and \mathcal{W}' are in fact webs. As we know, total spaces of both bundles $\mathcal{T}^*B/\text{Ker}_\lambda \mathcal{W}$ and $\mathcal{T}^*B/\text{Ker}_\lambda \mathcal{W}'$ carry natural Poisson structures, and one can consider Poisson structures as bivector (thus tensor) fields η and η' . Thus given a k -jet ι of an isomorphism between these manifolds, one can consider $\iota_*\eta$, which is a $k - 1$ -jet of a Poisson structure on $\mathcal{T}^*B/\text{Ker}_\lambda \mathcal{W}'$. Obviously, $\iota_*\eta$ coincides with the $k - 1$ -jet of η' . This immediately implies

Lemma 4.10. *Consider two Kronecker webs \mathcal{W} and \mathcal{W}' on B which have the same k -jet near $b \in B$. Then there is a canonically defined k -jet of an isomorphism ι between $\Phi(\mathcal{W})$ and $\Phi(\mathcal{W}')$, and this isomorphism identifies $k - 1$ -jets of Poisson structures $\eta_{\lambda_1, \lambda_2}$ (on Φ) and $\eta'_{\lambda_1, \lambda_2}$ (on Φ') near fibers of these bundles over b .*

Thus

Lemma 4.11. *In the conditions of Lemma 4.10 suppose that $k = 1$, and the web \mathcal{W}' is in fact real-analytic. Then the tensor field $\eta_{\lambda_1, \lambda_2}$ depends linearly on (λ_1, λ_2) on the fiber of Φ over b .*

In particular, the theorem follows from the case $k = 1$ of

Conjecture 4.12. Fix $k \geq 0$. Given a Kronecker web \mathcal{W} on an open subset $U \subset \mathbb{R}^n$, $0 \in U$, one can find a real-analytic Kronecker web \mathcal{W}' on $U' \subset U$, $0 \in U'$, such that k -jets of \mathcal{W} and \mathcal{W}' at 0 coincide.

In fact, in the case $k = 1$ we propose the following amplification:

Conjecture 4.13. Given a Kronecker web \mathcal{W} on an open subset $U \subset \mathbb{R}^n$, $0 \in U$, one can find $U' \subset U$, $0 \in U'$, and a diffeomorphism $U' \xrightarrow{f} V \subset \mathbb{R}^n$, $f(0) = 0$, such that 1-jet of $f_*\mathcal{W}$ (the transfer of \mathcal{W} via f) coincides with 1-jet of an appropriate translation-invariant¹⁷ Kronecker web.

Not only this conjecture implies the case $k = 1$ of Conjecture 4.12, but it would also directly imply the theorem we are proving. However, since these conjectures are not settled yet, one needs an alternative way to prove the theorem.

We are going to do this without the above extension-properties for webs by introducing integrability conditions on k -jets of prewebs. Since jets are not sensible to a change of the base field, we will be able to consider a k -jet of a C^∞ -web as a k -jet of a complex-analytic web, thus we will be able to extend $\eta_{\lambda_1, \lambda_2}$ to $(\lambda_1, \lambda_2) \in \mathbb{C}^2 \setminus (0, 0)$, then apply the arguments of Corollary 4.7.

¹⁶It is possible to define what is a k -jet of something *near a submanifold*, and the k -jet in question is near the fibers of these linear bundles over b .

¹⁷A Kronecker web \mathcal{W} on a vector space V is *translation-invariant* if the vector subspace $\mathcal{W}_v \subset \mathcal{T}_v^*V \oplus \mathcal{T}_v^*V = V^* \oplus V^*$ does not depend on $v \in V$.

Definition 4.14. Say that k -jet W of a vector subbundle of \mathcal{T}^*B is k -integrable near $b_0 \in B$ if there is a foliation \mathcal{F} such that the k -jet of the normal bundle to \mathcal{F} coincides with W .

Definition 4.15. Say that a k -jet \mathcal{W} of a preweb is a k -jet-web if for any $\lambda \in \mathbb{K}\mathbb{P}^1$ the k -jet $\text{Ker}_\lambda \mathcal{W}$ of a vector subbundle of \mathcal{T}^*B is k -integrable. Say that k -jet \mathcal{W} of a web near b is *Kronecker* if the relation \mathcal{W}_b in \mathcal{T}_b^*B is Kronecker.

Due to Remark 3.10, if a k -jet near b of preweb \mathcal{W} on B is Kronecker, then the preweb \mathcal{W} is Kronecker in a neighborhood of b . Thus the above definition of a Kronecker k -jet-web is compatible with taking a k -jet of a Kronecker web.

Consider a k -jet-web \mathcal{W} . It is a k -jet of a preweb, denote this preweb by \mathcal{W}' . Given $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$, the k -jet of $\mathcal{T}^*B / \text{Ker}_\lambda \mathcal{W}'$, $\lambda = \lambda_1 : \lambda_2$, is naturally identified with the k -jet of the cotangent bundle of the corresponding foliation \mathcal{F}_λ , in other words, one can construct a k -jet of an identification α of $\mathcal{T}^*B / \text{Ker}_\lambda \mathcal{W}'$ with $\mathcal{T}^*\mathcal{F}_\lambda$. Recall that a k -jet of a diffeomorphism acts on $k - 1$ -jets of tensor fields. $\mathcal{T}^*\mathcal{F}_\lambda$ carries a natural Poisson structure, thus it carries the tensor field which describes the bracket of the Poisson structure. Using the above identification α , one obtains a $k - 1$ -jet of a tensor field $\eta_{\lambda_1, \lambda_2}$ on $\mathcal{T}^*B / \text{Ker}_\lambda \mathcal{W}'$. Obviously, one can consider this $k - 1$ -jet as living on $\mathcal{T}^*B / \text{Ker}_\lambda \mathcal{W}$. It does not depend on the choice of \mathcal{W}' .

If $\mathbb{K} = \mathbb{C}$, and $k = 1$, then one obtains a 0-jet of a tensor field $\eta_{\lambda_1, \lambda_2}$ for any $(\lambda_1, \lambda_2) \in \mathbb{C}^2 \setminus (0, 0)$, and the same arguments as in the proof of Corollary 4.7 show that

Proposition 4.16. *Let $\mathbb{K} = \mathbb{C}$, consider a Kronecker 1-jet-web \mathcal{W} near $b_0 \in B$. The family of tensors fields $\eta_{\lambda_1, \lambda_2}$ on the total space of Φ is well defined on the fiber over b_0 of the projection $\Phi \rightarrow B$. This family depends linearly on λ_1, λ_2 .*

To finish the proof of the theorem, it is enough to show that given a Kronecker 1-jet-web over \mathbb{R} , one can consider it as a Kronecker 1-jet-web over \mathbb{C} . Recall that a jet of a preweb is just a collection of Taylor coefficients of a section of a bundle, thus a collection of numbers. Any collection of real numbers can be considered as a collection of complex numbers, thus a 1-jet of a preweb over \mathbb{R} can be considered as 1-jet of a preweb over \mathbb{C} . Since the condition of being Kronecker does not change when we change field of scalars, the only thing we need to prove is the integrability condition for complex $\lambda_1 : \lambda_2$.

Recall Frobenius integrability condition:

Proposition 4.17. *Consider a linear subbundle E of \mathcal{T}^*B . Suppose that for any small open subset $U \subset B$ and any section α of E over U its de Rham differential $d\alpha$ (which is a 2-form on B) can be written as $\sum \alpha_i \wedge \beta_i$ with α_i being sections of E and β_i being arbitrary differential forms on B . Then E is integrable.*

By constructing jet-solutions of ordinary differential equations, one can easily prove the following jet-analogue of Proposition 4.17:

Proposition 4.18. *Consider a k -jet of a linear subbundle E of \mathcal{T}^*B near $b \in B$. Suppose that for any open subset $U \subset B$ and for any k -jet α of a section of E near $b \in B$ its de Rham differential $d\alpha$ (which is a $k - 1$ -jet of a 2-form on B) can be written as $\sum \alpha_i \wedge \beta_i$ with α_i being $k - 1$ -jets of sections of E and β_i being arbitrary differential forms on B . Then E is k -integrable.*

In fact one can do more. Given a section α of E , consider the image of $d\alpha$ under projection $\Omega^2 B = \Lambda^2 \mathcal{T}^*B \rightarrow \Lambda^2(\mathcal{T}^*B/E)$. This defines a mapping δ from sections of E to sections of $\Lambda^2(\mathcal{T}^*B/E)$. A priori it is a differential operator of order 1, but in fact it has order 0, thus is a linear mapping between bundles. Indeed, if $\alpha|_b = 0$, then one can easily check that $\delta\alpha|_b = 0$.

Thus δ defines a section of the linear bundle $E^* \otimes \Lambda^2(\mathcal{T}^*B/E)$, call this section the *torsion* of E . Given a k -jet of a linear subbundle E , δ is defined as $k - 1$ -jet of a section of $E^* \otimes \Lambda^2(\mathcal{T}^*B/E)$. If a linear subbundle E_t depends smoothly on a parameter $t \in T$, then δ_t depends smoothly on t (i.e., it is a smooth section of the vector bundle $E_t^* \otimes \Lambda^2(\mathcal{T}^*B/E_t)$ over $B \times T$).

Suppose $k \geq 1$. Apply the above construction of δ to the linear subbundle $\text{Ker}_\lambda \mathcal{W} \subset \mathcal{T}^*B$ of a Kronecker preweb \mathcal{W} in the case $\mathbb{K} = \mathbb{C}$. Here $\lambda \in \mathbb{C}\mathbb{P}^1$, thus δ is a section of $(\text{Ker}_\lambda \mathcal{W})^* \otimes \Lambda^2(\mathcal{T}^*B/\text{Ker}_\lambda \mathcal{W})$ over $B \times \mathbb{C}\mathbb{P}^1$. Restrict this section to $\{b\} \times \mathbb{C}\mathbb{P}^1$, $b \in B$. We obtain a vector bundle over $\mathbb{C}\mathbb{P}^1$, and a regular section $\delta^{(b)}$ of this vector bundle. There may be only two different cases: either $\delta^{(b)} = 0$, or $\delta^{(b)}$ vanishes at a finite number of points of $\mathbb{C}\mathbb{P}^1$.

Proposition 4.19. *Consider a Kronecker 1-jet-web \mathcal{W} near $0 \in \mathbb{R}^n$ (and $\mathbb{K} = \mathbb{R}$). The complexification of this 1-jet-web is a 1-jet $\mathcal{W}_\mathbb{C}$ of a Kronecker preweb with $\mathbb{K} = \mathbb{C}$. Then $\mathcal{W}_\mathbb{C}$ is a Kronecker 1-jet-web.*

Proof. Arbitrarily extend \mathcal{W} to a preweb \mathcal{W}' in neighborhood of 0 in \mathbb{R}^n , and arbitrarily extend $\mathcal{W}_\mathbb{C}$ to a preweb $\mathcal{W}'_\mathbb{C}$ in a neighborhood of 0 in \mathbb{C}^n . Consider the torsion $\delta_\mathbb{C}$ of the subbundles $\text{Ker}_\lambda \mathcal{W}'_\mathbb{C}$ restricted to $\{0\} \times \mathbb{C}\mathbb{P}^1$, and the torsion δ of the subbundles $\text{Ker}_\lambda \mathcal{W}'$ restricted to $\{0\} \in \mathbb{R}\mathbb{P}^1$. Obviously, δ is a restriction of $\delta_\mathbb{C}$ from $\mathbb{C}\mathbb{P}^1$ to $\mathbb{R}\mathbb{P}^1$. On the other hand, δ vanishes, since 1-jet \mathcal{W} of \mathcal{W}' is a 1-jet-web. Thus $\delta_\mathbb{C}$ vanishes on $\mathbb{R}\mathbb{P}^1$, thus at infinitely many points of $\mathbb{C}\mathbb{P}^1$, thus $\delta_\mathbb{C} = 0$. This implies that the conditions of Proposition 4.18 are satisfied, which finishes the proof. \square

This finishes the proof of Theorem 4.8. \square

In the above proof we ignored the question of extension of jets of webs completely. Let us state

Conjecture 4.20. Given a Kronecker k -jet-web \mathcal{W} near $0 \in \mathbb{C}^n$, one can find $U \subset \mathbb{C}^n$, $0 \in U$, and a Kronecker web \mathcal{W}' on U such that k -jet of \mathcal{W}' at 0 coincides with \mathcal{W} .

Anyway, we proved the following

Corollary 4.21. *Given a Kronecker web \mathcal{W} , the total space of the vector bundle $\Phi(\mathcal{W})$ is equipped with a natural bihamiltonian structure.*

5. PAIRS OF SKEWSYMMETRIC FORMS

Recall the classification of pairs of skewsymmetric bilinear pairings from [11] (see also [12, 13]). For $k \in \mathbb{N}$ consider the identity $k \times k$ matrix I_k . For $\mu \in \mathbb{C}$ consider the Jordan block $J_{k,\mu}$ of size k and eigenvalue μ . The pair of matrices

$$H_1^{(\mu)} = \begin{pmatrix} 0 & J_{k,\mu} \\ -J_{k,\mu}^t & 0 \end{pmatrix}, \quad H_2^{(\mu)} = \begin{pmatrix} 0 & I_k \\ -I_k & 0 \end{pmatrix}$$

defines a pair of skewsymmetric bilinear pairings on vector space \mathbb{C}^{2k} . The limit case of $\mu \rightarrow \infty$ may be deformed to

$$H_1^{(\infty)} = \begin{pmatrix} 0 & I_k \\ -I_k & 0 \end{pmatrix}, \quad H_2^{(\infty)} = \begin{pmatrix} 0 & J_{k,0} \\ -J_{k,0}^t & 0 \end{pmatrix}.$$

Denote the pair $(H_1^{(\mu)}, H_2^{(\mu)})$ of skewsymmetric bilinear pairings by $\mathcal{J}_{2k,\mu}$, $k \in \mathbb{N}$, $\mu \in \mathbb{C}\mathbb{P}^1$.

Add to this list the so-called Kronecker pair \mathcal{K}_{2k-1} . This is a pair in a vector space \mathbb{C}^{2k-1} with a basis $(\mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_{2k-2})$. The only non-zero pairings are

$$(5.1) \quad (\mathbf{w}_{2l}, \mathbf{w}_{2l+1})_1 = 1, \quad (\mathbf{w}_{2l+1}, \mathbf{w}_{2l+2})_2 = 1,$$

for $0 \leq l \leq k-2$. Obviously, different pairs from this list are not isomorphic.

Theorem 5.1. ([11, 29]) *Any pair of skewsymmetric bilinear pairings on a finite-dimensional complex vector space can be decomposed into a direct sum of pairs of the pairings isomorphic to $\mathcal{J}_{2k,\mu}$, $k \in \mathbb{N}$, $\mu \in \mathbb{P}^1$, and \mathcal{K}_{2k-1} , $k \in \mathbb{N}$. The types of the components of this decomposition are uniquely determined.*

Though this simple statement was known for a long time (say, the preprint of [29] existed in 1973), we do not know whether it was published before it was used in [11]. The discussions in [8] and [32] come very close, but do not state this result.

Definition 5.2. Say that a pair of bilinear skewsymmetric forms $(,)_1$ and $(,)_2$ in a vector space V over \mathbb{C} is *Kronecker* if it has no Jordan blocks. The *rank* of a Kronecker pair is the number of Kronecker blocks in the decomposition of the pair into undecomposable components.¹⁸

Given a Kronecker pair of bilinear skewsymmetric forms in V let the *action subspace* of V be spanned by vector subspaces $\text{Ker}(,)_{\lambda_1, \lambda_2}$, $(\lambda_1, \lambda_2) \in \mathbb{C}^2 \setminus (0, 0)$. Here $(v, v')_{\lambda_1, \lambda_2} \stackrel{\text{def}}{=} \lambda_1 (v, v')_1 + \lambda_2 (v, v')_2$.

¹⁸Note a conflict in these notations: for a pair $(,)_{1,2}$ of rank r , any form has corank k .

Proposition 5.3. *The action subspace of a Kronecker pair of bilinear forms in V of rank r is isotropic with respect to $(\cdot, \cdot)_1$ and $(\cdot, \cdot)_2$, and has dimension $\frac{\dim V + r}{2}$. It is a maximal isotropic subspace for any form $(\cdot, \cdot)_{\lambda_1, \lambda_2}$, $(\lambda_1, \lambda_2) \in \mathbb{C}^2 \setminus (0, 0)$.*

Proof. This follows immediately from the explicit form of a Kronecker block. \square

Proposition 5.4. *The action subspace \mathcal{A} of a Kronecker pair of bilinear forms in V of rank r has a natural Kronecker linear relation of rank r .*

Proof. Since \mathcal{A} is isotropic with respect to $(\cdot, \cdot)_1$, $(\cdot, \cdot)_1$ induces a natural pairing of \mathcal{A} with V/\mathcal{A} , or a mapping $\alpha_1: \mathcal{A} \rightarrow (V/\mathcal{A})^*$. Similarly, $(\cdot, \cdot)_2$ induces a mapping $\alpha_2: \mathcal{A} \rightarrow (V/\mathcal{A})^*$. Consider the relation $\alpha_2^{-1}\alpha_1$ in \mathcal{A} . Looking on the explicit form of a Kronecker block of a pair of skewsymmetric bilinear forms, one can easily recognize in the relation $\alpha_2^{-1}\alpha_1$ a direct sum of Kronecker blocks. \square

Proposition 5.5. *Consider two families $(\cdot, \cdot)_{1,t}$, $(\cdot, \cdot)_{2,t}$, $t \in T$, of skewsymmetric bilinear forms in a vector space V , parameterized by a manifold T . Suppose that there is $r \in \mathbb{N}$ such that for any $t \in T$ the pair $(\cdot, \cdot)_{1,t}$, $(\cdot, \cdot)_{2,t}$ is Kronecker of rank r . Let \mathcal{A}_t be the action subspace of the pair $(\cdot, \cdot)_{1,t}$, $(\cdot, \cdot)_{2,t}$. Then \mathcal{A}_t depends smoothly on t .*

Proof. This follows immediately from the following

Lemma 5.6. *Consider a vector space V and numbers $n_1, \dots, n_k, N \in \mathbb{N}$. Consider the product X of Grassmannians $\prod_{i=1}^k \text{Gr}_{n_i}(V)$ and the subset $Z \subset X$ consisting of k -tuples of subspaces such that the linear span of such a k -tuple has dimension N . Consider the mapping*

$$\iota: Z \rightarrow \text{Gr}_N(V) : (V_1, \dots, V_k) \mapsto V_1 + \dots + V_k,$$

here V_i is a vector subspace of V of dimension n_i . Then the mapping ι is smooth.

Proof. Let $x_0 \in Z$, U_0 be a small neighborhood of x_0 in X . For $x \in X$ denote by $V_i(x)$ the i -th component of x , $V_i(x) \in \text{Gr}_{n_i}(V)$. Choose a basis $v_{il}(x)$ in $V_i(x)$, $i = 1, \dots, k$, $l = 1, \dots, n_i$, which depends regularly on $x \in U$. Pick up a basis $\{v_\alpha(x_0)\}_{\alpha \in A}$ out of vectors $v_{il}(x_0)$, $i = 1, \dots, k$, $l = 1, \dots, n_i$ (each α has a form il). Then in an open subset U_1 of U_0 the vectors $v_\alpha(x)$ remain linearly independent, thus on $U_1 \cap Z$ they span $V_1 + \dots + V_k$. This implies¹⁹ that $V_1(z) + \dots + V_k(z)$ depends regularly on $z \in Z$. \square

This finishes the proof of Proposition 5.5. \square

6. MICRO-KRONECKER BIHAMILTONIAN STRUCTURES

Recall that a bihamiltonian structure on M induces a pair $(\cdot, \cdot)_1$, $(\cdot, \cdot)_2$ of bilinear skewsymmetric forms in \mathcal{T}^*M . As a corollary, it also induces a pair of bilinear skewsymmetric forms in $\mathcal{T}^*M \otimes \mathbb{C}$, which we will denote by the same symbols.

¹⁹A similar argument can show that Z is a submanifold of X .

Definition 6.1. Say that a bihamiltonian structure on M is *micro-Kronecker* at $m \in M$ if the pair of bilinear skewsymmetric forms $(,)_1$ and $(,)_2$ in \mathcal{T}_m^*M has no Jordan blocks. The *rank* of the bihamiltonian structure at $m \in M$ is the number of Kronecker blocks. Say that a bihamiltonian structure on M is *micro-Kronecker of rank r* if it is micro-Kronecker at all the points $m \in M$ of the same rank r .

The above definition is almost identical to the definition of *completeness* in [4] (see also [25]), a similar but more restrictive definition of *homogeneity* was introduced in [10].

Proposition 6.2. *Consider a micro-Kronecker bihamiltonian structure of rank r on a manifold M . There is a foliation \mathcal{F} on M such that for any $m \in M$ the action subspace of \mathcal{T}_m^*M is the normal space to the leaf of \mathcal{F} through m .*

Proof. By Proposition 5.5 action subspaces form a subbundle of \mathcal{T}^*M . Consider $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$ and the Poisson structure $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ on M . This Poisson structure has corank r , consider the symplectic foliation $\tilde{\mathcal{F}}_{\lambda_1, \lambda_2}$. Then the kernel of the bilinear form $(,)_{\lambda_1, \lambda_2}$ in \mathcal{T}_m^*M is the normal space to the leaf of $\tilde{\mathcal{F}}_{\lambda_1, \lambda_2}$ through m . Now the statement follows from the following lemmas:

Lemma 6.3. *Consider a pair of skewsymmetric bilinear forms in V which has no Jordan blocks. Suppose that dimensions of all the Kronecker components of V are $\leq k$. If $\{\lambda_i\}, i \in K$, is a finite subset of $\mathbb{K}\mathbb{P}^1$ with k elements, then $V = \sum_{i \in K} \text{Ker}_{\lambda_i} W$.*

Proof. Direct corollary of the explicit form of a Kronecker block. \square

Lemma 6.4. *Consider vector subbundles E, E_1, \dots, E_k of \mathcal{T}^*M , such that $E = \sum E_i$. Suppose that E_i coincides with the normal bundle to a foliation \mathcal{F}_i on M , $i = 1, \dots, k$. Then there is a foliation \mathcal{F} on M such that E coincides with the normal bundle to \mathcal{F} . Leaves of \mathcal{F} can be described as intersections of leaves of foliations \mathcal{F}_i .*

Proof. This follows immediately for the Frobenius integrability criterion (Proposition 4.17). \square

This finishes the proof of Proposition 6.2. \square

Obviously, the foliation \mathcal{F} is canonically defined by the bihamiltonian structure. A (most important) particular case of Proposition 6.2 was announced in [25].

Definition 6.5. Given a micro-Kronecker bihamiltonian structure on a manifold M , call the foliation \mathcal{F} of Proposition 6.2 the *action foliation* of the bihamiltonian structure.

Definition 6.6. Given a micro-Kronecker bihamiltonian structure on M such that the action foliation \mathcal{F} is a fibration, let \mathcal{B}_M be the base of this fibration. If M is clear from the context, denote the base by \mathcal{B} .

From now on suppose that the foliation \mathcal{F} is in fact a fibration of M over the base \mathcal{B} . One can always achieve this by decreasing M .

Theorem 6.7. *The base of the action foliation of a micro-Kronecker bihamiltonian structure has a canonically defined structure of a Kronecker web.*

Proof. Indeed, consider $m \in M$ and the projection b of m to \mathcal{B} . The (co)differential $(db|_m)^*$ of the mapping of projection identifies the vector space $\mathcal{T}_b^*\mathcal{B}$ with the action subspace at m . However, by Proposition 5.4 the action subspace at m is equipped with a Kronecker linear relation, thus $\mathcal{T}_b^*\mathcal{B}$ is equipped with such a relation as well. To show that \mathcal{B} has a structure of a Kronecker preweb, it is enough to show that this relation in $\mathcal{T}_b^*\mathcal{B}$ does not depend on the choice of the point m over b .

Let m, m' be two point of M over $b \in \mathcal{B}$. Let $\mathcal{A}_m, \mathcal{A}_{m'}$ be the action subspaces in \mathcal{T}_m^*M and $\mathcal{T}_{m'}^*M$. Both \mathcal{A}_m and $\mathcal{A}_{m'}$ are identified with $\mathcal{T}_b^*\mathcal{B}$, thus one with the other.

Lemma 6.8. *For any $\lambda \in \mathbb{K}\mathbb{P}^1$ the identification between \mathcal{A}_m and $\mathcal{A}_{m'}$ sends $\text{Ker}_\lambda W_m \subset \mathcal{A}_m$ to $\text{Ker}_\lambda W_{m'} \subset \mathcal{A}_{m'}$.*

Proof. Let $\lambda = \lambda_1 : \lambda_2$. Consider the Poisson structure $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ on M . Then $\text{Ker}_\lambda W$ is the normal bundle to the symplectic foliation $\widetilde{\mathcal{F}}_\lambda$ for this Poisson structure. Consider the leaf L of this foliation which passes through m . This leaf contains the leaf of the action foliation \mathcal{F} through m , thus L is a preimage of a submanifold $\widetilde{L} \subset \mathcal{B}$. Thus $\text{Ker}_\lambda W_m \subset \mathcal{A}_m$ is the image of the normal space to \widetilde{L} under the (co)differential of the projection mapping.

By the same reason m' is in L , and $\text{Ker}_\lambda W_{m'} \subset \mathcal{A}_{m'}$ is also the image the same normal space. Thus the identification of \mathcal{A}_m and $\mathcal{A}_{m'}$ via projection to \mathcal{B} indeed sends $\text{Ker}_\lambda W_m$ to $\text{Ker}_\lambda W_{m'}$. \square

Lemma 6.9. *Consider vector spaces V and V' with Kronecker linear relations W and W' in V and V' correspondingly. Consider an isomorphism $\alpha: V \rightarrow V'$. If α sends $\text{Ker}_\lambda W$ to $\text{Ker}_\lambda W'$ for any $\lambda \in \mathbb{K}\mathbb{P}^1$, then $\alpha \oplus \alpha$ sends $W \subset V \oplus V$ to $W' \subset V' \oplus V'$.*

This lemma is equivalent to Theorem 7.1 proven in Section 7.

We had shown that the base \mathcal{B} of the action foliation has a canonically defined structure of a Kronecker preweb. Denote by $\widetilde{\mathcal{W}}_b$ the relation in $\mathcal{T}_b^*\mathcal{B}$, $b \in \mathcal{B}$.

To show that this preweb is in fact a web, it is enough to describe the λ -integrating foliation \mathcal{F}_λ of $\widetilde{\mathcal{W}}$. Fix $\lambda_1, \lambda_2 \in \mathbb{K}$, consider the Poisson structure $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ on M . Since this is a Poisson structure of constant rank, symplectic leaves form a foliation on M . This foliation depends only on $\lambda_1 : \lambda_2 \in \mathbb{K}\mathbb{P}^1$, denote this foliation $\widetilde{\mathcal{F}}_{\lambda_1:\lambda_2}$.

The normal space to this foliation at $m \in M$ is the kernel of the bilinear form $\lambda_1 (,)_1 + \lambda_2 (,)_2$ in \mathcal{T}_m^*M . In particular, the normal bundle to $\widetilde{\mathcal{F}}_{\lambda_1:\lambda_2}$ is contained in the action bundle of the bihamiltonian structure. Thus \mathcal{F} is a subfoliation of $\widetilde{\mathcal{F}}_{\lambda_1:\lambda_2}$. In particular, $\widetilde{\mathcal{F}}_{\lambda_1:\lambda_2}$ induces a ‘‘quotient’’ foliation $\mathcal{F}_{\lambda_1:\lambda_2}$ on the base \mathcal{B} of the foliation \mathcal{F} . Now one can immediately see that the normal space to the foliation

$\mathcal{F}_{\lambda_1:\lambda_2}$ at $b \in \mathcal{B}$ coincides with $\text{Ker}_\lambda \widetilde{\mathcal{W}}_b$, thus $\mathcal{F}_{\lambda_1:\lambda_2}$ is the $\lambda_1 : \lambda_2$ -integrating foliation of the preweb on \mathcal{B} . Thus the preweb structure on \mathcal{B} is indeed a Kronecker web. \square

A statement which is parallel to a particular case of Theorem 6.7 was announced in [25].

7. LATTICE OF KERNELS

Given a bisurjective relations W in V , one obtains a collection of vector subspaces $\text{Ker}_\lambda W$, $\lambda \in \mathbb{K}\mathbb{P}^1$, of the vector space V . The target of this section is to prove

Theorem 7.1. *Let $GR(V)$ denotes the set of all the vector subspaces of a vector space V . Associate to a bisurjective linear relation W in V a mapping of sets $\mathcal{K}_W: \mathbb{K}\mathbb{P}^1 \rightarrow GR(V)$, $\mathcal{K}_W(\lambda) = \text{Ker}_\lambda W$.*

If W and W' are two bisurjective linear relations in V such that $\mathcal{K}_W = \mathcal{K}_{W'}$ and W is Kronecker, then $W = W'$.

This theorem is a corollary of the following

Amplification 7.2. *In the conditions of Theorem 7.1 suppose that all the Kronecker blocks of W have dimensions k or less. Let $\Lambda \subset \mathbb{K}\mathbb{P}^1$ be a collection of $k + 2$ points. Then if $\mathcal{K}_W|_\Lambda = \mathcal{K}_{W'}|_\Lambda$, then $W = W'$.*

Proof. First, the following lemma follows immediately from the explicit description of Kronecker and Jordan blocks:

Lemma 7.3. *Consider a relation $W \subset V \oplus V$ which is a Kronecker block with $\dim V \leq k$, or is a Jordan block. Let $\{\lambda_i\}$, $1 \leq i \leq k$, be a subset of $\mathbb{K}\mathbb{P}^1$. Then $\text{Ker}_{\lambda_k} W \cap \sum_{i=1}^{k-1} \text{Ker}_{\lambda_i} W = 0$.*

By Theorem 3.7, any bisurjective relation W in V can be represented as $W = W_{\text{Kronecker}} \oplus W_{\text{Jordan}}$ with $W_{\text{Kronecker}}$ being Kronecker, and W_{Jordan} having no Kronecker blocks. Correspondingly, $\text{Ker}_\lambda W = \text{Ker}_\lambda W_{\text{Kronecker}} \oplus \text{Ker}_\lambda W_{\text{Jordan}}$. From Lemmas 6.3 and 7.3 it is clear that

$$\text{Ker}_\lambda W_{\text{Kronecker}} = \text{Ker}_\lambda W \cap \sum_{i=1}^l \text{Ker}_{\lambda_i} W$$

if l is large enough, thus $\text{Ker}_\lambda W_{\text{Kronecker}}$ is

1. a canonically defined subspace of $\text{Ker}_\lambda W$, and
2. can be found basing on $\{\text{Ker}_\mu W\}_{\substack{\mu \in \Lambda \\ \mu \neq \lambda}}$.

Moreover, $W_{\text{Kronecker}} = \sum_{\lambda \in \Lambda} \text{Ker}_\lambda W_{\text{Kronecker}}$, thus $W_{\text{Kronecker}}$ is determined by $\{\text{Ker}_\mu W\}_{\mu \in \Lambda}$ too (thus is canonically defined).

In particular, in the conditions of the amplification W' is Kronecker. Note that Lemmas 6.3, 7.3 imply that W' is Kronecker with the same rank and the same dimensions of Kronecker blocks as W . (We will not use the latter observation in what follows.)

Definition 7.4. Consider a linear mapping $f: V \rightarrow V'$ and a linear relation W in V . Denote by $f_!W$ the relation in V' given by the vector subspace $(f \oplus f)(W) \subset V' \oplus V'$.

Given a bisurjective linear relation W in V and $\Lambda \subset \mathbb{K}\mathbb{P}^1$, consider the collection of vector subspaces $\text{Ker}_\lambda W$, $\lambda \in \Lambda$. There is a natural mapping (of addition)

$$(7.1) \quad \alpha: \bigoplus_{\lambda \in \Lambda} \text{Ker}_\lambda W \rightarrow V.$$

In the conditions of the amplification this mapping is surjective by Lemma 6.3.

Given a vector space V and $\lambda = \lambda_1 : \lambda_2 \in \mathbb{K}\mathbb{P}^1$, denote by $\lambda \mathbf{1}$ the linear relation in V given by $\{(v_1, v_2) \mid \lambda_1 v_1 = \lambda_2 v_2\} \subset V \oplus V$. If $\lambda \neq \infty$, $\lambda \mathbf{1}$ is the graph of multiplication by λ in V .

Lemma 7.5. Consider a Kronecker relation W in V , and $\Lambda \subset \mathbb{K}\mathbb{P}^1$. Consider the linear relation W_0 in $V = \bigoplus_{\lambda \in \Lambda} \text{Ker}_\lambda W$ which is a direct sum of relations $\lambda \mathbf{1}$ in each of $\text{Ker}_\lambda W$. If dimensions of all the Kronecker blocks in V are less than $\text{card}(\Lambda) - 1$, then $W = \alpha_! W_0$, here α is from (7.1).

Proof. Due to Theorem 3.7, it is enough to prove this theorem in the case when W is a Kronecker block. We will use a Kronecker block of a particular form.

Let $k = \dim V - 1$. Recall that \mathcal{S} was introduced in Definition 2.2. Let V_0 be the symmetric power $\text{Sym}^{k-1} \mathcal{S}$, $V_1 = \text{Sym}^k \mathcal{S}$. Given $s \in \mathcal{S}$, the multiplication by s defines a linear operator $m_s: V_0 \rightarrow V_1$ and a dual operator $m_s^*: V_1^* \rightarrow V_0^*$. This defines a pencil of linear operators $(m_{s_1}^*, m_{s_2}^*)$ between V_1^* and V_0^* , thus a linear relation

$$\widetilde{W} = \{(v_1, v_2) \mid m_{s_1}^* v_1 = m_{s_2}^* v_2\} \subset V_1^* \oplus V_1^*,$$

which is a Kronecker block. We may assume that $V = V_1^*$, $W = \widetilde{W}$.

The above description of V as of $(\text{Sym}^k \mathcal{S})^*$ shows that it might be useful to restate the lemma in terms of the dual space to V . First of all, given a subspace $W \subset V \oplus V$, the orthogonal complement W^\perp is a subspace of $V^* \oplus V^*$, thus a relation in V^* . Given a surjective linear mapping $f: V \rightarrow V'$ and a relations W in V , W' in V' , obviously $W' = f_!W$ iff $(f^*)_! W'^\perp = W^\perp \cap \text{Im}(f^* \otimes f^*)$.

It is clear that α from (7.1) is surjective, and $\widetilde{W}^\perp = \text{Im}(m_{s_1}, -m_{s_2})$, here

$$(m_{s_1}, -m_{s_2}): V_0 \rightarrow V_1 \oplus V_1: p \mapsto (s_1 p, -s_2 p).$$

Consider the mapping α^* .

Given $p \in \mathcal{S}^*$, one can define linear functionals $\varepsilon_{p,l}$, $l = k$ or $l = k - 1$, on V_1 and on V_0 correspondingly. Indeed, one can naturally identify $\text{Sym}^l \mathcal{S}$ with the space of homogeneous polynomials of degree l on \mathcal{S}^* , and $\varepsilon_{p,l}$ is evaluation of a polynomial at $p \in \mathcal{S}^*$. Obviously, $m_s^* \varepsilon_{p,k} = \langle s, p \rangle \varepsilon_{p,k-1}$, here \langle, \rangle is the pairing between \mathcal{S} and \mathcal{S}^* . Thus the 1-dimensional subspace $\text{Ker}_{\lambda_1: \lambda_2} W$ is spanned by $\varepsilon_{p,k}$, here p is orthogonal to $\lambda_1 e_1 + \lambda_2 e_2$.

Thus we constructed a basis $\varepsilon_{p,k}$ in $\mathbf{V} = \bigoplus_{\lambda \in \Lambda} \text{Ker}_\lambda W$, here p runs through an appropriate subset $\Pi \subset \mathcal{S}^*$, $\text{card}(\Pi) = \text{card}(\Lambda)$. Thus \mathbf{V}^* can be identified with $\mathbb{K}^{\text{card}(\Lambda)}$. Given any subset $\Pi \subset \mathcal{S}^*$, consider the evaluation mapping $\varepsilon_\Pi: V_1 \rightarrow \mathbb{K}^{\text{card}(\Pi)}: \varphi \mapsto (\varphi(p))_{p \in \Pi}$. Now it should be obvious that in the above basis α^* is identified with ε_Π .

Now describe W_0^\perp . After a linear transformation in \mathcal{S} , one may assume that $0 \notin \Lambda$, $\infty \notin \Lambda$. Then in the above basis of \mathbf{V} the vector subspace $W_0 \subset \mathbf{V} \oplus \mathbf{V}$ is the graph of a diagonal matrix $\text{diag } \Lambda$. Thus W_0^\perp is the graph of a diagonal matrix $-\text{diag } \Lambda$.

Now the statement $(\alpha^*)_! \widetilde{W}^\perp = W_0^\perp \cap \text{Im}(\alpha^* \otimes \alpha^*)$ which we need to prove can be restated as

$$\varepsilon_{\Pi!} \text{Im}(m_{\mathbf{s}_1}, -m_{\mathbf{s}_2}) = \text{Graph}(-\text{diag } \Lambda) \cap \text{Im}(\varepsilon_\Pi \otimes \varepsilon_\Pi).$$

To describe the subspace $\mathbf{W} = \text{Graph}(-\text{diag } \Lambda) \cap \text{Im}(\varepsilon_\Pi \otimes \varepsilon_\Pi)$ it is enough to describe sequences of numbers $(\varphi_p)_{p \in \Pi}$ such that both $(\varphi_p)_{p \in \Pi}$ and $(\lambda_p \varphi_p)_{p \in \Pi}$ are values of homogeneous polynomials of degree k on Π (here λ_p is defined by $p \perp (\lambda_p \mathbf{e}_1 + \mathbf{e}_2)$, due to the above restrictions on Λ the number λ_p is correctly defined).

One may assume that the points in Π lie on the line $\langle \mathbf{e}_1, p \rangle = -1$, The function $\lambda_p \equiv \langle \mathbf{e}_2, p \rangle$ is a coordinate on this line. Restrictions of homogeneous polynomials of degree k on this line are polynomials of degree k or less in λ_p . Thus the above subspace \mathbf{W} is described by the following obvious

Lemma 7.6. *Consider a subset $\Pi_0 \subset \mathbb{K}$. Consider two polynomials φ_1, φ_2 of degree $\leq k$ in $\mathbb{K}[x]$ such that $\varphi_2(p) = p\varphi_1(p)$ if $p \in \Pi_0$. If $\text{card}(\Pi_0) \geq k+2$, then $\varphi_2 = x\varphi_1$.*

This lemma shows that \mathbf{W} coincides with the set of polynomials of the form $(\mathbf{s}_1\varphi, -\mathbf{s}_2\varphi)$, here $\varphi \in \text{Sym}^{k-1} \mathcal{S}$. Thus $\mathbf{W} = \varepsilon_{\Pi!} \text{Im}(m_{\mathbf{s}_1}, -m_{\mathbf{s}_2})$, hence $(\alpha^*)_! \widetilde{W}^\perp = W_0^\perp \cap \text{Im}(\alpha^* \otimes \alpha^*)$, which implies Lemma 7.5. \square

This finishes the proof of Amplification 7.2 and Theorem 7.1. \square

8. LAGRANGIAN FOLIATIONS

Here we recall more results of symplectic geometry which will be useful in Section 9. See [3] for details.

Definition 8.1. Given a bracket $\{, \}$ on M , say that a submanifold $L \subset M$ is *involutive* if $\{f, g\}|_L = 0$ for any functions f and g on M such that²⁰ both $f|_L$ and $g|_L$ are constant. Say that L is *Lagrangian* if it is involutive, and any submanifold $L_1 \subset L$ of codimension 1 or more is not involutive. Say that a foliation \mathcal{F} on M is *Lagrangian* if each leaf of \mathcal{F} is Lagrangian.

In the context of Example 0.6 the foliation on fibers of the projection π is Lagrangian. The 0-section of π is a Lagrangian submanifold. In fact, on a symplectic

²⁰In complex-analytic situation one needs to consider functions on open subsets of M .

manifold M any Lagrangian submanifold L has $\dim L = \frac{\dim M}{2}$, a coisotropic submanifold is Lagrangian if it has this dimension, and locally any Lagrangian foliation can be reduced to the foliation on fibers of the projection π of Example 0.6:

Definition 8.2. A *locally-affine* structure on a manifold L is a connection in $\mathcal{T}L$ with vanishing curvature and torsion.

Proposition 8.3. Consider a symplectic Poisson structure on M , and a Lagrangian foliation \mathcal{F} . Suppose that leaves of \mathcal{F} are fibers²¹ of a projection $\pi: M \rightarrow N$. Let L be a leaf of the foliation \mathcal{F} , $\pi(L) = \{n\} \subset N$. Then all the tangent spaces to L are identified with \mathcal{T}_n^*N , this identification provides L with a locally-affine structure.

Consider a section $s: N \rightarrow M$ of the projection π . If the $\text{Im } s \subset M$ is a Lagrangian submanifold, there is exactly one identification of a neighborhood of $\text{Im } s$ with an open subset of \mathcal{T}^*N which is compatible with the projection on N , with Poisson structures on M and \mathcal{T}^*N , and which sends $\text{Im } s$ to the 0-section of \mathcal{T}^*N .

The above identification sends the locally-affine structure on leaves of \mathcal{F} to the locally-affine structure on vector spaces \mathcal{T}_nN , $n \in N$.

Remark 8.4. Obviously, the connection of a locally-affine structure on a contractible set is isomorphic to the tautological connection in $\mathcal{T}U$ for an appropriate open subset U of a vector space. Thus another way to define a locally affine structure is to introduce identifications of open subsets U_i which cover L with open subsets of vector spaces, and require that the transition functions on $U_i \cap U_j$ correspond to affine mappings of these vector spaces.

Definition 8.5. Consider a small open connected subset U of a manifold L with a locally-affine structure. Then tangent spaces at different points $l \in U$ are identified with each other (such an identification may depend on a choice of homotopy type of a curve which connects these points). Call an open subset $U \subset L$ *simple* if the above identifications do not depend on the choice of homotopy class of connecting curves. If U is simple, the corresponding set of equivalence classes of tangent vectors has a vector space structure. Call this structure *the vector space* associated to a simple locally-affine structure on U .

Corollary 8.6. Consider a Lagrangian foliation \mathcal{F} with a base B . By Proposition 8.3 leaves of \mathcal{F} have a canonical locally-affine structure. The vector space associated to the leaf over $b \in B$ is canonically identified with \mathcal{T}_b^*B .

We will need an explicit construction of the identification of \mathcal{T}_mL with \mathcal{T}_n^*N , here $n = \pi(m)$. Given a Lagrangian submanifold L in a symplectic manifold M , the Hamiltonian mapping $H: \mathcal{T}_m^*M \rightarrow \mathcal{T}_mM$ reduces to a mapping $\mathcal{N}_m^*L \rightarrow \mathcal{T}_mL$, $m \in L$, which is a bijection. On the other hand, for any leaf L of any foliation there is a canonical flat connection on the normal bundle $\mathcal{N}L$ to L , thus on the dual bundle \mathcal{N}^*L . Identifying \mathcal{N}^*L with $\mathcal{T}L$, one obtains a flat connection on $\mathcal{T}L$.

²¹Locally any foliation can be represented in such a form.

Moreover, $\mathcal{N}_m^* L$ is identified with $\mathcal{T}_{\pi(m)}^* B$. Basing on these data, it is easy to construct the identification of Proposition 8.3.

Now extend this discussion to the case of Poisson structures of constant rank.

Proposition 8.7. *Any Lagrangian submanifold of a Poisson manifold M is contained in a symplectic leaf²² of M . If the Poisson structure on M is of constant rank, Lagrangian foliations on M coincide with smooth families of Lagrangian foliations, one per each symplectic leaf of M .*

Remark 8.8. If the Lagrangian foliation is a fibration $M \xrightarrow{\pi} B$, and Poisson structure on M is of constant rank, each symplectic leaf $S \subset B$ coincides with $\pi^{-1}\pi S$, thus the symplectic foliation $\tilde{\mathcal{F}}$ on M is a preimage of a foliation \mathcal{F}_0 on B . Due to Proposition 8.3, given a section s of π such that intersection of $\text{Im } s$ with each symplectic leaf S is Lagrangian in S , one can identify a neighborhood of $\text{Im } s$ with a neighborhood of the 0-section of $\mathcal{T}^*\mathcal{F}_0$. Moreover, $\mathcal{T}^*\mathcal{F}_0$ carries a natural Poisson structure, and the above identification is compatible with Poisson structures on $\mathcal{T}^*\mathcal{F}_0$ and M .

9. TWO FUNCTORS

Presently we have two constructions: by Corollary 4.21, given a Kronecker web \mathcal{W} on B of rank r , one can construct a bihamiltonian structure on the total space of $\Phi(\mathcal{W})$ (which has dimension $2 \dim B - r$). By Theorem 6.7, given a micro-Kronecker bihamiltonian structure M of dimension d and rank r , one can associate to small open subsets $U \subset M$ a Kronecker web structure on manifold \mathcal{B}_U (of dimension $\frac{d+r}{2}$). Investigate the relation of these two constructs.

Proposition 9.1. *Consider a Kronecker web \mathcal{W} of rank r on a manifold B .*

1. *If $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$, then the symplectic foliation of the Poisson bracket $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ on $\Phi(\mathcal{W})$ is the preimage of the integrating foliation $\mathcal{F}_{\lambda_1: \lambda_2}$ on B .*
2. *The bihamiltonian structure on $\Phi(\mathcal{W})$ is micro-Kronecker of rank r .*
3. *The action foliation on $\Phi(\mathcal{W})$ coincides with foliation on fibers of projection $\Phi(\mathcal{W}) \rightarrow B$.*
4. *The structure of Kronecker web on B induced by the bihamiltonian structure on $\Phi(\mathcal{W})$ coincides with the initial Kronecker web structure \mathcal{W} on B .*

Proof. The first statement follows from the definition of the Poisson bracket $\{, \}_{\lambda_1: \lambda_2}$ on $\Phi(\mathcal{W})$. If $\mathbb{K} = \mathbb{C}$, the second statement is a direct corollary of the first one. If $\mathbb{K} = \mathbb{R}$, the second statement follows from the following

Lemma 9.2. *Consider a pair of skewsymmetric bilinear forms $(,)_1$ and $(,)_2$ in a vector space V . Then $\dim \text{Ker}(\lambda_1 (,)_1 + \lambda_2 (,)_2)$ is constant for (λ_1, λ_2) inside an*

²²It is possible to define what is a symplectic leaf of an arbitrary Poisson structure. However, for the purpose of our discussion, it is enough to restrict attention to Poisson structures of constant rank.

open subset of \mathbb{K}^2 , call this common value r . Suppose that for any finite subset $\Lambda_0 \subset \mathbb{K}\mathbb{P}^1$ there is $\Lambda \subset \mathbb{K}\mathbb{P}^1 \setminus \Lambda_0$ such the vector subspaces $\text{Ker}(\lambda_1(\cdot, \cdot)_1 + \lambda_2(\cdot, \cdot)_2)$, $\lambda_1 : \lambda_2 \in \Lambda$, span a vector subspace of V of dimension $\frac{\dim V + r}{2}$. Then the pair $(\cdot, \cdot)_1$, $(\cdot, \cdot)_2$ is Kronecker.

Proof. Lemma follows immediately from the classification of Theorem 5.1. \square

The remaining statements of the proposition follow immediately from the first two statements. \square

Proposition 9.3. *Consider a micro-Kronecker bihamiltonian structure on a manifold M . Then*

1. *the fibers of action foliations have a natural locally-affine structure;*
2. *suppose that the action foliation is a fibration with a base \mathcal{B} , and the locally-affine structures on fibers are simple. Denote by E the vector bundle over \mathcal{B} which is associated²³ to the above bundle of locally-affine structures. This vector bundle is canonically isomorphic to $\Phi(\mathcal{B})$.*

Proof. Consider Hamiltonian mappings $H_{1,2}: \mathcal{T}_m^*M \rightarrow \mathcal{T}_mM$ of Poisson structures on M . Restrict these mappings to the action subspace $\mathcal{A}_m \subset \mathcal{T}_m^*M$. By Proposition 5.3 the action subspace is isotropic with respect to both pairings in \mathcal{T}_m^*M , thus the elements $H_{1,2}a$, $a \in \mathcal{A}_m$, are perpendicular to \mathcal{A}_m . By definition, \mathcal{A}_m^\perp coincides with the tangent space to the fiber L_m of the action foliation which passes through m . In particular, there are two mappings $\tilde{H}_{1,2}: \mathcal{A}_m \rightarrow \mathcal{T}_mL_m$. Note that these mappings form a pencil which corresponds to the linear relation in \mathcal{A}_m (as in Proposition 5.4 and in the proof of Theorem 6.7).

Since the statement of the proposition is local on M , one can suppose that the action foliation is a fibration. Let $\pi: M \rightarrow \mathcal{B}$ be the projection to the base of foliation, and $b = \pi(m)$. Then π^* induces a canonical isomorphism $\mathcal{T}_b^*\mathcal{B} \simeq \mathcal{A}_m$. Now \mathcal{B} is a Kronecker web, thus $\mathcal{T}_b^*\mathcal{B}$ carries a Kronecker relation, and this relation is compatible with the relation on \mathcal{A}_m w.r.t. the above isomorphism. On the other hand, let Φ_b be the fiber of $\Phi(\mathcal{B})$ over b . Then the linear relation in $\mathcal{T}_b^*\mathcal{B}$ can be described both by a pencil $\tilde{\mathcal{P}}_{1,2} = \tilde{H}_{1,2} \circ \pi^*: \mathcal{T}_b^*\mathcal{B} \rightarrow \mathcal{T}_mL_m$ and by a pencil $\mathcal{P}_{1,2}: \mathcal{T}_b^*\mathcal{B} \rightarrow \Phi_b$. Proposition 2.4 gives a canonical isomorphism between \mathcal{T}_mL_m and Φ_b , $b = \pi(m)$. As a corollary there is a canonical identification of tangent spaces to L_m at different points (since they all project to the same point of \mathcal{B}). This induces a flat connection on the tangent bundle to L_m .

What remains is to show that this connection on L_m is a locally-affine structure. Pick up any $\lambda \in \mathbb{P}^1$, say, $\lambda = 1 : 0$. Consider the corresponding Poisson structure $1\{\cdot, \cdot\}_1 + 0\{\cdot, \cdot\}_2$ of the pencil.

By Proposition 5.3, $\{\pi^*\varphi_1, \pi^*\varphi_2\}_1 = 0$ for any functions φ_1, φ_2 on \mathcal{B} . This implies that fibers of \mathcal{F} are coisotropic submanifolds of M w.r.t. $\{\cdot, \cdot\}_1$. On the other hand,

²³As in Definition 8.5.

if the rank of bihamiltonian structure on M is r , then symplectic leaves of M have dimension $\dim M - r$, and L_m has dimension $\frac{\dim M - r}{2}$. Since L_m has half the dimension of the symplectic leaf, and is coisotropic, it is Lagrangian. Thus \mathcal{F} is a Lagrangian foliation.

In particular, leaves of \mathcal{F} are equipped with locally-affine structures. Thus it is enough to prove that the constructed above connection on L_m coincides with the connection of this locally-affine structure. Recall that the connection on L_m can be described by mappings $\tilde{\mathcal{P}}_1: \mathcal{T}_b^* \mathcal{B} \rightarrow \mathcal{T}_m L_m$, in other words, $\mathcal{T}_b^* \mathcal{B} / \text{Ker} \left(\tilde{\mathcal{P}}_1 \right) \simeq \mathcal{T}_m L_m$. However, $\mathcal{T}_b^* \mathcal{B} / \text{Ker} \left(\tilde{\mathcal{P}}_1 \right)$ coincides with $\mathcal{T}_b^* F_b$, here F_b is the fiber of integrating foliation $\mathcal{F}_{1,0}$ which passes through $b \in \mathcal{B}$. Let $\tilde{F} = \pi^{-1} F_b$, recall that \tilde{F} is a symplectic leaf of the Poisson structure $\{, \}_1$ on M . The foliation \mathcal{F} allows a restriction $\mathcal{F}|_{\tilde{F}}$ to \tilde{F} , this restriction is a Lagrangian foliation on a symplectic manifold. Now it should be obvious that the the constructed above identification of $\mathcal{T}_b^* F_b$ coincides with the identification described in Section 8, which finishes the proof. \square

Remark 9.4. In the above proof we worked with two locally-affine structures on L_m : one constructed basing on the bihamiltonian structure, another based on the Poisson structure $\{, \}_1$. However, one could also consider another Poisson structure $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$. The fact that two locally-affine structures of the proof coincide shows that the locally-affine structures on L_m which correspond to the Poisson structure of the pencil $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ do not depend on $(\lambda_1, \lambda_2) \neq (0, 0)$.

10. CONJECTURE ON CLASSIFICATION

Conjecture 10.1. Consider a micro-Kronecker bihamiltonian structure on a manifold M . Suppose that the restriction of the action foliation to $U_0 \subset M$ is a fibration with a base \mathcal{B}_{U_0} , $m \in U_0$, and b is the projection of m to \mathcal{B} . Let \mathcal{W} be the Kronecker web on \mathcal{B}_{U_0} induced by the bihamiltonian structure on M . Let $m' \in \Phi(\mathcal{W})$ be the point on the 0-section of $\Phi(\mathcal{W})$ over b . Then the bihamiltonian structures on M and $\Phi(\mathcal{W})$ are locally isomorphic near m and m' . (In other words, there are neighborhoods U, U' of m and m' and a diffeomorphism $\alpha: U \rightarrow U'$ such that $\alpha(m) = m'$, α commutes with projections of U and U' to \mathcal{B} , and α identifies restrictions of bihamiltonian structures on M and $\Phi(\mathcal{W})$ to U and U' .)

This conjecture is a refinement of a conjecture from [10]. The case of analytic bihamiltonian structure on M of rank 1 is claimed in [12], the C^∞ -case of rank 1 is claimed in [31].

It looks like this conjecture would easily imply all the conjectures of [10]. (See how we prove Theorem 14.22.)

Remark 10.2. The conjecture says that to check that two micro-Kronecker bihamiltonian structures on M and M' are locally isomorphic, it is enough to check that the

structures of Kronecker webs on local bases of action foliations are locally isomorphic. By Amplification 7.2 to check an isomorphism of Kronecker webs on B and B' it is enough to restrict attention to a finite number of foliations on B and B' . It so happens that for webs which appear in problems of mathematical physics the latter problem is very easy (much easier than for generic Kronecker webs).

Remark 10.3. Yet another reformulation of the conjecture is that to check that two micro-Kronecker bihamiltonian structures on M and M' are locally isomorphic one should concentrate attention on *actions* (or first integrals) of these structures, and one can completely ignore the question of *phase variables*. Note that this enormously simplifies the question: say, usually first integrals may be explicitly written as polynomials in appropriate coordinates, while all one can say about phase variables is that they satisfy some partial differential equations, and that they may be expressed in terms of ϑ -functions.

11. ANTIINVOLUTIONS

Definition 11.1. An *involution* is a mapping $f: X \rightarrow X$ such that $f \circ f = \text{id}$. Say that an involution $\alpha: M \rightarrow M$ is an *antiinvolution* of a Poisson structure $\{, \}$ on a manifold M if $\{\alpha^*f, \alpha^*g\} = -\alpha^*\{f, g\}$ for any two functions f, g on M . An *antiinvolution* of a bihamiltonian structure on a manifold M is a mapping $M \rightarrow M$ which is an antiinvolution of both Poisson brackets $\{, \}_1, \{, \}_2$.

Proposition 11.2. *Let \mathcal{W} be a Kronecker web on a manifold B . The bihamiltonian structure on $\Phi(\mathcal{W})$ allows an antiinvolution.*

Proof. $\Phi(\mathcal{W})$ is a vector bundle over B . Let α be the multiplication by -1 in this bundle. To show that α is an antiinvolution, note that $\{, \}_1$ is isomorphic to the Poisson structure on $\mathcal{T}^*\mathcal{F}_{1,0}$ (here $\mathcal{F}_{1,0}$ is the integrating foliation on B), and this isomorphism is a mapping of vector bundles. Thus it is enough to show that multiplication by -1 is an antiinvolution on the cotangent bundle to a foliation. In turn, it is enough to prove this for the cotangent bundle to a manifold, which can be checked immediately in local coordinates. \square

Consider an antiinvolution ι of a micro-Kronecker bihamiltonian structure on a manifold M . Let $m \in M$ be a fixed point of ι . There is an ι -invariant open subset $U \ni m$ such that the restriction of the action foliation \mathcal{F} on U is a fibration, denote the base of this fibration by \mathcal{B} . Since the action foliation is determined by the *set* of linear combinations $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ of Poisson structures, and not by the parameterization (λ_1, λ_2) of this set, ι sends leaves of \mathcal{F} to leaves of \mathcal{F} . Thus ι induces an involution $\iota_{\mathcal{B}}$ of \mathcal{B} . Consider the submanifold²⁴ $\text{Fix}(\iota_{\mathcal{B}}) \subset \mathcal{B}$ of fixed points of $\iota_{\mathcal{B}}$.

²⁴Recall that the space of fixed point of an involution $i: M \rightarrow M$ on a manifold M is a submanifold $\text{Fix}(i) \subset M$, and for $f \in \text{Fix}(i)$ the tangent space to $\text{Fix}(i)$ at f coincides with the invariant subspace of the differential $i_*|_f$ of i . Indeed, consider an i -invariant Riemannian metric on M , then the exponential mapping of this mapping commutes with i .

Definition 11.3. The *defect* of antiinvolution ι at the fixed point $m \in M$ is the codimension of $\text{Fix}(\iota_{\mathcal{B}}) \subset \mathcal{B}$ at the image $b \in \mathcal{B}$ of m .

Obviously, the antiinvolution of Proposition 11.2 has defect 0. Now Conjecture 10.1 immediately implies

Conjecture 11.4. Consider a micro-Kronecker bihamiltonian structure on a manifold M , $m \in M$. There is a neighborhood $U \ni m$ and an antiinvolution α of the bihamiltonian structure on U such that $\alpha(m) = m$ and the defect of α at m is 0.

Antiinvolutions with defect 0 are important for us because of the following

Theorem 11.5. *Suppose that a bihamiltonian structure on a manifold M allows an antiinvolution ι such that $\iota(m) = m$, $m \in M$, and the defect of ι at m is 0. Then Conjecture 10.1 holds for m and M .*

The theorem is an immediate corollary of Propositions 11.7, 11.8 below.

Definition 11.6. Consider a micro-Kronecker bihamiltonian structure on M with the action foliation \mathcal{F} . Say that a submanifold $S \subset M$ is a *cross-section* if S is a transversal section of foliation \mathcal{F} , and for any $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$ and any symplectic leaf L of $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ the intersection $L \cap S$ is Lagrangian in L .

Proposition 11.7. *Given a micro-Kronecker bihamiltonian structure on M with a base \mathcal{B} of the action foliation, and a cross-section S , there is a local isomorphism of bihamiltonian structures M and $\Phi(\mathcal{B})$ which sends S to 0-section of $\Phi(\mathcal{B})$.*

Proof. Given a locally-affine structure L and a point $l \in L$, the exponential mapping of the connection on $\mathcal{T}L$ gives a canonical identification of a neighborhood of l in L and a neighborhood of 0 in $\mathcal{T}_l L$. Similarly, given a bundle $\mathcal{L} \rightarrow B$ with fibers having a locally-affine structure, a section s of this bundle, one obtains a canonical identification of a neighborhood of $\text{Im } s \subset \mathcal{L}$ with a neighborhood of 0-section in an appropriate vector bundle $\mathcal{E} \rightarrow B$. Obviously, fibers of \mathcal{E} are vertical tangent space of \mathcal{L} at points of $\text{Im } s$.

In particular, any submanifold $S \subset M$ which is transversal to leaves of \mathcal{F} , and such that $\dim S = \text{codim } \mathcal{F}$, gives a natural identification γ of a neighborhood of S with a neighborhood of 0-section of $\Phi(\mathcal{B})$ (here one can take S instead of the local base \mathcal{B}). We need to show that if S is a cross-section, then this identification is compatible with bihamiltonian structures on M and on $\Phi(\mathcal{B}_U)$.

It is enough to show that γ is compatible with $\{, \}_1$ (the same argument will be applicable to $\{, \}_2$). Consider a symplectic leaf $\tilde{L} \subset M$ for $\{, \}_1$. It is a preimage of a submanifold $L \subset \mathcal{B}_M$. The restriction foliation $\mathcal{F}|_{\tilde{L}}$ is well-defined and is a Lagrangian foliation. Moreover, $L \cap S$ is Lagrangian. Consider the identification of a neighborhood of $\tilde{L} \cap S$ in \tilde{L} with a neighborhood of 0-section of \mathcal{T}^*L (see Proposition 8.3). Since $\tilde{L} \cap S$ is Lagrangian w.r.t. $\{, \}_1$, this identification is compatible

with the Poisson structure $\{, \}_1$. Composing this identification with γ , one obtains an identification $\tilde{\gamma}$ of \mathcal{T}^*L with a submanifold of $\Phi(\mathcal{B}_M)$.

Obviously, $\tilde{\gamma}$ sends a fiber of \mathcal{T}^*L to a fiber of $\Phi(\mathcal{B}_M)$, and 0-section of \mathcal{T}^*L into a 0-section of $\Phi(\mathcal{B}_M)$. Due to Remark 9.4, $\tilde{\gamma}$ is compatible with affine structures on fibers of \mathcal{T}^*L and of $\Phi(\mathcal{B}_M)$, and since it sends 0 to 0, it is a linear mapping of vector bundles $\mathcal{T}^*L \xrightarrow{\tilde{\gamma}} \Phi|_L$. However, given $(\lambda_1, \lambda_2) \in \mathbb{K}^2 \setminus (0, 0)$, $\Phi|_L$ is canonically isomorphic to $\mathcal{T}^*\mathcal{F}_{\lambda_1, \lambda_2}$ (as in Section 4), and by Proposition 9.1 L is a leaf of the foliation $\mathcal{F}_{1,0}$. Thus there is a canonical isomorphism $i: \mathcal{T}^*L \rightarrow \Phi|_L$ which is compatible with the Poisson structure $\{, \}_1$ on Φ .

It is enough to show that $\tilde{\gamma} = i$. But this is a direct corollary of the proof of Proposition 9.3. \square

Proposition 11.8. *Given a micro-Kronecker bihamiltonian structure on M . If ι is an antiinvolution of M of defect 0, then $\text{Fix}(\iota)$ is a cross-section.*

Proof. This proposition immediately follows from the following

Lemma 11.9. *Consider a symplectic manifold M with a Lagrangian foliation \mathcal{F} and an antiinvolution $\iota: M \rightarrow M$ which sends each leaf of \mathcal{F} into itself. Then fixed points of ι form a Lagrangian submanifold of M which is transversal to leaves of \mathcal{F} .*

Proof. It is enough to restrict attention to a neighborhood of $\text{Fix}(\iota)$, thus one may assume that M is an open subset of \mathcal{T}^*B , and \mathcal{F} is the foliation on fibers of projection $M \rightarrow B$. Since the locally-affine structure on fibers of a Lagrangian foliation does not change if one multiplies the Poisson bracket by a number, the restriction of ι on any leaf of \mathcal{F} induces an affine transformation of this leaf (which is an open subset of \mathcal{T}_bB for an appropriate $b \in B$).

Recall that the set of fixed points of an involution ι of M forms a manifold $F \subset M$, and if $f \in F$, then $\mathcal{T}_fF \subset \mathcal{T}_fM$ coincides with the invariant subspace of $(d\iota)|_{\mathcal{T}_fM}$.

Lemma 11.10. *Consider a symplectic vector space V , and a Lagrangian subspace $l \subset V$. Consider a linear involution $i: V \rightarrow V$ such that $[iv_1, iv_2] = -[v_1, v_2]$ for any $v_1, v_2 \in V$. Suppose that $il = l$, and that i induces an identity mapping of V/l into itself. Then $i|_l$ is multiplication by -1 , and l is a complement to the invariant subspace of i in V , which is a Lagrangian subspace of V .*

Proof. Let V_1 is the vector subspace of fixed points of i , V_{-1} be the eigenspace of i with eigenvalue -1 . Obviously $V = V_1 \oplus V_{-1}$, and due to the conditions on i both V_1 and V_{-1} are isotropic. Thus both V_1 and V_{-1} are Lagrangian. Similarly, $l = l_1 \oplus l_{-1}$, $l_1 \subset V_1$, $l_{-1} \subset V_{-1}$. Let l'_λ be any complementary subspace to l_λ in V_λ , $\lambda \in \{1, -1\}$. Since the action of i in V/l is isomorphic to the action of i in $l'_1 \oplus l'_{-1}$, we see that $l'_{-1} = 0$, thus $l \supset V_{-1}$. Since l is Lagrangian, $l = V_{-1}$, which finishes the proof. \square

Applying this lemma to $V = \mathcal{T}_fM$, $l = \mathcal{T}_fL$ (here $f \in M$ is such that $if = f$, L is the leaf of \mathcal{F} through f) finishes the proof of Lemma 11.9. \square

This finishes the proof of Proposition 11.8. \square

12. REAL AND COMPLEX BIHAMILTONIAN STRUCTURES

Consider a real-analytic bihamiltonian structure $\{\cdot, \cdot\}_{1,2}$ on M . One can construct a *small complex-analytic* neighborhood $M_{\mathbb{C}} \supset M$ of the manifold M such that the brackets $\{\cdot, \cdot\}_{1,2}$ can be analytically continued on $M_{\mathbb{C}}$. If the bihamiltonian structure on M is micro-Kronecker, one may assume the same about $M_{\mathbb{C}}$ (possibly, after decreasing $M_{\mathbb{C}}$).

Proposition 12.1. *Let $m \in M \subset M_{\mathbb{C}}$, suppose that $U \subset M_{\mathbb{C}}$, $m \in U$, allows an antiinvolution $\iota: U \rightarrow U$ with $m = \iota m$ and defect 0 near m . Then there is a neighborhood U_1 of m in M which allows an antiinvolution $\iota': U' \rightarrow U'$ with $m = \iota' m$ and defect 0 near m .*

Proof. Recall that

Definition 12.2. An *antiholomorphic mapping* $C: N \rightarrow N'$ of complex analytic manifolds N and N' is such a mapping of sets that $\overline{C^*\varphi}$ is a holomorphic function on N if φ is a holomorphic function on N' .

Clearly, if C is antiholomorphic, and $Z \subset N'$ is a complex-analytic subvariety, then $C^{-1}Z$ is a complex analytic subvariety. Indeed, if $\varphi = 0$ is an equation of Z , then $\overline{C^*\varphi} = 0$ is an equation of $C^{-1}Z$. Similarly, one can transfer a Kronecker structure and a bihamiltonian structure via an antiholomorphic bijection. If $N_{\mathbb{C}}$ is a complexification of a real-analytic manifold N , then $N_{\mathbb{C}}$ is equipped with an antiholomorphic involution C such that $\text{Fix}(C) = N$. If N has a Kronecker or bihamiltonian structure, so has $N_{\mathbb{C}}$, and the structures on $N_{\mathbb{C}}$ are invariant w.r.t. C .

Consider the antiholomorphic involution C for $M_{\mathbb{C}}$, $\text{Fix}(C) = M$. Consider a cross-section σ of the projection $\pi_{\mathbb{C}}$ which passes through m . Then $C\sigma$ is also a cross-section which passes through m . Since fibers of \mathcal{F} have a locally-affine structure, the section $\frac{\sigma + C\sigma}{2}$ is well defined on an appropriate neighborhood of $\pi m \in \mathcal{B}$. Moreover, this section is a cross-section. Indeed, this immediately follows

Lemma 12.3. *Consider a Lagrangian foliation \mathcal{F} on a symplectic manifold M , and submanifolds S_1, S_2, S_3 of dimension $\frac{\dim M}{2}$ which are transversal to \mathcal{F} . Suppose that for any leaf L of \mathcal{F} the intersection $S_i \cap L$, $i = 1, 2, 3$, consists of one point p_i , and p_2 is the midpoint of the segment p_1p_3 in the locally affine structure on L . If L_1 and L_3 are Lagrangian, so is L_2 .*

Proof. Indeed, we may assume that $M = \mathcal{T}^*N$, leaves of \mathcal{F} are fibers of $\pi: M \rightarrow N$, S_1 is 0-section of \mathcal{T}^*N , $S_{2,2}$ are graphs of sections $\varepsilon_{2,3}$ of $\mathcal{T}^*N = \Omega^1N$, and $\varepsilon_3 = 2\varepsilon_2$.

Lemma 12.4. *Consider a section ε of \mathcal{T}^*B . The graph of ε (which is a submanifold of \mathcal{T}^*B) is Lagrangian iff $d\varepsilon = 0 \in \Omega^2B$.*

Application of this obvious lemma finishes the proof. □

Now the section $\frac{\sigma + C\sigma}{2}$ of π is C -invariant, thus it is a complexification of the section $\sigma_{\mathbb{R}}$ for M . Since the complexification of $\sigma_{\mathbb{R}}$ is a cross-section, so is $\sigma_{\mathbb{R}}$. □

13. ENDOMORPHISMS OF BIHAMILTONIAN STRUCTURES AND THE PRINCIPAL THEOREM

The following statement is widely known:

Proposition 13.1. *Suppose that (λ_1, λ_2) and (λ'_1, λ'_2) are two non-proportional vectors in \mathbb{K}^2 . Consider a bihamiltonian structure $\{, \}_1, \{, \}_2$ on a manifold M . Suppose that a function F on M is a Casimir function for the bracket $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$. Let $\chi \in \text{Vect } M$ be the Hamiltonian vector field²⁵ of F w.r.t. the bracket $\lambda'_1 \{, \}_1 + \lambda'_2 \{, \}_2$. Consider the flow α_t of χ in time $t \in \mathbb{K}$, as a mapping $\alpha_t: U_1 \rightarrow U_2$, here $U_{1,2}$ are open subsets of M . Consider $U_{1,2}$ with restriction bihamiltonian structures. Then α_t is an isomorphism of bihamiltonian structures.*

Proof. Indeed, the Hamiltonian vector field of any function w.r.t. a Poisson bracket $\{, \}$ preserves $\{, \}$. Thus the bracket $\lambda'_1 \{, \}_1 + \lambda'_2 \{, \}_2$ is preserved by χ , thus by α_t . On the other hand, due to the condition on F , χ is proportional to the Hamiltonian flow of F w.r.t. any bracket $\lambda''_1 \{, \}_1 + \lambda''_2 \{, \}_2$ (as far as $\lambda''_1 : \lambda''_2 \neq \lambda_1 : \lambda_2$), thus χ preserves the Poisson structure $\lambda''_1 \{, \}_1 + \lambda''_2 \{, \}_2$ as well. By linearity, it preserves $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$ too. \square

Definition 13.2. For $m \in M$ say that $v \in \mathcal{T}_m M$ is a *biflow* vector if on a small open subset $U \subset M$, $U \ni m$, the vector v can be represented as a value of χ at m , here χ is a vector field from Proposition 13.1.

Theorem 13.3. *Consider a micro-Kronecker bihamiltonian structure on a manifold M . Let $m \in M$, let L be the leaf of action foliation on M which passes through m . If $m' \in L$, then there are neighborhoods U, U' of m and m' and a diffeomorphism $\alpha: U \rightarrow U'$ which*

1. *sends the restriction bihamiltonian structure on U to the restriction bihamiltonian structure on U' ;*
2. *for $n \in U_1$ the point $\alpha(n)$ is on the same leaf of action foliation as n .*

Proof. Due to Proposition 13.1 it is enough to show that the span of biflow vectors in $\mathcal{T}_m M$ coincides with $\mathcal{T}_m L$. Decrease M so that the action foliation becomes a fibration $\pi: M \rightarrow \mathcal{B}$ with a base \mathcal{B} . Let $b = \pi(m)$.

Lemma 13.4. *Consider the pencil $\tilde{\mathcal{P}}_{1,2}: \mathcal{T}_b^* \mathcal{B} \rightarrow \mathcal{T}_m L$ from the proof of Proposition 9.3. A vector $v \in \mathcal{T}_m M$ is a biflow vector iff v can be written as $(\lambda'_1 \tilde{\mathcal{P}}_1 + \lambda'_2 \tilde{\mathcal{P}}_2) \alpha$ with $(\lambda_1 \tilde{\mathcal{P}}_1 + \lambda_2 \tilde{\mathcal{P}}_2) \alpha = 0$, $\lambda_1, \lambda_2, \lambda'_1, \lambda'_2 \in \mathbb{K}$, $(\lambda_1, \lambda_2) \neq (0, 0)$, $\alpha \in \mathcal{T}_b^* \mathcal{B}$.*

Proof. If f is a function on \mathcal{B} , and χ is the Hamiltonian flow of $f \circ \pi$ w.r.t. the Poisson bracket $\lambda'_1 \{, \}_1 + \lambda'_2 \{, \}_2$, then the value of χ at $m \in M$ coincides with $(\lambda'_1 \tilde{\mathcal{P}}_1 + \lambda'_2 \tilde{\mathcal{P}}_2) (df|_b)$. This implies the “only if” part of the lemma.

²⁵I.e., $\chi = \text{H}(dF)$, here H is the Hamiltonian mapping $\mathcal{T}^*(M) \rightarrow \mathcal{T}(M)$.

On the other hand, if $(\lambda_1 \tilde{\mathcal{P}}_1 + \lambda_2 \tilde{\mathcal{P}}_2) \alpha = 0$, then α is normal to the leaf of the integrating foliation $\mathcal{F}_{\lambda_1: \lambda_2}$ on \mathcal{B} . Decreasing \mathcal{B} , one can find a function f on \mathcal{B} such that f is constant on leaves of $\mathcal{F}_{\lambda_1: \lambda_2}$, and $df|_{\mathcal{B}} = \alpha$. This implies the “if” part of the lemma. \square

Lemma 13.5. *Consider a Kronecker relation in a vector space V , and the associated pencil $\mathcal{P}_{1,2}: V \rightarrow V'$. Then vectors $v' \in V'$ of the form $(\lambda'_1 \mathcal{P}_1 + \lambda'_2 \mathcal{P}_2)v$ span V' , here $v \in V$ and $(\lambda_1 \mathcal{P}_1 + \lambda_2 \mathcal{P}_2)v = 0$, $\lambda_1, \lambda_2, \lambda'_1, \lambda'_2 \in \mathbb{K}$, $(\lambda_1, \lambda_2) \neq (0, 0)$.*

Proof. We may assume $\lambda_1 = 1$, $\lambda_2 = 0$. Since $\mathcal{P}_1 V = V'$, it is enough to show that vectors $v \in V$ such that $(\lambda_1 \mathcal{P}_1 + \lambda_2 \mathcal{P}_2)v = 0$ for an appropriate $\lambda_1, \lambda_2 \in \mathbb{K}$ span V . But this is a corollary of Lemma 6.3. \square

Lemma 13.6. *In conditions of Proposition 13.1 assume that the bihamiltonian structure on M is micro-Kronecker, and \mathcal{F} is the action foliation. Then for any $m \in U_1$ points m and $\alpha_t m$ are on the same leaf of \mathcal{F} .*

Proof. It is enough to show that biflow vectors are tangent to leaves of action foliation. This is a statement of linear algebra, it follows directly from the explicit description of Kronecker blocks. \square

This finishes the proof of Theorem 13.3. \square

Theorem 13.7. *Consider a micro-Kronecker bihamiltonian structure on a manifold M and an antiinvolution α of M . Let Z be the set of fixed points of α of defect 0. Let U be the union of leaves of the action foliation of M which pass through points of Z . Then*

1. *The subset $U \subset M$ is open;*
2. *If $m \in U$, then Conjecture 10.1 holds for m and M .*

Proof. This is an immediate corollary of Theorems 11.5 and 13.3. \square

Proposition 11.2 and Theorem 13.7 show that in fact Conjectures 10.1 and 11.4 are equivalent to each other. However, the rôles of these conjecture are very dissimilar. As we will show in Section 14, for some particular bihamiltonian structures of mathematical physics Conjecture 11.4 is easy to verify by an explicit construction (see Theorem 14.14), thus for these structures Conjecture 10.1 *follows* from this construction.

On the other hand, as it was shown in [12], in the case an arbitrary bihamiltonian structure of rank 1 it is possible to prove Conjecture 10.1 using some “hard” cohomological statements. One can expect that a similar approach may succeed in the case of higher rank as well. But currently it is not clear how one could prove Conjecture 11.4 without a reference to Conjecture 10.1 (or the calculation of some cohomology which will immediately prove Conjecture 10.1).

This suggests that in some particular cases it is easier to directly deduce the statement of Conjecture 11.4, but in general case Conjecture 10.1 should be easier to tackle.

14. METHOD OF ARGUMENT TRANSLATION

Consider a Lie algebra \mathfrak{g} and a 2-cocycle c_2 of \mathfrak{g} . As in Example 0.10, such a pair induces a bihamiltonian structure $\{, \}_1, 2$ on \mathfrak{g}^* . Moreover, if c_2 is a coboundary of a 1-chain c_1 , one may consider c_1 as an element of \mathfrak{g}^* . Obviously, the Poisson structure $\{, \}_1$ is a Lie derivative of the Poisson structure $\{, \}_2$ in the direction of a parallel translation of \mathfrak{g}^* in the direction of $c_1 \in \mathfrak{g}^*$, and $\{, \}_1$ is translation-invariant. Thus $\{, \}_2 + \lambda \{, \}_1$ is a parallel translation of $\{, \}_2$ on λc_1 . Due to this observation consideration of the pair $\{, \}_1, 2$ when c_2 is a coboundary is often called the *method of argument translation*.

Moreover, if \mathfrak{g} is semisimple, then c_2 is automatically a coboundary.

Definition 14.1. Given $c_1 \in \mathfrak{g}^*$, let $\{, \}_1, 2$ be the bihamiltonian structure on \mathfrak{g}^* constructed based on 2-coboundary dc_1 . Say that this structure is *associated* to $c_1 \in \mathfrak{g}^*$.

Definition 14.2. Given a Lie algebra \mathfrak{g} over \mathbb{K} , let $\mathfrak{g}_{\mathbb{C}} \stackrel{\text{def}}{=} \mathfrak{g}$ if $\mathbb{K} = \mathbb{C}$, and $\mathfrak{g}_{\mathbb{C}} \stackrel{\text{def}}{=} \mathfrak{g} \otimes \mathbb{C}$ if $\mathbb{K} = \mathbb{R}$.

Lemma 14.3. Consider the Poisson structure $\{, \}_2$ on \mathfrak{g}^* and $\alpha \in \mathfrak{g}^*$. Let L be the symplectic leaf of $\{, \}_2$ through α . Then $\mathcal{N}_{\alpha}L = \text{Stab}_{\text{Ad}^*} \alpha$.

The proof of this lemma is a direct calculation.

Definition 14.4. The *rank* $\text{rk}(\mathfrak{g})$ of a Lie algebra \mathfrak{g} is $\min_{\alpha \in \mathfrak{g}^*} \dim \text{Stab}_{\text{Ad}^*} \alpha$. An element $\alpha \in \mathfrak{g}^*$ is *regular* if $\dim \text{Stab}_{\text{Ad}^*} \alpha = \text{rk}(\mathfrak{g})$, and *irregular* otherwise.

Obviously, regular elements form a (Zariski) open and dense subset of \mathfrak{g}^* .

Definition 14.5. Let $\alpha \in \mathfrak{g}^*$, $\beta \in \mathfrak{g}_{\mathbb{C}}^*$. Say that β is *compatible* with α if $\beta + \lambda\alpha$ is regular for any $\lambda \in \mathbb{C}$. Call a regular element $\alpha \in \mathfrak{g}^*$ *strongly regular* if there exists a compatible with α element of $\mathfrak{g}_{\mathbb{C}}^*$.

Definition 14.6. Say that \mathfrak{g} is *2-regular*, if the algebraic subvariety $\mathcal{I} \subset \mathfrak{g}_{\mathbb{C}}^*$ of irregular elements has codimension 2 or more.

Proposition 14.7. Suppose that there exists a strongly regular element α in $\mathfrak{g}_{\mathbb{C}}^*$. Then \mathfrak{g} is 2-regular.

Suppose that \mathfrak{g} is 2-regular, and α is a regular element of \mathfrak{g}^* . Then α is strongly regular, and the set of elements of \mathfrak{g}^* which are compatible with α is non-empty and Zariski open.

Proof. Let $\mathcal{I} \subset \mathfrak{g}_{\mathbb{C}}^*$ be the subvariety of irregular elements. Let π be the projection of $\mathfrak{g}_{\mathbb{C}}^*$ to $\mathfrak{g}_{\mathbb{C}}^*/\mathbb{K}\alpha$. The existence of an α -compatible element is equivalent to $\pi\mathcal{I} \neq \pi\mathfrak{g}_{\mathbb{C}}^*$.

If β is regular, then any non-zero scalar multiple of β is also regular. Thus one can consider a closed subvariety $\mathbb{P}\mathcal{I}$ of irregular elements in the projectivization $\mathbb{P}\mathfrak{g}^*$ of \mathfrak{g}^* . Given a strongly regular element α and a compatible element β , one obtains a line $l = \mathbb{P}\langle\alpha, \beta\rangle$ in $\mathbb{P}\mathfrak{g}^*$, and $l \cap \mathbb{P}\mathcal{I} = \emptyset$. Clearly, any nearby line l' will also not intersect $\mathbb{P}\mathcal{I}$. Thus the set of elements β which are compatible with α is Zariski open. Thus the intersection of this set with $\mathfrak{g}^* \subset \mathfrak{g}_{\mathbb{C}}^*$ is non-empty.

Since $l \cap \mathbb{P}\mathcal{I} = \emptyset$, $\mathbb{P}\mathcal{I}$ has codimension 2 or more, thus the same is true for \mathcal{I} . On the other hand, if \mathcal{I} has codimension 2 or more, then the projection of \mathcal{I} to $\mathfrak{g}^*/\mathbb{K}\alpha$ is not surjective, here $\alpha \in \mathfrak{g}^*$ is arbitrary. This implies that any regular element of \mathfrak{g}^* is strongly regular. \square

Proposition 14.8. *Suppose that $c_1 \in \mathfrak{g}^*$ is strongly regular. Then there is a dense open subset $U \subset \mathfrak{g}^*$ such that the restriction on U of the pair $\{, \}_{1,2}$ associated to c_1 is micro-Kronecker of rank $\text{rk}(\mathfrak{g})$.*

Proof. By Proposition 14.7, the set U of compatible with c_1 elements of $\mathfrak{g}_{\mathbb{C}}^*$ is Zariski open (thus dense). Show that U (or $U \cap \mathfrak{g}$ in the case $\mathbb{K} = \mathbb{R}$) satisfies the conditions of the proposition. One may assume that $\mathbb{K} = \mathbb{C}$.

We need to show that for $\beta \in U$ the symplectic leaf through β of $\lambda_1 \{, \}_1 + \lambda_2 \{, \}_2$, $(\lambda_1, \lambda_2) \neq (0, 0)$, has codimension $\text{rk}(\mathfrak{g})$. For $\lambda_2 = 0$ the normal space to this leaf coincides with $\text{Stab}_{\text{Ad}^*} c_1$, thus regularity of c_1 implies the statement. Thus we may assume $\lambda_2 = 1$. The Poisson structure $\lambda \{, \}_1 + \{, \}_2$ is a λc_1 -translation of $\{, \}_2$. If $\beta \in U$, then $\beta_1 = \lambda c_1 + \beta$ is regular.

Thus it is enough consider $\lambda = 0$. But the normal space to the leaf of symplectic foliation through β_1 is $\text{Stab}_{\text{Ad}^*} \beta_1$, which finishes the proof. \square

Proposition 14.9. *If \mathfrak{g} is reductive, then any regular element $\alpha \in \mathfrak{g}^*$ is strongly regular.*

Proof. Indeed [1], irregular elements of a semisimple Lie algebra form a Zariski closed subvariety of codimension 3. From this one can easily obtain the statement for reductive Lie algebras. \square

The above arguments are not new, see [4, 24].

The proposition cannot be inverted:

Example 14.10. (V. Serganova) For any vector space V there is a canonical symmetric pairing on $V \oplus V^*$. For any Lie algebra \mathfrak{g} this pairing on $\mathfrak{g} \oplus \mathfrak{g}^*$ is an invariant pairing on the Lie algebra $\mathfrak{G} = \mathfrak{g} \ltimes \text{ad}_{\mathfrak{g}}^*$, here $\text{ad}_{\mathfrak{g}}^*$ is the adjoint representation of \mathfrak{g} with trivial structure of Lie algebra.

Thus instead of studying dimension of stabilizers of elements of \mathfrak{G}^* , one can consider dimensions of stabilizers of elements of \mathfrak{G} . In the case of $\mathfrak{g} = \mathfrak{sl}_2$ it is easy to show that the set of irregular elements coincides with the radical of \mathfrak{G} , which has codimension 3.

Now show how one can refine the description of bihamiltonian structure on \mathfrak{g} by applying general machinery of this paper.

Definition 14.11. Say that a linear mapping $\iota: \mathfrak{g} \rightarrow \mathfrak{g}$ is an *antiinvolution* of \mathfrak{g} if ι is an involution of a vector space, and $[\iota X, \iota Y] = -\iota[X, Y]$ for any $X, Y \in \mathfrak{g}$. Say that an antiinvolution ι is *admissible*, if

1. The irregular elements in the vector subspace $\text{Fix}(\iota^*) \subset \mathfrak{g}^*$ of fixed points of ι in \mathfrak{g}^* form a subvariety of $\text{Fix}(\iota^*)$ of codimension 2 or more.
2. The subset $\mathcal{U}(\iota) \subset \text{Fix}(\iota^*)$ consisting of points $\alpha \in \text{Fix}(\iota^*)$ such that the Ad^* -orbit of α is transversal to $\text{Fix}(\iota^*)$ is not empty;

Say that $\alpha \in \mathfrak{g}^*$ is *admissible*, if α is regular, and there is an admissible antiinvolution ι such that $\iota\alpha = \alpha$.

Remark 14.12. Obviously, admissible elements exists only in 2-regular Lie algebras, and are strongly regular. Clearly, for an admissible antiinvolution ι the set $\mathcal{U}(\iota)$ is Zariski open in $\text{Fix}(\iota^*)$. Moreover, if α is admissible, and ι is the corresponding admissible antiinvolution, then elements $\beta \in \text{Fix}(\iota^*)$ which are compatible with α form a non-empty Zariski open subset of $\text{Fix}(\iota^*)$.

Theorem 14.13. *Suppose that $c_1 \in \mathfrak{g}^*$ is admissible. Consider the bihamiltonian structure $\{\cdot, \cdot\}_{1,2}$ associated to c_1 . Then there is an open subset $M \subset \mathfrak{g}^*$ such that for any $m \in M$ Conjecture 10.1 is satisfied.*

Proof. Consider an admissible antiinvolution ι such that $\iota^*c_1 = c_1$. Let U be the the Zariski open subset of $\mathfrak{g}_{\mathbb{C}}^*$ where $\{\cdot, \cdot\}_{1,2}$ is micro-Kronecker. Since ι is an antiinvolution of \mathfrak{g} , ι^* is an antiinvolution of the Poisson structure $\{\cdot, \cdot\}_2$ on \mathfrak{g}^* . Since ι^* preserves c_1 , and $\{\cdot, \cdot\}_1$ is the derivative of $\{\cdot, \cdot\}_2$ in the direction of c_1 , ι^* is an antiinvolution of $\{\cdot, \cdot\}_1$ as well. Thus U is ι^* -invariant, and $\iota^*|_U$ is an antiinvolution of a micro-Kronecker bihamiltonian structure.

Note that the same arguments as in the proof of Proposition 14.7 show that $U \cap \text{Fix}(\iota^*) \neq \emptyset$. Let $\tilde{U} = U \cap \mathcal{U}(\iota^*)$. This is a non-empty Zariski open subset of $\text{Fix}(\iota^*)$. Let $\beta \in \tilde{U}$. Show that the defect of ι^* at β is 0. Let U_1 be a neighborhood of β in U such that the action foliation \mathcal{F} of the bihamiltonian structure becomes a fibration $\pi: U_1 \rightarrow \mathcal{B}$. It is enough to show that for any function φ on \mathcal{B} the function $\varphi \circ \pi$ on U_1 is preserved by ι^* . In turn, it is enough to do the same for a large enough collection of functions φ on \mathcal{B} . Take as such collection functions φ which are constant on fibers of the integrating foliation \mathcal{F}_λ on \mathcal{B} , $\lambda \in \Lambda$ (for a sufficiently large $\Lambda \subset \mathbb{K}\mathbb{P}^1$). We may suppose $\Lambda \subset \mathbb{K}$.

On the other hand, φ being in the above subset is equivalent to $\varphi \circ \pi$ being constant on fibers of the symplectic foliation $\tilde{\mathcal{F}}_\lambda$ of the Poisson structure $\lambda\{\cdot, \cdot\}_1 + \{\cdot, \cdot\}_2$. Thus it is enough to show that ι^* preserves such functions. Since $\lambda\{\cdot, \cdot\}_1 + \{\cdot, \cdot\}_2$ is the result of translation of $\{\cdot, \cdot\}_2$ by λc_1 , it is enough to show this for $\lambda = 0$ and $\beta + \lambda c_1$, $\lambda \in \Lambda$, taken instead of β .

However, $\beta + \lambda c_1 \in \tilde{U}$ for λ in an open subset of \mathbb{K} , thus we can restrict our attention to a given $\beta \in \tilde{U}$ and $\lambda = 0$. Since $\beta \in \mathcal{U}(\iota^*)$, any Ad^* -orbit which passes near β intersects $\text{Fix}(\iota^*)$. Since ι^* sends an Ad^* -orbit to an Ad^* -orbit, this implies that ι^* preserves any Ad^* -orbit which passes near β . On the other hand, symplectic leaves of $\{, \}_2$ coincide with Ad^* -orbits in \mathfrak{g}^* . Thus on a neighborhood of β any function which is constant on symplectic leaves of $\{, \}_2$ is ι^* -invariant. Thus the defect of ι^* near β is indeed 0, and we are in conditions of Theorem 13.7. This implies the theorem for $M = U$. \square

Theorem 14.14. *Suppose that \mathfrak{g} is semisimple, and $c_1 \in \mathfrak{g}^*$ is regular and semisimple. Then there is a dense open subset $U \subset \mathfrak{g}^*$ such that the restriction on U of the pair $\{, \}_{1,2}$ associated to c_1 satisfies Conjecture 10.1.*

Proof. Identify \mathfrak{g}^* with \mathfrak{g} using the Killing form. Due to Proposition 12.1, it is enough to consider the case $\mathbb{K} = \mathbb{C}$.

Definition 14.15. Consider a semisimple Lie algebra \mathfrak{g} . Define the *Cartan antiinvolution* ι by its action on standard generators $e_i, f_i, h_i, i = 1, \dots, r$:

$$\iota(e_i) = f_i, \iota(f_i) = e_i, \iota(h_i) = h_i.$$

Lemma 14.16. *The Cartan antiinvolution of a semisimple Lie algebra is admissible.*

Proof. Identification of \mathfrak{g} with \mathfrak{g}^* allows one to consider ι instead of ι^* . Any antiinvolution sends an Ad -orbit to an Ad -orbit. Since $\text{Fix}(\iota) \supset \mathfrak{h}$, and an orbit of a regular element of \mathfrak{h} is transversal to \mathfrak{h} , ι satisfies the second condition of Definition 14.11. Thus to prove the lemma it is enough to show that irregular elements in $\text{Fix}(\iota)$ form a subvariety of codimension 2 or more. (Note that this statement is *not* true if one substitutes \mathfrak{h} instead of $\text{Fix}(\iota)$!)

Due to homogeneity of the set of regular elements, it is enough to prove this statement for an arbitrary vector subspace in $\text{Fix}(\iota)$ taken instead of $\text{Fix}(\iota)$. Recall that [5]:

Lemma 14.17. *There are numbers $a_i \neq 0, b_i \neq 0, c_i, i = 1, \dots, r$, such that for the elements $E = \sum a_i e_i, F = \sum b_i f_i, H = \sum c_i h_i$ the vector subspace $V = \langle E, F, H \rangle \subset \mathfrak{g}$ is a Lie subalgebra isomorphic to \mathfrak{sl}_2 . The adjoint action of V on \mathfrak{g} is a direct sum of r odd-dimensional irreducible representations of \mathfrak{sl}_2 .*

Since the action of any non-zero element of \mathfrak{sl}_2 in an odd-dimensional irreducible representation has 1-dimensional null-space, this shows that the stabilizer of any non-zero point of V has dimension r . But $\text{rk } \mathfrak{g} = r$, thus all the non-zero elements of V are regular. Moreover, conjugating V with elements of the Lie groups $\exp(\mathfrak{h})$ for \mathfrak{h} , one may assume that $a_i = b_i, i = 1, \dots, r$. Thus the subspace V is ι -invariant, let V_0 be the 2-dimensional invariant subspace $\langle H, E + F \rangle$ of $\iota|_V$.

Since $V_0 \subset \text{Fix}(\iota)$ intersects irregular elements on $\{0\}$, which is a subvariety of codimension 2, irregular elements in $\text{Fix}(\iota)$ form a submanifold of codimension 2 or more. This finishes the proof of Lemma 14.16. \square

Lemma 14.16 implies that any regular $c_1 \in \mathfrak{g}^*$ which is on an Ad^* -orbit of $\text{Fix}(\iota^*)$ is admissible. Since any regular semisimple element of \mathfrak{g} is conjugate to an element of \mathfrak{h} , it is on an Ad^* -orbit of an ι -invariant element, thus admissible.

Let us show that U can be taken Zariski open, thus dense. Recall that U is the union of leaves of \mathcal{F} (which is a foliation on a Zariski open subset U_0 of \mathfrak{g}) which intersect $\text{Fix}(\iota)$. Show that U contains a Zariski open subset.

There is a Zariski open subset U_1 of \mathfrak{g} such that U_1 is Ad -invariant, and Ad -invariant polynomials distinguish Ad -orbits in U_1 . Moreover, locally the fibers of the action foliation \mathcal{F} are intersections of a finite number of λc_1 -shifted Ad -orbits, $\lambda \in \Lambda$, $\text{card}(\Lambda) < \infty$. Taking a large enough finite collection p_i , $i \in I$, of invariant polynomials on \mathfrak{g} , we see that fibers of \mathcal{F} coincide with connected components of level sets of $p_{i,\lambda}$, $i \in I$, $\lambda \in \Lambda$, here $p_{i,\lambda}(X) \stackrel{\text{def}}{=} p_i(X + \lambda c_1)$. Let $U_2 = U_0 \cap \bigcap_{\lambda \in \Lambda} (U_1 - \lambda c_1)$, here $U_1 - \lambda c_1$ is the parallel translation of U_1 by $-\lambda c_1$. Obviously, U_2 is a union of fibers of foliation \mathcal{F} .

Let Π be the polynomial mapping of \mathfrak{g} to \mathbb{C}^N , $N = \text{card}(I) \text{card}(\Lambda)$, with components $p_{i,\lambda}$. Let $Y = \overline{\text{Im} \Pi}$, clearly ΠU_2 contains a Zariski open subset Y_0 of Y . Clearly, $\Pi|_{\Pi^{-1}Y_0}$ is a submersion to Y_0 . Since U contains an opens subset of \mathfrak{g} , $\Pi \text{Fix}(\iota)$ contains an open subset of Y , thus a Zariski open subset of Y . Thus one can assume that $Y_0 \subset \Pi \text{Fix}(\iota)$. Let $U_3 = \Pi^{-1}Y_0$, $Z = U_3 \cap \text{Fix}(\iota)$.

We obtain the following mappings: $Z \xrightarrow{i} U_3 \xrightarrow{\Pi} Y_0$, here U_3 is Zariski open on \mathfrak{g} , Z is Zariski closed in U_3 , Π is a polynomial mapping which is a submersion onto Y_0 , and $\Pi \circ i$ is surjective. Let U_4 be the union of connected components of fibers of Π which intersect Z . Since $U_4 \subset U$, it is enough to show that U_4 coincides with U_3 .

However, U_4 is obviously closed and open in U_3 . Since U_3 is connected (as a Zariski open subset of a vector space), $U_3 = U_4$. This finishes the proof of Theorem 14.14. \square

Amplification 14.18. *In Theorem 14.14 one can take \mathfrak{g} being reductive, and drop the condition of semisimplicity on c_1 .*

Proof. As the proof of Theorem 14.14 shows, it is enough to show that

Proposition 14.19. *Consider a reductive Lie algebra \mathfrak{g} and its Cartan antiinvolution ι . Then any element X of \mathfrak{g} is on Ad -orbit of ι -invariant element.*

Proof (V. Serganova). First of all, the statement for reductive algebras follows from the case of semisimple Lie algebras. For semisimple elements the statement is obvious (since \mathfrak{h} is ι -invariant). Consider the case when $X \in \mathfrak{g}$ is nilpotent.

One may assume that $X \neq 0$. If $\mathfrak{g} = \mathfrak{sl}_2$, then any nilpotent element is SL_2 -conjugate to $\begin{pmatrix} 1 & i \\ i & -1 \end{pmatrix}$ which is symmetric, thus ι -invariant. Thus to prove the proposition for the case of nilpotent X is enough to show

Lemma 14.20. *For any nilpotent $X \in \mathfrak{g}$ with a reductive Lie algebra \mathfrak{g} there is a subalgebra $\mathfrak{g}_0 \subset \mathfrak{g}$ such that $\mathfrak{g}_0 \simeq \mathfrak{sl}_2$, $\iota \mathfrak{g}_0 = \mathfrak{g}_0$, $\iota|_{\mathfrak{g}_0}$ is the Cartan antiinvolution of \mathfrak{sl}_2 , and X is conjugate to $X' \in \mathfrak{g}_0$. Here ι is the Cartan antiinvolution of \mathfrak{g} .*

Proof. It is enough to prove this for a semisimple \mathfrak{g} . The classification of nilpotent elements up to conjugation is well-known ([6], or it can be deduced from [5]):

Lemma 14.21. *For any nilpotent element $X \in \mathfrak{g}$, $X \neq 0$, there is a reductive subalgebra $\tilde{\mathfrak{g}}$ of \mathfrak{g} with $\text{rk}(\tilde{\mathfrak{g}}) = \text{rk}(\mathfrak{g})$ and a Cartan set of generators $\tilde{e}_i, \tilde{f}_i, i = 1, \dots, \text{rk}(\tilde{\mathfrak{g}}) - \dim Z(\tilde{\mathfrak{g}}), \tilde{h}_j, j = 1, \dots, \text{rk}(\tilde{\mathfrak{g}})$ of $\tilde{\mathfrak{g}}$ such that $X = \sum_{i=1}^{\text{rk}(\tilde{\mathfrak{g}})} \tilde{e}_i$.*

The Cartan subalgebra of $\tilde{\mathfrak{g}}$ is a Cartan subalgebra of \mathfrak{g} , thus after conjugation one can assume that \tilde{h}_j generate $\mathfrak{h} \subset \mathfrak{g}$. Then $\tilde{h}_j = \tilde{h}_j$ for any $j = 1, \dots, \text{rk}(\tilde{\mathfrak{g}})$. Since elements \tilde{e}_i are defined up to proportionality by coefficients a_{ij} given by $[\tilde{h}_j, \tilde{e}_i] = a_{ij}\tilde{e}_i$ (similarly for F_i), it is easy to see that $\iota\tilde{e}_i$ is proportional to \tilde{e}_i . Thus $\iota\tilde{\mathfrak{g}} = \tilde{\mathfrak{g}}$, and $\iota|_{\tilde{\mathfrak{g}}}$ is the Cartan antiinvolution of $\tilde{\mathfrak{g}}$.

Substituting $\tilde{\mathfrak{g}}$ instead of \mathfrak{g} , it follows that it is enough to prove the statement of Lemma 14.20 for $X = \sum_{i=1}^{\text{rk}(\mathfrak{g})} e_i$. In this case Lemma 14.17 implies that X is $\exp(\mathfrak{h})$ -conjugate to the element E of Lemma 14.17. Moreover, doing another $\exp(\mathfrak{h})$ -conjugation one can ensure that the vector space $\langle E, F, H \rangle$ of Lemma 14.17 is ι -invariant. \square

Now the proposition is proven for semisimple and for nilpotent elements $X \in \mathfrak{g}$. For an arbitrary $X \in \mathfrak{g}$, there is unique representation $X = X_{ss} + X_{\text{nil}}$ as a sum of commuting semisimple and nilpotent elements. Doing conjugation, we may assume $X_{ss} \in \mathfrak{h}$. Let $\mathfrak{g}_0 = \text{Stab}_{\text{ad}} X_{ss}$. Obviously, $\iota\mathfrak{g}_0 = \mathfrak{g}_0$, \mathfrak{g}_0 is a reductive Lie algebra, $\mathfrak{g}_0 \supset \mathfrak{h}$, and $\iota|_{\mathfrak{g}_0}$ is the Cartan antiinvolution of \mathfrak{g}_0 . Since $X_{\text{nil}} \in \mathfrak{g}_0$, we know that X_{nil} is G_0 -conjugate to an element of $\text{Fix}(\iota|_{\mathfrak{g}_0})$, here G_0 is the Ad-group of \mathfrak{g}_0 . Since X_{ss} is G_0 -invariant and ι -invariant, the above conjugation by $g \in G_0$ sends $X_{ss} + X_{\text{nil}}$ to $\text{Fix}(\iota|_{\mathfrak{g}_0}) \subset \text{Fix}(\iota)$. This finishes the proof of the proposition. \square

This finishes the proof of the amplification. \square

Finally, one can apply the accumulated information to prove one of conjectures of [10]:

Theorem 14.22. *Suppose that \mathfrak{g} is reductive, and $c_1 \in \mathfrak{g}^*$ is regular. Then there is a dense open subset $U \subset \mathfrak{g}^*$ such that the restriction on U of the pair $\{, \}_{1,2}$ is flat.*

Proof. Due to Theorem 14.14, it is enough to show that on a dense open subset of \mathfrak{g}^* the Kronecker web which is a (local) base of the action foliation is flat. This was proven in [25] (in less generality), here we reproduce a more general form of these arguments:

Lemma 14.23. *Consider a Lie algebra \mathfrak{g} , $\text{rk} \mathfrak{g} = r$, and r polynomials p_1, \dots, p_r on \mathfrak{g}^* . Let $U \subset \mathfrak{g}^*$ consists of points $\alpha \in \mathfrak{g}^*$ such that $dp_1|_{\alpha}, \dots, dp_r|_{\alpha}$ are linearly independent. Suppose that \mathfrak{g} is 2-regular, $U \neq \emptyset$, $\dim \mathfrak{g} = 2 \sum_{i=1}^r \deg p_i + r$, and that $c_1 \in \mathfrak{g}^*$ is regular. Consider the Kronecker structure on the local base \mathcal{B} of the action foliation \mathcal{F} of the bihamiltonian structure on \mathfrak{g}^* associated to c_1 . This Kronecker structure is flat on an open subset.*

Proof. Argue as in the end of the proof of Theorem 14.14. On a Zariski open subset U_0 of \mathfrak{g}^* the polynomials p_i , $i = 1, \dots, r$, locally distinguish Ad^* -orbits (thus symplectic leaves of $\{, \}_2$) on \mathfrak{g}^* . Thus on a Zariski open subset U_1 of \mathfrak{g}^* the polynomials $p_{i,\lambda}$, $i = 1, \dots, r$, $\lambda \in \Lambda \subset \mathbb{K}$, locally distinguish leaves of \mathcal{F} (here $p_{i,\lambda}(\alpha) \stackrel{\text{def}}{=} p_i(\alpha + \lambda c_1)$). Associate to a given $\alpha \in \mathfrak{g}^*$ the coefficients a_{ij} , $i = 1, \dots, r$, $j = 0, \dots, \deg p_i$, of polynomials $p_i(\alpha + \lambda c_1)$ in λ . This is a polynomial mapping $a: \mathfrak{g}^* \rightarrow \mathbb{K}^N$, $N = \sum_{i=1}^r \deg p_i + r$. We conclude that on U_1 connected components of fibers of $a|_{U_0}$ coincide with leaves of \mathcal{F} .

But the leaves of \mathcal{F} have codimension $\frac{\dim \mathfrak{g} + r}{2}$, thus in the conditions of the lemma the mapping $a|_{U_0}$ is a submersion, and leaves of \mathcal{F} are connected components of fibers of a . Thus on U_1 the manifold $a(U_1)$ may be considered as a base \mathcal{B} of the action foliation. Describe the structure of Kronecker web on $a(U_1)$.

Fix $\lambda \in \mathbb{K}$. The symplectic leaves of $\lambda \{, \}_1 + \{, \}_2$ are $-\lambda c_1$ -translations of Ad^* -orbits, thus on U_1 they coincide with level sets of $p_{i,\lambda}$, $l \in \mathbb{K}$. Thus the projections of these leaves to \mathcal{B} may be described by equations $\sum_j c_{ij} \lambda^j = C_i$. Thus fibers of integrating foliations \mathcal{F}_λ on \mathcal{B} are parallel planes in \mathbb{K}^N , which finishes the proof of the lemma. \square

To finish the proof of the theorem, it is enough to recall [5] that reductive Lie algebras satisfy the lemma. \square

Corollary 14.24. *In conditions of Theorem 14.22 there is an open dense subset $U \subset \mathfrak{g}^*$ such that the restriction on U of the bihamiltonian structure is locally isomorphic to a direct product of several copies of the structure of Example 0.9.*

Proof. It is enough to show that a micro-Kronecker translation-invariant bihamiltonian structure can be represented as a product of structures of Example 0.9. This a direct corollary of Theorem 5.1. \square

15. APPENDIX ON KRONECKER DECOMPOSITIONS

We know that any Kronecker relation W in a vector space V can be decomposed into a direct sum of Kronecker blocks W_i in subspaces $V_i \subset V$. However, this decomposition is not unique. Here we sketch the degree of arbitrariness of this decomposition.

Given such a decomposition $V = \bigoplus V_i$, consider the following objects:

Definition 15.1. The *isotypic component* $\mathcal{I}_k(V)$ of type k of a decomposition $V = \bigoplus V_i$ is the sum of subspaces V_i of dimension k . The *isotypic filtration* F_k of a decomposition $V = \bigoplus V_i$ is $F_k V = \sum_{l \leq k} \mathcal{I}_l(V)$.

Say that a vector subspace $S \subset V$ is a *k-isotypic block* if all the Kronecker blocks of V have dimension k .

Theorem 15.2. *The isotypic filtration of a Kronecker relation in V does not depend on the choice of a decomposition $V = \bigoplus V_i$ into Kronecker blocks.*

Proof. Start with

Definition 15.3. Given a decomposition $V = \bigoplus V_i$ of a relation W in V into Kronecker blocks and $\lambda \in \mathbb{K}\mathbb{P}^1$, let λ -filtration be the filtration $F_k \text{Ker}_\lambda W$ of $\text{Ker}_\lambda W$ by $\text{Ker}_\lambda W \cap F_k V$.

Due to Lemma 6.3, if λ_i , $i \geq 0$, is a sequence of different elements of $\mathbb{K}\mathbb{P}^1$, then $F_k V = \sum_{i=1}^k F_k \text{Ker}_{\lambda_i} W$. Thus it is enough to show that the λ -filtration in $\text{Ker}_\lambda W$ does not depend on the choice of decomposition into Kronecker blocks.

On the other hand, Lemmas 6.3 and 7.3 taken together imply that $F_k \text{Ker}_{\lambda_0} W = \text{Ker}_{\lambda_0} W \cap \sum_{i=1}^k \text{Ker}_{\lambda_i} W$. Thus $F_k \text{Ker}_{\lambda_0} W$ does not depend on the choice of decomposition. \square

Remark 15.4. Note that this theorem implies the statement of Theorem 3.7 about uniqueness of the collection of dimension of Kronecker blocks in the decomposition of a given Kronecker relation.

Obviously, a choice of a decomposition of an isotypic component into Kronecker blocks is extremely non-unique if the component has more than one block.

Proposition 15.5. Consider a Kronecker relation W in V , suppose that all the Kronecker blocks of W have the same dimension. Fix $\lambda_0 \in \mathbb{K}\mathbb{P}^1$. Given a decomposition $W = \bigoplus W_i$ into Kronecker blocks, one obtains a decomposition $\text{Ker}_{\lambda_0} W = \bigoplus \text{Ker}_{\lambda_0} W_i$ into a direct sum of 1-dimensional subspaces.

Given an arbitrary decomposition $\text{Ker}_{\lambda_0} W = \bigoplus Y_i$ into one-dimensional subspaces, one can find a decomposition $V = \bigoplus V_i$ into Kronecker blocks such that $\text{Ker}_{\lambda_0} V_i = Y_i$. The subspace V_i are uniquely determined by the subspace Y_i .

Proof. Suppose that dimensions of Kronecker blocks of V are equal to k . Let $\Lambda \subset \mathbb{K}\mathbb{P}^1$ has $k+1$ element, let $K_\lambda = \text{Ker}_\lambda W$, $\lambda \in \Lambda$. By Amplification 7.2 the collection $\{K_\lambda\}_{\lambda \in \Lambda}$ uniquely determines W . Express possible decompositions of W into Kronecker blocks in terms of this collection.

Suppose that $\Lambda = \Lambda_0 \cup \{\tilde{\lambda}\}$, $\tilde{\lambda} \notin \Lambda_0$. By lemmas 6.3, 7.3, $V = \bigoplus_{\lambda \in \Lambda_0} K_\lambda$, denote by π_λ , $\lambda \in \Lambda_0$, the projection of V on K_λ according to this decomposition. As one can easily check, $\pi_\lambda K_{\tilde{\lambda}} = K_\lambda$ if $\lambda \in \Lambda_0$ and V is a Kronecker block with $\dim V = k$. Hence $\pi_\lambda K_{\tilde{\lambda}} = K_\lambda$ for an arbitrary k -isotypic V . We see that projections π_λ identify all the K_λ with $K_{\tilde{\lambda}}$, thus one with another.

Assume $\lambda_0 \in \Lambda_0$. Due to the above identifications, a choice of a basis $v_i^{(\lambda_0)}$ in K_{λ_0} induces bases $v_i^{(\lambda)}$ in each of the subspaces K_λ , $\lambda \in \Lambda_0$, thus a basis in V . Let V_i is spanned by $v_i^{(\lambda)}$, $\lambda \in \Lambda_0$.

Lemma 15.6. Consider a vector space \tilde{V} , $\dim \tilde{V} = k$, and one-dimensional subspaces $T_\lambda \subset \tilde{V}$, $\lambda \in \Lambda \subset \mathbb{K}\mathbb{P}^1$, $\text{card}(\Lambda) = k+1$. Suppose that each collection of k subspaces out of $\{T_\lambda\}$ spans the whole vector space. Then there is one and only one Kronecker relation \tilde{W} in \tilde{V} such that $\text{Ker}_\lambda \tilde{W} = T_\lambda$, $\lambda \in \Lambda$,

Proof. The “only one” part follows from Amplification 7.2. On the other hand, if W_0 is a Kronecker block in V_0 , $\dim V_0 = k$, then there is (exactly one up to proportionally) linear mapping f from V_0 to \tilde{V} such that $f(\text{Ker}_\lambda W) = T_\lambda$, $\lambda \in \Lambda$. Since f is invertible, putting $\tilde{W} = f_! W_0$ finishes the proof. \square

Apply the lemma to $\tilde{V} = V_i$, $T_\lambda = V_i \cap K_\lambda$, $\lambda \in \Lambda$. By the construction of the basis $v_i^{(\lambda)}$, T_λ is one-dimensional, thus the conditions of the lemma apply. This gives a Kronecker-block linear relation \tilde{W}_i in V_i . Then $\tilde{W} = \bigoplus \tilde{W}_i$ is a Kronecker linear relation in V with all the Kronecker blocks having dimension k , and $\text{Ker}_\lambda \tilde{W} = K_\lambda$ for $\lambda \in \Lambda$. By Amplification 7.2, $W = \tilde{W}$, thus $\bigoplus \tilde{W}_i$ is the required decomposition of W into a direct sum of Kronecker blocks. \square

Due to Theorem 15.2, the only arbitrariness in the choice of k -isotypic block is the choice of an appropriate complement of $F_{k-1}V$ in $F_k V$. Study which complements may appear as isotypic blocks.

To simplify notations we may assume that $V = F_k V$ (call such W a *relation of type $\leq k$*). This assumption holds until the end of this section.

Definition 15.7. Given a k -isotypic block $S \subset V$ of a Kronecker relation W of type $\leq k$ in V , let λ -pivot space of S be $S \cap \text{Ker}_\lambda W$, here $\lambda \in \mathbb{K}\mathbb{P}^1$.

Lemma 15.8. Suppose that W is a relation of type $\leq k$ in V . Then a λ -pivot space is a complement to $F_{k-1} \text{Ker}_\lambda W$ in $\text{Ker}_\lambda W$.

Proof. This follows directly from decomposability into Kronecker blocks. \square

Definition 15.9. Consider 3 subspaces V, V', V'' of a vector space W . Say that $V' \equiv V'' \pmod{V}$ if $\dim V' = \dim V''$ and images of V' and V'' in W/V coincide.

Definition 15.10. Consider a finite subset $\Lambda \subset \mathbb{K}\mathbb{P}^1$ and a Kronecker relation W in V of type $\leq k$. Say that a collection of vector subspaces $S_\lambda \subset \text{Ker}_\lambda W$, $\lambda \in \Lambda$, is *admissible* if there is a k -isotypic block S such that $S_\lambda = S \cap \text{Ker}_\lambda W$ for any $\lambda \in \Lambda$.

Say that $\{S_\lambda\}_{\lambda \in \Lambda}$ is *l -admissible* if there is a k -isotypic block S such that $S_\lambda \equiv S \cap \text{Ker}_\lambda W \pmod{F_l \text{Ker}_\lambda W}$ for any $\lambda \in \Lambda$.

In particular, a collection $\{S_\lambda\}_{\lambda \in \Lambda}$ is $k-1$ -admissible if S_λ is a complement to $F_{k-1} \text{Ker}_\lambda W$ in $\text{Ker}_\lambda W$.

Definition 15.11. Say that a sequence v_1, \dots, v_l of elements of V forms a *W -chain* if for any two consecutive elements v, \tilde{v} of the sequence $0, v_1, \dots, v_l, 0$ the pair $(v, \tilde{v}) \in W$.

Consider the pencil $\mathcal{P}_1, \mathcal{P}_2: V \rightarrow V'$ which corresponds to W as in Section 2. It is clear that v_1, \dots, v_l forms a W -chain iff $v_1 + \lambda v_2 + \dots + \lambda^{l-1} v_l$ is in the kernel of $\lambda \mathcal{P}_1 - \mathcal{P}_2$ (here we consider λ as a new variable, thus the relation holds over $\mathbb{K}[\lambda]$). Each Kronecker block S_i of dimension l in V with a basis $\mathbf{f}_1^{(i)}, \dots, \mathbf{f}_l^{(i)}$ (as in Example 3.3) gives a W -chain $\mathbf{f}_1^{(i)}, \dots, \mathbf{f}_l^{(i)}$.

Definition 15.12. Given two W -chains v_1, \dots, v_l and v'_1, \dots, v'_m , $m < l$, define the n -th elementary operation as a change of v_i to $v_i + Cv'_{i-n}$. Here $C \in \mathbb{K}$, $0 \leq n \leq l - m$, and we extend the sequence v'_i to $i \leq 0$ and $i > m$ by 0. The elementary operation of the first kind is the 0-th elementary operation, elementary operation of the second kind is the $l - m$ -th elementary operation.

Note that an elementary operation transforms a W -chain into a W -chain, and that the operations of the first kind do not change v_l , while operations of the second kind do not change v_1 .

Remark 15.13. Consider a k -isotypic block S of V . Taking a W -chain v_i corresponding to a Kronecker block of S , and a W -chain v'_i corresponding to a Kronecker block of $F_{k-1}V$, one can perform elementary operations using these chains. These operations will change the chain v_i . The following lemma implies that this change corresponds to a change of the k -isotypic block S :

Lemma 15.14. *Suppose that vectors $v_{ij} \in L$, $1 \leq i \leq k$, $j \in J$, are linearly independent, span a complement S to $F_{k-1}V$ in V , and for any $j \in J$ the sequence v_{ij} , $1 \leq i \leq k$, is a W -chain in V . Then S is a k -isotypic block in V .*

Proof. Consider again the pencil $\mathcal{P}_1, \mathcal{P}_2: V \rightarrow V'$. It is clear that $\mathcal{P}_1 S = \mathcal{P}_2 S$ as subspaces in V' . If we prove that $\mathcal{P}_1(F_{k-1}V)$ (which coincides with $\mathcal{P}_2(F_{k-1}V)$) does not intersect $\mathcal{P}_1 S$, then one can split S and $\mathcal{P}_1 S$ into direct summands, which will prove the lemma.

Suppose that $\mathcal{P}_1(F_{k-1}V)$ does intersect $\mathcal{P}_1 S$. Consider an arbitrary k -isotypic block S' in V . Conditions of the lemma imply that $S \equiv S' \pmod{F_{k-1}V}$. This implies that $\mathcal{P}_1(F_{k-1}V)$ intersects $\mathcal{P}_1 S'$, which is a contradiction. \square

Definition 15.15. Given a k -isotypic block S of V , call the modifications of S resulting from elementary operations of Remark 15.13 the elementary operations over k -isotypic blocks.

Obviously:

Lemma 15.16. *Using elementary operations of the first kind one can change $S \cap \text{Ker}_{1;0} W$ to become an arbitrary complement to $F_{k-1} \text{Ker}_{1;0} W$ in $\text{Ker}_{1;0} W$ without changing $S \cap \text{Ker}_{0;1} W$. Similarly, the operations of the second kind will do the same with $S \cap \text{Ker}_{0;1} W$.*

This implies

Lemma 15.17. *Fix $\lambda', \lambda'' \in \mathbb{K}\mathbb{P}^1$, $\lambda' \neq \lambda''$. Suppose that $V = F_k V$. Then any $k - 1$ -admissible pair of subspaces $(S_{\lambda'}, S_{\lambda''})$ is admissible.*

Moreover, one can improve this statement by considering subsets $\Lambda \subset \mathbb{K}\mathbb{P}^1$ with more than two elements. Also, one can describe the degree of arbitrariness in the choice of an isotypic block S with given intersections $S \cap \text{Ker}_\lambda W$ for $\lambda \in \Lambda$. Start with the following

Proposition 15.18. *Consider two²⁶ k -isotypic blocks S and S' in V . There is a sequence of elementary operations which transforms S into S' .*

Proof. By Lemma 15.16, one may suppose that $S \cap \text{Ker}_{1;0} W = S' \cap \text{Ker}_{1;0} W$. Consider W -chains which form bases in S and S' . Denote these chains in S by $v_{i,j}$, in S' by $v'_{i,j}$ (here j enumerates chains, and i vectors inside a chain). By Proposition 15.5 we may assume that $v_{1,j} = v'_{1,j}$. Let $p_j = \sum_i v_{i,j} \lambda^{i-1}$ be the polynomial in λ which corresponds to the W -chain $v_{\bullet,j}$, similarly introduce p'_j . Fix j . Obviously, $p'_j - p_j$ can be written as λq , and the polynomial q corresponds to an appropriate W -chain \tilde{v}_i . Moreover, all the vectors \tilde{v}_i are in $F_{k-1}V$.

Decompose $F_{k-1}V$ into direct sum of Kronecker components, consider projections of \tilde{v}_i to these components. Clearly, for any such projection π_α the vectors $\pi_\alpha \tilde{v}_i$ form a W -chain. Now the proposition follows from the following

Lemma 15.19. *Consider a Kronecker block \widetilde{W} in \widetilde{V} and a \widetilde{W} -chain v_i which is a basis of \widetilde{V} . Let v'_i be an arbitrary \widetilde{W} -chain in \widetilde{V} . Let $p = \sum_i v_i \lambda^{i-1}$, $p' = \sum_i v'_i \lambda^{i-1}$. Then there is a polynomial $q \in \mathbb{K}[\lambda]$ such that $p' = qp$.*

Proof. Write v'_j in the basis v_i and compare the coefficients using the definition of a W -chain. □

This finishes proof of the proposition. □

From this proposition one can immediately deduce

Theorem 15.20. *Let S be a k -isotypic block in V , and $\Lambda \subset \mathbb{K}\mathbb{P}^1$.*

1. *Let S' be another k -isotypic block in V . Suppose that*

$$S \cap \text{Ker}_\lambda W \equiv S' \cap \text{Ker}_\lambda W \pmod{F_l \text{Ker}_\lambda W} \text{ for } \lambda \in \Lambda.$$

If $\text{card}(\Lambda) = k - l$, then

$$S \cap \text{Ker}_\lambda W \equiv S' \cap \text{Ker}_\lambda W \pmod{F_l \text{Ker}_\lambda W} \text{ for any } \lambda.$$

2. *Suppose that for $\lambda \in \Lambda$ a vector subspace $S'_\lambda \subset \text{Ker}_\lambda W$ is fixed, and $S'_\lambda \equiv S \cap \text{Ker}_\lambda W \pmod{F_{l+1} \text{Ker}_\lambda W}$ for $\lambda \in \Lambda$. Then if $\text{card}(\Lambda) = k - l$, then there exists another k -isotypic block S' in V such that*

$$S'_\lambda \equiv S' \cap \text{Ker}_\lambda W \pmod{F_l \text{Ker}_\lambda W}.$$

This theorem gives a complete description of the arbitrariness in the choice of the k -isotypic block in V . Say, consider a subset $\{\lambda_i\}$ of $\mathbb{K}\mathbb{P}^1$. Subspaces $S \cap \text{Ker}_{\lambda_1} W$ and $S \cap \text{Ker}_{\lambda_2} W$ (which may be arbitrary complements to $F_{k-1} \text{Ker}_{\lambda_i} W$ in $\text{Ker}_{\lambda_i} W$, $i = 1, 2$) completely determine $S \pmod{F_{k-2}}$. In particular, they determine $S \cap \text{Ker}_{\lambda_3} W \pmod{F_{k-2} \text{Ker}_{\lambda_3} W}$. Choosing an arbitrary subspace S_{λ_3} of $\text{Ker}_{\lambda_3} W$ with the same reduction $\pmod{F_{k-2} \text{Ker}_{\lambda_3} W}$ and requiring $S \cap \text{Ker}_{\lambda_2} W = S_{\lambda_3}$ completely

²⁶These blocks would correspond to different decomposition of V into direct sums of Kronecker blocks.

determines $S \bmod F_{k-3}$ and so on. Together with Proposition 15.5 this describes all possible decompositions of V into Kronecker blocks.

REFERENCES

1. J. Frank Adams, *Lectures on Lie groups*, W. A. Benjamin, Inc., New York-Amsterdam, 1969.
2. V. I. Arnol'd, *Mathematical methods of classical mechanics*, Springer-Verlag, New York, 199?, Translated from the 1974 Russian original by K. Vogtmann and A. Weinstein, Corrected reprint of the second (1989) edition.
3. V. I. Arnol'd and A. B. Givental', *Symplectic geometry*, Current problems in mathematics. Fundamental directions, Vol. 4 (Moscow), Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1985, pp. 5–139, 291.
4. A. V. Bolsinov, *Compatible Poisson brackets on Lie algebras and the completeness of families of functions in involution*, Izv. Akad. Nauk SSSR Ser. Mat. **55** (1991), no. 1, 68–92.
5. N. Bourbaki, *Groupes et algèbres de Lie*, Herman, Paris, 1975, Ch7-8.
6. E. B. Dynkin, *Maximal subgroups of the classical groups*, Trudy Moskov. Mat. Obšč. **1** (1952), 39–166.
7. A. S. Fokas and B. Fuchssteiner, *On the structure of symplectic operators and hereditary symmetries*, Lett. Nuovo Cimento (2) **28** (1980), no. 8, 299–303.
8. F. R. Gantmacher, *The theory of matrices. Vols. 1, 2*, Chelsea Publishing Co., New York, 1959, Translated by K. A. Hirsch.
9. I. M. Gel'fand and I. Ja. Dorfman, *Hamiltonian operators and algebraic structures associated with them*, Funktsional. Anal. i Prilozhen. **13** (1979), no. 4, 13–30, 96.
10. Israel M. Gelfand and Ilya Zakharevich, *Webs, Lenard schemes, and the local geometry of bihamiltonian Toda and Lax structures*, Archived as math.DG/9903080. To appear in Selecta Math.
11. ———, *Spectral theory for a pair of skew-symmetrical operators on S^1* , Func. Anal. Appl. **23** (1989), no. 1, 85–93.
12. ———, *Webs, Veronese curves, and bihamiltonian systems*, J. of Func. Anal. **99** (1991), 150–178.
13. ———, *On the local geometry of bihamiltonian structures*, The Gelfand mathematical seminar, 1990–1992 (Boston), Birkhäuser, 1993, pp. 51–112.
14. ———, *The spectral theory for a pencil of skewsymmetrical differential operators of the third order*, Comm. Pure Appl. Math. **47** (1994), no. 8, 1031–1041.
15. Victor Guillemin and Shlomo Sternberg, *Geometric asymptotics*, American Mathematical Society, Providence, R.I., 1977, Mathematical Surveys, No. 14.
16. A. A. Kirillov, *Local Lie algebras*, Uspehi Mat. Nauk **31** (1976), no. 4(190), 57–76.
17. Y. Kosmann-Schwarzbach and F. Magri, *Lax-Nijenhuis operators for integrable systems*, J. Math. Phys. **37** (1996), no. 12, 6173–6197.
18. Peter D. Lax, *Almost periodic solutions of the KdV equation*, SIAM Rev. **18** (1976), no. 3, 351–375.
19. Franco Magri, *A simple model of the integrable Hamiltonian equation*, Journal of Mathematical Physics **19** (1978), no. 5, 1156–1162.
20. ———, *On the geometry of soliton equations*, Preprint, 1988.
21. ———, *On the geometry of soliton equations*, Acta Appl. Math. **41** (1995), no. 1-3, 247–270, Geometric and algebraic structures in differential equations.
22. Henri McKean, private communication, 1990.
23. A. S. Miščenko and A. T. Fomenko, *Euler equation on finite-dimensional Lie groups*, Izv. Akad. Nauk SSSR Ser. Mat. **42** (1978), no. 2, 396–415, 471.

24. Andriy Panasyuk, *Symplectic realizations of bihamiltonian structures*, preprint, 1998.
25. ———, *Veronese webs for bihamiltonian structures of higher corank*, Banach Center Publications, Poisson geometry, Proceedings of the conference dedicated to the memory of Stanislaw Zakrzewski, Warsaw 1998 (Warszawa) (Pawel Urbański and Janusz Grabowski, eds.), Instytut Matematyczny PAN, 1999, to appear.
26. Marie-Hélène Rigal, *Géométrie globale des systèmes bihamiltoniens réguliers de rang maximum en dimension 5*, C. R. Acad. Sci. Paris Sér. I Math. **321** (1995), no. 11, 1479–1484.
27. ———, *Tissus de Veronese en dimension 3*, Séminaire Gaston Darboux de Géométrie et Topologie Différentielle, 1994–1995 (Montpellier) (Montpellier), Univ. Montpellier II, Montpellier, 1995, pp. iv, 63–68.
28. ———, *Systèmes bihamiltoniens en dimension impaire*, Ann. Sci. École Norm. Sup. (4) **31** (1998), no. 3, 345–359.
29. Robert C. Thompson, *Pencils of complex and real symmetric and skew matrices*, Linear Algebra Appl. **147** (1991), 323–371.
30. Francisco-Javier Turiel, *Classification locale d'un couple de formes symplectiques Poisson-compatibles*, Comptes Rendus des Seances de l'Academie des Sciences. Serie I. Mathematique **308** (1989), no. 20, 575–578.
31. ———, *C^∞ -équivalence entre tissus de Veronese et structures bihamiltoniennes*, C. R. Acad. Sci. Paris Sér. I Math. **328** (1999), no. 10, 891–894.
32. H. W. Turnbull and A. C. Aitken, *An introduction to the theory of canonical matrices*, Dover Publications Inc., New York, 1961.
33. Alan Weinstein, *The local structure of Poisson manifolds*, J. Differential Geom. **18** (1983), no. 3, 523–557.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, 231 W. 18 AVE, COLUMBUS, OH, 43210

E-mail address: ilya@math.ohio-state.edu