

Σ_1^1 in every real in a Σ_1^1 class of reals is Σ_1^1

Leo Harrington
Department of Mathematics
University of California, Berkeley
Berkeley CA 94270

Richard A. Shore*
Department of Mathematics
Cornell University
Ithaca NY 14853

Theodore A. Slaman†
Department of Mathematics
University of California, Berkeley
Berkeley CA 94270

July 11, 2016

Abstract

We first prove a theorem about reals (subsets of \mathbb{N}) and classes of reals: If a real X is Σ_1^1 in every member G of a nonempty Σ_1^1 class \mathcal{K} of reals then X is itself Σ_1^1 . We also explore the relationship between this theorem, various basis results in hyperarithmetic theory and omitting types theorems in ω -logic. We then prove the analog of our first theorem for classes of reals: If a class \mathcal{A} of reals is Σ_1^1 in every member of a nonempty Σ_1^1 class \mathcal{B} of reals then \mathcal{A} is itself Σ_1^1 .

1 Introduction

We work in Cantor space $2^{\mathbb{N}}$ and call its members $X \subseteq \mathbb{N}$, *reals*. We think of members of Baire space $\mathbb{N}^{\mathbb{N}}$ as functions $F : \mathbb{N} \rightarrow \mathbb{N}$ (coded as real consisting of pairs of numbers). We use the standard normal form theorems for reals and classes of reals as follows: A real X is Σ_1^1 (in a real G) if it is of the form $\{n \mid \exists F \forall x R(F \upharpoonright x, x, n)\}$ for a recursive (in G) predicate R . A class \mathcal{K} of reals is Σ_1^1 (in G) if it is of the form $\{X \mid \exists F \forall x R(X \upharpoonright x, F \upharpoonright x, x)\}$

*Partially supported by NSF Grant DMS-1161175. The last two authors began their work on this paper at a workshop of the Institute for Mathematical Sciences of the National University of Singapore which also partially supported them.

†Partially supported by NSF Grant DMS-1301659 and by the Institute for Mathematical Sciences of the National University of Singapore.

for a recursive (in G) predicate R . A real or class of reals is Δ_1^1 (or hyperarithmetical) (in G) if it and its complement are Σ_1^1 (in G). Our first main theorem is the following:

Theorem 2.1. *If a real X is Σ_1^1 in every member G of a nonempty Σ_1^1 class K of reals then X is itself Σ_1^1 .*

While the statement of this theorem and certainly the proof we provide in the next section seem to have little to do with either results of hyperarithmetical theory or model theory they are all, in fact, connected along a couple of paths. Indeed, we were thinking about related matters when we proved the theorem.

A basis theorem in recursion theory typically says that every nonempty class of some sort contains a member with some property. For example, the classes may be arbitrary Σ_1^1 classes \mathcal{K} of reals. One, the Gandy Basis Theorem (see Sacks [1990, III.1.5]), says that every nonempty Σ_1^1 class of reals contains one Z such that $\omega_1^Z = \omega_1^{CK}$. (For any Z , ω_1^Z is the least ordinal not recursive, or equivalently not Δ_1^1 in Z ; ω_1^{CK} is ω_1^Z for Z recursive (or Δ_1^1 .) Another, the Kreisel Basis Theorem (see Sacks [1990, III. 7.2]), says that if a real X is not hyperarithmetical (i.e. Δ_1^1) then \mathcal{K} also contains a real Z in which X is not Δ_1^1 . An equivalent version is that if X is Δ_1^1 in every member of \mathcal{K} then X is Δ_1^1 .

Our first theorem is the generalization of the Kreisel basis theorem where Δ_1^1 is replaced by Σ_1^1 . (To see that it implies the result of Kreisel note that it says that if X and \bar{X} are both Σ_1^1 (i.e. Δ_1^1) in every member of \mathcal{K} then they are both Σ_1^1 (and so Δ_1^1)). Our theorem also implies the basis result of Gandy: As Kleene's O is not Σ_1^1 there is a $Z \in \mathcal{K}$ in which O is not Σ_1^1 . By classical results of Spector (see Sacks [1990, II. 7.7]), this implies that $\omega_1^Z = \omega_1^{CK}$. (See Theorem 2.9.)

Sacks also provides results of hyperarithmetical theory as corollaries to Kreisel's theorem and others that, as he points out, can be viewed as omitting types theorems in ω -logic. They are also immediate consequences of our theorem as we indicate in the next section. We discuss these and other related results in next section after we prove our theorem.

After his proof, Sacks [1990, p. 75] says of this connection that "The recursion theorist winding his way through a Σ_1^1 set is a brother to the model theorist threading his way through a Henkin tree." Our proof, which requires no knowledge of either hyperarithmetical theory or model theory, shows that there is another sibling traipsing (or perhaps treading carefully) through a forcing construction.

Our theorem should have been a classical one of hyperarithmetical theory. It also has analogs, both recent and classical, in other settings. When we told Stephen Simpson the result he remarked that Andrews and Miller [2015, Proposition 3.6] had recently proven the analogous result for Π_1^0 classes in place of Σ_1^1 classes. We rephrase it in our terminology as follows:

Theorem 1.1 (Andrews and Miller). *Let P be a nonempty Π_1^0 class. If X is Π_1^0 in every member of P then X is Π_1^0 . (Or, equivalently, if X is Σ_1^0 in every member of P*

then X is Σ_1^0 .)

Their proof is a forcing proof similar to ours but using Π_1^0 classes instead of Σ_1^1 ones.

At the level of Σ_2^1 classes, a standard basis theorem gives the analogous result (as pointed out to us by John Steel). The classical result (see Moschovakis [1980, 4E.5]) is that the Δ_2^1 reals are a basis for the Σ_2^1 classes of reals. Thus if \mathcal{K} is Σ_2^1 it contains a Δ_2^1 real G and, of course, any real X which is Σ_2^1 in G via Θ is itself Σ_2^1 . ($X = \{n \mid \exists G(\Psi(G) \ \& \ \Theta(G, n))\}$ where Ψ is the Σ_2^1 formula saying G satisfies its Δ_2^1 definition.) Similar basis results hold at higher levels of the projective hierarchy assuming various set theoretic axioms. (See Moschovakis [1980, 5A.4 and 6C.6].)

About the only facts about Σ_1^1 reals and classes that we use in our proof are the the standard normal form theorems mentioned at the beginning of this Introduction.

Our second main theorem is one analogous to Theorem 2.1 but at the level of classes of real.

Theorem 3.1. *If a class \mathcal{A} of reals is Σ_1^1 in every member of a nonempty Σ_1^1 class \mathcal{B} of reals then it is Σ_1^1 .*

Our proof of this theorem requires some familiarity with effective descriptive set theory. We give some of the basic facts needed and the proof in §3.

2 The Proof for Reals

We now give the promised forcing style proof of our main theorem.

Theorem 2.1. *If a real X is Σ_1^1 in every member G of a nonempty Σ_1^1 class \mathcal{K} of reals then X is itself Σ_1^1 .*

Proof. We use the language of Gandy-Harrington forcing. Forcing conditions are nonempty Σ_1^1 classes \mathcal{L} of reals with set containment as extension. We view the Σ_1^1 formulas $\varphi(G, n)$ as of the form $\exists F \forall x R(G \upharpoonright x, F \upharpoonright x, x, n)$ with R recursive. We say that $\mathcal{L} \Vdash \varphi(G, n)$ if $(\forall Z \in \mathcal{L})(\varphi(Z, n))$. If, as usual, we say $\mathcal{L} \Vdash \neg \varphi(G, n)$ if $(\forall \hat{\mathcal{L}} \subseteq \mathcal{L})(\hat{\mathcal{L}} \not\Vdash \varphi(G, n))$, this definition is then equivalent to $(\forall Z \in \mathcal{L})(\neg \varphi(Z, n))$. The point here is that if there is a $Z \in \mathcal{L}$ such that $\varphi(Z, n)$ then $\hat{\mathcal{L}} = \mathcal{L} \cap \{Z \mid \varphi(Z, n)\}$ is a nonempty extension of \mathcal{L} which obviously forces $\varphi(G, n)$.

We now list all the Σ_1^1 formulas $\Theta_k(G, n)$. These are the formulas that could potentially define the reals Σ_1^1 in any G . We consider an X which is a candidate for being Σ_1^1 in every $G \in \mathcal{K}$. We build a sequence \mathcal{L}_k of conditions beginning with $\mathcal{L}_0 = \mathcal{K} = \{G \mid \exists F_0 \forall x R_{m_0}(G \upharpoonright x, F_0 \upharpoonright x, x)\}$ as well as initial segments γ_k (of length at least k) of our intended G and $\delta_{i,k}$ of witnesses F_i (of length at least k) showing that $G \in \mathcal{L}_k$. More precisely, each \mathcal{L}_k will be of the form $G \supset \gamma_k \ \& \ \forall i \leq k \exists F_i \supset \delta_{i,k} \forall x R_{m_i}(G \upharpoonright x, F_i \upharpoonright x, x)$

for some recursive R_{m_i} (independent of k). Thus, if we successfully continue our construction keeping \mathcal{L}_k nonempty for each k then the $F_i = \lim_k \delta_{i,k}$ for $i \leq k$ will witness that $G = \lim_k \gamma_k$ is in every \mathcal{L}_k as we guarantee that $R_{m_i}(\gamma_k \upharpoonright x, \delta_{i,k} \upharpoonright x, x)$ holds for every $i, x < k$ and every k .

We begin with $\gamma_0 = \emptyset = \delta_{0,0}$ and R_{m_0} as specified by \mathcal{K} . So our G will at least be in \mathcal{K} as desired. Suppose we have defined γ_j and $\delta_{i,j}$ for $j, i \leq k$ and wish to define \mathcal{L}_{k+1} , γ_{k+1} and $\delta_{i,k+1}$ for $i \leq k+1$ so as to prevent X from being Σ_1^1 in G via Θ_k . We ask if there is an $m \in \omega$ and a nonempty $\mathcal{L} \subseteq \mathcal{L}_k$ such that

1. $m \notin X$ and $\mathcal{L} \Vdash \Theta_k(G, m)$ or
2. $m \in X$ and $\mathcal{L} \Vdash \neg\Theta_k(G, m)$.

Suppose there is such an \mathcal{L} of the form $\exists F_{k+1} \forall x R_{m_{k+1}}(G \upharpoonright x, F_{k+1} \upharpoonright x, x)$. As $\mathcal{L} \subseteq \mathcal{L}_k$ is nonempty we can choose $\gamma_{k+1} \supset \gamma_k$ and $\delta_{i,k+1} \supset \delta_{i,k}$ for $i \leq k$ and some $\delta_{k+1,k+1}$ all of length at least $k+1$ such that \mathcal{L}_{k+1} as given by $G \supset \gamma_{k+1}$ & $(\forall i \leq k+1)(\exists F_i \supset \delta_{i,k+1})(\forall x R_{m_i})(G \upharpoonright x, F_i \upharpoonright x, x)$ is a nonempty subclass of \mathcal{L} (and so, in particular, $R_{m_i}(\gamma_{k+1} \upharpoonright x, \delta_{i,k+1} \upharpoonright x, x)$ for every $i, x \leq k+1$). We can now continue our induction.

Note that if we can successfully define nonempty \mathcal{L}_k in this way for every k then we build a $G = \lim_k \gamma_k$ and $F_i = \lim_k \delta_{i,k}$ for each i such that $\forall x R_{m_i}(G \upharpoonright x, F_i \upharpoonright x, x)$. In particular $\forall x R_{m_0}(G \upharpoonright x, F_0 \upharpoonright x, x)$ and so $G \in \mathcal{K}$. Similarly, $G \in \mathcal{L}_k$ for every $k > 0$. If X is $\Sigma_1^1(G)$ as assumed, then $X = \{n \mid \Theta_k(G, n)\}$ for some k . We consider the construction at stage $k+1$ and the \mathcal{L} chosen at that stage. If we were in case (1) then as $\mathcal{L} \Vdash \Theta_k(G, m)$ and $G \in \mathcal{L}_{k+1}$, $\Theta(G, m)$ is true but $m \notin X$ for a contradiction. Similarly, if we were in case (2), as $\mathcal{L} \Vdash \neg\Theta_k(G, m)$ and $G \in \mathcal{L}_{k+1}$, $\neg\Theta(G, m)$ is true but $m \in X$ again for a contradiction.

Thus we can assume that there is some first stage $k+1$ at which there are no m and $\mathcal{L} \subseteq \mathcal{L}_k$ as required in the construction. In this case we claim that X is Σ_1^1 as desired. Indeed, we claim that X is defined as a Σ_1^1 real by $m \in X \Leftrightarrow (\exists Z \in \mathcal{L}_k)\Theta_k(Z, m)$. To see this suppose first that $(\exists Z \in \mathcal{L}_k)\Theta_k(Z, m)$. Then \mathcal{L} as defined by \mathcal{L}_k & $\Theta_k(G, m)$ is a nonempty Σ_1^1 class such that $\mathcal{L} \Vdash \Theta_k(G, m)$ and so we would have $m \in X$ as desired by the assumed failure of (1) at stage $k+1$ of the construction. On the other hand, if $(\forall Z \in \mathcal{L}_k)(\neg\Theta_k(Z, m))$ then $\mathcal{L}_k \Vdash \neg\Theta_k(G, m)$ and so by the failure of (2) at stage $k+1$ of the construction, $m \notin X$ as desired. \square

As usual, we may relativize the Theorem to any real C .

Our theorem easily implies several basic results of hyperarithmetic theory without any appeal to the theory of hyperarithmetic sets as used, for example, in Sacks [1990]. Many of them can also be seen as consequences of type omitting theorems for certain classes of generalized logics. These type omitting arguments are also immediate consequences of our Theorem. We presented two basis theorems of this sort in the introduction and note here

that the proof of Kreisel's in Sacks [1990] uses several deep facts about hyperarithmetical reals and Σ_1^1 classes.

Following his proof of the Kreisel basis theorem Sacks [1990] gives as a corollary a result of Kreisel about the intersections of all ω -models of various theories of second order arithmetic from which follow some previous specific results. We state that result now along with some similar ones earlier in Sacks's presentation. These can all be seen as type omitting arguments. After stating them, we explain a general setting which includes them all and give the relevant type omitting theorems as Corollaries of Theorem 2.1.

Theorem 2.2 (Sacks [1990, III. 4.10]). *The intersection of all ω -models of Δ_1^1 comprehension is HYP, the class of all hyperarithmetical sets or equivalently the class of all Δ_1^1 sets.*

More generally, we have the following result of Kreisel.

Theorem 2.3 (Sacks [1990, III.7.3]). *Let K be a Π_1^1 set of axioms in the language of analysis (i.e. second order arithmetic). If a real X belongs to every countable ω -model of K then X is Δ_1^1 .*

A similar result is the following.

Theorem 2.4 (Sacks [1990, III.4.13]). *The intersection of all ω -models of Σ_1^1 choice downward closed under many-one reducibility is also HYP.*

In all of these results it is easy to see that the class of models described is Σ_1^1 and, of course, every member X of such a model is recursive in it and so any real in every such model is Σ_1^1 but these models are all trivially closed under complementation. So they all follow from our Theorem.

Moving to the type omitting point of view we, somewhat more generally, consider two sorted logics $(\mathcal{N}, \mathcal{M}, \dots)$ in the usual sense of having two types of variables one ranging over the elements of \mathcal{N} and the other over those of \mathcal{M} in addition to the usual apparatus of function, relation and constant symbols of ordinary first order logic. While formally merely a version of first order logic gotten by adding on predicates for N and M , this logic can be turned into a much stronger one (\mathcal{N} -logic) by requiring that all models have their first sort (with some functions and relations on it as given in the structure) isomorphic to some given countable first order structure. The most common example of these logics is ω -logic where we require that \mathcal{N} be isomorphic to the ordinal ω or the standard model \mathbb{N} of arithmetic (depending on the language intended). Again, the most common examples are given by classes of ω -models of fragments T of second order arithmetic as mentioned above. Here, in addition to requiring that \mathcal{N} be the standard model of arithmetic we intend that the elements of \mathcal{M} are subsets of \mathcal{N} and the membership relation \in between members of \mathcal{N} and those of \mathcal{M} is in the language (with the usual axiom of extensionality so that the elements of \mathcal{M} may be identified with true subsets of $\mathcal{N} = \mathbb{N}$). As being an \mathcal{N} model, or even also satisfying some Π_1^1 theory T , is clearly Σ_1^1 in \mathcal{N} , we immediately

get all the results from Sacks [1990] mentioned above as a corollaries of our theorem. Indeed, we have the following generalization of Kreisel's result in Sacks [1990, III.7.3]:

Theorem 2.5. *If T is a Π_1^1 set of sentences in the two sorted language of $(\mathcal{N}, \mathcal{M}, \dots)$ and \mathcal{N} is a countable structure for the appropriate sublanguage (restricted to the first sort), T has an \mathcal{N} -model and p is a n -type (i.e. a complete consistent set of formulas $\varphi(x)$ with n free variables in the language of $(\mathcal{N}, \mathcal{M}, \dots)$) which is not Σ_1^1 in \mathcal{N} , then there is an \mathcal{N} -model of T not realizing p . (Note that, as types are complete sets of formulas, p being Σ_1^1 (in \mathcal{N}) is equivalent to its being Δ_1^1 (in \mathcal{N})).*

Proof. Being an \mathcal{N} model of T is a Σ_1^1 in \mathcal{N} property and so by our Theorem (relativized to \mathcal{N}) there is even an \mathcal{N} -model $(\mathcal{N}, \mathcal{M}, \dots)$ of T in which p is not even Σ_1^1 . (Of course, any type realized in $(\mathcal{N}, \mathcal{M}, \dots)$ is recursive in the complete diagram of $(\mathcal{N}, \mathcal{M}, \dots)$ and so hyperarithmetic in $(\mathcal{N}, \mathcal{M}, \dots)$.) \square

Viewing our theorem as a type omitting argument suggests that we should be able to omit any countable sequence of types (reals) of the appropriate sort rather than just one. A simple modification of our proof gives the expected result.

Theorem 2.6. *If \mathcal{K} is a nonempty Σ_1^1 class reals and X_n a countable sequence of reals none of which is Σ_1^1 , then there is a $G \in \mathcal{K}$ such that no X_n is Σ_1^1 in G . Similarly if no X_n is Δ_1^1 , then there is a $G \in \mathcal{K}$ such that no X_n is Δ_1^1 in G .*

Proof. Repeat the proof of the Theorem but at step $k + 1 = \langle n, j \rangle$ of the construction replace X by X_n and Θ_k by Θ_j . If we successfully pass through all steps k then the previous argument shows that no X_n is Σ_1^1 in $G \in \mathcal{K}$. On the other hand, if the construction terminates at step $k + 1 = \langle n, j \rangle$ then the previous argument shows that X_n is defined as a Σ_1^1 real by $m \in X_n \Leftrightarrow (\exists Z \in \mathcal{L}_k)\Theta_j(Z, m)$ for a contradiction. For the Δ_1^1 version, simply consider the sequence Y_n where $Y_n = X_n$ if X_n is not Σ_1^1 and Y_n is the complement if X_n otherwise (i.e. X_n is not Π_1^1). As now no Y_n is $\Sigma_1^1(G)$, no X_n is $\Delta_1^1(G)$. \square

This version of our Theorem also extends the analog of the result actually given by Andrews and Miller [2015, Proposition 3.6].

Of course, we can relativize this theorem as well to any real C . To give a somewhat different example of a such type omitting argument application of this last theorem we provide one for nonstandard models of ZFC for which we have uses elsewhere.

Corollary 2.7. *For every real C and reals X_n not Δ_1^1 in C , there is a countable ω -model of ZFC containing C but not containing any X_n whose well founded part consists of the ordinals less than ω_1^C , the first ordinal not recursive in C .*

Proof. Being a countable ω -model of ZFC containing (a set isomorphic to) C (under the isomorphism taking the ω of the model to true ω) is clearly a Σ_1^1 in C property. Now apply Theorem 2.6 adding on a new real $X_0 = O^C$ (i.e. Kleene's O relativized to C) to

the list. It supplies a countable ω -model of ZFC containing C but not containing any of the X_n . As it contains C it contains every ordering recursive in C and so order types for every ordinal less than ω_1^C . On the other hand, if there were an ordinal in the model isomorphic to ω_1^C then, by standard results of hyperarithmetic theory, O^C would be in the model as well.

Finally, we point out that the complexity of the G of Theorem 2.6 (and hence of Corollary 2.7 as well) can be as low as possible. \square

Theorem 2.8. *If \mathcal{K} is a nonempty Σ_1^1 class reals and X_n a countable sequence of reals uniformly Δ_1^1 (recursive) in O none of which is Σ_1^1 , then there is a $G \in \mathcal{K}$ with $G \Delta_1^1$ (recursive) in O such that no X_n is Σ_1^1 in G . Indeed, G can be chosen to be of strictly smaller hyperdegree than O , i.e. O is not Δ_1^1 in G . As in Theorem 2.6, if we assume only that the X_n are not Δ_1^1 then we may conclude that none are Δ_1^1 in G .*

Proof. Suppose we are at step $k = \langle n, j \rangle$ of the construction. We know that either there is an $m \in X_n$ such that $(\forall Z \in \mathcal{L}_k)(\neg \Theta_k(Z, m))$ or an $m \notin X_n$ such that $(\exists Z \in \mathcal{L}_k)(\Theta_k(Z, m))$. As the X_n are uniformly Δ_1^1 (recursive) in O , and the rest of the conditions considered in the construction are either Σ_1^1 or Π_1^1 , O can hyperarithmetically (recursively) decide which case to apply. As choosing the $\gamma_{k+1} \supset \gamma_k$ and $\delta_{i,k+1} \supset \delta_{i,k}$ for $i \leq k$ and so \mathcal{L}_{k+1} now only require finding ones for which the corresponding Σ_1^1 class \mathcal{L}_{k+1} is nonempty, this step is also recursive in O . Of course, as we can add O onto the list of X_n , we then guarantee that O is not Σ_1^1 in G and so, of course, not Δ_1^1 in G as required. \square

Note that by a result of Spector's (see Sacks [1990, Theorem II.7.6(ii)]) $\omega_1^{CK} < \omega_1^A$ implies that O is Δ_1^1 in A (indeed there is a pair of Σ_1^1 formulas $\varphi(X, n)$ and $\theta(X, n)$ which define O and its complement for any X with $\omega_1^X > \omega_1^{CK}$), we have the Kleene and Gandy basis theorem for Σ_1^1 classes as well.

Theorem 2.9 (Kleene and Gandy Basis Theorems). *Every nonempty Σ_1^1 class of reals \mathcal{K} contains an element A recursive in and of strictly smaller hyperdegree than O . In particular, one with $\omega_1^A = \omega_1^{CK}$.*

3 The Proof for Classes of Reals

In this section we prove our result for classes of reals.

Theorem 3.1. *If a class \mathcal{A} of reals is Σ_1^1 in every member of a nonempty Σ_1^1 class \mathcal{B} of reals then it is Σ_1^1 .*

The proof relies on several basic and important results of effective descriptive set theory. To ease reading the proof, we state the most important ones now. We state lightface versions without parameters. Relativizations to individual real parameters are

routine. (Note that, when ordinals or lengths of well-ordered relations are involved, relativization to Z includes replacing ω_1^{CK} by ω_1^Z .) We don't need the full boldface versions. These facts can be found in basic books on effective descriptive set theory such as Moschovakis [1980], higher recursion theory such as Sacks [1990] or Hinman [1978] or even reverse mathematics such as Simpson [2009].

Proposition 3.2 (Codes). *We can code Δ_1^1 classes of reals \mathcal{V} as either Δ_1^1 reals C (Δ_1^1 codes) or as numbers e by coding the Δ_1^1 code C as a number e (hyperarithmetical codes for Δ_1^1 reals). In either case, the property of being a code is Π_1^1 and membership of a real Z in the set coded by C or e is a Δ_1^1 relation given that C and e are codes. Similarly, membership of a number n in a Δ_1^1 real with hyperarithmetical code e is a Δ_1^1 relation. We can pass in a Δ_1^1 way between these types of codes and the syntactic ones given by the formulas required in our definition of Δ_1^1 reals and classes given that all the objects are, in fact, codes. In this situation we often abuse notation by writing $Z \in C$ to denote the assertion that Z is in the class coded by C . When C and D are both codes, we use $D \subseteq C$ to denote the assertion that $\forall Z (Z \in D \rightarrow Z \in C)$ and similarly for $D \supseteq C$. These relations are then all Π_1^1 . These facts also imply that the predicate Z is $\Delta_1^1(X)$ is Π_1^1 . (We also use $C \supseteq \mathcal{A}$ for an arbitrary class \mathcal{A} of reals to mean that every real in the set coded by C is in \mathcal{A} .)*

Proposition 3.3 (Representation Theorem). *If \mathcal{V} is a Π_1^1 class then there is a Δ_1^1 function \mathcal{F} such that $Z \in \mathcal{V} \Leftrightarrow \mathcal{F}(Z) \in WO$ where WO is the class of reals Z which, viewed as a set of pairs of numbers, represents a well ordering. If $Z \in WO$, we write $|Z|$ for the ordinal represented by Z .*

Proposition 3.4 (Bounding). *If \mathcal{V} is a Π_1^1 class of reals, \mathcal{F} is as in Proposition 3.3 then \mathcal{V} is Δ_1^1 if and only if there is a bound $< \omega_1^{CK}$ on the order types of $\mathcal{F}(Z)$ for $Z \in \mathcal{V}$. Moreover, if \mathcal{V} contains only Δ_1^1 reals and \mathcal{G} is a Δ_1^1 subset of \mathcal{V} then such a bound (expressed as either a real or a number coding a recursive well-ordering) for $\{\mathcal{F}(Z) \mid Z \in \mathcal{G}\}$ (or even any Σ_1^1 subset of the Δ_1^1 well orderings) can be found in a Δ_1^1 way from the codes (or indices) for \mathcal{F} , \mathcal{G} and \mathcal{V} . As a consequence we may, in this case, divide \mathcal{V} into an increasing, continuous sequence $\langle \mathcal{V}_i \mid i < \omega_1^{CK} \rangle$ of uniformly Δ_1^1 sets given by $\mathcal{V}_i = \{Z \in \mathcal{V} \mid |\mathcal{F}(Z)| < i\}$.*

Remark 3.5. While we have not found an explicit statement in our references of the uniformity described in this bounding theorem, it can easily be deduced from the uniform version of the analogous theorem for sets of numbers (as in e.g. Sacks [1990, II.3.4]) by translating the real codes for ordinals $< \omega_1^{CK}$ to numbers in O of at least as large a rank given by Sacks [1990, I.4.3].

Proposition 3.6 (Gandy-Harrington Forcing). *We can define a general notion of forcing whose conditions are Σ_1^1 classes ordered by inclusion as extension. A simplified version of the proof of Theorem 2.1 that leaves out the diagonalization requirements shows that we may construct a generic G in any given Σ_1^1 class meeting any countable collection*

of dense sets. Thus we may use this forcing notion in any of the common ways. As usual, we will be interested in forcing over countable standard models of fragments of ZFC containing various specified reals. In addition to the typical results about forcing such as forcing equals truth, we note that, by the arguments in the proof of Theorem 2.1, a Π_1^1 sentence $\varphi(G)$ about the generic G is forced by a condition (Σ_1^1 set) \mathcal{P} if and only if $\forall Z \in \mathcal{P}(\varphi(Z))$. We also note that if $\langle G_0, G_1 \rangle$ is generic then both G_0 and G_1 are generic. (See Miller [1995, §30] for more about this forcing notion and Lemma 30.3 there for this last particular fact.) Absoluteness considerations will also play a role in our applications of this forcing.

As a notational convenience in proving our theorem, we can, by the Gandy basis theorem (Theorem 2.9) and the fact that $\omega_1^B = \omega_1^{CK}$ is a Σ_1^1 predicate (of B), assume without loss of generality that $\omega_1^B = \omega_1^{CK}$ for every $B \in \mathcal{B}$. (Note $\omega_1^B = \omega_1^{CK} \Leftrightarrow \forall e(\{e\}^B$ is a well-ordering $\rightarrow \exists i \exists f(f$ is an isomorphism of $\{i\}$ and $\{e\}^B)$.)

We begin with some crucial approximations to our class \mathcal{A} and an analysis of their properties.

Notation 3.7. We let $\mathcal{D}_B = \{C \mid C \text{ is a } \Delta_1^1(B) \text{ code \& } C \supseteq \mathcal{A}\}$ and $\mathcal{A}_B = \{A \mid (\forall C \in \mathcal{D}_B)(A \in C)\}$. Similarly, we let $\mathcal{A}_0 = \{A \mid A \text{ is a member of every } \Delta_1^1 \text{ class containing } \mathcal{A}\}$. For $B \in \mathcal{B}$, we let $\psi_B(Z)$ be a $\Sigma_1^1(B)$ formula defining \mathcal{A} .

Lemma 3.8. For $B \in \mathcal{B}$, \mathcal{D}_B is $\Pi_1^1(B)$ and \mathcal{A}_B is $\Sigma_1^1(B)$.

Proof. Fix $B \in \mathcal{B}$. For any real C , $C \in \mathcal{D}_B$ if and only if C is a $\Delta_1^1(B)$ code and $\forall Z(\psi_B(Z) \rightarrow Z \in C)$. As ψ_B is $\Sigma_1^1(B)$, both conjuncts here are $\Pi_1^1(B)$ by Proposition 3.2 and so \mathcal{D}_B is $\Pi_1^1(B)$ as required. The second claim now follows directly from the definition of \mathcal{A}_B and Proposition 3.2. (Rephrase the definition of \mathcal{D}_B in terms of number codes to make the quantifier count work.) \square

Lemma 3.9. For any reals A and B in \mathcal{B} with $\omega_1^{A,B} = \omega_1^{CK}$, $A \in \mathcal{A} \Leftrightarrow A \in \mathcal{A}_B$.

Proof. Clearly $A \in \mathcal{A}$ implies that $A \in \mathcal{A}_B$ for every $B \in \mathcal{B}$ by the definition of \mathcal{A}_B . For the other direction suppose $A \notin \mathcal{A}$. Let $\mathcal{F} \in \Delta_1^1(B)$ be as in Proposition 3.3 for the $\Pi_1^1(B)$ class $\mathcal{V} = \{Z \mid \neg \psi_B(Z)\}$ which is the complement of \mathcal{A} . Thus $\mathcal{F}(A)$ is a well ordering which is $\Delta_1^1(A, B)$ and so, by hypothesis, less than some $i < \omega_1^{CK}$. Thus the (obviously Σ_1^1) set $\{Z \mid \mathcal{F}(Z) < i\}$ is equal to the Π_1^1 set $\{Z \mid \neg \psi_B(Z) \ \& \ (i+1) \not\leq \mathcal{F}(Z)\}$ and so is $\Delta_1^1(B)$. It is disjoint from \mathcal{A} and has A as a member by our choice of i . The $\Delta_1^1(B)$ code C for its complement is then a member of \mathcal{D}_B not containing A as a member. This C is then a witness that $A \notin \mathcal{A}_B$ as required. \square

Lemma 3.10. \mathcal{A}_0 is Σ_1^1 .

Proof. Consider the real $J = \{e \mid e \text{ is a hyperarithmetic index for a } \Delta_1^1 \text{ code of a superset of } \mathcal{A}\} = \{e \mid e \text{ is a hyperarithmetic index for a real in } \mathcal{D}_B\}$. This real J is $\Pi_1^1(B)$ in every

$B \in \mathcal{B}$ by Lemma 3.8 and Proposition 3.2. So by Theorem 2.1 (formally applied to the complements) is Π_1^1 . Thus \mathcal{A}_0 which is the intersection of the sets coded by indices in J is Σ_1^1 : $Z \in \mathcal{A}_0 \Leftrightarrow \forall e(e \in J \rightarrow Z \text{ is in the set coded by } e)$. \square

Lemma 3.11. *If $B, C \in \mathcal{B}$ and $\omega_1^{B,C} = \omega_1^{CK}$, then $\mathcal{A}_B = \mathcal{A}_C$.*

Proof. If not, then we have, without loss of generality, an $A \in \mathcal{A}_C - \mathcal{A}_B$. So there is a code $D \in \mathcal{D}_B$ with $A \notin D$ and $A \in \mathcal{A}_C$. Now the nonempty class $\mathcal{W} = \{Z | Z \in \mathcal{A}_C \ \& \ Z \notin D\}$ is $\Sigma_1^1(B, C)$ by Lemma 3.8 and Proposition 3.2. Thus by the Gandy basis theorem (relative to B, C) (Theorem 2.9) there is a W in \mathcal{W} and so in $\mathcal{A}_C - \mathcal{A}_B$ with $\omega_1^{W,B,C} = \omega_1^{CK}$. Lemma 3.9, however, tells us that $W \in \mathcal{A}_B \Leftrightarrow W \in \mathcal{A} \Leftrightarrow W \in \mathcal{A}_C$ for a contradiction. \square

Lemma 3.12. *If $B, C \in \mathcal{B}$ and $\omega_1^{B,C} = \omega_1^{CK}$, then for every $X \in \mathcal{D}_B$ there is a $Y \in \mathcal{D}_B$ and a $Z \in \mathcal{D}_C$ such that $Y \subseteq X$ and $Z \subseteq Y \subseteq X$, i.e. Y and Z are codes for the same set.*

Proof. By Proposition 3.4, there is a uniformly Δ_1^1 continuous increasing sequences $\mathcal{D}_{B,i}$ ($i < \omega_1^{CK} = \omega_1^B$) with union \mathcal{D}_B . We can then set $\mathcal{A}_{B,i} = \{A | \forall C \in \mathcal{D}_{B,i}(A \in C)\}$. This sequence is clearly nested and continuous with intersection \mathcal{A}_B . As $\mathcal{D}_{B,i}$ and all its members are $\Delta_1^1(B)$, the $\mathcal{A}_{B,i}$ are also uniformly $\Delta_1^1(B)$ by Proposition 3.2 as we can convert to number codes. Similarly, we have $\mathcal{D}_{C,i}$ and $\mathcal{A}_{C,i}$ ($i < \omega_1^{CK} = \omega_1^C$). By Lemma 3.11 we know that for each $i < \omega_1^{CK}$ and $Z \in \overline{\mathcal{A}_{B,i}}$ there is a $j < \omega_1^{CK}$ such that $Z \in \overline{\mathcal{A}_{C,j}}$. By Proposition 3.4, there is a $k < \omega_1^{CK}$ such that for every $Z \in \overline{\mathcal{A}_{B,i}}$, $Z \in \overline{\mathcal{A}_{C,k}}$ and we can get k uniformly $\Delta_1^1(B, C)$. Of course, the analogous fact switching B and C is also true. Iterating and interleaving these $\Delta_1^1(B, C)$ functions starting with any $i < \omega_1^{CK}$ produces a $\Delta_1^1(B, C)$ increasing sequence of $k < \omega_1^{CK}$. By Proposition 3.4, this sequence has a bound and hence a supremum $l < \omega_1^{CK}$ and $\mathcal{A}_{B,l} = \mathcal{A}_{C,l}$.

Now consider any $X \in \mathcal{D}_{B,i}$ so $X \supseteq \mathcal{A}_{B,l}$ for any $l > i$ in ω_1^{CK} . We may now choose one such that $\mathcal{A}_{B,l} = \mathcal{A}_{C,l}$. As $\mathcal{A}_{B,l} \in \Delta_1^1(B)$ and contains \mathcal{A} , there is a code $Y \in \mathcal{D}_B$ for it. Similarly there is a code $Z \in \mathcal{D}_C$ for $\mathcal{A}_{C,l}$. As $\mathcal{A}_{B,l} = \mathcal{A}_{C,l}$, these are then the desired Y and Z . \square

Lemma 3.13. *For every $B \in \mathcal{B}$, $\mathcal{A}_B = \mathcal{A}_0$.*

Proof. Fix $B \in \mathcal{B}$. Clearly, it suffices to prove that $\forall X \in \mathcal{D}_B \exists Y \in \mathcal{D}_B (X \supseteq Y \ \& \ \mathcal{V} = \{Z | Z \in Y\} \in \Delta_1^1)$. (As this says there is, for each $X \in \mathcal{D}_B$, a Δ_1^1 code V for a Δ_1^1 class \mathcal{V} contained in the class coded by X and containing \mathcal{A} (as $Y \in \mathcal{D}_B$). This code shows that $\mathcal{A}_0 \subseteq \mathcal{A}_B$ by definition. On the other hand, $\mathcal{A}_B \subseteq \mathcal{A}_0$ for every B .)

Fix an $X \in \mathcal{D}_B$. Consider now the class $\mathcal{W} = \{C \in \mathcal{B} | (\forall Y \in \mathcal{D}_B)(X \supseteq Y \rightarrow \{Z | Z \in Y\} \notin \Delta_1^1(C))\}$. By Proposition 3.2, this class is $\Sigma_1^1(B)$. If it were nonempty then, by the Gandy basis theorem (relative to B) (Theorem 2.9), it would have a member C with $\omega_1^{B,C} = \omega_1^{CK}$. This would provide a counterexample to Lemma 3.12 and so \mathcal{W} is empty.

We now work with a countable standard model which contains B and satisfies a fragment of ZFC sufficient to guarantee the absoluteness of Σ_1^1 formulas. Note, for example, that all reals Δ_1^1 in B (and so all in \mathcal{D}_B) are in this model.

Let $G \in \mathcal{B}$ be a Gandy-Harrington generic over this model as in Proposition 3.6. As $G \notin \mathcal{W}$, there is a $Y \in \mathcal{D}_B$ such that $X \supseteq Y$ and $\{Z \mid Z \in Y\} \in \Delta_1^1(G)$. Fix a specific Δ_1^1 definition of this class from G , i.e. $\Sigma_1^1(G)$ formulas φ and θ such that $\forall Z(\varphi(G, Z) \leftrightarrow \neg\theta(G, Z))$, $\forall Z(Z \in Y \rightarrow \varphi(G, Z))$ and $\forall Z(Z \notin Y \rightarrow \theta(G, Z))$. As G is generic we have a Σ_1^1 \mathcal{P} forcing these sentences. Now consider the Σ_1^1 class $\mathcal{Q} = \{\langle C, C' \rangle \mid C, C' \in \mathcal{P} \ \& \ \exists Z(\varphi(C, Z) \ \& \ \theta(C', Z) \vee \varphi(C', Z) \ \& \ \theta(C, Z))\}$. If \mathcal{Q} is nonempty then there is a Gandy-Harrington generic $\langle C, C' \rangle \in \mathcal{Q}$. Each of C and C' is in \mathcal{P} and Gandy-Harrington generic by Proposition 3.6. Thus any Z witnessing that $\langle C, C' \rangle \in \mathcal{Q}$ would be a counterexample to one of the sentences above forced by \mathcal{P} and hence true of C and C' . Thus \mathcal{Q} is empty and so $Z \in Y \Leftrightarrow (\forall C \in \mathcal{P})(\neg\theta(C, Z))$ and $Z \notin Y \Leftrightarrow (\forall C \in \mathcal{P})(\neg\varphi(C, Z))$ and $\{Z \mid Z \in Y\}$ is Δ_1^1 as required. \square

We now prove our theorem on Σ_1^1 classes.

Proof of Theorem 3.1: We claim that $A \in \mathcal{A}$ if and only if $A \in \mathcal{A}_0$ (which is Σ_1^1 by Lemma 3.10) and one of the following two Σ_1^1 statements hold for a Σ_1^1 formula ψ that we will define below:

- (1) $\omega_1^A = \omega_1^{CK}$ or
- (2) $\omega_1^A > \omega_1^{CK} \rightarrow \psi(A)$.

Now $A \in \mathcal{A} \rightarrow A \in \mathcal{A}_0$ by the definition of \mathcal{A}_0 . So we may assume that $A \in \mathcal{A}_0$ and show that $A \in \mathcal{A} \Leftrightarrow$ (1) or (2) holds. If (1) holds then by the Gandy basis theorem (Theorem 2.9) (relative to A) we may choose a $B \in \mathcal{B}$ with $\omega_1^{A,B} = \omega_1^{CK}$. Now by Lemma 3.9, $A \in \mathcal{A} \Leftrightarrow A \in \mathcal{A}_B$ while $A \in \mathcal{A}_B \Leftrightarrow A \in \mathcal{A}_0$ by Lemma 3.13. Thus, in this case, $A \in \mathcal{A} \Leftrightarrow A \in \mathcal{A}_0$ as required.

Assume then that (1) fails and so the hypothesis of (2) holds. We now must argue that we have a Σ_1^1 formula $\psi(A)$ that, under these assumptions, is equivalent to $A \in \mathcal{A}$. As mentioned just before Theorem 2.9, there is a pair if Σ_1^1 formulas $\varphi(X, n)$ and $\theta(X, n)$ which define O and its complement for any X with $\omega_1^X > \omega_1^{CK}$. By the Kleene basis theorem (Theorem 2.9) there is a recursive index computing a $B \in \mathcal{B}$ from O . By the hypothesis of our theorem there is a $\Sigma_1^1(B)$ formula $\psi_B(Z)$ defining \mathcal{A} . Thus there is a Σ_1^1 formula $\hat{\psi}(X, Z)$ which defines \mathcal{A} from any X with $\omega_1^X > \omega_1^{CK}$. We now take our desired ψ to be $\hat{\psi}(A, A)$. \square

As a final comment, we point out that if we had only wanted to prove Theorem 3.1 in the Δ_1^1 case we would have a simple proof along the lines of the last paragraph of the proof of Lemma 3.13. This argument also gives a proof of the analog for classes of reals of the Δ_1^1 case of Theorem 2.6.

Theorem 3.14. *If \mathcal{B} is a nonempty Σ_1^1 class of reals and \mathcal{X}_n a countable sequence of classes of reals none of which is Δ_1^1 , then there is a $G \in \mathcal{B}$ such that no \mathcal{X}_n is Δ_1^1 in G .*

Proof. Note that if $\mathcal{X}_m \not\subseteq \Delta_1^1(B)$ for every $B \in \mathcal{B}$ then any $G \in \mathcal{B}$ works for \mathcal{X}_m . Thus we may assume that for every n there is a $B_n \in \mathcal{B}$ and φ_n and θ_n Σ_1^1 formulas with two free real variables which, with B_n for the first variable, define \mathcal{X}_n and its complement. Let $G \in \mathcal{B}$ be a Gandy-Harrington generic over a countable standard model of a sufficient fragment of ZFC containing the B_n . We claim no \mathcal{X}_n is $\Delta_1^1(G)$. If not, let φ and θ be $\Sigma_1^1(G)$ formulas defining some \mathcal{X}_n and its complement. Let \mathcal{P} be a condition which forces that $(\forall Z)(\varphi_n(B_n, Z) \rightarrow \varphi(G, Z))$, $(\forall Z)(\theta_n(B_n, Z) \rightarrow \theta(G, Z))$ and $(\forall Z)(\varphi(G, Z) \leftrightarrow \neg\theta(G, Z))$. Now consider the Σ_1^1 class $\mathcal{Q} = \{\langle C, C' \rangle \mid C, C' \in \mathcal{P} \ \& \ \exists Z(\varphi(C, Z) \ \& \ \theta(C', Z) \vee \varphi(C', Z) \ \& \ \theta(C, Z))\}$. If \mathcal{Q} is nonempty then there is a Gandy-Harrington generic $\langle C, C' \rangle \in \mathcal{Q}$. Each of C and C' is in \mathcal{P} and Gandy-Harrington generic by Proposition 3.6. Thus any Z witnessing that $\langle C, C' \rangle \in \mathcal{Q}$ would be a counterexample to one of the sentences above forced by \mathcal{P} and hence true of C and C' . Thus \mathcal{Q} is empty and so $Z \in \mathcal{X}_n \Leftrightarrow (\forall C \in \mathcal{P})(\neg\theta(C, Z))$ and $Z \notin \mathcal{X}_n \Leftrightarrow (\forall C \in \mathcal{P})(\neg\varphi(C, Z))$ and so \mathcal{X}_n is Δ_1^1 for the desired contradiction. \square

Corollary 3.15. *Any class \mathcal{A} of reals which is Δ_1^1 in every member of a Σ_1^1 class \mathcal{B} of reals is Δ_1^1 .*

We do not know if the full analog of Theorem 2.6 for classes of reals, i.e. Theorem 3.14 with Δ_1^1 replaced by Σ_1^1 , is also true.

4 Bibliography

Andrews, U. and Miller, J. S. [2015], Spectra of theories and structures, Proceedings of the American Mathematical Society **143**, 1283-1298.

Hinman, P. G. [1978], *Recursion Theoretic Hierarchies*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin.

Miller, A. W. [1995], *Descriptive Set Theory and Forcing*, Lecture Notes in Logic, Springer-Verlag, Berlin.

Moschovakis, Y. N. [1980], *Descriptive Set Theory*, Studies in Logic and the Foundations of Mathematics **100**, North-Holland, Amsterdam.

Sacks, G. E. [1990], *Higher Recursion Theory*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin.

Simpson, S. G. [2009], *Subsystems of Second Order Arithmetic 2nd ed.*, *Perspectives in Logic*, ASL and Cambridge University Press, New York.