

DETERMINACY FROM STRONG REFLECTION

JOHN STEEL AND STUART ZOBLE
SEPTEMBER 16, 2008

ABSTRACT. The Axiom of Determinacy holds in the inner model $L(\mathbb{R})$ assuming Martin's Maximum for partial orderings of size c .

1. INTRODUCTION

A theorem of Neeman gives a particularly elegant sufficient condition for a set B of reals to be determined: B is determined if there is a triple (M, τ, Σ) which *captures* B in the sense that M is a model of a sufficient fragment of set theory, τ is a forcing term in M with respect to the collapse of some Woodin cardinal δ of M to be countable, and Σ is an $\omega + 1$ -iteration strategy for M such that

$$B \cap N[g] = i(\tau)_g,$$

whenever $i : M \rightarrow N$ is an iteration map by Σ , and g is generic over N for the collapse of $i(\delta)$. The core model induction, subject of the forthcoming book [16], is a method pioneered by Woodin for constructing such triples (M, τ, Σ) by induction on the complexity of the set B . It seems to be the only generally applicable method for making fine consistency strength calculations above the level of one Woodin cardinal. We employ this method here to establish that the Axiom of Determinacy holds in the inner model $L(\mathbb{R})$ from consequences of the maximal forcing axiom $\text{MM}(c)$, or Martin's Maximum for partial orderings of size c . The particular consequences we use are the saturation of the nonstationary ideal on ω_1 , and the simultaneous reflection principle $\text{WRP}_{(2)}(\omega_2)$ asserting that for any stationary subsets S and T of $[\omega_2]^\omega$ there is an ordinal $\delta < \omega_2$ so that $S \cap [\delta]^\omega$ and $T \cap [\delta]^\omega$ are both stationary in $[\delta]^\omega$.

Theorem 1. $\text{WRP}_{(2)}(\omega_2)$ plus NS saturated implies $\text{AD}^{L(\mathbb{R})}$.

Corollary 2. $\text{MM}(c)$ implies $\text{AD}^{L(\mathbb{R})}$.

2000 *Mathematics Subject Classification.* Primary 03E45; Secondary 03E35.

Key words and phrases. Stationary Reflection, Nonstationary Ideal, Determinacy.

This theorem, obtained in late 2000, builds on Woodin's proof of PD from the same hypotheses (9.85 of [25]), and represents the first proof of the consistency of the Axiom of Determinacy from Forcing Axioms. The first author subsequently obtained the same conclusion in [22] from a single failure of square (and hence from PFA) building on Woodin's theorem that PFA together with an inaccessible gives AD in the Solovay model. Unlike that proof, which relies on covering lemmas to produce the models required for the induction step, we use the generic embedding derived from the saturated ideal. This has its precedents in the first author's proof of Δ_2^1 determinacy from a presaturated ideal on ω_1 together with a measurable cardinal, and in Woodin's proof, via the core model induction, of $\text{AD}^{L(\mathbb{R})}$ from an ω_1 -dense ideal on ω_1 .

While our theorem represents the best known lower bound for the consistency strength of $\text{MM}(c)$, this principle is believed to be much stronger. In Chapter 6 we discuss some results suggesting that the arguments here cannot take us much farther than $\text{AD}^{L(\mathbb{R})}$, and some extensions of the main theorem which could plausibly yield an equiconsistency result at the level of ω^2 Woodin cardinals from modified hypotheses.

Acknowledgement. Some of the arguments presented here have their origin in [26], written under the supervision of Hugh Woodin. The second author would like to thank him for his guidance and support.

2. FRAMEWORK OF THE INDUCTION

Let us recall some terminology from [22]

Definition 3. Let $U \subseteq \mathbb{R}$, and $k < \omega$. Let N be countable and transitive, and suppose $\delta_0, \dots, \delta_k, S$, and T are such that

- (a) $N \models \text{ZFC} \wedge \delta_0 < \dots < \delta_k$ are Woodin cardinals,
- (b) $N \models S, T$ are trees which project to complements after the collapse of δ_k to be countable, and
- (c) there is an $\omega_1 + 1$ -iteration strategy Σ for N such that whenever $i: N \rightarrow P$ is an iteration map by Σ and P is countable, then $p[i(S)] \subseteq U$ and $p[i(T)] \subseteq \mathbb{R} \setminus U$.

Then we say that N is a *coarse* (k, U) -Woodin mouse, as witnessed by $S, T, \Sigma, \delta_0, \dots, \delta_k$.

Definition 4. W_α^* denotes the following assertion. If $U \subseteq \mathbb{R}$, and there are scales $\vec{\phi}$ and $\vec{\psi}$ on U and $\mathbb{R} \setminus U$ respectively such that $\vec{\phi}^*, \vec{\psi}^* \in J_\alpha(\mathbb{R})$, where $\vec{\phi}^*$ and $\vec{\psi}^*$ are the sequences of prewellorders associated to the scales, then for all $k < \omega$ and $x \in \mathbb{R}$ there are N, Σ such that

- (1) $x \in N$, and N is a coarse (k, U) -Woodin mouse, as witnessed by Σ , and
- (2) $\Sigma \upharpoonright \text{HC} \in J_\alpha(\mathbb{R})$.

Our core model induction will show that

$$V[g] \models \forall \alpha W_\alpha^*,$$

whenever $g \subset \text{Col}(\omega, \omega_1)$ is V -generic. From W_α^* we get a version of mouse capturing by fine-structural mice. Let us recall the relevant definitions from [22]. To any Σ_1 formula $\theta(v)$ we associate formulae $\theta^k(v)$ for $k \in \omega$, such that θ^k is Σ_k , and for any γ and any real x ,

$$J_{\gamma+1}(\mathbb{R}) \models \theta[x] \Leftrightarrow \exists k < \omega J_\gamma(\mathbb{R}) \models \theta^k[x].$$

Our fine-structural witnesses are as follows.

Definition 5. Suppose $\theta(v)$ is a Σ_1 formula (in the language of set theory expanded by a name for \mathbb{R}), and z is a real; then a (θ, z) -witness is an ω -sound, $(\omega, \omega_1, \omega_1 + 1)$ -iterable z -mouse \mathcal{N} in which there are $\delta_0 < \dots < \delta_9$, S , and T such that \mathcal{N} satisfies the formulae expressing

- (a) ZFC,
- (b) $\delta_0, \dots, \delta_9$ are Woodin,
- (c) S and T are trees on some $\omega \times \eta$ which are absolutely complementing in $V^{\text{Col}(\omega, \delta_9)}$, and
- (d) For some $k < \omega$, $p[T]$ is the Σ_{k+3} -theory (in the language with names for each real) of $J_\gamma(\mathbb{R})$, where γ is least such that $J_\gamma(\mathbb{R}) \models \theta^k[z]$.

Definition 6. W_α is the assertion: if $\theta(v)$ is Σ_1 , $z \in \mathbb{R}$, and $J_\alpha(\mathbb{R}) \models \theta[z]$, then there is a (θ, z) -witness \mathcal{N} whose associated iteration strategy, when restricted to countable iteration trees, is in $J_\alpha(\mathbb{R})$.

We have

Lemma 7. *Assume W_α^* holds; then*

- (a) $J_\alpha(\mathbb{R}) \models \text{AD}$, and
- (b) W_α holds if α is a limit ordinal.

See [22] for a proof of (b), which is essentially Woodin's mouse set theorem for $L(\mathbb{R})$. Part (a) is an easy exercise for our intended reader.¹ We note that W_α easily implies other forms of capturing by fine-structural mice, and in particular:

¹It follows directly from Neeman's theorem in [12], but one doesn't need that much firepower. The results of Martin-Steel [9], together with Woodin's genericity iterations (see [20]), give it easily.

Lemma 8. *Assume W_α holds; then if a is countable transitive, and $b \subseteq a$, b is ordinal definable from parameters in $a \cup \{a\}$ over some $J_\gamma(\mathbb{R})$, where $\gamma < \alpha$, then there is a a -premouse \mathcal{M} such that $b \in \mathcal{M}$, and $J_\alpha(\mathbb{R}) \models \mathcal{M}$ is ω_1 -iterable.*

In our proof of $W_{\beta+1}^*$, we get the capturing mice we need in $V[g][G]$, where $G \subset (P(\omega_1)/NS)^V$ is generic over $V[g]$. We then use an inductively maintained resemblance between $V[g]$ and $V[g][G]$ to find these mice in $V[g]$. This leads us to a second induction hypothesis.

Definition 9. I_α is the assertion: whenever $h \times G$ is $\text{Col}(\omega, \omega_1) \times (P(\omega_1)/NS)^V$ -generic over V , there is a Σ_1 embeddings

$$\pi: J_\alpha(\mathbb{R})^{V[h]} \rightarrow J_\alpha(\mathbb{R})^{V[G][h]}$$

such that $\pi \upharpoonright \omega\alpha$ is the identity.

An easy consequence of Lemma 7(a) and our induction hypotheses together is

Lemma 10. *Assume I_α holds. Let $g \times G$ be $\text{Col}(\omega, \omega_1) \times (P(\omega_1)/NS)^V$ -generic over V , and suppose $V[g] \models W_\alpha^*$; then AD holds in $J_\alpha(\mathbb{R})^{V[g]}$, and in $J_\alpha(\mathbb{R})^{V[G][g]}$.*

We shall see later that AD holds in $J_\alpha(\mathbb{R})^{V[G]}$ as well; see 39.

As mentioned above, we shall be proving that W_α^* holds in $V[g]$, by induction on α . Clearly, the only stages which matter are the *critical* ones, where

Definition 11. An ordinal β is *critical* just in case there is some set $U \subseteq \mathbb{R}$ such that U and $\mathbb{R} \setminus U$ admit scales in $J_{\beta+1}(\mathbb{R})$, but U admits no scale in $J_\beta(\mathbb{R})$.

Once again, we are identifying a scale with the *sequence* of its prewellorderings here. Clearly, we need only show that $W_{\beta+1}^*$ holds whenever β is critical, in order to conclude that W_α^* holds for all α . It follows from [23] that if β is critical, then $\beta + 1$ is critical. Moreover, if β is a limit of critical ordinals, then β is critical if and only if $J_\beta(\mathbb{R})$ is not an admissible set. Letting β be critical, we then have the following possibilities

- (1) $\beta = \eta + 1$, for some critical η ;
- (2) β is a limit of critical ordinals, and either
 - (a) $\text{cof}(\beta) = \omega$, or
 - (b) $\text{cof}(\beta) > \omega$, but $J_\beta(\mathbb{R})$ is not admissible;
- (3) $\alpha = \sup(\{\eta < \beta \mid \eta \text{ is critical}\})$ is such that $\alpha < \beta$, and either
 - (a) $[\alpha, \beta]$ is a Σ_1 gap, or

(b) $\beta - 1$ exists, and $[\alpha, \beta - 1]$ is a Σ_1 gap.

We shall call (3) the *admissible case*, because it corresponds precisely to crossing a Σ_1 gap whose initial ordinal is admissible.

The hypothesis I_α is not used as an input in the arguments of the admissible case but is produced as an output. On the other hand, it is used in the inadmissible case. I_α originates in [26], where it is part of a proof that the saturation of NS and $WRP_{(2)}(\omega_2)$ imply W_α^* for α the first admissible over the reals. Our argument here follows the overall structure of [26] pretty closely. What we add are some techniques for getting past admissible ordinals using hybrid strategy mice. These techniques were also used in [22].

A final remark on the organization of our proof. It might perhaps be more natural to think of ourselves as proving that W_α^* holds in V , for all α . As in any core model induction, given a critical ordinal β , the first step toward $W_{\beta+1}^*$ in V is to find a (hybrid) mouse operator J which codes up truth at the level of the first pointclass $\sum_n^{J_\beta(\mathbb{R})}$ having the scale property. In order to prove $W_{\beta+1}^*$, we then need to capture truth over $J_\beta(\mathbb{R})$ in full, and for this we need to construct the “ k -many $-J$ -Woodins” operators M_k^J , for all k . These “successor steps” are where core model theory (relativised to J) comes in. The core model theory in our argument requires that J first be extended to $H(\omega_3)$, in a way that is consistent with its images $\pi(J)$ under NS -generic ultrapower maps. The extension to $H(\omega_2)$ is equivalent to an extension to $H(\omega_1)^{V[g]}$. It is at this point that we must consider W_γ^* in $V[g]$, where $\gamma < \pi(\beta)$ for some NS -generic π . Extending J involves showing that $\pi(\beta)$ is independent of the NS -generic, and W_γ^* and holds in $V[g]$ at all $\gamma \leq \pi(\beta)$. This subinduction in $V[g]$ leading to an extension of J is also where I_γ is used.

3. THE SUCCESSOR STEP

The successor step in a core model induction is the step from a model operator J to the *one J -Woodin* operator. There are two important sorts of model operators for which one needs to make this step, the *mouse operators* and the *hybrid mouse operators*. We shall consider only mouse operators in this section, but the proof works without much change for model operators in general. We shall consider hybrid mouse operators in the last section of the paper.

Our proof builds on the proofs of the following theorems.

Theorem 12 (Steel, [18]). *Assume there is a measurable cardinal and a presaturated ideal on ω_1 ; then Δ_2^1 determinacy holds.*

Theorem 13 (Woodin, [25][9.85]). *(Woodin) Assume $\text{WRP}_{(2)}(\omega_2)$ and that NS is saturated; then PD holds, and continues to hold in the universe after ω_2 is collapsed.*

Roughly, the proof of the first theorem supplies the core model theory we need, and the proof of the second shows how to integrate the core model theory into a core model induction.

Woodin's argument shows inductively that $H(\omega_3)$ is closed under the $M_n^\#$ mouse operator for each $n < \omega$. We shall give the proof in somewhat greater generality. Roughly, rather than doing the first ω steps of a core model induction, we will be doing the general successor step. For that, we need to have as data an operator J such that $H(\omega_3)$ is closed under J ; we then show that $H(\omega_3)$ is closed under $M_1^{J,\#}$. ($M_1^\#(a) = M_1^{J,\#}(a)$, for $J(\mathcal{P}) = \text{rud}(\mathcal{P})$.) Here we shall just consider the case that J is a *first order mouse operator*. In the last section we shall be forced to consider more general J .

Since it requires only a little additional work, we shall assume only $\text{WRP}_{(2)}(\omega_2)$, there is a presaturated ideal on ω_1 , and $2^{\omega_1} \leq \omega_2$. Woodin has shown that $\text{WRP}_{(2)}(\omega_2)$ and the saturation of NS together imply that $2^{\omega_1} \leq \omega_2$; see 30 below. We believe that in fact $\text{WRP}_{(2)}(\omega_2)$ together with a presaturated ideal on ω_1 should be enough for our argument, but have not checked that carefully. See 31 below.

The properties of our initial mouse operator J that make the general successor step possible are that it

- (1) J condenses well,
- (2) J relativises well,
- (3) J determines itself on generic extensions, and
- (4) I -generic embeddings move J to itself.

Here I is our presaturated ideal. We now explain these properties. The reader should keep in mind the example $J_n(b) = M_n^\#(b)$, which has all of them. That it has property (4) is one of the main things our induction will show.

We need to consider premeice over some transitive set b , with a distinguished parameter $a \in b$. In this context, we shall always in this paper assume that b is *selfwellordered*, that is, equipped with a wellorder that is (uniformly over all b under consideration) rudimentary in a .² The language \mathcal{L}_0 of such relativized premeice is the language of premeice, together with an additional constant symbols \dot{b} and \dot{a} for the set thrown in at the bottom and its distinguished element.

²We are essentially working with b 's which are sets of ordinals, and sweeping some codings under the rug.

Iterations of a relativized premouse M are always by extenders on its coherent sequence, all of which have critical points above \dot{b}^M . A relativised premouse M is countably iterable if whenever $\pi : N \rightarrow M$ is \mathcal{L}_0 -elementary with N countable then N is $\omega_1 + 1$ iterable. Fix b transitive. Any two sound countably iterable premice over b which project to b are comparable (see [20]). (Their possibly different parameters are irrelevant at this point.) The lower part closure of b is defined as the union of all b -premise N which are countably iterable, sound, and satisfy $\rho_\omega(M) = b$. $Lp(b)$ can be regarded as a countably iterable b -premise in its own right, over any $a \in b$. We sometimes write $Lp^a(b)$ when we want to think of it this way, but we shall drop the superscript a when it is safe to do so.

Definition 14. Let $\nu \geq \omega_1$ be regular, and $a \in H(\nu)$. Let φ be an rQ -sentence of \mathcal{L}_0 . Suppose that for any transitive, selfwellordered $b \in H(\nu)$ such that $a \in b$, there is a countably iterable premouse M over b with parameter a such that $M \models \varphi$; then we set

$$J_\varphi(b) = Lp^a(b) \upharpoonright \gamma,$$

where γ is least such that $Lp^a(b) \models \varphi$. We call the map $b \mapsto J_\varphi(b)$ a (first order) (ν, a) mouse operator. We say that J_φ is defined on the $H(\nu)$ -cone above a .

Remark 15. In some contexts, the countable iterability requirement we have imposed above is too onerous. For example, if $H(\omega_1)$ is closed under sharps, then ω_1 -iterability is enough to identify the true M_1^\sharp . We don't need full $\omega_1 + 1$ iterability. This fact will be important for us when we consider mice in $V[h]$, for h generic over $\text{Col}(\omega, \omega_2)$. Our hypotheses are consistent with $V = L[A]$ for some $A \subseteq \omega_3$, so they do not imply that M_1^\sharp is $\omega_1 + 1$ iterable in $V[h]$. (An iteration to make A generic will provide a counterexample.) We do have to consider countable mice in $V[h]$, in order to show the mouse operators in smaller models behave well. However, we don't need to consider mouse operators in $V[h]$, so we can stick with the $\omega_1 + 1$ iterability requirement of 14.

An important property of first order mouse operators is that they *condense well*, in the sense of the following lemma.

Lemma 16. *Let J be a first order mouse operator with parameter a . Let $b \in \text{dom}(J)$, and let $\pi : \mathcal{M} \rightarrow J(b)$ be rQ -elementary, with $\pi(c) = b$, and $\pi \upharpoonright TC(a \cup \{a\})$ being the identity; then $c \in \text{dom}(J)$, and $\mathcal{M} = J(c)$.*

One theme of this paper is that simultaneous reflection can be used to lift closure under certain operations from $P(\omega_1)$ to $P(\omega_2)$. In [25],

Woodin gives a proof that under $\text{WRP}_{(2)}(\omega_2)$, closure of $P(\omega_1)$ under sharps entails closure of $P(\omega_2)$ under sharps. His proof of Theorem 10 involved analogous arguments for the $M_n^\#$ operation. The following is a straightforward generalization to first order mouse operators. There is a related argument in [27], where it is shown that under $\text{WRP}_{(2)}(\omega_2)$, ω_1 -Universally Baire self-justifying systems are ω_2 -Universally Baire. What is key to all the arguments is that the function being extended from $P(\omega_1)$ to $P(\omega_2)$ condenses well.

Definition 17. For regular cardinals $\kappa < \lambda$, we say that *Mouse Reflection holds at (κ, λ)* iff for every $a \in H(\kappa)$, every (κ, a) -mouse operator can be extended to a (λ, a) -mouse operator. If $\lambda = \kappa^+$, we say that *Mouse Reflection holds at κ* .

Lemma 18. $\text{WRP}_{(2)}(\omega_2)$ implies *Mouse Reflection at ω_2* .

Proof. Let $J = J_\varphi$ be a first order mouse operator with parameter a defined on the $H(\omega_2)$ cone above a . Fix a transitive, selfwellordered b in $H(\omega_3)$ such that $a \in b$. We must show that there is a countably iterable b -premouse with parameter a that satisfies φ .

For $\sigma \in [b]^\omega$ such that $a \in \sigma$, let b_σ be the transitive collapse of σ , and let a_σ be the collapse of a .

Claim. For club many $\sigma \in [b]^\omega$, there is a countably iterable b_σ -premouse \mathcal{M} , with parameter a_σ , such that $\mathcal{M} \models \varphi$.

Proof. If not, then one-set stationary reflection for $[b]^\omega$ gives us an $X \subseteq b$ such that $|X| = \omega_1$, $TC(a \cup \{a\}) \subseteq X$, and for stationary many $\sigma \in [X]^\omega$ the conclusion of the claim fails. But let b_X be the collapse of X , and note that a is fixed by this collapse. Let $\mathcal{N} = J_\varphi(b_X)$. It is clear that for club many $\sigma \in [X]^\omega$, there is a countable $Y \prec \mathcal{N}$ such that $Y \cap X = \sigma$. For such σ , the collapse of Y is an \mathcal{M} as in the claim. Contradiction. \square

For $c \in \text{dom}(J)$, let $T^J(c) = \{\langle \psi, t \rangle \mid t \in c^{<\omega} \wedge \psi \in \mathcal{L}_0 \wedge J(c) \models \psi[t]\}$. Since $J(c)$ projects to c , it is coded by $T^J(c)$. Our goal is to define the appropriate theory to be $T^J(b)$. To this end, for $t \in [b]^{<\omega}$ and $\psi(v)$ an \mathcal{L}_0 formula, put

$$S_{\psi,t} = \{\sigma \in [b]^\omega \mid t_\sigma \in T^J(b_\sigma)\}.$$

Claim. For any ψ, t , one of $S_{\psi,t}$ and $S_{\neg\psi,t}$ contains a club in $[b]^\omega$.

Proof. Otherwise we can find $X \subseteq b$ such that $|X| = \omega_1$, $t \in X$, $TC(a \cup \{a\}) \subseteq X$, and both $S_{\psi,t}$ and $S_{\neg\psi,t}$ are stationary in $[X]^\omega$. Let b_X be the transitive collapse of X , and t_X the image of t under the collapse. The collapse fixes a . Arguing as in the first claim, we

see that if $J(b_X) \models \psi[t_X]$, then $S_{\psi,t}$ contains a club in $b_X]^\omega$, and if $J(b_X) \models \neg\psi[t_X]$, then $S_{\neg\psi,t}$ contains a club in $[b_X]^\omega$. In either case, we have a contradiction. \square

Now we put

$$\langle \psi, t \rangle \in T \Leftrightarrow S_{\psi,t} \text{ contains a club in } [b]^\omega.$$

It is easy to see that T is the theory with parameters of a countably iterable b -premouse \mathcal{M} with parameter a satisfying φ . \square

Remark 19. We believe that with more work, one can show that simultaneous reflection for pairs of stationary subsets of ω_2 implies Mouse Reflection at ω_2 .

Definition 20. Let J be a (ν, a) -mouse operator, and let \mathcal{M} be a b -premouse with parameter a ; then we say \mathcal{M} is *J -level-closed* iff whenever $\eta < \xi < \nu$ and ξ is a cardinal of \mathcal{M} , then $J(\mathcal{M}|\xi) \trianglelefteq \mathcal{M}$.

For any (ν, a) -mouse operator J and set b in its domain there is a corresponding minimal J -level-closed premouse $L^J(b)$, obtained by concatenating extender sequences. A condensation argument shows that $L^J(b)$ is a bone-fide b -premouse with parameter a . It has ordinal height $o(L^J(b)) = \nu$, and it is countably iterable.

If they are defined on the $H(\nu)$ cone above a , then J^\sharp , J^* , and J_n^w are themselves (ν, a) -mouse operators. The reader should see [22] and [16] for background.

Definition 21. A (ν, a) -mouse operator J *relativizes well* iff

- (1) there is a formula $\theta(u, v, w, z)$ such that whenever $b, c \in \text{dom}(J)$, $b \in c$, and N is a transitive model of ZFC^- such that $J(c) \in N$, then $J(b) \in N$ and $J(b)$ is the unique $x \in N$ such that $N \models \theta[x, a, b, J(c)]$, and
- (2) if $b \in \text{dom}(J)$ and η is a cutpoint of $J(b)$, then $J(J(b)|\eta)$ is not a proper initial segment of $J(b)$.

We shall only be dealing with operators that relativize well. Clause (2) is used in the proof of

Lemma 22. *Suppose that J is a (ν, a) mouse operator that relativizes well; then for all $b \in \text{dom}(J)$, $J(b)$ is ν -iterable.*

Proof. We show that $J(b)$ is iterable by the strategy of choosing the unique cofinal branch b of \mathcal{T} such that $Q(b, \mathcal{T}) \trianglelefteq J(\mathcal{M}(\mathcal{T}))$. The usual reflection argument shows that this works. \square

If J relativizes well, then a J -level-closed premouse that satisfies ZFC^- is fully closed under J . Because of this, we shall sometimes say “ J -closed” when we mean “ J -level-closed”. Note that if $n \leq \omega$ and the M_n^\sharp operator is total on $H(\nu)$, then it is a $(\nu, 0)$ -mouse operator that relativizes well. Another very useful property of the M_n^\sharp operator is that it determines itself on generic extensions, in the following sense.

Definition 23. Let J be a (ν, a) -mouse operator; we say that J *determines itself on generic extensions* iff for all b in the $H(\omega_1)$ -cone over a , and all g that are \mathbb{P} -generic over $J(b)$ for some $\mathbb{P} \in \text{rud}(b)$, we have that $J(b)[g] = J(\langle b, g \rangle)$.

Notice here that $J(b)[g]$ can be regarded as a premouse over $\langle b, g \rangle$, because the forcing is small with respect to extenders on the sequence of $J(b)$. Definition 23 requires that so regarded, $J(b)[g]$ is just $J(\langle b, g \rangle)$. It is clear that if the M_n^\sharp operator is defined on $H(\omega_1)$, then it determines itself on generic extensions.

It is shown in [21][section 4] that under AD, every mouse operator on $H(\omega_1)$ determines itself on generic extensions in some $H(\omega_1)$ cone.

Although we stated definition 23 in terms of generic extensions of countable models that exist in V , condensation leads to extendibility beyond V :

Lemma 24. *Let $a \in H(\omega_1)$, and let $J = J_\varphi$ be a first order (ν, a) -mouse operator, where $\nu \geq \omega_1$ is regular. Suppose that $J \upharpoonright H(\omega_1)$ relativizes well and determines itself on generic extensions. Let h be V -generic over some partial order of size $< \nu$; then $V[h]$ satisfies “for any c in the $H(\nu)$ cone over a , there is a c -premouse \mathcal{M} with parameter a such that $\mathcal{M} \models \varphi$, and \mathcal{M} is ν -iterable.*

Proof. Let θ be the formula witnessing that J relativizes well. Let $\tau \in H(\nu)$ be a term such that $c = \tau_h$. Let b be in the $H(\nu)^V$ cone over a with $\tau \in b$.

The canonical re-arrangement of $J(b)[h]$ is a $\langle b, h \rangle$ -premouse with parameter a satisfying φ . For if not, we can find in V a countable elementary submodel N of V such that forcing over N yields g , where $\bar{J}(\bar{b})[g]$ does not re-arrange to a premouse satisfying φ . But $\bar{J} \subseteq J \upharpoonright H(\omega_1)$ because J condenses well. Since J determines itself on generic extensions, we have a contradiction.

Let \mathcal{N} be the canonical re-arrangement of $J(b)[h]$. Let θ be the formula witnessing that $J \upharpoonright H(\omega_1)$ relativizes well. Let $M \models \text{ZFC}^-$ be transitive, with $b, h \in M$. A Lowenheim-Skolem argument like that in the last paragraph shows that there is a unique c -premouse \mathcal{M}

satisfying φ such that for some (equivalently all) transitive $S \models \mathbf{ZFC}^-$, $S \models \theta[\mathcal{M}, a, c, \mathcal{N}]$. This is our desired \mathcal{M} . \square

If $\nu = \omega_1$ in $V[h]$, the premouse \mathcal{M} in the conclusion of 24 may not be $\omega_1 + 1$ -iterable in $V[h]$. See remark 15 above. It is, however, definable in $V[h]$ from $J \upharpoonright V$ and c , uniformly over all $V[h]$. Thus we write

$$\mathcal{M} = J^h(c)$$

for the c -premouse satisfying φ and obtained from J as above. If $\nu > \omega_1$ in $V[h]$, we get

Corollary 25. *Under the hypotheses of 24, if $\nu > \omega_1$ in $V[h]$, then J^h is a (ν, a) mouse operator extending J , and J^h relativizes well, and determines itself on generic extensions.*

Definition 26. Let I be a presaturated ideal on ω_1 , and suppose $2^{\omega_1} = \omega_2$. Let $a \in H(\omega_1)$, and let J be a (ω_3, a) mouse operator that relativizes well and determines itself on generic extensions. We say that J is *I -absolute* iff whenever $\pi: V \rightarrow \text{Ult}(V, G)$ is a generic embedding associated to some (I^+, \subseteq) -generic G , and J^G is the extension of J to $V[G]$ given by corollary 25, then

$$\pi(J) \upharpoonright H(\omega_1)^{V[G]} = J^G \upharpoonright H(\omega_1)^{V[G]}.$$

If J is I -absolute, then in fact $\pi(J) \subseteq J^G$ in full, by a simple Lowenheim-Skolem argument based on the condensation property of J .

Definition 27. Let J be a (ν, a) -mouse operator, and let $b \in \text{dom}(J)$; then

- (1) $J^\sharp(b)$ is the minimal active, countably iterable, J -level-closed b -premouse, if there is one.
- (2) $J^*(b) = (J^\sharp)^\sharp(b)$, if it exists.
- (3) For $n \geq 1$, $J_n^w(b) = M_n^{J, \sharp}(b)$ is the minimal active, countably iterable, J -level-closed b -premouse satisfying “there are n Woodin cardinals above $o(b)$ ”, if there is one. We put $J^w(b) = J_1^w(b)$.

If they are defined on the $H(\nu)$ cone above a , then J^\sharp , J^* , and J_n^w are themselves (ν, a) -mouse operators. The reader should see [22] and [16] for background.

The successor step in a core model induction is the step from J -closure to J^w -closure. We are ready now to execute it, in the case that our given J is a first order mouse operator.

Theorem 28. *Suppose that I is a presaturated ideal on ω_1 , $2^{\omega_1} = \omega_2$, and Mouse Reflection holds at ω_2 . Let $a \in H(\omega_1)$, and suppose that J*

is a first order (ω_3, a) -mouse operator that relativizes well, determines itself on generic extensions, and is I -absolute; then J^w is a first order (ω_3, a) -mouse operator that relativizes well, determines itself on generic extensions, and is I -absolute.

Proof. We first take a smaller step.

Claim 1. J^\sharp is an (ω_3, a) mouse operator that relativizes well, determines itself on generic extensions, and is I -absolute.

Proof. Let G be V -generic over (I^+, \subseteq) , and

$$\pi: V \rightarrow M = \text{Ult}(V, G)$$

be the generic embedding. We have that M is closed under ω -sequences in $V[G]$, $\pi(\omega_1^V) = \omega_2^V$, $\pi(\omega_3^V) = \omega_3^V = \omega_2^{V[G]}$, and

$$V[G] \models \{\alpha \mid \pi(\alpha) = \alpha\} \text{ is stationary in } \omega_2.$$

By corollary 25, there is in $V[G]$ a unique $(\omega_2^{V[G]}, a)$ -mouse operator J^G extending J . Because J is I -absolute, $V[G] \models \pi(J) \upharpoonright H(\omega_1) = J^G \upharpoonright H(\omega_1)$. It follows from this that

$$\pi(J) \subseteq J^G.$$

For take any $c \in \text{dom}(\pi(J))$; then $c \in \pi(H(\omega_3)) \subseteq H(\omega_2)^{V[G]}$, so $c \in \text{dom}(J^G)$. If $J^G(c) \neq \pi(J)(c)$, then by a simple Skolem hull argument in $V[G]$, using condensation for J^G in $V[G]$ and for $\pi(J)$ in M , we get a countable b such that $J^G(b) \neq \pi(J)(b)$. This is a contradiction.

Now let b be in the $H(\omega_1)$ -cone over a of V . We wish to show that in V , $J(b)^\sharp$ exists and is countably iterable. This is basically a well known result of Kunen, but we sketch a proof for completeness. Setting $\theta = \omega_3^V$, we have in $V[G]$

$$\pi: L_\theta^J(b) \rightarrow L_\theta^J(b),$$

with a stationary set of fixed points. We leave it to the reader to show that ω_1^V and ω_2^V are inaccessible cardinals in $L^J(b)$. Let $\kappa_0 = \omega_1^V$, and let

$$U_0 = \{A \subseteq \kappa_0 \mid A \in L^J(b) \wedge \kappa_0 \in \pi(A)\}.$$

Let $\mu = (\kappa_0^+)^{L^J(b)}$, and

$$\mathcal{M}_0 = (L_\mu^J(b), \in, U_0).$$

\mathcal{M}_0 is an amenable structure by an argument of Kunen. Let \mathcal{M}_α be the α -th iterate of \mathcal{M}_0 by U_0 and its images. We show by induction on $\alpha < \omega_1^{V[G]}$ that \mathcal{M}_α has the form $(L_{\mu_\alpha}^J(b), \in, U_\alpha)$, where μ_α is a cardinal of $L_\theta^J(b)$. At the same time we define maps $i_{\beta,\alpha}: L_\theta^J(b) \rightarrow L_\theta^J(b)$ extending the iteration map from \mathcal{M}_β to \mathcal{M}_α , and “realization maps”

$$\pi_\alpha: L_\theta^J(b) \rightarrow L_\theta^J(b),$$

with $\pi_0 = \pi$, such that for all $\beta < \alpha$

$$\pi_\beta = \pi_\alpha \circ i_{\beta,\alpha}.$$

If α is a limit ordinal, then $i_{\beta,\alpha}$ is the direct limit map, and $\pi_\alpha(i_{\beta,\alpha}(x)) = \pi_\beta(x)$ for all x . Note that π_α embeds the direct limit into $L_\theta^J(b)$, and thus the direct limit does indeed have the form $L_\theta^J(b)$, by the fact that J condenses well. We let

$$i_{\alpha,\alpha+1}: L_\theta^J(b) \rightarrow \text{Ult}(L_\theta^J(b), U_\alpha)$$

be the ultrapower map, and

$$\pi_{\alpha+1}(i_{\alpha,\alpha+1}(f)(\kappa_\alpha)) = \pi_\alpha(f)(\kappa_\alpha),$$

where $\kappa_\alpha = \text{crit}(U_\alpha) = i_{0,\alpha}(\kappa_0)$. This works so long as U_α is the ultrafilter derived from π_α , that is, $\text{crit}(\pi_\alpha) = \kappa_\alpha$, and for $X \subseteq \kappa_\alpha$ in $L_\theta^J(b)$,

$$(*) X \in U_\alpha \Leftrightarrow \kappa_\alpha \in \pi_\alpha(X).$$

Again, the fact that J condenses well then yields that $\text{Ult}(L_\theta^J(b), U_\alpha) = L_\theta^J(b)$, as desired.

We omit the proof that $\text{crit}(\pi_\alpha) = \kappa_\alpha$. To see (*), let $\nu < \mu_0$, $W = U_0 \cap L_\nu^J(b)$, and $f: \kappa_0 \rightarrow P(\kappa_0) \cap L_\nu^J(b)$. Let c, τ be such that $\pi(c) = c$, and for all $\xi < \kappa_0$, $f(\xi) = \tau^{L_\theta^J(b)}[c](\xi) \cap \kappa_0$. Then

$$L_\theta^J(b) \models \forall \xi < \kappa_0 ((\tau[c](\xi) \cap \kappa_0 \in W) \leftrightarrow \kappa_0 \in \tau[c](\xi)).$$

This fact is preserved by $i_{0,\alpha}$, and from that, we easily get (*).

So for b in the $H(\omega_1)$ cone over a of V , $V[G] \models J^\sharp(b)$ exists, and is ω_2 iterable. (The iterations that do not drop are linear, so we can go to $\omega_2^{V[G]}$. This uses 24 for iterations that do drop.) Let $\mathcal{M} = J^\sharp(b)^{V[G]}$, and let h be V -generic over $\text{Col}(\omega, \omega_2)$ and such that $G \in V[h]$. Our proof shows that \mathcal{M} is definable in $V[h]$ from b and J^V : it is the unique putative $J(b)^\sharp$ that is linearly ω_1 -iterable by its last extender in a way that moves the $L^J(b)$ to itself. It follows that $\mathcal{M} \in V$, and it is easy to see that it is countably iterable in V . So for b in the $H(\omega_1)$ cone over a of V , $V \models J^\sharp(b)$ exists.

Now let $b \in H(\omega_2)^V$. Since $b \in H(\omega_1)^M$, we have an \mathcal{N} such that $M \models \mathcal{N} = \pi(J)^\sharp(b)$. Since J is I -absolute, this gives

$$V[G] \models \mathcal{N} = (J^G)^\sharp(b),$$

with \mathcal{N} being ω_2 -iterable in $V[G]$. As in the last paragraph, this gives that \mathcal{N} is definable from J^V and b in $V^{\text{Col}(\omega, \omega_2)}$, so $\mathcal{N} \in V$, and $V \models \mathcal{N} = J^\sharp(b)$.

Finally, J^\sharp can be extended to $H(\omega_3)^V$ by Mouse Reflection at ω_2 .

It is easy to see that J^\sharp relativizes well, determines itself on generic extensions, and is I -absolute. Indeed, the proof that it is I -absolute is part of our proof that that J^\sharp is defined on $H(\omega_2)^V$. \square

Claim 2. J^* is an (ω_3, a) mouse operator that relativizes well, determines itself on generic extensions, and is I -absolute.

Proof. $J^* = (J^\sharp)^\sharp$, so we can just use the proof of claim 1, with J^\sharp replacing J . \square

We are ready to prove that J^w an (ω_3, a) mouse operator that relativizes well, determines itself on generic extensions, and is I -absolute. We show first that it is an (ω_1, a) mouse operator. The proof is parallel to that in the step from J to J^\sharp , but now the core model theory is too involved to be reproduced, so we must just quote it.

Let b be in the $H(\omega_1)$ cone over a . Let $C \subseteq \omega_2$ code $\langle H(\omega_2), I \rangle$. We work in the model

$$N = L_{\omega_2}^{J^\sharp}[C].$$

Because $J^*(C)$ exists, $N \models \text{ZFC}$, and letting Ω be the critical point of the last extender of $J^*(C)$, Ω behaves in N enough like a measurable cardinal that the core model theory of [18], relativised to J , goes through. Let $K^{c,J}(b)$ be the result of the J -relativized K^c -construction over b of length Ω . (See [16].³) It is enough to show that $K^{c,J}(b)$ reaches an active level \mathcal{P} satisfying “there is a Woodin cardinal”. That is because the first such \mathcal{P} is countably iterable in N , and hence countably J -iterable⁴ in V because $H(\omega_2) \subseteq N$. It follows that \mathcal{P} can be re-arranged as a countably iterable putative $M_1^{J, \sharp}(b)$.

³ J is fed into the model being constructed as a model operator, meaning that if the current model in the construction is \mathcal{P} , and we are not adding an extender, then the next model is the core of \mathcal{Q} , where $\mathcal{Q} = J(\mathcal{P})$ unless some proper initial segment of $J(\mathcal{P})$ projects strictly across $o(\mathcal{P})$, in which case \mathcal{Q} is the first such initial segment of $J(\mathcal{P})$. Strictly speaking, the levels of $K^{c,J}(b)$ are not b -premise, but hybrid b -premise relative to the model operator F^J , where $F^J(\mathcal{P})$ is the \mathcal{Q} we just described. See [16].

⁴This means that the iterations move F^J correctly. See [16].

But if the $K^{c,J}(b)$ construction does not reach such a \mathcal{P} , then in N , $K^J(b)$ exists, and has the basic properties of the unrelativized K from [18]. Since I is a presaturated ideal in N , the argument of [18][section 7] leads to a contradiction. This shows that J^w is an (ω_1, a) mouse operator. It is easy to see that it relativizes well and determines itself on generic extensions, using those properties of J .

We now extend J^w to an I -absolute (ω_2, a) operator. Let b be in the $H(\omega_2)$ cone over a . Let

$$\pi: V \rightarrow M = \text{Ult}(V, G)$$

be the generic embedding associated to $G \subseteq I^+$. Let \mathcal{P} be such that

$$M \models \mathcal{P} = M_1^{\pi(J), \#}(b).$$

Claim. \mathcal{P} is in V , and countably iterable in V .

Proof. Let h be V -generic over $\text{Col}(\omega, \omega_2)$, and such that $G \in V[h]$. In $V[h]$, \mathcal{P} is ω_1 -iterable by the following strategy Σ : pick the unique cofinal branch b of \mathcal{T} such that $\mathcal{Q}(b, \mathcal{T}) \sqsubseteq (J^\#)^h(\mathcal{M}(\mathcal{T}))$. To see that Σ works, note that it works in $V[G]$, because M and $V[G]$ have the same reals, and $\pi(J) \subseteq J^G$. But $H(\omega_2)^{V[G][h]} = H(\omega_1)^{V[h]}$, and $(J^\#)^G$ determines $(J^\#)^h$ on $H(\omega_1)^{V[h]}$. A simple Lowenheim Skolem argument for the forcing from $V[G]$ to $V[h]$ shows that if Σ fails in $V[h]$, it fails in $V[G]$. Thus Σ works in $V[h]$. Note that Σ is definable from $J^\#$, which is in V .

If \mathcal{Q}, Λ have in $V[h]$ the properties of \mathcal{P}, Σ just described, then working in $J^*(\langle \mathcal{P}, \mathcal{Q} \rangle)$, where both \mathcal{P} and \mathcal{Q} are $\omega_1 + 1$ -iterable, we can compare them. This shows $\mathcal{Q} = \mathcal{P}$. It follows that \mathcal{P} is definable in $V[h]$ from b and $(J^*)^V$, and thus $\mathcal{P} \in V$ by the homogeneity of $\text{Col}(\omega, \omega_2)$. For a similar reason, $\Sigma \upharpoonright V \in V$, and witnesses that \mathcal{P} is countably iterable (in fact ω_3 -iterable) in V . \square

Thus J^w extends to an (ω_2, a) mouse operator, and our proof showed that it is I -absolute. By Mouse Reflection at ω_2 , J^w extends to an (ω_3, a) mouse operator. The extension relativises well and determines itself on generic extensions, because these properties depend only on $J \upharpoonright H(\omega_1)$. It is I -absolute, because this property depends only on $J \upharpoonright H(\omega_2)$.

This completes the proof of theorem 28. \square

We have at once

Corollary 29. *Suppose there is a presaturated ideal on ω_1 , and $2^{\omega_1} = \omega_2$, and Mouse Reflection holds at ω_2 ; then for all $n < \omega$ and all $b \in H(\omega_3)$, $M_n^\sharp(b)$ exists, and is countably iterable. Thus PD holds, and continues to hold in $V[H]$ whenever H is V -generic for a poset of size $\leq \omega_2$.*

4. AN ASIDE ON $2^{\omega_1} = \omega_2$

We digress briefly in order to prove the following theorem, due to Woodin and implicit in [25] (see Thm. 9.82 for example).

Lemma 30. *Assume $\text{WRP}_{(2)}(\omega_2)$ and NS saturated; then*

$$2^\omega = 2^{\omega_1} = \delta_2^1 = \omega_2.$$

Proof. A theorem of Todorćević (see Thm 6.4 of [6]) gives $2^\omega \leq \omega_2$ under $\text{WRP}(\omega_2)$. The idea is that there is always an injection from 2^ω into any club subset C of $[\omega_2]^\omega$, and under $\text{WRP}(\omega_2)$ there is such a club of size ω_2 , namely

$$C = \bigcup_{\delta < \omega_2} C_\delta$$

where each C_δ be a club of size ω_1 in $[\delta]^\omega$. Now, NS saturated gives closure of $P(\omega_1)$ under the dagger operation and $\text{WRP}_{(2)}(\omega_2)$ lifts this closure to $P(\omega_2)$ as in Lemma 18 below. For $B \subseteq \omega_2$, we write $L[B, U]$ for the minimal model with one measurable cardinal over B described by B^\dagger . We construct a function $B : \omega_2 \rightarrow P(\omega_1)$ as follows. Let $B(0) \subset \omega_1$ be such that $\omega_1^{L[B]}$ is ω_1 . Given $B \upharpoonright \gamma$, let $(X_\xi \mid \xi < \omega_1)$ enumerate those nonstationary sets $X \in L[B \upharpoonright \gamma, U]$ which are stationary in $L[B \upharpoonright \gamma, U]$. Let $B(\gamma)$ code a sequence of clubs disjoint from the X_ξ as well as a surjection from ω_1 to γ . Every subset of ω_2 in the final $L[B, U]$ is in $L[B \upharpoonright \gamma, U]$ for some $\gamma < \omega_2$, so $L[B, U]$ thinks that NS is saturated, and computes ω_2 correctly. Since $L[B, U]$ has a measurable cardinal, it follows from 3.17 of [25] that $\delta_2^1 = \omega_2$ in $L[B, U]$, and thus $2^\omega = \omega_2$ in V , and we are left to show that $2^{\omega_1} \leq \omega_2$.

For this, let G be V -generic over (NS, \subseteq) , and let $j : V \rightarrow M \simeq \text{Ult}(V, G)$ be the generic embedding. We have $\omega_2^V = \omega_1^{V[G]}$ and $\omega_3^V = \omega_2^{V[G]}$ by the ω_2 -c.c. Thus

$$j(\omega_2^V) = \omega_2^M \leq \omega_2^{V[G]} = \omega_3^V.$$

But j is continuous at ω_2 , so

$$j(\omega_2^V) < \omega_3^V.$$

But $M \models 2^\omega = \omega_2$, and M contains all the reals of $V[G]$, so CH holds in $V[G]$. On the other hand, if V thinks that f is an injection of ω_3

into $P(\omega_1)$, then $V[G]$ thinks that f enumerates ω_2 distinct subsets of ω_1^V , contrary to CH. \square

The hypothesis that $2^\omega \leq \omega_2$ is not needed for corollary 29, and probably not for theorem 28, though we have not checked the latter carefully. The following little lemma lets us drop the hypothesis.

Lemma 31. *Suppose I is a presaturated ideal on ω_1 and J is an (ω_3, a) -mouse operator, where $a \in H(\omega_2)$. Suppose that $J^\#$ is an (ω_2, a) -mouse operator; then there is a $B \subset \omega_2$ such that*

- (1) $H(\omega_2)^{L^J(B)}$ is fully elementary in $H(\omega_2)$
- (2) $\bar{I} = I \cap L^J(B) \in L^J(B)$ and
- (3) $L^J(B) \models \bar{I}$ is a presaturated ideal on ω_1 .

Proof. We inductively fold in all of the necessary data into B , which we regard as a function from ω_2 to $P(\omega_1)$. Let $B(0)$ code a , and be such that $\omega_1^{L[B(0)]} = \omega_1$. Let $\delta < \omega_2$, and suppose $B \upharpoonright \delta$ has been defined. We let $B(\delta)$ code W_δ and I_δ , where I_δ in turn codes

$$I \cap J^\#(B \upharpoonright \delta)$$

and $W_\delta \in [P(\omega_1)]^{\omega_1}$ has the following property: if $c \in \mathcal{M}^\#(B \upharpoonright \delta)$ and $\phi(x, y)$ is a formula of the language of set theory so that

$$H(\omega_2)^V \models \phi[c, d]$$

for some d , then there is such a d coded (in some simple way) by an element of W_δ .

Finally, we let

$$\bar{I} = \bigcup \delta < \omega_2 I_\delta.$$

It is easy to check (1)-(3) of the lemma. \square

5. I_α AND COHEN FORCING

In this section, we discuss our resemblance hypothesis I_α , and we prove some lemmas on Cohen forcing over models of AD that are relevant to its formulation.

5.1. The role of I_α . In section 3, we proved PD by a "cycling" argument, first closing $H(\omega_1)$ under a mouse operator, then lifting this closure to $H(\omega_2)$ and finally to $H(\omega_3)$. One idea of [26] is that the main obstacle to continuing this "cycling" argument through the levels of $L(\mathbb{R})$ is the lack of homogeneity of the forcing $P(\omega_1)/NS$. In order to make the $H(\omega_1)$ to $H(\omega_2)$ step, we need at least enough homogeneity that for J the current mouse operator on $H(\omega_1)^V$, $\pi(J) \upharpoonright H(\omega_2)^V$ is independent of the NS-generic embedding π . J will be definable

at some level of $L(\mathbb{R})^V$, and $\pi^G(J)$ at some level of $L(\mathbb{R})^{V[G]}$. This makes it natural to relate the $L(\mathbb{R})$ of the homogeneous extension by $Col(\omega, \omega_1)$ that of an extension by $P(\omega_1)/NS \times Col(\omega, \omega_1)$. This latter model $V^{P(\omega_1)/NS \times Col(\omega, \omega_1)}$ is a Cohen extension of $V^{P(\omega_1)/NS}$, and a *ccc* extension of $V^{Col(\omega, \omega_1)}$ if NS is saturated. The author of [26] believed that the existence of an embedding between the $L(\mathbb{R})$ of these models together with our hypotheses would yield $AD^{L(\mathbb{R})}$. That is, he conjectured the following weakening of our main theorem:

Assume $WRP_{(2)}(\omega_2)$ and NS saturated. Suppose that whenever $G \subset P(\omega_1)/NS$ is V -generic and $g \subset Col(\omega, \omega_1)$ is $V[G]$ -generic there is an embedding

$$\bar{\pi} : L(\mathbb{R})^{V[g]} \rightarrow L(\mathbb{R})^{V[G][g]}$$

which is Σ_1 and fixes ordinals. Then AD holds in $L(\mathbb{R})$.

On the other hand, it was also known that if the induction were to succeed in proving AD in $L(\mathbb{R})$, and that this persists after collapsing ω_2 , then we would have such an embedding. This follows easily from results of Foreman and Magidor ([3]) and Woodin ([2],[25]) that we shall use later in a similar way to prove I_α . We therefore give the argument here. Let us first recall the Foreman-Magidor results.

Definition 32. ([3]) A partial ordering \mathbb{P} is *reasonable* iff for all ordinals α , $[\alpha]^\omega \cap V$ is stationary in $V^\mathbb{P}$.

All proper posets are reasonable, and thus all *ccc* posets are reasonable.

Definition 33. ([2]) A set B of reals is κ -*universally Baire* just in case there are trees T and U on some $\omega \times \gamma$ such that $p[T] = B$, and whenever \mathbb{P} has cardinality $< \kappa$, then $V^\mathbb{P} \models p[T] = \mathbb{R} \setminus p[U]$. We call T and U κ -*absolute complements* in this situation.

Of course, we can speak of κ -universally Baire relations on reals as well. If B is κ universally Baire, and T, U are κ -absolute complements such that $p[T] = B$, and G is V -generic for a poset of size $< \kappa$, then we write B^G for $p[T] \cap V[G]$. The notation is justified because if T^*, U^* is another such complementing pair, then $p[T] = p[T^*]$ holds in $V[G]$. Foreman and Magidor note that if B is an equivalence relation on \mathbb{R} , then B^G is an equivalence relation on $\mathbb{R}^{V[G]}$ ([3][3.3]). It is also easy to see that if B is a prewellorder of \mathbb{R} , then B^G is a prewellorder of $\mathbb{R}^{V[G]}$. Foreman and Magidor show

Theorem 34. ([3][3.4]) *Let B be a κ -universally Baire prewellorder of \mathbb{R} , and let G be V generic over a reasonable poset of size $< \kappa$; then for every $x \in \mathbb{R}^{V[G]}$ there is a $y \in \mathbb{R}^V$ such that $B^G(x, y)$ and $B^G(y, x)$.*

That is, every B^G -equivalence class has a representative in V . In fact, [3][3.4] shows that if B is a thin equivalence relation, then every B^G equivalence class has a representative in B . One can show that the equivalence relation generated by a prewellorder with the Baire property is thin.

Theorem 35. *Assume NS saturated and $WRP_{(2)}(\omega_2)$. Assume $AD^{L(\mathbb{R})}$ holds in $V[g]$ whenever $g \subset Col(\omega, \omega_1)$ is V -generic. Suppose G and g are the factors of generic filter on $P(\omega_1)/NS \times Col(\omega, \omega_1)$. Then there is a fully elementary embedding*

$$\pi : L(\mathbb{R})^{V[g]} \rightarrow L(\mathbb{R})^{V[G][g]},$$

and any such embedding satisfies $\pi \upharpoonright \theta^{L(\mathbb{R})} = id$. Moreover, $AD^{L(\mathbb{R})}$ holds in V and in the universe after collapsing ω_2 .

Proof. By Lemma 30, we have $2^{\omega_1} = \omega_2$. Thus from the perspective of $V[g]$ the algebra $\mathbb{B} = (P(\omega_1)/NS)^V$ has size ω_1 .

Claim 1. In $V[g] \models (P(\omega_1)/NS)^V$ has the countable chain condition.

Proof. Otherwise there is a condition $p \in Col(\omega, \omega_1)$ which forces that some \dot{f} enumerates an antichain of length ω_2^V . On cardinality grounds there must be a condition $q \leq p$ which decides ω_2 of the values of \dot{f} , a contradiction as two of these values must therefore be compatible by saturation of NS in V . \square

Now, since $2^{\omega_1} = \omega_2$ and $P(\omega_2)$ is closed under sharps, we have $\mathbb{R}^\#$ in $V[g]$. By 9.83 of [25] then we have that $\mathbb{R}^\#$ exists and $L(\mathbb{R}) \models AD$ in the universe after collapsing ω_2 . This uses $WRP_{(2)}(\omega_2)$. Now we may assume there is $h \subset Col(\omega, \omega_2)$ which is $V[G][g]$ generic and so that $V[G][g][h] = V[\bar{h}]$ for some $\bar{h} \subset Col(\omega, \omega_2)$. The argument of 5.2 of [2] now shows that there are definable trees S, T in $V[g]$ such that $p[S] = \mathbb{R}^\#$ and $p[T] = \mathbb{R} \setminus \mathbb{R}^\#$ in $V[\bar{h}]$. These trees belong to $V, V[g]$, and $V[G][g]$ by homogeneity and it can be argued that $p[S] = \mathbb{R}^\#$ in the sense of each model. Thus there is a fully elementary embedding⁵

$$\pi : L(\mathbb{R})^{V[g]} \rightarrow L(\mathbb{R})^{V[G][g]}.$$

We also have that from the perspective of $V[g]$, every set of reals in $L(\mathbb{R})$ is ω_1^+ -Universally Baire and hence \mathbb{B} -Universally Baire as $2^{\omega_1} = \omega_2$.

⁵The argument of this paragraph can be recast using mice; essentially we are crossing the weak gap $(\delta_1^2, \theta)^{L(\mathbb{R})}$, and we can do it using the techniques of Lemma 66 below.

Now suppose π is such an embedding and fix α less than the θ of $L(\mathbb{R})^{V[g]}$. There is a prewellordering $\preceq \in L(\mathbb{R})^{V[g]}$ (with associated equivalence relation \simeq) of length α in $V[g]$, together with S, T such that

- (1) $p[S] = \preceq$ in $V[g]$
- (2) $p[S] = \pi(\preceq)$ in $V[G][g]$
- (3) $p[S] = \mathbb{R} \setminus p[T]$ in $V[g][G]$.

Since $\pi(\alpha)$ is the length of $\pi(\preceq)$ we must show that $p[S]^{V[G][g]}$ has length α . This follows from the Foreman-Magidor theorem 34. \square

The preceding remarks suggest that we add the existence of approximations to such an embedding to the induction hypothesis. This was how [26] came to formulate I_α . In the next section we shall use W_α^* and I_α to get $I_{\alpha+\omega}$. If α is not in the range of π_{NS} this will give $W_{\alpha+\omega}^*$ easily. Otherwise the arguments of section 3 will produce the required witnessing structures.

5.2. $L(\mathbb{R})$ and Cohen Forcing. Recall that the ordinal height of the transitive structure $J_\alpha(\mathbb{R})$ is $\omega\alpha$, and that the new sets of reals appearing in $J_{\alpha+1}(\mathbb{R})$ are precisely the new sets which are first order definable over $J_\alpha(\mathbb{R})$, that is

$$P(\mathbb{R}) \cap J_{\alpha+1}(\mathbb{R}) = P(\mathbb{R}) \cap \Sigma_\omega^1(J_\alpha(\mathbb{R})).$$

We say that α begins a gap if there is no $\beta < \alpha$ with $J_\beta(\mathbb{R})$ a Σ_1 elementary (with real parameters) submodel of $J_\alpha(\mathbb{R})$.

Lemma 36. *Suppose $h \subset \text{Col}(\omega, \kappa)$ is V -generic and α begins a gap in $L(\mathbb{R})^{V[h]}$. Suppose $J_\alpha(\mathbb{R})^{V[h]} \models \text{AD}$; then there is a unique pair α_0, ψ such that α_0 that begins a gap in $L(\mathbb{R})$, and*

$$\psi: J_{\alpha_0}(\mathbb{R}) \rightarrow J_\alpha(\mathbb{R})^{V[h]}$$

is Σ_1 elementary.

Proof. There are uniformly Σ_1 definable functions

$$f_\alpha: [\omega\alpha]^{<\omega} \times \mathbb{R} \rightarrow J_\alpha(\mathbb{R})$$

which are surjective. Using these we define a Σ_1 function F as follows. Given a real x , decode a sequence (x_0, \dots, x_n) of reals and a real y and suppose there is a finite sequence t such that

$$J_\alpha(\mathbb{R}) \models \phi_{y(0)}((x_0, \dots, x_n), f_\alpha(t, \hat{y}))$$

where $\hat{y}(n) = y(n+1)$ and $(\phi_k \mid k < \omega)$ enumerates Σ_1 formulae with two free variables. Let t^* be the Brouwer-Kleene least such t and set

$F(x) = f_\alpha(t^*, \hat{y})$. F is a uniformly Σ_1 partial map and if α begins a gap then F as defined over $J_\alpha(\mathbb{R})$ is surjective. Let $M = F[\mathbb{R}^V]$ where F is computed in $J_\alpha(\mathbb{R})$ of $V[g]$. By the homogeneity of $\text{Col}(\omega, \kappa)$, $\mathbb{R} \cap M = \mathbb{R}^V$. Because $(\Sigma_1^{J_\alpha(\mathbb{R})})^{V[h]}$ has the scale property, $M \prec_{\Sigma_1} J_\alpha(\mathbb{R})^{V[h]}$. It follows that $M \simeq J_{\alpha_0}(\mathbb{R})^V$ for some ordinal α_0 , and the inverse of the collapse is the desired map ψ . \square

Lemma 37. *Suppose α is an inadmissible limit ordinal which begins a gap.*

- (1) *There is a surjective function $f : \mathbb{R} \rightarrow J_\alpha(\mathbb{R})$ which is Δ_1 definable over $J_\alpha(\mathbb{R})$ from a real z_0 .*
- (2) *If $A \in P(\mathbb{R}) \cap J_{\alpha+1}(\mathbb{R})$ then A is projective in a set $D \in P(\mathbb{R})$ which is Δ_1 definable over $J_\alpha(\mathbb{R})$ from a real.*
- (3) *If $\underline{\Delta}_{2k+1}(J_\alpha(\mathbb{R}))$ -determinacy holds then the pointclasses*

$$\underline{\Pi}_{2k+2}(J_\alpha(\mathbb{R})) \text{ and } \underline{\Sigma}_{2k+3}^{J_\alpha(\mathbb{R})}$$

have the scale property.

Proof. Inadmissibility of $J_\alpha(\mathbb{R})$ together with a Skolem hull argument gives a $\underline{\Sigma}_1^{J_\alpha(\mathbb{R})}$ map $g : \mathbb{R} \rightarrow \omega\alpha$ which is cofinal. Using the uniform Σ_1 Skolem function this can be turned into the desired map f . For (2) note that every such A can be obtained from a Δ_1 set of the form

$$D = \{(x, x_1, \dots, x_k) \in \mathbb{R}^{k+1} \mid J_\alpha(\mathbb{R}) \models \phi(x, f(x_1), \dots, f(x_k), f(r))\}$$

by taking projections and complements, for some Σ_0 formula ϕ and real r . Part (3) follows from the second periodicity theorem. See [23] for further details. \square

Let $g \times G$ be V -generic over the product $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$. In order to relate $L(\mathbb{R})^{V[g]}$ to $L(\mathbb{R})^{V[g][G]}$, we must at the same time relate $L(\mathbb{R})^{V[G]}$ to $L(\mathbb{R})^{V[G][g]}$. Now $V[G][g]$ comes from $V[G]$ by adding just one Cohen real. The following lemma, due to Woodin and H. Friedman, establishes the level-by-level relationship between the two $L(\mathbb{R})$'s that we need. See [8] for other results in this vein.

Lemma 38. *Suppose $J_\alpha(\mathbb{R}) \models \text{AD}$ and α begins a gap. Suppose $g \subset \text{Col}(\omega, \omega)$ is Cohen generic over V ; then there exists a unique Σ_1 elementary embedding*

$$j : J_\alpha(\mathbb{R}) \rightarrow J_\alpha(\mathbb{R})^{V[g]}$$

such that

$$j \upharpoonright \alpha = \text{identity}.$$

Furthermore, if all $\underline{\Sigma}_1^{J_\alpha(\mathbb{R})}$ sets of reals have the Baire property, then j is Σ_2 -elementary.

Proof. The reader of [8][section 5] will easily adapt it so as to obtain this localization. Nevertheless, we sketch a proof for the sake of completeness.

We think of the reals as ω^ω . For $p \in \omega^{<\omega}$ let N_p denote the neighborhood determined by p . Let \mathcal{B} denote the σ -algebra of sets of reals in $L(\mathbb{R})$ which have the Baire property. Let \mathbb{B} denote the quotient algebra \mathcal{B}/I where I is the ideal of meager sets. Clearly the map

$$\pi : Col(\omega, \omega) \rightarrow \mathbb{B}$$

be defined by $\pi(p) = [N_p]$ is a dense embedding so if $g \subset Col(\omega, \omega)$ is V -generic then g induces an ultrafilter U_g on \mathbb{B} . That is, a set A is in U_g if and only if $A \cap N_p$ is comeager in N_p for some $p \in g$.

In $V[g]$ we form the ultrapower $Ult(J_\alpha(\mathbb{R}), U_g)$ using functions belonging to $J_\alpha(\mathbb{R})$. We may assume that these functions are total with domain \mathbb{R} . We first show that $Ult(J_\alpha(\mathbb{R}), U_g)$ is well-founded. Assume a condition p forces that $([f_n])$ is a decreasing sequence in the ultrapower. For $s \in \omega^{<\omega}$ define a condition $p_s \in Col(\omega, \omega)$, a set A_s , and a function f_s all in $J_\alpha(\mathbb{R})$ such that

- (a) $p \subseteq p_\emptyset$
- (b) $\{p_{s \smallfrown n} \mid n \in \omega\}$ is a maximal antichain below p_s
- (c) $A_s \subset N_s$ is comeager in N_s ,
- (d) $p_s \Vdash_{Col(\omega, \omega)} \dot{f}_{lh(s)} = f_s$.
- (e) $s \subset t, s \neq t$, and $x \in A_t$ implies $f_t(x) \in f_s(x)$.

By the Baire Category Theorem

$$\bigcap_{n < \omega} \bigcup_{lh(s)=n} A_s \neq \emptyset$$

so there are $x, h \in \omega^\omega$ so that $x \in A_{h \smallfrown n}$ for every $n < \omega$ and hence $\{f_{h \smallfrown n}(x) \mid n < \omega\}$ is an \in -decreasing sequence, giving the desired contradiction. Thus the ultrapower has a transitivity M .

Los' theorem for Σ_0 formulae comes from almost-everywhere uniformization:

Claim. Suppose $A \subset \mathbb{R} \times \mathbb{R}$ and $A \in J_\alpha(\mathbb{R})$; then there is a continuous function f and a comeager set $D \subseteq \text{dom}(f)$ such that if $x \in D$ and there is y such that $(x, y) \in A$ then $(x, f(x)) \in A$.

This standard result is proved by unfolding the Banach-Mazur game. It yields

Claim. For any functions $f_1, \dots, f_n \in J_\alpha(\mathbb{R})$ and Σ_0 formula ϕ ,

$$Ult(J_\alpha(\mathbb{R}), U_g) \models \phi([f_1], \dots, [f_n])$$

if and only if

$$\{r \in \mathbb{R} \mid J_\alpha(\mathbb{R}) \models \phi(f_1(r), \dots, f_n(r))\} \in U_g.$$

This follows by the usual induction. For the existential step, we may assume that $\beta < \alpha$ begins a gap and $J_\beta(\mathbb{R}) \models \exists Y \phi(Y, f(x))$, for all x in some set $B \in U_g$. Let $F \in J_\alpha(\mathbb{R})$ map \mathbb{R} onto $J_\beta(\mathbb{R})$. Let A denote the set of (x, y) such that $\phi(F(y), f(x))$ holds in $J_\beta(\mathbb{R})$, let h^* uniformize this A as in the claim, and set $h(x) = F(h^*(x))$. By induction, $\phi([h], [f])$ holds in the ultrapower, as desired.

It follows at once that the ultrapower map $j : J_\alpha(\mathbb{R}) \rightarrow M$ is Σ_1 elementary. We leave it to the reader to check that the standard terms for reals correspond to continuous functions on \mathbb{R} , and thus $\mathbb{R}^M = \mathbb{R}^{V[g]}$. It follows that $M = J_\gamma(\mathbb{R})^{V[g]}$, for some γ .

In $J_\alpha(\mathbb{R})$ any well-ordered union of meager sets is meager. It follows that for any ordinal η and function $f : \mathbb{R} \rightarrow \eta$ in $J_\alpha(\mathbb{R})$ there is a dense set $D \subset \text{Col}(\omega, \omega)$ such that f is constant on a comeager subset of N_p for any $p \in D$. It follows that $\gamma = \alpha$ and that $j \upharpoonright \omega\alpha$ is the identity.

Finally, suppose all $\sum_1^{J_\alpha(\mathbb{R})}$ sets of reals have the Baire property. We show that Los' theorem holds for Σ_1 formulae, so that j is Σ_2 elementary, as desired. So suppose ϕ is Σ_1 , and

$$J_\alpha(\mathbb{R}) \models \exists Y \phi(Y, f(x)),$$

for comeager many x in N_p , where $p \in g$. Say this holds for all $x \in B$, where B is Borel and comeager in N_p . Let us assume α is a limit ordinal for simplicity; otherwise we use Jensen's S -hierarchy. For $\beta < \alpha$, put

$$x \in B_\beta \Leftrightarrow J_\beta(\mathbb{R}) \models \exists Y \phi(Y, f(x)).$$

The prewellordering $x_1 \leq x_2$ iff $\mu\beta(x_1 \in B_\beta) \leq \mu\beta(x_2 \in B_\beta)$ is Σ_1 over $J_\alpha(\mathbb{R})$, so it has the Baire property, and thus by Kuratowski-Ulam some B_β is nonmeager in N_p , and thus comeager on some N_q with $p \subseteq q$. By density, there is such a β, q with $q \in g$. We can now find the desired witness function h for ϕ, f so that $h \in J_{\beta+1}(\mathbb{R})$. \square

The following lemma shows one way we shall use these results.

Lemma 39. *Assume I_α holds, and that α begins a gap in $V^{\text{Col}(\omega, \omega_1)}$. Let $h \times H$ be $\text{Col}(\omega, \omega_1) \times (P(\omega_1)/NS)^V$ -generic over V , and suppose $V[h] \models W_\alpha^*$; then*

- (1) *there is a unique Σ_1 embedding from $J_\alpha(\mathbb{R})^{V[H]}$ to $J_\alpha(\mathbb{R})^{V[H][h]}$ that fixes all ordinals,*
- (2) *there is a unique ordinal α_0 that begins a gap in V , and Σ_1 embedding from $J_{\alpha_0}(\mathbb{R})^V$ to $J_\alpha(\mathbb{R})^{V[h]}$.*

Thus AD holds in $J_{\alpha_0}(\mathbb{R})^V$, in $J_\alpha(\mathbb{R})^{V[h]}$, in $J_\alpha(\mathbb{R})^{V[H]}$, and in $J_\alpha(\mathbb{R})^{V[H][h]}$.

Proof. We have AD in $J_\alpha(\mathbb{R})^{V[h]}$ and $J_\alpha(\mathbb{R})^{V[H][h]}$ by Lemma 10. Moreover, the embedding given by I_α shows that α begins a gap in $V[H][h]$.

$V[H][h]$ is a $\text{Col}(\omega, \omega)$ extension of $V[H]$, so Lemma 36 gives a unique Σ_1 embedding from $J_\gamma(\mathbb{R})^{V[H]}$ to $J_\alpha(\mathbb{R})^{V[H][h]}$, for some γ that begins a gap in $V[H]$. But then AD holds in $J_\gamma(\mathbb{R})^{V[H]}$, so we can apply our lemma 38 to see that $\gamma = \alpha$, and the embedding is the identity on the ordinals.

Finally, α_0 and the embedding of (2) are just what we get when we apply 36 to the $\text{Col}(\omega, \omega_1)$ -extension from V to $V[h]$. \square

5.3. The Baire property for $\sum_1^{J_\alpha(\mathbb{R})}$. We present a sufficient condition for the pointclass $\Gamma = \sum_1^{J_\alpha(\mathbb{R})}$ to have the Baire property. The condition is implicit in Solovay's theorems on the Baire property for Σ_2^1 sets. We assume that α begins a gap, $J_\alpha(\mathbb{R}) \models \text{AD}$, and $\text{cof}(\alpha) > \omega$.⁶ We define

$$C_\Gamma(x) = \mathbb{R} \cap OD_x^{<\alpha},$$

that is, for reals x, y we put $y \in C_\Gamma(x)$ if there is $\beta < \alpha$ such that y is ordinal definable over $J_\beta(\mathbb{R})$ from the parameter x .⁷ Similarly

$$C_\Gamma(a) = P(a) \cap OD_{a \cup \{a\}}^{J_\alpha(\mathbb{R})}$$

for a countable transitive set a . Because we have W_α ,

$$C_\Gamma(a) = \text{Lp}^\Gamma(a)$$

is the result of stacking all mice projecting to a and having ω_1 -iteration strategies belonging to $J_\alpha(\mathbb{R})$. A transitive set M is Γ -closed if $a \in M$ implies $C_\Gamma(a) \in M$. The model $L^\Gamma[x]$ is the minimal transitive model of height ω_1 which contains x and is Γ -closed. Finally, we say that ω_1 is Γ -inaccessible to reals iff $C_\Gamma(x)$ is countable for all reals x , or equivalently,

$$\omega_1^{L^\Gamma[x]} < \omega_1$$

for all reals x .

The classical argument of Solovay in the case $\Gamma = \Sigma_2^1$ yields

Lemma 40. *Assume $J_\alpha(\mathbb{R}) \models \text{AD}$, α begins a gap, $\text{cof}(\alpha) > \omega$ and ω_1 is Γ -inaccessible to reals where $\Gamma = \sum_1^{J_\alpha(\mathbb{R})} \cap P(\mathbb{R})$; then*

- (1) Γ has the Baire Property.
- (2) ω_1 is Γ -inaccessible to reals in $V[g]$ whenever g is Cohen generic over V .

⁶In the case that $\text{cof}(\alpha) = \omega$ or α is a successor, the facts that the class of sets with the Baire property is closed under countable unions is enough for us.

⁷If $J_\alpha(\mathbb{R})$ is admissible, then C_Γ is the largest thin Γ set, but we don't know whether this is true in general.

Proof. Let A be a set in Γ . Let x be a real such that A is Σ_1 -definable over $J_\alpha(\mathbb{R})$ from x . Let N be a rank initial segment of $L^\Gamma[x]$ containing its reals. There are comeager many Cohen generics over N so if each one lands in $\mathbb{R} \setminus A$ we're finished. Assume therefore that there is g_0 which is Cohen generic over N with $g_0 \in A$. Let $\beta < \alpha$ be least such that $J_\beta(\mathbb{R}) \models g_0 \in A$. Let $\beta_1 = \beta + \omega$ and let T be the tree of the scale on the universal $\Sigma_1(J_{\beta_1}(\mathbb{R}))$ set. Note that N is a rank initial segment of $L[T, N]$. Now assume toward a contradiction that there exists a sequence of open dense sets $D_n \subseteq \mathbb{R}$ such that

$$g \in \bigcap_{n \in \omega} D_n \Rightarrow g \in (\neg A)^{J_\beta(\mathbb{R})}.$$

Then for any real z coding N , the model $J_{\beta_1}(\mathbb{R})$ satisfies the sentence asserting the existence of g_0 , $\{D_n \mid n \in \omega\}$ and β satisfying

- (1) g_0 is Cohen generic over the model coded by z
- (2) $g_0 \in A^{J_\beta(\mathbb{R})}$
- (3) $g \in \bigcap_{n \in \omega} D_n \Rightarrow g \in (\neg A)^{J_\beta(\mathbb{R})}$

This is a Σ_1 sentence and so it holds in $L[T, z]$ by absoluteness. Thus for every g which is $L[T, z]$ -generic, $g \in (\neg A)^{J_\beta(\mathbb{R})}$. Now, g_0 is generic over $L[T, N]$. Let G be generic over $L[T, N][g_0]$ for $Col(\omega, N)$. Then g_0 is generic over $L[T, N][G]$. But G codes N which is a contradiction. Thus by the Baire Property there is a condition p such that $g \in A^{J_\beta(\mathbb{R})}$ for comeager many g below p . Let O be the union of all neighborhoods which have this property. We claim that $A \setminus O$ is meager. Otherwise there is an $L^\Gamma[x, y]$ generic g which lands in $A \setminus O$ where y codes the open set O . But $A \setminus O$ is Σ_1 so the preceding analysis produces a q such that comeager many g below q land in $A \setminus O$, a contradiction as $N_q \subseteq O$. For part (2) note that we interpret Γ in $V[g]$ as $(\Sigma_1(J_\alpha(\mathbb{R})) \cap P(\mathbb{R}))^{V[g]}$. Let τ be a standard term for a real. It suffices to show that $C_\Gamma(\tau_g)$ is countable. Assume otherwise. Thus we may assume that there is a single Σ_1 formula $\psi(x, y, z)$ and a condition p which forces that $\{r_\eta \mid \eta < \omega_1\}$ is a distinct sequence of reals where

$$r_\eta = \{n < \omega \mid J_\alpha(\mathbb{R})^{V[g]} \models \psi(\eta, n, \tau_g)\}.$$

Let z be a real coding τ and let $\phi(\eta, q, n)$ be the Σ_1 formula which asserts that $q \leq p$, z codes a term τ and

$$q \Vdash_{Col(\omega, \omega)} \psi(\eta, n, \tau).$$

Setting $r_\eta^* = f[\{(q, n) \mid \phi(\eta, q, n)\}]$ where $f : Col(\omega, \omega) \times \omega \rightarrow \omega$ is a fixed bijection we see that each r_η^* belongs to $C_\Gamma(z)$ and that they are distinct, contradicting our hypotheses. \square

6. THE INDUCTION STEP IN THE INADMISSIBLE CASE

We now handle the induction step in the uncountable cofinality, inadmissible case. This was the case labelled (2)(b) in case breakdown 2. The countable cofinality inadmissible case (2)(a) is similar but simpler, so we omit it. The general successor case (1) involves no ideas beyond those in section 3, so again, we omit it.

We assume for the rest of this section that NS is saturated, and $\text{WRP}_{(2)}(\omega_2)$ holds. We fix for the remainder of the section an α such that I_α holds, and for some (equivalently all) g generic over $\text{Col}(\omega, \omega_1)$:

- (1) $V[g] \models W_\alpha^*$, and
- (2) $V[g] \models \alpha$ has uncountable cofinality, and $J_\alpha(\mathbb{R})$ is inadmissible.

Our goal is to prove that $I_{\alpha+1}$ holds, and that $W_{\alpha+1}^*$ holds in $V[g]$.

6.1. The map $\pi_{NS} \upharpoonright \alpha_0$. If G is any V -generic over $P(\omega_1)/NS$, then we let

$$\pi_G: V \rightarrow \text{Ult}(V, G)$$

be the canonical embedding into the generic ultrapower. One byproduct of the proof of our main theorem is that $\pi_G \upharpoonright \Theta^{L(\mathbb{R})}$ is independent of G . As we go along, we are showing that $\pi_G(\gamma)$ is independent of G for certain γ . In particular, for any $\text{Col}(\omega, \omega_1)$ -generic g , let α_0 be the ordinal beginning a gap in V that is given by 39, and

$$\tau^g: J_{\alpha_0}(\mathbb{R})^V \rightarrow J_\alpha(\mathbb{R})^{V[g]}$$

the unique Σ_1 map. By homogeneity, $\tau^g \upharpoonright (\mathbb{R} \times \omega\alpha_0)$ is independent of g , and so we may omit the superscript g when we are only concerned with this part of τ^g . Of course, $J_{\alpha_0}(\mathbb{R})$ is the Σ_1 hull of $(\mathbb{R} \times \omega\alpha_0)$. We have

Lemma 41. *Let G be $P(\omega_1)/NS$ -generic over V ; then*

- (a) $\pi_G \upharpoonright \omega\alpha_0 = \tau \upharpoonright \omega\alpha_0$,
- (b) $\pi_G(\alpha_0) \geq \alpha$, and
- (c) if $\pi_G(\alpha_0) = \alpha$, then for all $P(\omega_1)/NS$ generic H , $\pi_H(\alpha_0) = \alpha$.

Proof. Let g be such that $g \times G$ is $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$ generic. Let π witness I_α , and let

$$\psi: J_\alpha(\mathbb{R})^{V[G]} \rightarrow J_\alpha(\mathbb{R})^{V[G][g]}$$

be the Cohen ultrapower map. ψ is Σ_1 elementary, and the identity on ordinals. We have then

$$\tau^g = \pi^{-1} \circ \psi \circ (\pi_H \upharpoonright J_{\alpha_0}(\mathbb{R})^V),$$

because both sides yield Σ_1 embeddings of $J_{\alpha_0}(\mathbb{R})^V$ into $J_{\alpha_0}(\mathbb{R})^{V[h]}$. Since ψ and π are the identity on ordinals, we have (a).

For (b), let β be least such that $\pi_G(\beta) \geq \alpha$, and suppose $\alpha_0 < \beta$ toward contradiction. Let γ end the gap in V that begins with α_0 . If $\beta \leq \gamma$, then $[\pi_G(\alpha_0), \pi_G(\beta)]$ is a gap in $V[G]$ by the elementarity of π_G , and α is inside it, so α does not begin a gap in $V[G]$, contradiction. Suppose $\gamma < \beta$. Let x be a real in V and $\varphi(v)$ a Σ_1 formula such that $J_{\gamma+1}(\mathbb{R})^V \models \varphi[x]$, but $J_{\alpha_0}(\mathbb{R}) \not\models \varphi[x]$. Using $\tau^g = \pi^{-1} \circ \psi \circ (\pi_G \upharpoonright J_{\alpha_0}(\mathbb{R})^V)$, we see that $J_{\alpha}(\mathbb{R})^{V[g]} \models \varphi[x]$. But this contradicts the elementarity of τ^g .

For (c), we show first that all $(\sum_1^{J_{\alpha}(\mathbb{R})})^{V[G]}$ sets have the Baire property. Let $\Gamma = (\sum_1^{J_{\alpha}(\mathbb{R})})^{V[G]}$. We wish to show that in $V[G]$, ω_1 is Γ -inaccessible to reals. Letting $\Gamma_0 = (\sum_1^{J_{\alpha_0}(\mathbb{R})})^V$, it is enough to show that in V , ω_1 is Γ_0 -inaccessible to reals. If not, there is in V a real x such that every countable ordinal is the order type of a wellorder in $C_{\Gamma_0}(x)$. Applying π_G to this fact, we have $y \in C_{\Gamma}(x)^{V[G]}$ coding a wellorder of ω of order type ω_1^V . Applying $\pi^{-1} \circ \psi$ to this fact, we see that $y \in V[g]$, and y is ordinal definable in $V[g]$ from x . But then $y \in V$, a contradiction. \square

Having shown $\pi_G(\gamma)$ independent of G , we may use $\pi_{NS}(\gamma)$ to denote the common value of all $\pi_G(\gamma)$. So at the moment, $\pi_{NS} \upharpoonright \alpha_0$ is determined, and it is the Skolem hull map τ .

Let us say that a pointclass has the Baire property just in case all sets of reals belonging to it have the Baire property. Our proof of the last lemma gives

Lemma 42. *Let $g \times G$ be $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$ -generic; then*

- (1) *if $\pi_H(\alpha_0) = \alpha$, then $(\sum_1^{J_{\alpha_0}(\mathbb{R})})^V$ has the Baire property, and*
- (2) *in any case, each of the pointclasses*
 - (a) $(\sum_1^{J_{\alpha}(\mathbb{R})})^{V[G]}$,
 - (b) $(\sum_1^{J_{\alpha}(\mathbb{R})})^{V[G][g]}$, and
 - (c) $(\sum_1^{J_{\alpha}(\mathbb{R})})^{V[g]}$*has the Baire property in its respective model.*

Proof. We showed (1) in proving the last lemma.

(2)(a) follows from (1) in the case that $\pi_G(\alpha_0) = \alpha$ by elementarity. But if $\pi_G(\alpha_0) > \alpha$, then AD in $J_{\alpha_0}(\mathbb{R})^V$ gives us AD, and hence the Baire property, for all sets in $J_{\alpha+1}(\mathbb{R})^{V[G]}$. So this gives (2)(a) in any case.

We get (2)(b) from (2)(a) and part (2) of Lemma 40.

For (2)(c), we show that ω_1 is Γ -inaccessible to reals in $V[g]$, where $\Gamma = (\Sigma_1^{J_\alpha(\mathbb{R})})^{V[g]}$. If not, there is a real $x \in V[g]$ such that every countable ordinal has a counting that is $\text{OD}(x)$ over some $J_\beta(\mathbb{R})^{V[g]}$, where $\beta < \alpha$. Applying π^g , we have that ω_1 is not $(\Sigma_1^{J_\alpha(\mathbb{R})})^{V[G][g]}$ -inaccessible to reals in $V[G][g]$, contrary to (2)(b). \square

6.2. The plan. The following diagram helps illustrate the steps of our proof.

$$\begin{array}{ccc}
 J_{\alpha_0}(\mathbb{R}) & \xrightarrow{\tau^g} & J_\alpha(\mathbb{R})^{V[g]} \\
 \pi_G \downarrow & \searrow & \downarrow \pi \\
 J_\alpha(\mathbb{R})^{V[G]} & \xrightarrow{\psi} & J_\alpha(\mathbb{R})^{V[G][g]} \\
 & & \downarrow j \\
 & & J_{\alpha^*}(\mathbb{R})^{V[G][g][h]}
 \end{array}$$

Here $g \times G$ is an arbitrary $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$ generic, and $h \subset \text{Col}(\omega, \omega_2^V)$ is $V[G][g]$ -generic. Of course, $V[G][g][h]$ could be reorganized as $V[\bar{h}]$ for another $\text{Col}(\omega, \omega_2^V)$ -generic. $J_\alpha(\mathbb{R})^{V[G]}$ and $J_\alpha(\mathbb{R})^{V[G][g]}$ are connected via the Cohen ultrapower map ψ . The map π is given by I_α , and τ^g is the Σ_1 hull map of 36. It may be that α_0 is a discontinuity point of π_G , so in general, π_G is only Σ_1 elementary as a map from $J_{\alpha_0}(\mathbb{R})$ to $J_\alpha(\mathbb{R})^{V[G]}$. If $\pi_G(\alpha_0) = \alpha$, then the map is fully elementary.

Our first burden is to define an appropriate level α^* of the $L(\mathbb{R})$ of $V^{\text{Col}(\omega, \omega_2)}$ and to construct j . We shall show that α^* and $j \upharpoonright \omega_\alpha$ are independent of g, G and h , using the fact that α has a “name” provided by any witness to its inadmissibility in $V[g]$. We will use j and α^* to show that π is in fact Σ_2 -elementary.

The proof then breaks into cases. If $\pi_G(\alpha_0) = \alpha$ for some (equivalently all) G , then α_0 is inadmissible and begins a gap in $L(\mathbb{R})^V$, so we have from [22] a natural mouse operator J on $H(\omega_1)^V$ coding up Σ_1 truth over $J_{\alpha_0}(\mathbb{R})$. We can extend J to $H(\omega_2)$ using π_G and π ; at this point we use j , and the fact that π is Σ_2 -elementary, to see that $\pi_G(J) \upharpoonright H(\omega_2)^V$ is independent of G . By Mouse Reflection, J extends to $H(\omega_3)^V$. We can now use the core model theory of Theorem 28 to get the mice needed for $W_{\alpha+1}^*$ in $V[g]$, and to prove $I_{\alpha+1}$.

If $\pi_G(\alpha_0) > \alpha$, then since $V \models W_{\alpha_0}^*$, $V[G] \models W_{\alpha+1}^*$, so $V[G][g] \models W_{\alpha+1}^*$ by the elementarity of the Cohen ultrapower. So we don't need core model theory to get the new mice, but we do need to get them

in $V[g]$, rather than in $V[G][g]$. This is done with an induction that shows at the same time that π and j are fully elementary.

6.3. Lifting $J_\alpha(\mathbb{R})^{V[G][g]}$ to $V^{\text{Col}(\omega, \omega_2)}$.

Lemma 43. *There is a function $l: \omega_1 \rightarrow (\alpha_0 + 1)$ such that whenever G is $P(\omega_1)/NS$ -generic over V , then $\alpha = [l]_G$.*

Proof. Let $\{S_\xi, l_\xi \mid \xi < \omega_1\}$ be such that $\{S_\xi \mid \xi < \omega_1\}$ is a maximal antichain in $P(\omega_1)/NS$, $l_\xi: \omega_1 \rightarrow (\alpha_0 + 1)$, and S_ξ forces l_ξ to represent α in the generic ultrapower. Define $l(\eta) = l_\xi(\eta)$ where ξ is least such that $\eta \in S_\xi$. It's easy to see that l is as desired. \square

We fix l as in the lemma. By lemma 42, we may assume that for all β , $\sum_1^{J_{l(\beta)}(\mathbb{R})}$ has the Baire property.

Let F be the uniform Σ_1 function for levels of $L(\mathbb{R})$, as described in the proof of 36.

We use “ $\forall^* \sigma$ ” to abbreviate “for a club of $\sigma \in [\omega_2]^\omega$ ”. If $\sigma \in [\omega_2]^\omega$, then $otp(\sigma)$ is the image of σ under transitive collapse, that is, its order-type. For $a \subseteq \sigma^{<\omega}$, a^σ is the image of a under the collapse of σ .

Lemma 44. *There is a term for an ordinal α^* such that $V^{\text{Col}(\omega, \omega_2)} \models \alpha^*$ begins a gap, and for any $p \in \text{Col}(\omega, \omega_2)$, standard term for a real τ , and Σ_1 formula ϕ , the following are equivalent:*

- (1) $p \Vdash_{\text{Col}(\omega, \omega_2)} J_{\alpha^*}(\mathbb{R}) \models \phi(F(\tau))$
- (2) $\forall^* \sigma \ p^\sigma \Vdash_{\text{Col}(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models \phi(F(\tau^\sigma))$

Proof. We define a name for a structure

$$\dot{\mathcal{M}} = (\dot{M}, \dot{\in}_{\mathcal{M}}, \dot{=}_{\mathcal{M}})$$

as follows. For $p \in \text{Col}(\omega, \omega_2)$ and $\tau, \bar{\tau}$ standard terms for reals

- (a) $(p, \tau) \in \dot{M}$ if and only if

$$\forall^* \sigma \ p^\sigma \Vdash_{\text{Col}(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models F(\tau^\sigma) \text{ exists,}$$

- (b) $(p, (\tau, \bar{\tau})) \in \dot{\in}_{\mathcal{M}}$ if and only if

$$\forall^* \sigma \ p^\sigma \Vdash_{\text{Col}(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models F(\tau^\sigma) \in F(\bar{\tau}^\sigma),$$

- (c) $(p, (\tau, \bar{\tau})) \in \dot{=}_{\mathcal{M}}$ if and only if

$$\forall^* \sigma \ p^\sigma \Vdash_{\text{Col}(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models F(\tau^\sigma) = F(\bar{\tau}^\sigma).$$

By $(\tau, \bar{\tau})$ above we really mean the term for the ordered pair. Notice that because $\sum_1^{J_{l(\beta)}(\mathbb{R})}$ has the Baire property, we can understand $p^\sigma \Vdash_{\text{Col}(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models \varphi(\tau)$ to mean that $\varphi(\tau^h)$ is true in $J_{l(\sigma \cap \omega_1)}(\mathbb{R})$, for comeager many generics h such that $p^\sigma \in h$. We prove the following Los-type assertion.

Claim 45. For a condition p , terms τ_1, \dots, τ_k , and a Σ_1 formula ϕ the following are equivalent.

- (1) $p \Vdash_{Col(\omega, \omega_2)} \tau_1, \dots, \tau_k \in \dot{M} \wedge \dot{\mathcal{M}} \models \phi(\tau_1, \dots, \tau_k)$
- (2) $\forall^* \sigma \ p^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{h(\sigma \cap \omega_1)}(\mathbb{R}) \models \phi(F(\tau_1^\sigma), \dots, F(\tau_k^\sigma))$.

Proof. Note that $p \Vdash_{Col(\omega, \omega_2)} \tau \in \dot{M}$ if and only if $(p, \tau) \in \dot{M}$. The atomic cases follow easily. We handle negation as follows. Assume $\phi(\tau_1, \dots, \tau_k)$ is $\neg\psi(\tau_1, \dots, \tau_k)$, ψ is Σ_1 , and the equivalence above holds for $\psi(\tau_1, \dots, \tau_k)$. Assume first that $p \Vdash_{Col(\omega, \omega_2)} \dot{\mathcal{M}} \models \phi(\tau_1, \dots, \tau_k)$. Assume toward a contradiction that there is a stationary set $A_0 \subset [\omega_2]^\omega$ such that for $\sigma \in A$ we have

$$\neg p^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models \phi(F(\tau_1^\sigma), \dots, F(\tau_k^\sigma)).$$

Thus by refining p^σ and pressing down we find a q below p and a stationary subset $A \subset A_0$ such that $\sigma \in A$ implies

$$q^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models \psi(F(\tau_1^\sigma), \dots, F(\tau_k^\sigma)).$$

Similarly we get a stationary set B and a condition r below q such that for $\sigma \in B$ we have

$$r^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models \phi(F(\tau_1^\sigma), \dots, F(\tau_k^\sigma)).$$

We now find an ordinal $\gamma < \omega_2$ above r so that $A \cap [\gamma]^\omega$ and $B \cap [\gamma]^\omega$ are both stationary in $[\gamma]^\omega$. Let $(\sigma_\xi \mid \xi < \omega_1)$ be a continuous, exhaustive chain in $[\gamma]^\omega$ and let $\bar{A} = \{\xi \mid \sigma_\xi \in A\}$ and similarly define \bar{B} . Let $G_A, G_B \subset P(\omega_1)/NS$ be V -generic with $\bar{A} \in G_A$ and $\bar{B} \in G_B$. Let $g \subset Col(\omega, \gamma)$ be generic over both $V[G_A]$ and $V[G_B]$ with $r \in g$. Thus

$$J_\alpha(\mathbb{R})^{V[G_A][g]} \models \psi(F((\tau_1 \upharpoonright \gamma)_g), \dots, F((\tau_k \upharpoonright \gamma)_g))$$

and

$$J_\alpha(\mathbb{R})^{V[G_B][g]} \models \phi(F((\tau_1 \upharpoonright \gamma)_g), \dots, F((\tau_k \upharpoonright \gamma)_g)).$$

By hypothesis there are Σ_1 embeddings

$$\bar{\pi}_A : J_\alpha(\mathbb{R})^{V[g]} \rightarrow J_\alpha(\mathbb{R})^{V[G_A][g]}$$

and

$$\bar{\pi}_B : J_\alpha(\mathbb{R})^{V[g]} \rightarrow J_\alpha(\mathbb{R})^{V[G_B][g]}.$$

Thus $J_\alpha(\mathbb{R})^{V[g]}$ satisfies

$$\psi(F((\tau_1 \upharpoonright \gamma)_g), \dots, F((\tau_k \upharpoonright \gamma)_g)) \wedge \neg\psi(F((\tau_1 \upharpoonright \gamma)_g), \dots, F((\tau_k \upharpoonright \gamma)_g)),$$

which is the desired contradiction. The other direction of the negation case follows similarly.

We now treat the unbounded existential case. For the nontrivial direction suppose for a club of $\sigma \in [\omega_2]^\omega$ that

$$p^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models \exists x \phi(x, F(\tau^\sigma)).$$

For simplicity we assume there is only one parameter. For a real z , the set

$$\{x \mid \phi(F(x), F(z))\},$$

as interpreted in a level of $L(\mathbb{R})$ beginning a gap, is the projection of the tree of the Σ_1 -scale on this set. We let $lw(z)$ denote witness obtained from the leftmost branch of this tree. A key point is that there is a Σ_1 formula ψ so that $\psi(u, z)$ holds if and only if $u = lw(z)$. We define a term $lw(\tau)$ as follows. For a condition q and a pair $n, m \in \omega$ we put the term $(q, (n, m))$ (abusing notation) in $lw(\tau)$ if and only if for a club of σ ,

$$q^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models lw(\tau^\sigma)(n) = m.$$

We need to see that for a club of $\sigma \in [\omega_2]^\omega$

$$p^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models lw(\tau)^\sigma = lw(\tau^\sigma).$$

We will then have

$$p^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models \phi(F((lw(\tau)^\sigma), F(\tau^\sigma)))$$

for each such σ as desired. Assume otherwise. We extract a condition q below p , integers n, m_1, m_2 , an ordinal $\gamma < \omega_2$ above q and stationary sets $A, B \subset [\gamma]^\omega$ such that for $\sigma \in A$ we have

$$q^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models lw(\tau^\sigma)(n) = m_1$$

and for $\sigma \in B$ we have

$$q^\sigma \Vdash_{Col(\omega, otp(\sigma))} J_{l(\sigma \cap \omega_1)}(\mathbb{R}) \models lw(\tau^\sigma)(n) = m_2.$$

As in the negation case we get generics $G_A, G_B \subset P(\omega_1)/NS$ and $g \subset Col(\omega, \gamma)$ such that

$$J_\alpha(\mathbb{R})^{V[G_A][g]} \models lw((\tau \upharpoonright \gamma)_g)(n) = m_1$$

and

$$J_\alpha(\mathbb{R})^{V[G_B][g]} \models lw((\tau \upharpoonright \gamma)_g)(n) = m_2.$$

Using $\bar{\pi}$ we get a contradiction. This completes the proof of the claim. \square

Now let $h \subset \text{Col}(\omega, \omega_2)$ be V -generic. In $V[h]$ we form a structure

$$(M, E) = (\dot{M}_h / \dot{=}^h, \dot{e}_h / \dot{=}^h).$$

An easy argument which uses the fact that the club filter on $P([\omega_2]^\omega)$ is countably complete shows that M is wellfounded, and so there is an isomorphism

$$i : (M, E) \rightarrow (N, \in),$$

for some transitive set N . The construction of the function F we have used is such that for any real r there is a real \hat{r} obtained from r so that $F(\hat{r}) = r$ and r, \hat{r} are Turing equivalent. Thus for any standard term for a real τ , there is a term $\hat{\tau}$ so that

$$\hat{\tau}_h = \hat{\tau}.$$

It follows that $\mathbb{R}^N = \mathbb{R}^{V[h]}$. By claim 45, N satisfies the sentence asserting that it is a level of $L(\mathbb{R})$ and thus $N = J_{\alpha^*}(\mathbb{R})$ in $V[h]$, for some ordinal α^* (which would seem to depend on h). We finish the proof of lemma 44 by showing that the following are equivalent for a Σ_1 formula ϕ , a condition $p \in \text{Col}(\omega, \omega_2)$ and a standard term for a real τ .

- (1) $p \Vdash \dot{M} \models \phi(\tau)$
- (2) $p \Vdash J_{\alpha^*}(\mathbb{R}) \models \phi(F(\tau))$

Note that $\dot{M} \models F(\hat{\tau}) = \tau$ so $J_{\alpha^*}(\mathbb{R}) \models F(i(\hat{\tau}) = i(\tau))$. Since $i(\hat{\tau}) = \tau$ we see that $i = F$, which gives the equivalence above. The equivalence shows that $J_{\alpha^*}(\mathbb{R})^{V[h]}$ is Σ_1 -generated by its reals, so that α^* begins a gap in $V[h]$. \square

If h is $\text{Col}(\omega, \omega_2)$ -generic over V , then we write $\alpha^*(h)$ for the ordinal given by lemma 44. We shall eventually show $\alpha^*(h)$ is independent of h .

Lemma 46. *Let g be $\text{Col}(\omega, \omega_1)$ -generic over V , and suppose that $g \in V[h]$, where h is $\text{Col}(\omega, \omega_2)$ -generic over V ; then there is a Σ_2 -elementary embedding*

$$\pi_2 : J_\alpha(\mathbb{R})^{V[g]} \rightarrow J_{\alpha^*(h)}(\mathbb{R})^{V[h]}.$$

Proof. We consider first the case that $h = g \times k$, where k is $\text{Col}(\omega, \omega_2^V)$ generic over $V[g]$. The proof of 44 generalizes trivially from $\text{Col}(\omega, \omega_2)$ to $\text{Col}(\omega, \omega_1) \times \text{Col}(\omega, \omega_2)$, and shows that the ordinal

$$\alpha^* = \alpha^*(h) = \alpha^*(g \times k)$$

that begins a gap in $V[g][k]$ and satisfies the natural variant of the Los property recorded in the statement of 44. We show that $J_\alpha(\mathbb{R})^{V[g]}$ Σ_2 -embeds into $J_{\alpha^*}(\mathbb{R})^{V[g][k]}$.

Since $J_\alpha(\mathbb{R})^{V[g]}$ is pointwise Σ_1 -definable from its reals, it is enough to show that whenever x is a real in $V[g]$ and $\varphi(v)$ is a Σ_2 formula, then

$$J_\alpha(\mathbb{R})^{V[g]} \models \varphi[x] \Leftrightarrow J_{\alpha^*}(\mathbb{R})^{V[g][k]} \models \varphi[x].$$

We show this first for $\varphi(v)$ a Σ_1 formula. Suppose $J_{\alpha^*}(\mathbb{R})^{V[g][k]} \models \varphi[x]$. Let $x = \rho^g$, where ρ is a standard $\text{Col}(\omega, \omega_1)$ term for a real. Let

$$(p, q) \Vdash J_{\alpha^*(g \times k)}(\mathbb{R}) \models \varphi(\rho^g),$$

with $(p, q) \in g \times k$. Applying 44, we have a club C of $\sigma \in [\omega_1 \times \omega_2]^\omega$ such that, letting $\sigma_0 = \{\xi \mid \exists \nu \langle \xi, \nu \rangle \in \sigma\}$ and $\sigma_1 = \{\nu \mid \exists \xi \langle \xi, \nu \rangle \in \sigma\}$, we have

$$(p, q^\sigma) \Vdash J_{I(\sigma_0 \cap \omega_1)}(\mathbb{R}) \models \varphi[(\rho^{\sigma_0})^g],$$

where the forcing is $\text{Col}(\omega, \sigma_0 \cap \omega_1) \times \text{Col}(\omega, \text{otp}(\sigma_1))$. Thus the forcing is essentially Cohen forcing. Note that σ_0 is transitive, $p^{\sigma_0} = p$, and $\rho^{\sigma_0} = \rho \cap \sigma_0$, for club many σ . We have by $\text{WRP}_{(2)}(\omega_2)$ a $\gamma < \omega_2$ such that C is club in $[\omega_1 \times \gamma]^\omega$, and $q \in \gamma^{<\omega}$. Now let G be such that $g \times G$ is $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$ -generic over V . (It doesn't matter whether $G \in V[h]$.) Los theorem for the generic ultrapower by G gives

$$V[G] \models (p, q) \Vdash J_\alpha(\mathbb{R}) \models \varphi[\rho^g],$$

which gives

$$V[G][g] \models q \Vdash J_\alpha(\mathbb{R}) \models \varphi[x],$$

where the forcing is $\text{Col}(\omega, \gamma)$, and hence essentially Cohen forcing. Since $\Sigma_1^{J_\alpha(\mathbb{R})}$ has the Baire property in $V[G][g]$, we get that

$$J_\alpha(\mathbb{R})^{V[G][g]} \models \varphi[x],$$

and hence

$$J_\alpha(\mathbb{R})^{V[g]} \models \varphi[x],$$

as desired.

The proof just given works equally well for Π_1 formulae φ , and so it gives the desired equivalence for Σ_1 formulae. That in turn gives us a unique Σ_1 elementary embedding $\pi_2: J_\alpha(\mathbb{R})^{V[g]} \rightarrow J_{\alpha^*}(\mathbb{R})^{V[g][k]}$. We want to show π_2 is Σ_2 -elementary. For that, it suffices to show that Σ_2 formulae about a real x go down, and since α^* begins a gap in $V[g][k]$, we may assume the outer quantifier is witnessed by a real. Let $x = \rho^g = \mu^{g \times k}$. We have

$$J_{\alpha^*}(\mathbb{R})^{V[g][k]} \models \varphi[\rho^g, \nu^{g \times k}],$$

where ν is a canonical term for a real, and $\varphi(\nu)$ is Π_1 . Let (p, q) force this. As before, we get a $\gamma < \omega_2$ such that for club many $\sigma \in [\omega_1 \times \gamma]^\omega$,

$$(p, q^\sigma) \Vdash J_{I(\sigma_0 \cap \omega_1)}(\mathbb{R}) \models \varphi[(\rho^{\sigma_0})^{\dot{g}}, (\nu^\sigma)^{\dot{g} \times k}].$$

Again, let G be such that $g \times G$ is $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$ generic over V . We have

$$V[G] \models (p, q) \Vdash J_\alpha(\mathbb{R}) \models \varphi[\rho^{\dot{g}}, \nu^{\dot{g} \times k}],$$

which gives

$$V[G][g] \models q \Vdash J_\alpha(\mathbb{R}) \models \varphi[x, \hat{\nu}],$$

where $\hat{\nu}$ is a $\text{Col}(\omega, \gamma)$ term obtained from ν and g . Let k_0 be $\text{Col}(\omega, \gamma)$ -generic over $V[G][g]$, and let $z = \nu^{g \times k_0}$. We have

$$V[G][g][k_0] \models J_\alpha(\mathbb{R}) \models \varphi[x, z].$$

But then

$$V[g][k_0] \models J_\alpha(\mathbb{R}) \models \varphi[x, z]$$

because $g \times k_0$ is equivalent to a $\text{Col}(\omega, \omega_1)$ generic over V , and we have I_α . Since $\sum_1^{J_\alpha(\mathbb{R})}$ has the Baire property in $V[g]$, 38 gives

$$V[g] \models J_\alpha(\mathbb{R}) \models \exists z \varphi[x, z],$$

as desired.

Thus we have a unique Σ_2 elementary

$$\pi_2: J_\alpha(\mathbb{R})^{V[g]} \rightarrow J_{\alpha^*(g \times k)}(\mathbb{R})^{V[g][k]}.$$

Let φ, x be a witness to the inadmissibility of α in $V[g]$; that is, $\varphi(u, v)$ is Σ_1 , $x \in \mathbb{R} \cap V[g]$, and α is the least ordinal β such that $J_\beta(\mathbb{R})^{V[g]} \models \forall u \in \mathbb{R} \varphi[x]$. Since π_2 is Σ_2 elementary, φ, x witness the inadmissibility of $\alpha^*(g \times k)$ in $V[g][k]$, so that φ, x gives us a “name” for $\alpha^*(g \times k)$.

We now show $\alpha^*(h)$ is independent of h . It is enough to show $\alpha^*(h_1) = \alpha^*(h_2)$ whenever h_1 and h_2 are $\text{Col}(\omega, \omega_2)$ -generic, and $V[h_1] = V[h_2]$. We may re-arrange h_i so that $h_i = g_i \times k_i$, where $g_i \times k_i$ is $\text{Col}(\omega, \omega_1) \times \text{Col}(\omega, \omega_2)$ -generic. Let φ_i, x_i be an inadmissibility witness for α in $V[g_i]$. Since $\sum_1^{J_\alpha(\mathbb{R})}$ has the Baire property in $V[g_i]$, we have that in $V[g_1][g_2]$, both φ_1, x_1 and φ_2, x_2 are inadmissibility witnesses for α . Let $g_3 = g_1 \times g_2$, and let k_3 on $\text{Col}(\omega, \omega_2)$ be such that $V[g_3][k_3] = V[h_1] = V[h_2]$. By what we just showed, both φ_1, x_1 and φ_2, x_2 are inadmissibility witnesses for $\alpha^*(g_3 \times k_3)$ in $V[h_1] = V[h_2]$. It follows that $\alpha^*(h_1) = \alpha^*(g_3 \times k_3) = \alpha^*(h_2)$, as desired.

This finishes the proof of lemma 46. \square

Henceforth we write α^* for the common value of all $\alpha^*(h)$.

Lemma 47. *Let $g \times G$ be $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$ generic over V , and suppose $g, G \in V[h]$, where h is $\text{Col}(\omega, \omega_2)$ generic over V ; then there is a unique Σ_1 elementary*

$$j: J_\alpha(\mathbb{R})^{V[G][g]} \rightarrow J_{\alpha^*}(\mathbb{R})^{V[h]}.$$

Moreover, $\pi_2 = j \circ \pi$.

Proof. In $V[h]$, $J_{\alpha^*}(\mathbb{R}) \models \text{AD}$, and α^* begins a gap. Moreover, $V[h] = V[k]$ where k is $\text{Col}(\omega, \omega_2^V)$ generic over $V[G][g]$. Thus by lemma 36, there is a unique γ beginning a gap in $V[G][g]$ and a unique Σ_1 -elementary

$$j: J_\gamma(\mathbb{R})^{V[G][g]} \rightarrow J_{\alpha^*}(\mathbb{R})^{V[h]}.$$

It will be enough to show $\gamma = \alpha$. (We then get $\pi_2 = j \circ \pi$ because the maps are determined by their restrictions to reals.)

To see $\gamma = \alpha$, in $V[g]$ let B be the prewellordering associated to a $\Sigma_1^{J_\alpha(\mathbb{R})}$ norm on a universal $\Sigma_1^{J_\alpha(\mathbb{R})}$ set of reals. B has length α . In $V[h]$, let C be the set with the same definition, but over $J_{\alpha^*}(\mathbb{R})$. Since $\Sigma_1^{J_{\alpha^*}(\mathbb{R})}$ has the scale property in $V[h]$, there are definable-in- $V[h]$ trees T, U projecting to C and its complement. By homogeneity, T and U are in V . Because π_2 exists, T and U project to B and its complement in $V[h]$. It follows that

$$V[g] \models B \text{ is } \omega_2\text{-universally Baire.}$$

But G is generic over $V[h]$ for a reasonable forcing, so by the Foreman-Magidor theorem 34, every level of C with a representative in $V[G][g]$ has a representative in $V[g]$. The order type of $C \cap V[G][g]$ is just γ , and the order type of $B = C \cap V[g]$ is α , so $\gamma = \alpha$. \square

We now consider the case that α is in the range of a generic embedding.

Lemma 48. *Suppose $\pi_G(\alpha_0) = \alpha$ for some $P(\omega_1)/NS$ generic; then*

- (1) $J_{\alpha_0}(\mathbb{R})$ is inadmissible in V , and
- (2) $\pi_G(\alpha_0) = \alpha$ for all $P(\omega_1)/NS$ generic G .

Proof. Let g be such that $g \times G$ is $\text{Col}(\omega_1, \omega) \times P(\omega_1)/NS$ generic. Let h on $\text{Col}(\omega, \omega_2)$ be such that $g, G \in V[h]$. In $V[g]$, let x be a real and φ a Σ_1 formula such that

$$J_\alpha(\mathbb{R}) \models \forall y \in \mathbb{R} \varphi[y, x],$$

and α is least such that this is true. So φ, x witness the inadmissibility of α in $V[g]$. Because π_2 is Σ_2 elementary, φ and x witness the inadmissibility of α^* in $V[h]$. But then our Σ_1 map j from 47 shows that φ, x define a total function over $J_\alpha(\mathbb{R})$ in $V[G][g]$, and the map π given

by I_α shows this function has range cofinal in α . Thus φ, x witness the inadmissibility of α in $V[G][g]$. The Cohen map is Σ_2 , so φ, x witness the inadmissibility of α in $V[G]$. Finally, π_G is fully elementary, and this gives us part (1).

For (2), let φ, x witness the inadmissibility of α_0 in V . Let $g \times G$ be any $\text{Col}(\omega_1, \omega) \times P(\omega_1)/NS$ generic, with $\alpha = \pi_G(\alpha_0)$. Then φ, x witnesses inadmissibility of α in $V[G]$, hence $V[G][g]$, and hence $V[g]$. If $k \times H$ is any other generic, and $\xi = \pi_H(\alpha_0)$, then φ, x witness inadmissibility for ξ in $V[k]$. Since $x \in V$ and $\text{Col}(\omega, \omega_1)$ is homogeneous, $\xi = \alpha$, as desired. \square

Lemma 49. *Suppose $\alpha = \pi_G(\alpha_0)$ for some G ; then $I_{\alpha+1}$ holds, and $W_{\alpha+1}^*$ holds in all $V[g]$, for g on $\text{Col}(\omega, \omega_1)$.*

Proof. Let φ, x witness the inadmissibility of α_0 in V . Let $g \times G$ be $\text{Col}(\omega_1, \omega) \times P(\omega_1)/NS$ generic. Let h on $\text{Col}(\omega, \omega_2)$ be such that $g, G \in V[h]$. We have that φ, x witnesses inadmissibility for α in $V[g]$, and for α^* in $V[h]$. Working in $V[h]$, we now obtain a function J on $H(\omega_1)$ with parameter x such that J codes up $\Sigma_1^{J_{\alpha^*}(\mathbb{R})}$ truth. Here we just follow the construction of [22]. Given a countable transitive set b such that $x \in b$, $J(b)$ is the minimal b -premouse with parameter x , call it \mathcal{M} , such that

$$J_{\alpha^*}(\mathbb{R}) \models \mathcal{M} \text{ is } \omega_1\text{-iterable,}$$

and

$$\mathcal{M} \models \psi.$$

Here ψ asserts roughly that the Σ_1 function defined by φ, x is defined at all reals of the form τ_g , where g is $\text{Col}(\omega, b)$ generic over \mathcal{M} and τ is a term that is rudimentary in b .

More precisely, if $l \subset \text{Col}(\omega, b)$ is \mathcal{M} -generic, then there is a real $z(l, b)$ which simply codes the pair (l, b) . Thus $\mathcal{M}[l]$ is a $z(l, b)$ -mouse. Let $\sigma \in \mathcal{M}$ be term so that whenever l is such a generic

$$\{(\sigma_l)_i \mid i > 0\} = \{\rho_l \mid \rho \in L_1(b)\}$$

and $(\sigma_l)_0 = x$. We let the sentence ψ assert that whenever l is generic there is a γ so that $\mathcal{M}(z(l, b)) \upharpoonright \gamma$ is a (ϕ_n^*, σ_l) -witness, in the sense referred to in W^* , where $\phi_n^*(v)$ is the Σ_1 formula: there is an ordinal ξ for which $\omega\xi + n$ exists and

$$J_\xi(\mathbb{R}) \models \forall i > 0 \phi(v_i, v_0).$$

Since in $V[h]$, φ, x witnesses inadmissibility for $J_{\alpha^*}(\mathbb{R})$, and $J_{\alpha^*}(\mathbb{R})$ thinks that every Σ_1 truth about a real is captured by an ω_1 -iterable

premouse, we have that $J(b)$ exists for all b such that $x \in b$. J is not quite a mouse operator in $V[h]$, because that requires $\omega_1 + 1$ -iterability. However, ω_1 -iterability in a model of **AD** is enough for comparison, so $J(b)$ is Σ_1 -definable over $J_{\alpha^*}(\mathbb{R})$ from Letting $J_0 = J \upharpoonright V$, we then have in V that J_0 is a first order mouse operator on $H(\omega_3)$, and $J = J_0^h$. It is not hard to see that J_0 relativises well, condenses well, and determines itself on generic extensions. (See [22].)

Our lemmas 47 on the existence of j and the Cohen ultrapower map ψ imply that J is NS-absolute. For let $k \times H$ be $\text{Col}(\omega, \omega_1) \times P(\omega_1)/NS$ generic over V , and both in $V[l]$, where l is $\text{Col}(\omega, \omega_2)$ generic. Let θ be the Σ_1 formula defining $J_0 \upharpoonright H(\omega_1)^V$ from x over $J_{\alpha_0}(\mathbb{R})$. Then θ defines J^l over $J_{\alpha^*}(\mathbb{R})^{V[l]}$, because $j \circ \psi \circ \pi_H$ exists. Thus θ defines $J^l \upharpoonright V[H] = J^H$ over $J_{\alpha}(\mathbb{R})^{V[H]}$, because $j \circ \psi$ exists. So $\pi_H(J_0) \subseteq J^H$.

So in V , J_0 is an (ω_3, x) mouse operator that relativises and condenses well, determines itself on generic extensions, and is NS-absolute. By theorem 28, setting

$$J_{n+1} = (J_n)^w,$$

we have the same properties for J_n , for all $n < \omega$.

This gives $W_{\alpha_0+1}^*$ in V , with the witnessing mice being the $J_n(b)$. Going back to g, G , and h , we also have that $W_{\alpha+1}^*$ holds in $V[g]$, with the witnessing mice being those of the form $J_n^g(b)$ for $b \in H(\omega_1)^{V[g]}$. For details on why these mice suffice as witnesses, see [22]; the key is that Σ_n truth over $J_{\alpha}(\mathbb{R})$ reduces to (Σ_n^1) -in- J_0 truth by inadmissibility.

We also get $I_{\alpha+1}$. For let

$$\pi: J_{\alpha}(\mathbb{R})^{V[g]} \rightarrow J_{\alpha}(\mathbb{R})^{V[G][g]}$$

be the Σ_1 map given by I_{α} . It suffices to show π is fully elementary. This is because the J_n 's code truth over $J_{\alpha}(\mathbb{R})$. More precisely, fix $n < \omega$. There is a recursive function t such that for any Σ_n formula φ and any b in the $H(\omega_1)$ -cone over x of V , we have

$$J_{\alpha_0}(\mathbb{R})^V \models \varphi[b, x] \Leftrightarrow J_n(b) \models t(\varphi)[b].$$

But then for any $b \in H(\omega_1)^{V[G]}$,

$$J_{\alpha}(\mathbb{R})^{V[G]} \models \varphi[b, x] \Leftrightarrow \pi_G(J_n)(b) \models t(\varphi)[b].$$

Since $J_{\alpha+1}(\mathbb{R})^{V[G]} \models \text{AD}$, the Cohen ultrapower map ψ is fully elementary, and $\pi_G(J_n)^g = J_n^{g \times G}$, for any $b \in H(\omega_1)^{V[G][g]}$,

$$J_{\alpha}(\mathbb{R})^{V[G][g]} \models \varphi[b, x] \Leftrightarrow J_n^{g \times G}(b) \models t(\varphi)[b].$$

Finally, for $b \in H(\omega_1)^{V[g]}$,

$$J_{\alpha}(\mathbb{R})^{V[g]} \models \varphi[b, x] \Leftrightarrow J_n^g \models t(\varphi)[b].$$

Since $J_n^g \subseteq J_n^{g \times G}$, this gives us that the inclusion map on the reals is fully elementary from $J_\alpha(\mathbb{R})^{V[g]}$ to $J_\alpha(\mathbb{R})^{V[G][g]}$. That implies π is fully elementary, and hence $I_{\alpha+1}$. \square

We will be done with the inadmissible, uncountable cofinality case when we show:

Lemma 50. *Suppose $\alpha < \pi_G(\alpha_0)$ for some G ; then $I_{\alpha+1}$ holds, and $W_{\alpha+1}^*$ holds in all $V[g]$, for g on $\text{Col}(\omega, \omega_1)$.*

Proof. Let $g \times G$ be $\text{Col}(\omega_1, \omega) \times P(\omega_1)/NS$ generic. Let h on $\text{Col}(\omega, \omega_2)$ be such that $g, G \in V[h]$. Let π be given by I_α, π_2 by lemma 46, ψ by the Cohen ultrapower, and j by lemma 47. We have that π and j are Σ_1 elementary, while ψ and π_2 are Σ_2 elementary. Let φ, x witness inadmissibility for α in $V[g]$, $V[G]$, and $V[G][g]$, and for α^* in $V[h]$.

Just as before, working in $V[h]$, we obtain a function J_0 on $H(\omega_1)$ with parameter x such that J_0 codes up $\Sigma_1^{J_{\alpha^*}(\mathbb{R})}$ truth. $J_0(b)$ is the least b -premouse with parameter x satisfying a certain sentence φ , and being ω_1 -iterable in $J_{\alpha^*}(\mathbb{R})$. The difference now is just that the parameter x may only be in $V[g]$. Since $V[h]$ is a homogeneous extension of $V[g]$, we do have in $V[g]$ and $V[G][g]$ (ω_2, x) mouse operators K_0, K_1 such that $J_0 = K_i^h$. K_i is just the restriction of J_0 to the $H(\omega_2)$ of its model, so we write J_0 for K_i .

Claim. In $V[h]$, there are functions J_n on the $H(\omega_1)$ cone over x such that for any b in this cone

- (a) $J_{n+1}(b) \models$ “I am $J_n^w(b)$ ”, and
- (b) $J_{n+1}(b)$ is ω_1 iterable in $V[h]$ by a J_n -guided iteration strategy.

Remark. So J_{n+1} is J_n^w , except that $\omega_1 + 1$ strategies have to be replaced in $V[h]$ by absolutely definable ω_1 strategies.

Proof.

We begin with J_1 , though our method works in general.

Subclaim 1. In $V[g]$, for any b in the $H(\omega_1)$ cone over x , $J_0^w(b)$ exists, and has a J_0 -guided ω_2 -iteration strategy.

Proof. Fix b . We have b in $V[G][g]$. Because j exists, $J_0 \upharpoonright V[G][g]$ has the same definition over $J_\alpha(\mathbb{R})$ in $V[G][g]$ that J_0 has over $J_{\alpha^*}(\mathbb{R})$ in $V[h]$. But $\pi_G(\alpha_0) > \alpha$, and hence

$$V[G] \models W_{\alpha+1}^*,$$

so that

$$V[G][g] \models W_{\alpha+1}^*,$$

by the elementarity of the Cohen ultrapower. Hence in $V[G][g]$ we can find \mathcal{M} such that

- (a) $\mathcal{M} \models$ “ I am $J_0^w(b)$ ”, and
- (b) $V[G][g] \models$ “ \mathcal{M} has a J_0 -guided ω_1 -iteration strategy”.

Using condensation for J_0 in $V[G][g]$ and the fact that it determines itself on generic extensions, we see by a standard Lowenheim-Skolem argument that \mathcal{M} has a J_0 -guided ω_1 -iteration strategy in $V[h]$. This fact defines \mathcal{M} in $V[h]$ from $J_0 \upharpoonright V[g]$ and b , so $\mathcal{M} \in V[g]$. It is easy to see that $V[g] \models \mathcal{M} = J_0^w(b)$, and that \mathcal{M} J_0 -guided ω_2 -iteration strategy in $V[g]$. \square

Subclaim 2. In $V[h]$, for any b in the $H(\omega_1)$ cone over x , there is a b premouse \mathcal{M} with parameter x such that

- (a) $\mathcal{M} \models$ “ I am $J_0^w(b)$ ”, and
- (b) $V[h] \models$ “ \mathcal{M} has a J_0 -guided ω_1 -iteration strategy”.

Proof. We go back to V , so that we can use Mouse Reflection.

Let τ be a standard term such that $\tau_g = x$. Thus $\tau \in V$, and $\tau \subset \omega \times \omega_1$. Working in V , we can find for any b in the $H(\omega_2)$ cone over τ a countably iterable b -premouse \mathcal{M} with parameter τ such that $J_0^w(b)$ is the canonical re-arrangement of $\mathcal{M}[g]$ as a premouse over $\langle b, g \rangle$ with parameter x . (The hierarchy of \mathcal{M} is defined by induction. See [24] or [14] for the details of this method of inverting generic extensions of mice.) Writing $\mathcal{M} = I(b)$, we can summarize this as

$$I(b)[g] = J_0^w(b),$$

for all b in the $H(\omega_2)^V$ -cone over τ . I is a first order mouse operator with parameter $\langle \tau, p \rangle$ in V , its sentence being “it is forced in $\text{Col}(\omega, \omega_1)$ by p that my canonical re-arrangement as a premouse over $\langle b, g \rangle$ with parameter τ_g thinks it is $J_0^w(b)$.” (Here p is an appropriately chosen fixed condition.)

By Mouse Reflection, I extends in V to an $(\omega_3, \langle p, \tau \rangle)$ mouse operator, which we also call I . Given now b in the $H(\omega_1)^{V[h]}$ cone over x , we can find an \mathcal{M} as in subclaim 2 as follows. Let $V[h] = V[g][k]$, where k is $\text{Col}(\omega, \omega_2)$ generic over $V[g]$, and let $b = \sigma_g$. Let c be in the domain of I , with $\tau, \sigma \in c$. Then $I(c)[g][k]$ can be re-arranged as a premouse over $\langle c, g, k \rangle$ with parameter $x = \tau_g$. Let \mathcal{N} be this re-arrangement. It is easy to see

- (a) $\mathcal{N} \models$ “ I am $J_0^w(\langle c, g, k \rangle)$ ”, and
- (b) $V[g][k] \models$ “ \mathcal{N} has a J_0 -guided ω_1 -iteration strategy”.

The reason is simply that we can reflect any failure of (a) or (b) into the $H(\omega_1)$ cone over x of $V[g]$, where I does indeed determine J_0^w by the method whereby we obtained \mathcal{N} .

Since J_0^w relativises well, we can use \mathcal{N} to find a \mathcal{M} over b satisfying subclaim 2. \square

This gives us the function J_1 as required by our claim. Notice also that the proof has shown that the $H(\omega_1)$ cones over x of $V[g]$ and $V[G][g]$ are closed under J_1 , and that

$$J_1 = J_0^w$$

holds in $V[g]$ and $V[G][g]$ as well. This is what we need to obtain J_2 by the same method. We get J_1^w on $H(\omega_1)^{V[G][g]}$ using $W_{\alpha+1}^*$ there. The J_1 -closure of $V[h]$ and homogeneity of $\text{Col}(\omega, \omega_2)$ give us J_1^w on $H(\omega_1)^{V[g]}$ in $V[g]$. We get I in V such that $I(b)[g] = J_1^w(\langle b, g \rangle)$. I extends to an (ω_3, τ) operator in V , where $\tau_g = x$. Going back to $V[h]$, we can convert I into the desired J_1^w -like function J_2 .

We leave any further detail to the reader. This proves the claim. \square

We can now finish the proof of lemma 50. Let J_n be the function in $V[h]$ we constructed. In $V[g]$, the $J_n(b)$ for b in the $H(\omega_1)$ cone over x collectively witness $W_{\alpha+1}^*$. This is because, as in the proof of 49, we can reduce Σ_n truth about b in $J_\alpha(\mathbb{R})^{V[g]}$ to Σ_1 truth in $J_n(b)$. We get $I_{\alpha+1}$ because, as a byproduct of our construction, $J_n \cap H(\omega_1)^{V[g]}$ is contained in $((J_0)^{w, \dots, w})^{V[G][g]}$. The former captures truth over $J_\alpha(\mathbb{R})^{V[g]}$, and the latter captures truth over $J_\alpha(\mathbb{R})^{V[G][g]}$. This implies that π is fully elementary, as desired. \square

7. THE ADMISSIBLE CASES

Let us fix $g \subset \text{Col}(\omega, \omega_1)$ which is V -generic, and a critical ordinal γ in $V[g]$ of type (3). That is, letting α be the strict sup of the critical ordinals $< \gamma$, we have $\alpha < \gamma$. We assume that W_α^* holds in $V[g]$, and we wish to show that $W_{\gamma+1}^*$ holds in $V[g]$. The analysis of scales in $L(\mathbb{R})$ shows that α begins a Σ_1 gap $[\alpha, \beta]$, and $J_\alpha(\mathbb{R})$ is admissible. The possibilities are that $\alpha = \beta = \gamma - 1$ (the admissible empty gap), that $\alpha < \beta = \gamma - 1$ (the strong gap), or that $\alpha < \beta = \gamma$ (the weak gap). But for the most part, we do not need to distinguish these three cases here. We also assume $\text{WRP}_{(2)}(\omega_2)$, and that NS is saturated. Our overall plan is:

Step 1. Working in $V[g]$, we construct a mouse N and iteration strategy Σ^g which code up truth at the end of the gap $[\alpha, \beta]$. N will be a mouse over some real parameter z .

Step 2. Letting $\tau_g = z$, we show that N and Σ^g yield a mouse N_τ over τ and an ω_2 -iteration strategy Σ for N_τ , both in V , via the equations $N_\tau[g] = N$, and $\Sigma = \Sigma^g \upharpoonright V$.

Step 3. We show that Σ extends to act on $H(\omega_3)$.

Step 4. We then further extend Σ so that it acts on all trees in the $H(\omega_1)$ of $V[g][h]$, where $h \subset \text{Col}(\omega, \omega_2^V)$ is $V[g]$ -generic. At the same time we find versions $[\alpha^H, \beta^H]$ of our gap $[\alpha, \beta]$ in $V[g][H]$, whenever $H \in V[g][h]$, along with appropriately elementary embeddings from $J_\beta(\mathbb{R})^{V[g]}$ to $J_{\beta^H}(\mathbb{R})^{V[g][H]}$.

Step 5. We proceed as in the inadmissible case, but using Σ -mice with Woodin cardinals to witness $W_{\gamma+1}^*$ in $V[g]$. As before, the proof breaks into cases, according to whether $\alpha \in \text{ran}(\pi_{\text{NS}})$ or not.

Steps 1 and 2 follow [22] closely. The only difference here is that we want N to have ω Woodin cardinals, so that we can lift the gap $[\alpha, \beta]$ to $V[g][h]$, for h generic over $\text{Col}(\omega, \omega_2)$, using an \mathbb{R} -genericity iteration over $V[g][h]$. We now proceed to details.

Definition 51. Let Γ be the pointclass $\Sigma_1^{J_\alpha(\mathbb{R})}$. What is called the envelope of Γ , or $\text{ENV}(\Gamma)$, is the class of all $A \subseteq \mathbb{R}$ which are countably captured by Γ in that there is a real x such that for any countable $\sigma \subseteq \mathbb{R}$, $A \cap \sigma$ is $OD^{<\alpha}(\sigma, x)$. The analysis of scales from [23] shows that if $\alpha = \beta$ or $[\alpha, \beta]$ is strong, then

$$\text{ENV}(\Gamma) = J_{\beta+1}(\mathbb{R}) \cap P(\mathbb{R}),$$

and if $[\alpha, \beta]$ is weak, then

$$\text{ENV}(\Gamma) = J_\beta(\mathbb{R}) \cap P(\mathbb{R}).$$

If $\alpha = \beta$ or $[\alpha, \beta]$ is strong, put

$$e(\Gamma) = \{A \subseteq \mathbb{R} \mid A \text{ is ordinal definable over } J_\beta(\mathbb{R})\}.$$

If $[\alpha, \beta]$ is weak, put

$$e(\Gamma) = \{A \subseteq \mathbb{R} \mid \exists \xi < \beta \text{ (} A \text{ is ordinal definable over } J_\xi(\mathbb{R})\text{)}\}.$$

So

$$\text{ENV}(\Gamma) = \bigcup_{z \in \mathbb{R}} e(\Gamma)(z)$$

is the boldface pointclass associated to $e(\Gamma)$.

A 0-suitable premouse is a minimal premouse N with one Γ Woodin, called δ^N . Such an N is A -iterable if it has a partial iteration strategy moving the $Col(\omega, \delta^N)$ term relation for A correctly. The reader should see [22] or [24] for full definitions. We have the following basic result of Woodin.

Theorem 52. (*Woodin*) *For any countable transitive set X , and A such that $A \in e(\Gamma)(z)$ for some $z \in X$, there is a 0-suitable, A -iterable premouse over X .*

No full proof of this key lemma has ever been written up. There is part of a proof in [16]. The paper [24] outlines a proof in the weak gap case.

If $[\alpha, \beta]$ is weak, we let z_0 be a real parameter such that for some finite set F of ordinals, $\langle z_0, F \rangle$ satisfies a non-reflecting Σ_n type, where n is least such that $\rho_n(J_\beta(\mathbb{R})) = \mathbb{R}$. We let F_0 be the Brouwer-Kleene least such F , and let

$$J_\beta(\mathbb{R}) = \bigcup_n H_n$$

be the decomposition given in [23]. Thus in particular each H_n collapses to a member of $J_\beta(\mathbb{R})$. If $[\alpha, \beta]$ is not weak, let z_0 be a real such that for some Σ_1 formula $\varphi(v)$, we have

$$J_{\beta+1}(\mathbb{R}) \models \varphi[z_0] \text{ but } J_\beta(\mathbb{R}) \not\models \varphi[z_0].$$

Let ρ be a standard $Col(\omega, \omega_1^V)$ term for a real such that $\rho_g = z_0$, and let p_0 force all the properties of ρ we have enumerated so far. For $p \in Col(\omega, \omega_1^V)$ such that $p_0 \subseteq p$, let $g_p(n) = p(n)$ if $n \in \text{dom}(p)$, and $g_p(n) = g(n)$ otherwise. Let τ be a term for a real such that τ_g codes ρ_g and g in some natural way. It is easy to do this so that

- (a) $z_0 \leq_T \tau_g$, and
- (b) for all p , $\tau_g \equiv_T \tau_{g_p}$.

Put $z = \tau_g$. For any $A \in e(\Gamma)(z)$, put

$$B(A) = \{ (y, t) \in \mathbb{R} \times \mathbb{R} \mid y \text{ codes a countable, transitive } X \text{ such that} \\ z \in X, \text{ and } t \text{ codes } \text{Th}_\omega^N(X \cup \{X, \tau_A^N\}), \\ \text{for some (all) 0-suitable, } A\text{-iterable } X\text{-premouse } N \}.$$

Here τ_A^N is the standard $Col(\omega, \delta^N)$ -term capturing A . “Some” is equivalent to “all” in the definition above because A -iterability yields

an approximation to the comparison process which suffices to determine the theory in question. Note that $B(A) \in e(\Gamma)(z)$, because $e(\Gamma)(z)$ is closed under real quantification. By the scale analysis of [23], we have a self-justifying system $\mathcal{A} = \{A_i \mid i < \omega\}$ such that

- (1) each A_i is in $e(\Gamma)(z)$,
- (2) if $\alpha = \beta$ or $[\alpha, \beta]$ is strong, then for each $n < \omega$, $\text{Th}_n^{J_\beta(\mathbb{R})}(\mathbb{R}) \in \mathcal{A}$
- (3) if $[\alpha, \beta]$ is weak, then for all n , $\text{Th}_\omega^{H_n}(\mathbb{R} \cup \{z, F_0\}) \in \mathcal{A}$, and
- (4) for any n , $B(\langle A_i \mid i \leq n \rangle) \in \mathcal{A}$, where we regard $\langle A_i \mid i \leq n \rangle$ as a set of reals via some natural coding.

It is easy to also arrange that there is a fixed term $\dot{\mathcal{A}}$ such that

- (5) for all $p \supseteq p_0$, $\dot{\mathcal{A}}^{g_p} = \dot{\mathcal{A}}^g$.

Let X be countable transitive, with $z \in X$. Let N^n be a 0-suitable, $\langle A_i \mid i \leq n \rangle$ -iterable mouse over X . As in [22] we can simultaneously compare all the N^n to get a 0-suitable N over X such that N is $\langle A_i \mid i \leq n \rangle$ -iterable for all n . But then condensation for term relations implies that N has a unique fullness preserving (ω_1, ω_1) iteration strategy which moves all the term relations $\tau_{A_i}^N$ correctly.⁸ Put

$$Q(X) = \text{Hull}^N(X \cup \{X\} \cup \{\tau_A^N \mid A \in \mathcal{A}\}).$$

Condensation for term relations of a self-justifying system implies that $Q(X)$ has all the properties of N , namely, it is 0-suitable, and has a unique fullness preserving (ω_1, ω_1) iteration strategy which moves all the term relations $\tau_{A_i}^{Q(X)}$ correctly. Moreover, $Q(X)$ is “sound”, in that

$$Q(X) = \text{Hull}^{Q(X)}(X \cup \{X\} \cup \{\tau_A^{Q(X)} \mid A \in \mathcal{A}\}).$$

Let

$$N_0 = Q(V_\omega \cup \{z\}),$$

and

$$N_{k+1} = Q(N_k),$$

and

$$N = \bigcup_{k < \omega} N_k.$$

Put $\delta_k^N = \delta^{N_k}$. We regard N as a premouse over z in the natural way. Note that because N_k is suitable, and hence Γ -full, no level of N_{k+1} projects across $o(N_k)$, and thus the δ_k are all Woodin in N .

Lemma 53. *There is a unique \mathcal{A} -guided strategy for N in $V[g]$.*

⁸See [22]. The strategy chooses the limit over n of branches b_n moving all $\tau_{A_i}^N$ for $i \leq n$ correctly.

Proof. As in [22], there is a unique \mathcal{A} -guided iteration strategy Σ_0 for N_0 . Let

$$i: N_0 \rightarrow S_0$$

be an iteration map by Σ_0 . We can let i act on all of N , giving rise to

$$i: N \rightarrow S.$$

Put also $S_m = i(N_m)$, for all m . We do not yet know that S is even wellfounded, but in fact

Claim 54. For all m , $S_{m+1} = Q(S_m)$.

Proof. We prove it for $m = 0$. Let

$$W_k = \text{Th}_\omega^{N_1}(N|\delta \cup \{\tau_{A_0}^{N_1}, \dots, \tau_{A_k}^{N_1}\}),$$

where $\delta = \delta_0^N$. Note $W_k \in N_0$ because N_0 is Γ -full. Let

$$B = B(\langle A_0, \dots, A_k \rangle).$$

Now N_0 satisfies the sentence “it is forced in $Col(\omega, \delta)$ that if y codes $N|\delta$ and t codes W_k , then $(y, t) \in \tau_B^{N_0}$.” Thus the same sentence is true of $i(\delta)$, $i(W_k)$, and $i(\tau_B^{N_0})$ in S_0 . But $i(\tau_B^{N_0}) = \tau_B^{S_0}$, and so

$$\begin{aligned} \text{Th}_\omega^{S_1}(S|i(\delta) \cup \{\tau_{A_0}^{S_1}, \dots, \tau_{A_k}^{S_1}\}) &= i(W_k) \\ &= \text{Th}_\omega^{Q(S_0)}(S|i(\delta) \cup \{\tau_{A_0}^{Q(S_0)}, \dots, \tau_{A_k}^{Q(S_0)}\}). \end{aligned}$$

It follows that there is a natural isomorphism between

$$\text{Hull}_\omega^{S_1}(S|i(\delta) \cup \{\tau_{A_0}^{S_1}, \dots, \tau_{A_k}^{S_1}\})$$

and

$$\text{Hull}_\omega^{Q(S_0)}(S|i(\delta) \cup \{\tau_{A_0}^{Q(S_0)}, \dots, \tau_{A_k}^{Q(S_0)}\}).$$

Moreover, these isomorphisms commute with the inclusion maps on the hulls, because they are determined by the $i(W_k)$. Finally, S_1 and $Q(S_0)$ are the unions of the respective sequences of hulls, as k varies. (In the case of S_1 , this is because $N_1 = Q(N_0)$, and i came from an iteration based on N_0 .) Thus $S_1 \cong Q(S_0)$. The proof for $m > 0$ is the same. \square

But now $S_1 = Q(S_0)$ has a unique iteration strategy Σ_1 for trees above S_0 . Letting $i: S \rightarrow T$ come from an iteration of S_1 by this strategy, and $T_m = i(S_m)$ we get $T_{m+1} = Q(T_m)$ for all $m \geq 1$ by repeating the proof of the claim above. We can then move on to iterating T_2 above T_1 , etc. Clearly, this describes an iteration strategy for N acting on normal trees.⁹ \square

⁹In fact our strategy applies to trees of the form: a stack of normal trees below the first Woodin, then a stack of normal trees between the first and second Woodin, then a stack between the second and third, etc.

N is a mouse over τ_g , but it can be re-arranged as a mouse over τ_{g_p} whenever $p \supseteq p_0$. The re-arranged mouse has the same universe and extender sequence; it just has a different (but Turing equivalent) real distinguished at the bottom. What is more, we have a fixed term \dot{N} such that for all $p \supseteq p_0$, \dot{N}_{g_p} is the re-arrangement of N as a mouse over τ_{g_p} . This is because of the symmetry in the construction of N , and in particular, because $\dot{A}_g = \dot{A}_{g_p}$ for all such p . This enables us to build in V a premouse N_τ over τ such that $N_\tau[g] = N$. We construct $N_\tau|\alpha$ by induction on α , maintaining that

$$(N_\tau|\alpha)[g] = N|\alpha,$$

along with the correspondence of projecta and parameters. α is active in N_τ iff it is active in N , and if so,

$$E_\alpha^{N_\tau} = E_\alpha^N \cap N_\tau|\alpha.$$

$N_\tau|(\alpha + 1) \in V$ because by induction, $N_\tau|\alpha \in V$, and because E_α^N is independent of g . $E_\alpha^{N_\tau}$ is an extender over N_τ because g was generic over V , and the forcing is small. The reader can find all the details of this construction in [24]. Let Σ^g be the unique iteration strategy for N given by the Lemma. Iterating N_τ is the same as iterating $N_\tau[g] = N$, because the forcing is small, and thus we can regard Σ^g as a strategy for N_τ . Moreover Σ , which denotes $\Sigma^g \upharpoonright V$, is in V by the symmetry in our construction of Σ^g . Since Σ^g condenses well, Σ condenses well. We have finished steps 1 and 2 of the general plan.

We now execute step 3. Here we use $\text{WRP}_{(2)}(\omega_2)$ in V to extend our ω_2 -iteration strategy to an ω_3 -iteration strategy. In fact, simultaneous stationary reflection for pairs of subsets of ω_2 is enough.

Lemma 55. *Let M be a premouse of cardinality $\leq \omega_1$, and let Σ be any ω_2 iteration strategy for M which condenses well. Suppose that whenever $S, T \subseteq \omega_2$ are stationary and consist of ordinals of countable cofinality, there is a $\nu < \omega_2$ such that S and T are stationary in ν . Then there is a unique ω_3 iteration strategy Ω for M such that*

- (1) $\Sigma \subseteq \Omega$, and
- (2) Ω condenses well.

Proof. We omit the easy proof that there is at most one such Ω . Fix η large. Let \mathcal{T} be an iteration tree on M with $\text{lh}(\mathcal{T}) < \omega_3$. We say $\langle X_\alpha \mid \alpha < \omega_2 \rangle$ is a \mathcal{T} -chain iff

- (a) $X_\alpha \prec V_\eta$, for all $\alpha < \omega_2$,
- (b) $\alpha < \beta \Rightarrow X_\alpha \subsetneq X_\beta$, and $X_\lambda = \bigcup_{\alpha < \lambda} X_\alpha$ for limit λ ,
- (c) $M, \mathcal{T} \in X_0$, and
- (d) $|X_\alpha| = \omega_1$, and $X_\alpha \cap \omega_2 \in \omega_2$, for all $\alpha < \omega_2$.

Given a \mathcal{T} -chain \vec{X} , we let $\pi_\alpha: H_\alpha \cong X_\alpha$ with H_α transitive, let $\pi_{\alpha,\beta} = \pi_\beta^{-1} \circ \pi_\alpha$, and let $\mathcal{T}_\alpha = \pi_\alpha^{-1}(\mathcal{T})$. We say that \vec{X} is Σ -good iff each \mathcal{T}_α is by Σ , and in that case, we set

$$b_\alpha = \Sigma(\mathcal{T}_\alpha)$$

for all $\alpha < \omega_2$. There is of course no reason that we should have $b_\alpha \in H_\alpha$.

Claim 56. Let \vec{X} be a Σ -good \mathcal{T} -chain, and let $\gamma < \omega_2$ with $\text{cof}(\gamma) = \omega_1$; then for club many $\alpha < \gamma$, $\pi_{\alpha,\gamma}{}''b_\alpha \subseteq b_\gamma$.

Proof. We take cases on the cofinality of the length of \mathcal{T} . Suppose first $\text{cof}(\text{lh}(\mathcal{T})) = \omega$. Then for all sufficiently large $\alpha < \gamma$, $\text{ran}(\pi_{\alpha,\gamma})$ is cofinal in b_γ , and thus applying condensation to the support-closed subtree of $\mathcal{T}_\gamma \widehat{\ } b_\gamma$ determined by $\text{ran}(\pi_{\alpha,\gamma})$, we get that $\pi_{\alpha,\gamma}^{-1}{}''b_\gamma = \Sigma(\mathcal{T}_\alpha) = b_\alpha$. So the desired club is just a tail below γ . Suppose $\text{cof}(\text{lh}(\mathcal{T})) = \omega_1$. Then $\text{cof}(\text{lh}(\mathcal{T}_\xi)) = \omega_1$, for all ξ . Also, for all $\alpha < \gamma$, $\pi_{\alpha,\gamma}{}''b_\alpha$ is cofinal in $\text{lh}(\mathcal{T}_\gamma)$. Since \mathcal{T}_γ has at most one cofinal branch, we get $\pi_{\alpha,\gamma}{}''b_\alpha \subseteq b_\gamma$. Finally, suppose $\text{cof}(\text{lh}(\mathcal{T})) = \omega_2$. As in case two, $\text{cof}(\text{lh}(\mathcal{T}_\xi)) = \text{cof}(X_\xi \cap \omega_2)$, for all ξ , but now, $\alpha < \gamma \Rightarrow \pi_{\alpha,\gamma}$ is discontinuous at $\text{lh}(\mathcal{T}_\alpha)$. Fixing γ with $\text{cof}(\gamma) = \omega_1$, we can find club many $\alpha < \gamma$ such that $\text{ran}(\pi_{\alpha,\gamma}) \cap b_\gamma$ is cofinal in $\sup \pi_{\alpha,\gamma}{}''\text{lh}(\mathcal{T}_\alpha)$. For any such α , condensation for the support-closed subtree of $\mathcal{T}_\gamma \widehat{\ } b_\gamma$ determined by $\text{ran}(\pi_{\alpha,\gamma})$, implies that $\pi_{\alpha,\gamma}^{-1}{}''b_\gamma = \Sigma(\mathcal{T}_\alpha) = b_\alpha$. \square

Let \vec{X} be a Σ -good \mathcal{T} -chain. We say \vec{X} is *coherent* if and only if whenever $\alpha < \gamma < \omega_2$, then $\pi_{\alpha,\gamma}{}''b_\alpha \subseteq b_\gamma$. In this case, we say \vec{X} *justifies* b , where

$$b = \bigcup_{\alpha < \omega_2} b_\alpha.$$

It is easy to see

Claim 57. \mathcal{T} has at most one branch b which is justified by some coherent \mathcal{T} -chain.

Proof. If \vec{X} and \vec{Y} are Σ -good \mathcal{T} -chains, then for club many $\alpha < \omega_2$, $X_\alpha \cap \omega_2 = Y_\alpha \cap \omega_2$. Thus for club many $\alpha < \omega_2$, $\mathcal{T}_\alpha^{\vec{X}} = \mathcal{T}_\alpha^{\vec{Y}}$ and $b_\alpha^{\vec{X}} = b_\alpha^{\vec{Y}}$. \square

So we define

$$\Omega(\mathcal{T}) = b \Leftrightarrow b \text{ is justified by some coherent } \mathcal{T}\text{-chain.}$$

Claim 58. If \mathcal{T} is by Ω , then every \mathcal{T} -chain is Σ -good.

Proof. Let \mathcal{T} be of minimal length such that the claim is false. Suppose first that $lh(\mathcal{T})$ is a limit ordinal. Let \vec{X} be a \mathcal{T} -chain. If $\alpha < \gamma < \omega_2$, then \mathcal{T}_α is the collapse of a support-closed subtree of \mathcal{T}_γ , so since Σ condenses well, we have that \mathcal{T}_γ is not by Σ for all sufficiently large $\gamma < \omega_2$. Using a surjective map $f: \omega_2 \rightarrow lh(\mathcal{T})$ with $f \in X_0$, and a Fodor argument, we can fix $\xi < lh(\mathcal{T})$ such that for stationary many $\alpha < \omega_2$, $\pi_\alpha^{-1}(\mathcal{T} \upharpoonright \xi)$ is not by Σ . But \vec{X} is a $\mathcal{T} \upharpoonright \xi$ -chain, contrary to the minimality of $lh(\mathcal{T})$. Thus $lh(\mathcal{T}) = \lambda + 1$ for some λ . It is clear that λ must be a limit ordinal. Let $b = \Omega(\mathcal{T} \upharpoonright \lambda)$, and let \vec{X} be a \mathcal{T} -chain. Let \vec{Y} be a $\mathcal{T} \upharpoonright \lambda$ -chain which justifies b . There are club many $\alpha < \omega_2$ such that $X_\alpha \cap lh(\mathcal{T}) = Y_\alpha \cap lh(\mathcal{T})$, and for such α , $(\pi_\alpha^{\vec{X}})^{-1}(b) = (\pi_\alpha^{\vec{Y}})^{-1}b = b_\alpha^{\vec{Y}}$. Thus for club many α , $\mathcal{T}_\alpha^{\vec{X}}$ is by Σ . Condensation implies this is true for all α . This contradiction completes the proof. \square

Claim 59. $\Sigma \subseteq \Omega$, and Ω condenses well.

Proof. If \mathcal{T} is by Σ , then in any \mathcal{T} -chain, we have $\mathcal{T}_\alpha = \mathcal{T}$ for all $\alpha < \omega_2$, so every \mathcal{T} -chain justifies $\Sigma(\mathcal{T})$. For condensation, suppose $\Omega(\mathcal{T}) = b$, and $\mathcal{U} \hat{=} c$ is the collapse of a support-closed subtree of $\mathcal{T} \hat{=} b$, and $\Omega(\mathcal{U}) = d$ where $d \neq c$. It is easy to see that there is a single \vec{X} , with $\mathcal{T}, \mathcal{U}, b, c, d \in X_0$, which justifies both b and d . But this gives a failure of condensation for Σ . \square

Claim 60. Suppose \mathcal{T} is by Ω , and $lh(\mathcal{T}) < \omega_3$; then there is a b such that $\Omega(\mathcal{T}) = b$.

Proof. Fix any $\xi < lh(\mathcal{T})$, and any \mathcal{T} -chain \vec{X} . Since \vec{X} is Σ -good, we have $b_\alpha = \Sigma(\mathcal{T}_\alpha)$ for $\alpha < \omega_2$. We claim that exactly one of the following holds:

- (1) for ω -club many $\alpha < \omega_2$, $\pi_\alpha^{-1}(\xi) \in b_\alpha$,
- (2) for ω -club many $\alpha < \omega_2$, $\pi_\alpha^{-1}(\xi) \notin b_\alpha$.

It is clear that not both can hold. Suppose both fail. Let S be the stationary set of α of cofinality ω where $\xi \in \text{ran}(\pi_\alpha)$ and (1) fails, and T the stationary set of α of cofinality ω where (2) fails. By our stationary reflection hypothesis, we can fix γ of cofinality ω_1 such that both S and T are stationary in γ . Note $\xi \in \text{ran}(\pi_\gamma)$. If $\pi_\gamma^{-1}(\xi) \in b_\gamma$, then by the first claim, $\pi_\alpha^{-1}(\xi) \in b_\alpha$ for club-in- γ many α , so T was not stationary in α , contradiction. Similarly, if $\pi_\gamma^{-1}(\xi) \notin b_\gamma$, then The first claim implies S is not stationary in γ , a contradiction. So at least one of (1) and (2) holds. It also implies that the ω -clubs of (1) and (2) can be taken to be fully club in ω_2 . Define b by:

$$\xi \in b \Leftrightarrow \text{for club many } \alpha < \omega_2, \pi_\alpha^{-1}(\xi) \in b_\alpha.$$

Taking a diagonal intersection, we can find a club $C \subseteq \omega_2$ such that for $\alpha \in C$, $\pi_\alpha "b_\alpha \subseteq b$. But then $\langle X_\alpha \mid \alpha \in C \rangle$ is a coherent \mathcal{T} -chain which justifies b . \square

This completes the proof of Lemma 55. \square

Applying Lemma 55, let us use Σ to denote the unique ω_3 iteration strategy for N_τ which condenses well, and extends $\Sigma^g \upharpoonright V$. Proceeding to step 4, we need to further extend Σ so that it acts on all trees in $H(\omega_1)^{V[g][h]}$, whenever h is $V[g]$ -generic over some poset in $H(\omega_2)^{V[g]}$. These extensions of Σ will be mutually consistent. At the same time, we will be showing that the gap $[\alpha, \beta]$ of $V[g]$ has counterparts in every $V[g][h]$. The following little lemma will be useful.

Lemma 61. *Let Γ be an iteration strategy for S which condenses well. Let $\pi: R \rightarrow S$ be sufficiently elementary that the pullback strategy Γ^π for R exists; then Γ^π also condenses well.*

Proof. Let \mathcal{T} be a tree according to Γ^π , and \mathcal{U} a support-closed subtree of \mathcal{T} corresponding to those $\mathcal{M}_\alpha^\mathcal{T}$ such that $\alpha \in X$, and let $\bar{\mathcal{U}}$ be the collapse of \mathcal{U} . It is easy to see the lifted tree $\pi\bar{\mathcal{U}}$ is the collapse of the support-closed subtree of $\pi\mathcal{T}$ corresponding to those $\mathcal{M}_\alpha^{\pi\mathcal{T}}$ such that $\alpha \in X$. Since Γ condenses well, $\pi\bar{\mathcal{U}}$ is according to Γ , and hence $\bar{\mathcal{U}}$ is according to Γ^π . \square

We need to use hybrid strategy mice. Suppose Ω is an iteration strategy for some structure M , and Ω condenses well. Let A be transitive, with $M \in A$. We obtain a hybrid Ω -premouse by adding extenders with critical points above A to a coherent sequence we are building, and at the same time closing the model we are building under Ω , and giving it a predicate for Ω . The construction can only go on as long as all (non-dropping) iteration trees according to Ω we construct are in the domain of Ω . (M may or may not be a fine-structural premouse, but in any case, it is convenient to only close under Ω on non-dropping trees.) We refer the reader to [22] for a brief discussion of such Ω -hybrids, and to [16] for a more thorough one.

Definition 62. Let Ω be an $|A|^+$ -iteration strategy for M which condenses well, where A is transitive and $M \in A$; then $P_n^\Omega(A)^\sharp$ is the minimal $|A|^+$ -iterable hybrid Ω -mouse over A which is active, and satisfies “there are n Woodin cardinals”.

We note that the iterations referred to here all leave A , and hence M , fixed. It is part of iterability that they must move Ω to itself. One can hope to construct such iterable hybrid mice in a K^c construction,

because Ω condenses well, and hence Ω will condense to itself under realizing maps. The iterability demand we have made for $P_n^\Omega(A)^\sharp$ in the definition above is the minimal one which guarantees uniqueness, granted that $H(|A|^+)$ is closed under P_{n-1}^Ω -sharps. We shall never consider a putative $P_n^\Omega(A)$ -sharp unless we already know $H(|A|^+)$ is closed under P_{n-1}^Ω -sharps. In practice, we often have more iterability than the minimal demand. Our core-model-induction proof that $H(\omega_3)$ of V is closed under the M_n^\sharp operators generalizes routinely to hybrid mouse operators, and gives:

Lemma 63. *Assume NS is saturated, and $\text{WRP}_{(2)}(\omega_2)$ holds. Let $S \in H(\omega_1)$, and let Ω be an ω_3 -iteration strategy for S which condenses well. Then for all transitive $A \in H(\omega_3)$ such that $S \in A$, and all $n < \omega$, $P_n^\Omega(A)^\sharp$ exists and is ω_3 -iterable.*

This is proved exactly as was theoremjtojw, so we omit the details.

Lemma 63 is the place core model theory gives us new mice. We shall eventually apply it in the case that $\alpha \in \text{ran}(\pi_G)$ for some NS-generic G . In that case, we can take our real z_0 to be in V , and N_τ to be in V , and countable there. (We don't actually need τ at all, N could be a z_0 -premouse.) We shall then apply lemma 63 with $S = N$ and $\Omega = \Sigma$.

Before we take cases on whether $\alpha \in \text{ran}(\pi_G)$, however, we do some further preliminary work related to I_γ .

Lemma 64. *For all $A \in H(\omega_3)$, $P_0^\Sigma(A)^\sharp$ exists and is ω_3 -iterable.*

Proof. We show first that $P_0^\Sigma(A)^\sharp$ exists for all such $A \in H(\omega_2)$, then extend this to $A \in H(\omega_3)$ using simultaneous reflection. Let $G \subset P(\omega_1)/NS$ be V -generic and let

$$i: V \rightarrow M \subseteq V[G]$$

be the generic embedding. Since $N_\tau \in H(\omega_2)^V$, we have $N_\tau \in M$, and $i \upharpoonright N_\tau \in M$. So inside M , we can form the $(i \upharpoonright N_\tau)$ -pullback of $i(\Sigma)$, which we denote by $i(\Sigma)^i$. From the point of view of M , $i(\Sigma)^i$ is an ω_3 iteration strategy for N_τ , and by Lemma 61, it condenses well in M .

Claim 65. $i(\Sigma)^i$ agrees with Σ on all trees in the intersection of the two domains.

Proof. We first consider trees in $H(\omega_2)^V$, all of which are in both domains. Let $\mathcal{T} \in H(\omega_2)^V$ be a tree according to Σ . Note that $i \upharpoonright N_\tau \in M$. In M the copied tree $i\mathcal{T}$ on $i(N_\tau)$ is satisfied to be the collapse of a support closed subtree $i(\mathcal{T})$. Since $i(\Sigma)$ condenses well in M , $i\mathcal{T}$ is according to $i(\Sigma)$. Hence \mathcal{T} is according to the pullback $i(\Sigma)^i$. Now let \mathcal{U} be a tree in V of size ω_2^V in V which is according

to both Σ and $i(\Sigma)^i$, and is of limit length. Let

$$b = i(\Sigma)^i(\mathcal{U})$$

and

$$c = \Sigma(U)$$

and

$$b \neq c.$$

Note that $cf(lh(\mathcal{U}))$ must be countable in $V[G]$. In V , we can write $\mathcal{U} = \bigcup_{\alpha < \omega_2} \mathcal{U}_\alpha$, where this is an increasing continuous chain of support-closed subtrees, each of size ω_1 . Going to $V[G]$, where $cf(lh(\mathcal{U}))$ is countable, we see that

$$b \cap \mathcal{U}_\alpha \text{ is cofinal in } \mathcal{U}_\alpha$$

and

$$c \cap \mathcal{U}_\alpha \text{ is cofinal in } \mathcal{U}_\alpha,$$

for all sufficiently large α , so we may assume all α . Let $\bar{\mathcal{U}}_\alpha$ denote the collapse of \mathcal{U}_α , \bar{b}_α the collapse of $b \cap \mathcal{U}_\alpha$, \bar{c}_α the collapse of $c \cap \mathcal{U}_\alpha$. Fix α such that $\bar{b}_\alpha \neq \bar{c}_\alpha$. Now

$$\bar{c}_\alpha = \Sigma(\bar{\mathcal{U}}_\alpha)$$

because Σ condenses well in V . On the other hand,

$$\bar{b}_\alpha = i(\Sigma)^i(\bar{\mathcal{U}}_\alpha),$$

because $i(\Sigma)^i$ condenses well in M . Since Σ and $i(\Sigma)^i$ must agree at $\bar{\mathcal{U}}_\alpha$ by the first part, we are done. \square

Now fix a transitive $A \in H(\omega_2)$ such that $N_\tau \in A$. Let $L^\Sigma[A]$ be the minimal model of height ω_3 which has A as a member and is closed under Σ , and is expanded by a predicate for Σ . In M , we can form $L^{i(\Sigma)^i}[A]$ in parallel fashion. By claim 65, these two models are the same. So $L^\Sigma[A] \in M$. But $i(NS)$ is saturated in M , so by the same argument that shows that the existence of a saturated ideal implies 0^\sharp exists, we get some $P \in M$ such that

$$M \models P = P_0^\Sigma(A)^\sharp.$$

Being $P_0^\Sigma(A)^\sharp$ is a first order property, combined with linear iterability by the last (and only) extender in a way that moves Σ to itself, for iterations of length $< \omega_3^M = \omega_3^V$. But now let h be V -generic for $Col(\omega, \omega_2^V)$, and such that $G \in V[h]$. The required iterability of P is upward absolute, that is

$$V[h] \models P \text{ is } \omega_3^V \text{ iterable,}$$

so since ω_3^V is still uncountable in $V[h]$,

$$V[h] \models P = P_0^\Sigma(A)^\sharp.$$

By the homogeneity of $Col(\omega, \omega_2)$, $P \in V$, and

$$V \models P = P_0^\Sigma(A)^\sharp,$$

and we are done in the case $A \in H(\omega_2)$. We now use $WRP_{(2)}(\omega_2)$, via hybrid mouse reflection at ω_2 , to show that $P_0^\Sigma(A)^\sharp$ exists, for all transitive $A \in H(\omega_3)$ such that $N_\tau \in A$. Without loss of generality, let us assume $A \subseteq \omega_2$. Let ϕ be a formula in the language of set theory together with a predicate symbol $\dot{\Sigma}$ and constant symbol \dot{A} , and let $\vec{\alpha} \in \omega_2^{<\omega}$. For $\sigma \prec H(\omega_2)$ countable, let

$$\pi_\sigma: M_\sigma \rightarrow H_{\omega_2}$$

be the transitive collapse, and $A_\sigma = \pi_\sigma^{-1}(A)$, $N_\sigma = \pi_\sigma^{-1}(N_\tau)$, and $\vec{\alpha}_\sigma = \pi_\sigma^{-1}(\vec{\alpha})$. Note that for such σ , the pullback strategy Σ^{π_σ} is a full ω_3 iteration strategy for N_σ , and it condenses well. Using our saturated ideal, we then have that

$$P_0^{\Sigma^{\pi_\sigma}}(A_\sigma)^\sharp \text{ exists,}$$

and is ω_3 iterable. We put

$$\begin{aligned} (\phi, \vec{\alpha}) \in P_0^\Sigma(A)^\sharp &\Leftrightarrow \text{for club many } \sigma \in P_{\omega_1}(H_{\omega_2}) \\ &(\phi, \vec{\alpha}_\sigma) \in P_0^{\Sigma^{\pi_\sigma}}(A_\sigma)^\sharp. \end{aligned}$$

(Here we identify the structure $P_0^\Sigma(A)^\sharp$ with its theory with parameters from ω_2 .) In order to see that this definition works, we must show that every $(\phi, \vec{\alpha})$ is decided on a club. So suppose neither $(\phi, \vec{\alpha})$ nor $(\neg\phi, \vec{\alpha})$ is in $P_0^\Sigma(A)^\sharp$ according to the definition above. As usual, we find a transitive $X \prec H_{\omega_2}$ with $|X| = \omega_1$ such that both sets are stationary in $P_{\omega_1}(X)$. Without loss of generality, assume $\vec{\alpha}, N_\tau \in X$, and

$$(\phi, \vec{\alpha}) \in P_0^\Sigma(A \cap X)^\sharp.$$

It is then easy to see that for club many $\sigma \in P_{\omega_1}(X)$, $(\phi, \vec{\alpha}_\sigma) \in P_0^{\Sigma^{\pi_\sigma}}(A_\sigma)^\sharp$. That is because for club many $\sigma \prec X$, $\sigma = Z \cap X$ for some $Z \prec V_\eta$ with $P_0^\Sigma(A \cap X)^\sharp \in Z$. Letting $\pi \supseteq \pi_\sigma$ be the collapse of Z , we get that

$$\pi^{-1}(P_0^\Sigma(A \cap X)^\sharp) = P_0^{\Sigma^{\pi_\sigma}}(A_\sigma)^\sharp.$$

To see this, note $\pi^{-1}(\Sigma) \subseteq \Sigma^{\pi_\sigma}$ by our argument in the first part of the proof of 64. So the strategy predicate in $\pi^{-1}(P_0^\Sigma(A \cap X)^\sharp)$ denotes Σ^{π_σ} . Moreover, iterates S of $\pi^{-1}(P_0^\Sigma(A \cap X)^\sharp)$ embed into iterates S^* of $P_0^\Sigma(A \cap X)^\sharp$, and the strategy predicate of S^* denotes a fragment of Σ , so the strategy predicate of S denotes a fragment of Σ^{π_σ} . So we

have shown $(\phi, \vec{\alpha}) \in P_0^\Sigma(A)^\sharp$, or $(\neg\phi, \vec{\alpha}) \in P_0^\Sigma(A)^\sharp$. This easily gives that our $P_0^\Sigma(A)^\sharp$ has the first order properties required of $P_0^\Sigma(A)^\sharp$.

We must see that its strategy predicate denotes Σ , and that linear iterates of it move Σ correctly. Let I be a linear iteration of length $< \omega_3$ of P , with last model Q such that $\dot{\Sigma}^Q \not\subseteq \Sigma$. We can find

$$\pi: H \rightarrow V_\eta$$

such that M is countable transitive, and everything relevant is in $\text{ran}(\pi)$. Because $\text{ran}(\pi) \cap \omega_2$ meets the clubs definable over V_η from elements of $\text{ran}(\pi)$, we get

$$\pi^{-1}(P) = P_0^{\Sigma^\pi}(\pi^{-1}(A))^\sharp.$$

Also, $\pi^{-1}(\Sigma) = \Sigma^\pi \cap H$. So $\dot{\Sigma}^{\pi^{-1}(N)} \not\subseteq \Sigma^\pi$. This contradicts the fact that linear iterations of $P_0^{\Sigma^\pi}(\pi^{-1}(A))^\sharp$ do move Σ^π to itself, by definition. \square

We shall now use genericity iterations of N_τ to lift $J_\beta(\mathbb{R})^{V[g]}$ and Σ^g to $V[g][h]$, for any h generic over $V[g]$ for a poset in $H(\omega_2)^{V[g]}$. To this end, recall our self-justifying system $\mathcal{A} = \{A_i \mid i < \omega\}$ in $V[g]$. For $A \in \mathcal{A}$ and δ a Woodin cardinal of N , we have the $Col(\omega, \delta)$ -term $\tau_{A,\delta}^N$, whose images in Σ^g -iterations always capture A . Since $N = N_\tau[g]$, we have $\tau_{A,\delta}^N = \rho_g$ for some $Col(\omega, \omega_1^V)$ -term ρ . Let $\sigma_{A,\delta}$ be the canonical $Col(\omega, \omega_1^V) \times Col(\omega, \delta)$ term such that for all generics $k \times l$,

$$(\sigma_{A,\delta})_{k \times l} = (\rho_k)_l.$$

Thus

$$(\sigma_{A,\delta})_{g \times l} = (\tau_{A,\delta}^N)_l,$$

for l being $Col(\omega, \delta)$ generic over $N_\tau[g]$.

Lemma 66. *Let $h \subset \mathbb{P}$ be $V[g]$ -generic where $\mathbb{P} \in (H(\omega_3^V))^{V[g]}$. Then in $V[g][h]$ there are*

- (1) sets $A_i^* \subseteq \mathbb{R}$ such that

$$(HC^{V[g]}, \in, A_i)_{i < \omega} \prec (HC^{V[g][h]}, \in, A_i^*)_{i < \omega},$$

- (2) an ordinal β^h and embedding

$$\pi: J_\beta(\mathbb{R})^{V[g]} \rightarrow J_{\beta^h}(\mathbb{R})^{V[g][h]}$$

such that π is fully elementary if $\alpha = \beta$ or $[\alpha, \beta]$ is strong, and π is Σ_n -elementary for n least such that $\rho_n(J_\beta(\mathbb{R})^{V[g]}) = \mathbb{R}$ otherwise, and

- (3) a unique ω_3^V -iteration strategy $\Sigma^h \in V[g][h]$ for N which extends $\Sigma^g \cup \Sigma$ and condenses well.

Proof. We begin with (1). A_i^* comes from interpreting the images of $\tau_{A_i}^{N_\tau}$ under genericity iterations. Note first

Claim 67. In V , let $M \in H(\omega_3)$ be any non-dropping Σ -iterate of N_τ , and let $k < \omega$. Let $x \in \mathbb{R} \cap V[g][h]$; then there is (in V) a Σ -iteration map $i: M \rightarrow P$ with $\text{crit}(i) > \delta_k^M$ such that for any $\text{Col}(\omega, \delta_k^M)$ -generic l over P such that $l \in V[g][h]$ and $g, h \in P[l]$, we have that $x \in P[l][f]$, for some $f \in V[g][h]$ such that f is $\text{Col}(\omega, \delta_{k+1}^P)$ -generic over P .

Proof. Let $x = \sigma_{g \star h}$. Working in V , we use the standard genericity iteration for the ω_2^V -generator version of the extended algebra of M at δ_{k+1}^M to make σ generic. We get in V an $i: M \rightarrow P$ with $\text{crit}(i) > \delta_k^M$ such that for any $\text{Col}(\omega, \delta_k^M)$ -generic l over P , there is f as in our claim with $\sigma \in P[l][f]$. So if $g, h \in P[l]$, then $x \in P[l][f]$. The important thing to note is that the genericity iteration yielding i terminates. This follows from the fact that $P^\Sigma(C)^\sharp$ exists, where $C \in H(\omega_3)$ codes up σ and the iteration from N_τ to M . \square

Claim 68. Let $i: N_\tau \rightarrow P$ and $j: N_\tau \rightarrow Q$ be non-dropping Σ -iterations of N_τ (in V), and let δ and μ be Woodin cardinals of N_τ . Let $A \in \mathcal{A}$ be in our self-justifying system from $V[g]$. Let $x \in \mathbb{R}^{V[g][h]}$ be such that

$$x \in P[g][l_0] \cap Q[g][l_1],$$

where l_0, l_1 are generic over $P[g], Q[g]$ for the collapses of $i(\delta)$ and $j(\mu)$, respectively. Then

$$x \in i(\sigma_{A,\delta})_{g \times l_0} \Leftrightarrow x \in j(\sigma_{A,\mu})_{g \times l_1}.$$

Proof. If not, we have $(p, q) \in g \star h$ such that

$$(p, q) \Vdash \phi(\check{N}_\tau, \check{\Sigma}),$$

where ϕ in the language for forcing over V expresses the failure of our claim in $V[g][h]$. Here ϕ also involves check-names for $\sigma_{A,\delta}$ and $\sigma_{A,\mu}$, which we have suppressed. In V , let

$$\pi: H \rightarrow V_\eta,$$

where H is transitive and of size ω_1 , $\omega_1 \in H$, and everything relevant is in $\text{ran}(\pi)$. We can find

$$\bar{h} \in V[g]$$

so that

$$g \star \bar{h} \text{ is } \text{Col}(\omega, \omega_1) \star \pi^{-1}(\dot{\mathbb{P}})\text{-generic over } H,$$

and $\pi^{-1}(q) \in \bar{h}$. Note that by condensation for Σ ,

$$\pi^{-1}(\Sigma) \subseteq \Sigma.$$

But then, the fact that $H[g][\bar{h}] \models \phi[N_\tau, \pi^{-1}(\Sigma)]$ yields a Σ^g -iteration of N which fails to move one of the term relations for A correctly. This is a contradiction. \square

Motivated by these claims, working in $V[g][h]$ we put for $x \in \mathbb{R}$ and $A \in \mathcal{A}$,

$$x \in A^h \Leftrightarrow \exists i \exists \delta \exists l (i: N_\tau \rightarrow P \text{ is a } \Sigma\text{-iteration and} \\ l \text{ is } P[g]\text{-generic and } x \in i(\sigma_{A,\delta})_{g \times l}).$$

It is easy to see that

$$A^h \cap V[g] = A,$$

because the iteration given by 67 can be taken in $H(\omega_2)^V$ in this case, and such iterations correspond to iterations by Σ^g , which moves $\tau_{A,\delta}^N$ correctly. Note \mathcal{A} is closed under real quantification. Fixing i , we have a j such that

$$V[g] \models \forall \vec{x} \in \mathbb{R}^{<\omega} (A_j(\vec{x}) \Leftrightarrow \exists y A_i(\vec{x}, y)).$$

But this fact is coded into the first order theory over N_τ of the term relations $\sigma_{A_i,\delta}$ and $\sigma_{A_j,\mu}$. More precisely, given $\delta < \mu$ Woodins of N_τ , there is a $p \in g$ which forces over N_τ the statement “whenever k is $N_\tau[\dot{g}]$ -generic over $Col(\omega, \delta)$ and $\vec{x} \in N_\tau[\dot{g}][k]$, then $\vec{x} \in (\sigma_{A_j,\delta})_{\dot{g} \times k}$ if and only if 1 forces in $Col(\omega, \mu)$ over $N_\tau[\dot{g}][k]$ “there is a y such that $(\vec{x}, y) \in (\sigma_{A_i,\mu})_{\dot{g} \times t}$, where t is the re-arrangement of $k \times \dot{G}$.”” These first order facts are preserved by our genericity iterations of N_τ , and those are sufficiently numerous by 67, and coherent in how they move the $\sigma_{A,\nu}$ by 68, that we get

$$V[g][h] \models \forall \vec{x} \in \mathbb{R}^{<\omega} (A_j^h(\vec{x}) \Leftrightarrow \exists y A_i^h(\vec{x}, y)).$$

We leave any further calculation here to the reader.¹⁰ Also, for any i there is a j such that

$$V[g] \models \forall \vec{x} (A_i(\vec{x}) \Leftrightarrow \neg A_j(\vec{x})).$$

Fixing such i, j , we then have

$$V[g][h] \models \forall \vec{x} (A_i^h(\vec{x}) \Leftrightarrow \neg A_j^h(\vec{x})).$$

Generalizing slightly, we get that for any formula ϕ in the language of our two structures, there is a $j = j_\phi$ such that for all \vec{x} in $V[g]$,

$$((\text{HC}^{V[g]}, \in, A_i)_{i < \omega} \models \phi[\vec{x}]) \Leftrightarrow A_j(\vec{x}),$$

¹⁰See [21] for a similar argument. It was to make this argument possible that we moved to an N with ω Woodins, rather than just one.

and for all \vec{x} in $V[g][h]$

$$((\text{HC}^{V[g][h]}, \in, A_i^h)_{i < \omega} \models \phi[\vec{x}]) \Leftrightarrow A_j^*(\vec{x}).$$

Since $A_j = A_j^h \cap V[g]$, we are done with part (1). Part (2) of the theorem follows easily from part (1), and the fact that the A_i code the appropriate fragments of the theory of $J_\beta(\mathbb{R})$. It is routine then use the A_i^h to construct a structure over $\mathbb{R} \cap V[g][h]$ into which $J_\beta(\mathbb{R})^{V[g]}$ embeds with the required degree of elementarity. One need only show the structure over $\mathbb{R} \cap V[g][h]$ one gets is well-founded. The proof of this is a reflection argument very similar to the proof of Claim 55, so we omit it. We let β^h be such that ω^{β^h} is the height of this structure.

Finally, we turn to (3). By part (2), β^h ends a gap $[\alpha^h, \beta^h]$ in $V[g][h]$, and the A_i^h constitute a self-justifying system which seals this gap. We claim that the A_i^h guide an iteration strategy Σ^h for N_τ , or equivalently for $N = N_\tau[g]$, and that $\Sigma^g \cup \Sigma \subseteq \Sigma^{g,h}$. This is again a simple reflection argument along the lines of the proof of Claim 55, and so again, we omit it. Being guided by a self-justifying system, Σ^h condenses well. \square

Remark 69. An earlier version of this paper, posted on the first author's webpage, had at this point an argument purporting to show that the pullback strategy $i(\Sigma)^i$ used in the proof of Lemma 64 codes that same sort of gap in the $L(\mathbb{R})$ of $\text{Ult}(V, G)$ that Σ^G does in $V[g][G]$. Trevor Wilson found a serious gap in this argument, namely, its assumption that the theory coded into $i(\Sigma)^i$ describes a wellfounded model. Fortunately, we don't need this argument.

Remark 70. We eventually get $\beta^G = \beta$, but only after we have shown $W_{\gamma+1}^*$ holds in $V[g]$. That is because the Foreman-Magidor argument requires a universally Baire prewellorder of length β in $V[g]$.

Corollary 71. *Let $G \subset (P(\omega_1)/NS)^V$ be generic over $V[g]$. Then*

$$V[g][G] \models \forall A \in H(\omega_1) \forall n (P_n^{\Sigma^G}(A)^\sharp \text{ exists and is } \omega_1\text{-iterable}).$$

Proof. Suppose first that $\alpha \in \text{ran } \pi_G$. We may then suppose that $N_\tau = N$ is in V , and countable there. We can then repeat the inductive proof of theorem 28, showing that $H(\omega_3)^V$ is closed under the P_n^Σ operator, for all n . This easily yields the corollary.

Suppose next that $\alpha \notin \text{ran}(\pi_G)$, and let α_0 be least such that $\alpha < \pi_G(\alpha_0)$. As in the inadmissible case, α_0 begins a gap in V . Thus $\pi_G(\alpha_0)$ begins a gap in $\text{Ult}(V, G)$, so $\beta^G < \pi(\alpha_0)$. It follows that in $\text{Ult}(V, G)$, $J_{\beta^G+\omega}(\mathbb{R}) \models \text{AD}$. Standard results on the existence of iterable models with Woodin cardinal under determinacy imply that $J_{\beta^G+\omega}(\mathbb{R})^{V[G]} \models \forall A \in H(\omega_1) \forall n (P_n^{\Sigma^G}(A)^\sharp \text{ exists and is } \omega_1\text{-iterable}).$ (\square)

See [21, theorem 10.1].) But then the same is true in $J_{\beta^{G+\omega}}(\mathbb{R})^{V[G][g]}$, by the elementarity of the Cohen ultrapower. \square

Now let h be $Col(\omega, \omega_2^V)$ -generic over $V[g]$, and $G \in V[g][h]$ be NS -generic over $V[g]$. Note that the extension from $V[g][G]$ to $V[g][h]$ is by a partial order which, in $V[g][G]$, is of size ω_1 and collapses ω_1 . So $V[g][g]$ -to- $V[g][h]$ is a homogeneous extension. We shall show the mice $P_n^{\Sigma^G}(A)^\sharp$ given by 71 are definable from A in $V[g][h]$, thus in $V[g]$ when $A \in V[g]$. Definability comes from lifting their strategies to $V[g][h]$, and that comes from lifting the operators themselves to $V[g][h]$. To do that, we need to use simultaneous reflection in V , so we must consider the P_n^Σ -sharp operators on $H(\omega_3)^V$. The following lemma does the job.

Lemma 72. *For all $n < \omega$,*

- (1) $V \models$ for all transitive $A \in H(\omega_3)$ such that $N_\tau \in A$, $P_n^\Sigma(A)^\sharp$ exists and is ω_3 -iterable,
- (2) if $A \in H(\omega_3)^V$ and P is such that $V \models$ “ $P = P_n^\Sigma(A)^\sharp$ is ω_3 -iterable”, then $V[g][h] \models$ “ $P = P_n^{\Sigma^h}(A)^\sharp$ is ω_1 -iterable”, and
- (3) $V[g][h] \models$ for all countable transitive A such that $N \in A$, $P_n^{\Sigma^h}(A)^\sharp$ exists and is ω_1 -iterable.

Proof. By induction on n . We have already proved (1) when $n = 0$. Part (2) is trivial in this case, since the iterations of P are all linear iterations by its unique extender, and hence are all in V . For part (3), Note that the P_0^Σ -sharp operator determines itself on $V[g][h]$. More precisely, the P_0^Σ -sharp operator on $H(\omega_3)^V$ determines the $P_0^{\Sigma^h}$ -sharp operator on $H(\omega_1)^{V[g][h]}$. For let A be countable transitive in $V[g][h]$, and say $A = \rho_{g \times h}$. Let $B \in V$ be the transitive closure of $\{\rho, N_\tau\}$. We have an ω_3^V -iterable $P = P_0^\Sigma(B)^\sharp$ in V . But then $P[g \times h]$ exists in $V[g][h]$, and we can obtain $P_0^{\Sigma^h}(A)^\sharp$ from it. This is because the determination of Σ^h from Σ we gave (via \mathbb{R} -genericity iterations which create a self-justifying system guiding Σ^h) is sufficiently local that if $M \models \text{ZFC}$ and $\Sigma \cap M \in M$ and $g, h \in M$, then $\sigma^h \cap M \in M$ and is uniformly-in- M definable over M from $\Sigma \cap M, g$, and h . Iterations of $P_0^{\Sigma^h}(A)^\sharp$ correspond to iterations of P as in (2). The latter stretch Σ into Σ , so the former stretch Σ^h into Σ^h .

Now suppose (1) through (3) hold for $n = k$. We consider (1) for $n = k + 1$. We first consider the case $A \in H(\omega_2)^V$. In $V[g]$, let B be the first admissible set over $\{A, g\}$, so that $N \in B$. By Corollary 71 we have P in $V[g][G]$ such that

$$V[g][G] \models P = P_{k+1}^{\Sigma^G}(B)^\sharp,$$

in the sense that P has the first order properties, and is $\omega_1^{V[g][G]}$ -iterable via a strategy which moves Σ^G to itself. We claim that

$$V[g][h] \models P = P_{k+1}^{\Sigma^h}(B)^\sharp,$$

in the sense that P is $\omega_1^{V[g][h]}$ iterable in $V[g][h]$ via a strategy which moves Σ^h to itself. The iteration strategy for P in $V[g][h]$ is the one guided by the Q -structures provided by (3) for $n = k$. Let Γ be this strategy, and suppose Γ fails in $V[g][h]$. Let \mathcal{O}^h be the $P_k^{\Sigma^h}$ -sharp operator of $V[g][h]$, and let $\mathcal{O} = \mathcal{O}^h \upharpoonright V[g][G]$. So \mathcal{O} is defined on $A \in H(\omega_2)^{V[g][G]}$ with $N \in A$. We have that $\mathcal{O} \in V[g][G]$, because the extension to $V[g][h]$ is homogeneous, and \mathcal{O}^h is definable in $V[g][h]$ from Σ . Moreover, \mathcal{O} determines the full \mathcal{O}^h in $V[g][h]$ via the process we have described. So Γ is definable in $V[g][h]$ from \mathcal{O} , and

$V[g][G] \models$ “it is forced that the strategy for P determined by \mathcal{O} fails”.

From the point of view of $V[g][G]$, the forcing in question is just $Col(\omega, \omega_1)$. But now, working in $V[g][G]$, let

$$\pi: S \rightarrow V_\eta,$$

where S is countable transitive, with everything relevant in its range. Let l be S -generic for the collapse of ω_1^S , with $l \in V[g][G]$. It is then easy to see that $\pi^{-1}(\mathcal{O})$ is contained in the $P_k^{\Sigma^G}$ -sharp operator of $V[g][G]$, and what it determines on $S[l]$ is also contained in the $P_k^{\Sigma^G}$ -sharp operator of $V[g][G]$. Since P did have a strategy in $V[g][G]$ guided by this operator, we have a contradiction, proving our claim.

Now we can invert the extension leading from A to B , getting a Σ^h premouse Q over A such that

$$P = \text{canonical re-arrangement of } Q[g] \text{ as a premouse over } B.$$

By the homogeneity of the forcing and the definability of P in $V[g][h]$, we get that inductively that all levels of Q are in V , and that all trees to which Σ^h is applied in such levels are in V . Thus Q is a Σ -premouse in V . The iteration strategy for P in $V[g][h]$ induces a strategy for Q in V , and this strategy is in V by homogeneity, and it witnesses

$$Q = P_{k+1}^\Sigma(A)^\sharp$$

in V , and that Q is ω_3 -iterable in V . We now use simultaneous reflection to extend the P_{k+1}^Σ -sharp operator to $H(\omega_3)$ in V , just as we did in the $n = 0$ case. For $A \subseteq \omega_2$, the key definition is

$$\begin{aligned} (\phi, \vec{\alpha}) \in P_{k+1}^\Sigma(A)^\sharp &\Leftrightarrow \text{for club many } \sigma \in P_{\omega_1}(H_{\omega_2}) \\ &(\phi, \vec{\alpha}_\sigma) \in P_{k+1}^{\Sigma^{\pi_\sigma}}(A_\sigma)^\sharp. \end{aligned}$$

Here we use Lemma 49 to see that for each such $\sigma \in P_{\omega_1}(\omega_2)$, $P_{k+1}^{\Sigma^{\pi\sigma}}(A_\sigma)^\sharp$ exists, and is ω_3 -iterable. Just as in the $n = 0$ case, we get that everything is decided on a club, so that the definition yields a structure with the first order properties of $P_{k+1}^\Sigma(A)^\sharp$. An argument parallel to that in the $n = 0$ case shows that this structure interprets $\dot{\Sigma}$ as Σ , and that it is ω_3 -iterable in a way that moves Σ to itself. Here one uses the corresponding properties of the $P_{k+1}^{\Sigma^{\pi\sigma}}(A_\sigma)^\sharp$, and the fact that Σ collapses into its pullbacks under Skolem hulls. This finishes the proof of (1) at $k + 1$. We leave the straightforward proofs of (2) and (3) at $k + 1$ to the reader. \square

We have finally done what we set out to do in this section.

Lemma 73. *The following holds in $V[g]$. For all transitive $A \in H(\omega_1)$ such that $N \in A$, and all $n < \omega$, $P_n^{\Sigma^g}(A)^\sharp$ exists and is ω_1 -iterable. Hence $W_{\gamma+1}^*$ holds in $V[g]$.*

Proof. This follows at once from (3) of 72, and the homogeneity of $Col(\omega, \omega_1)$. Every set in $J_\gamma(\mathbb{R})^{V[g]}$ is (boldface) projective in Σ^g . So the $P_n^{\Sigma^g}(A)^\sharp$ are the desired coarse capturing mice. \square

Lemma 74. *$I_{\gamma+1}$ holds.*

Proof. We have shown that the $P_n^{\Sigma^G}$ -sharp operator of $V[g][G]$, when restricted to $HC^{V[g]}$, is just the $P_n^{\Sigma^g}$ -sharp operator of $V[g]$. Projective-in- Σ truth is coded into these operators, so we get

$$(HC^{V[g]}, \in, \Sigma^g) \prec (HC^{V[g][G]}, \in, \Sigma^G).$$

But Σ^g codes truth at the bottom of the Levy hierarchy over $J_\gamma(\mathbb{R})^{V[g]}$, and Σ^G codes truth at the bottom of the Levy hierarchy over $J_{\gamma^G}(\mathbb{R})^{V[g][G]}$, where $\gamma^G = \beta^G$ if our gap was weak, and $\gamma^G = \beta^G + 1$ otherwise. (Truth is coded via \mathbb{R} -genericity iterations which determine self-justifying systems at the end of these gaps, as in our argument.) So we get from the line displayed above an embedding

$$\pi: J_{\gamma+1}(\mathbb{R})^{V[g]} \rightarrow J_{\gamma^G+1}(\mathbb{R})^{V[g][G]}$$

which is Σ_1 elementary. But in $V[g]$ we have an ω_1 Universally Baire prewellorder of length γ , so we can use the Foreman-Magidor argument to show $\gamma = \gamma^G$, and $\pi = \text{identity}$. \square

Repeating the relevant arguments ω times gives $W_{\gamma+\omega}^*$ in $V[g]$ as well as $I_{\gamma+\omega}$.

8. CONCLUDING REMARKS

Like many of the well-known consequences of $\text{MM}(c)$, our hypotheses follow from the Strong Reflection Principle of Todorćević, denoted $\text{SRP}(\omega_2)$, which asserts that for every projective stationary subset S of $[\omega_2]^\omega$ there is an ordinal $\delta < \omega_2$ so that $S \cap [\delta]^\omega$ contains a club in $[\delta]^\omega$. Thus, our main theorem gives $\text{AD}^{L(\mathbb{R})}$ from $\text{SRP}(\omega_2)$ as well. While this represents the best known lower bound for the strength of $\text{SRP}(\omega_2)$, and even $\text{MM}(c)$, these principles are almost certainly much stronger.¹¹ Moreover, in a precise sense, our arguments can not take us much farther. In section 9.5 of [25] Woodin defines principles $\text{SRP}^*(\omega_2)$ and $\text{WRP}_{(2)}^*(\omega_2)$ and shows that the latter is a consequence of the former if NS is saturated (see Lemma 9.93 of [25]). $\text{SRP}^*(\omega_2)$ asserts the existence of a normal fine ideal $I \subset P([\omega_2]^\omega)$ with the following two properties: (1) For every $T \in P(\omega_1) \setminus NS$ the set

$$S_T = \{\sigma \in [\omega_2]^\omega \mid \sigma \cap \omega_1 \in T\}$$

is I -positive, and (2) for every $S \subset [\omega_2]^\omega$ which satisfies $S \cap S_T \notin I$ for every $T \in P(\omega_1) \setminus NS$, there is $\gamma < \omega_2$ such that $S \cap [\gamma]^\omega$ contains a club in $[\gamma]^\omega$. $\text{WRP}_{(2)}^*(\omega_2)$ asserts the existence of a normal fine ideal I with property (1) above so that any pair $S, T \notin I$ simultaneously reflect to stationary sets in some $[\gamma]^\omega$. Woodin obtains $\text{SRP}^*(\omega_2)$ together with the saturation of NS in a \mathbb{P}_{max} extension of a determinacy model whose existence is equiconsistent with ω^2 Woodin cardinals.

Theorem 75. (Woodin; 9.102 of [25]) *The following are equiconsistent.*

- (1) $F \cap L(F, \mathbb{R})$ is an ultrafilter where F is the club filter on $[\mathbb{R}]^\omega$ and AD holds in the model $L(F, \mathbb{R})$
- (2) There exists a set of Woodin cardinals of order type ω^2

Moreover, if G is \mathbb{P}_{max} generic over $L(F, \mathbb{R})$ as in (1) then

$$L(F, \mathbb{R})[G] \models \text{SRP}^*(\omega_2) \text{ and } NS \text{ saturated} .$$

Woodin remarks in 9.98 of [25] that his proof of Theorem 10 also proves PD from NS saturated and $\text{WRP}_{(2)}^*(\omega_2)$. The same is true of our argument.

Corollary 76. *Assume NS saturated, $\text{WRP}_{(2)}^*(\omega_2)$ holds, and $2^\omega \leq \omega_2$. Then $L(\mathbb{R}) \models AD$.*

¹¹They can be obtained via forcing from a supercompact cardinal (see [4]) or from $\text{AD}_{\mathbb{R}} + \Theta$ regular (see 10.88 of [25]).

Proof. This amounts to checking that $\text{WRP}_{(2)}^*(\omega_2)$ can serve in the place of $\text{WRP}_{(2)}(\omega_2)$ in every $H(\omega_2)$ to $H(\omega_3)$ lifting argument. For example in Lemma 14 we would show that the sets S_t are measured by the filter dual to I (as opposed to the club filter). One therefore gets $2^{\omega_1} = \omega_2$ as in Lemma 8 and the rest of the proof is the same as in the proof of the Main Theorem. \square

Very likely the $2^\omega \leq \omega_2$ hypothesis can be dropped, although we haven't thought this through.¹² Thus the consistency strength of NS saturated together with $\text{WRP}_{(2)}^*(\omega_2)$ and $2^\omega \leq \omega_2$ is somewhere in the interval

$$(\omega \text{ Woodins}, \omega^2 \text{ Woodins}]$$

and we have reason to believe that the following conjecture is true.

Conjecture 77. The following are equiconsistent.

- (1) There exists a set of Woodin cardinals of order type ω^2
- (2) NS is saturated and $\text{WRP}_{(2)}^*(\omega_2)$ holds.
- (3) NS is saturated and $\text{SRP}^*(\omega_2)$ holds.

The first step is to prove that $K(\mathbb{R}) \models \text{AD}$. We leave this for another time.

REFERENCES

- [1] Feng, Q., Jech, T., *Projective Stationary Sets and Strong Reflection Principles*, Journal of the London Mathematical Society, 58 (2), 271-283, 1999
- [2] Feng, Q., Magidor, M., Woodin, H., *Universally Baire sets of reals*, MSRI Pub. 26, 1992, pp. 203-242
- [3] Foreman, M., Magidor, M., *Large cardinals and definable counterexamples to the continuum hypothesis*, Ann. Pure Appl. Logic 76 (1995), no. 1, 47-97
- [4] Foreman, M., Magidor, M., Shelah, S., *Martin's Maximum, Saturated Ideals, and Nonregular Ultrafilters*, Annals of Mathematics 127 (1998), 1-47
- [5] Hauser, K., *Generic Relativizations of Fine Structure*, Archive for Mathematical Logic 39, 2000, no. 4, 227-251
- [6] Jech, T., *Stationary Sets*, in: Handbook of Set Theory, vol. 1, M. Foreman and A. Kanamori eds., Springer 2010, pp. 93-129.
- [7] Kechris, A., Woodin, H., *Equivalence of Partition Properties and Determinacy*, Proc. Natl. Acad. Sci. USA 80 (1983), 1783-1786.
- [8] A.S. Kechris and W.H. Woodin, *Generic codes for uncountable ordinals*, in: Games, scales, and Suslin cardinals, The Cabal Seminar, volume I, Kechris et. al. eds., Cambridge University Press, 2008, pp. 379-397.
- [9] Martin, D., Steel, J., *A proof of projective determinacy*, J. Amer. Math. Soc. 2 (1989), no. 1, 71-125.
- [10] Mitchell, W., Steel, J., *Fine Structure and Iteration Trees*, Lecture Notes in Logic, Vol. 3, Springer Verlag, 1991

¹²The point is that we use $2^{\omega_1} = \omega_2$ at various points and the hypotheses NS saturated and $\text{WRP}_{(2)}^*(\omega_2)$ give all of Lemma 8 except Todorćević's bound on 2^ω .

- [11] Moschovakis, Y., *Descriptive Set Theory*, Studies in Logic and the Foundations of Mathematics, 100. North-Holland Publishing Co., Amsterdam-New York, 1980
- [12] Neeman, I., *Optimal proofs of determinacy*, Bull. Symbolic Logic 1 (1995), no. 3, 327–339
- [13] Neeman, I., *Itay Optimal proofs of determinacy II*, J. Math. Log. 2 (2002), no. 2, 227–258
- [14] R.D. Schindler and J.R. Steel, *The self-iterability of $L[E]$* , Journal of Symbolic Logic, vol. 74 (2009), pp. 751–779.
- [15] Schindler, R., Steel, J., *The Strength of AD*, unpublished manuscript.
- [16] Schindler, R., Steel, J., *The Core Model Induction*, unpublished manuscript.
- [17] Steel, J., *Scales on Σ_1^1 sets*, Cabal seminar 79–81, 72–76, Lecture Notes in Math., 1019, Springer, Berlin, 1983.
- [18] Steel, J., *The core model iterability problem*, Lecture Notes in Logic, 8. Springer-Verlag, Berlin, 1996.
- [19] Steel, J., *A Theorem of Woodin on Mouse Sets*, to appear in: Ordinal definability and recursion theory: The Cabal seminar, volume III, Kechris et. al. eds., Cambridge University Press.
- [20] J.R. Steel, *An outline of inner model theory*, in Handbook of Set Theory, vol. 3, M. Foreman and A. Kanamori eds., Springer 2010, pp. 1595–1684.
- [21] J.R. Steel, *Derived models associated to mice*, in Computational Prospects of Infinity, Chong et. al. eds., Part 1, Tutorials, pp. 105–193, Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap., 14, World Sci. Publ., Hackensack, NJ, 2008.
- [22] Steel, J., *PFA implies AD in $L(\mathbb{R})$* , J. Symbolic Logic 70 (2005), no. 4, 1255–1296.
- [23] Steel, J., *Scales in $L(\mathbb{R})$* , Cabal seminar 79–81, 107–156, Lecture Notes in Math., 1019, Springer, Berlin, 1983
- [24] J.R. Steel, *Scales in $K(\mathbb{R})$* , in: Games, scales, and Suslin cardinals, The Cabal seminar, volume I, Kechris et. al. eds., Cambridge University Press, 2008, pp. 176–208.
- [25] Woodin, H., *The axiom of determinacy, forcing axioms, and the nonstationary ideal*, de Gruyter Series in Logic and its Applications, 1. Walter de Gruyter & Co., Berlin, 1999
- [26] Zoble, S., *Stationary Reflection and the Determinacy of Inductive Games*, Doctoral Dissertation, University of California at Berkeley, 2000
- [27] Zoble, S., *Stationary reflection and the universal Baire property*, Fund. Math. 191 (2006), no. 1, 45–56

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT BERKELEY
E-mail address: `steel@math.berkeley.edu`

DEPARTMENT OF MATHEMATICS, WESLEYAN UNIVERSITY
E-mail address: `azoble@wesleyan.edu`