

## Outline

Holomorphic functions

- Cauchy-Goursat theorem
- Morera's theorem
- Holomorphic vs. real analytic functions

Conformal maps

- Riemann mapping theorem
- Important examples: disc to upper half-plane and vice versa, upper half-plane to a sector, Schwarz-Christoffel integral
- Automorphisms of the disk; Schwarz lemma, Blaschke factors

The extended/compactified complex plane

Classification of singularities

- Removable singularities
- Poles
- Essential singularities; Casorati-Weierstrass

Local and global behavior of holomorphic functions

- Open mapping theorem
- Maximum modulus principle
- Liouville's theorem

## Problems

**5.2.16 – Spring 1978 12** Prove that the uniform limit of a sequence of complex analytic functions is complex analytic. Is the analogous theorem true for real analytic functions?

**5.4.13 – Fall 1990 13** Suppose that  $f$  is analytic on the open upper half-plane and satisfies  $|f(z)| \leq 1$  for all  $z$ ,  $f(i) = 0$ . How large can  $|f(2i)|$  be under these conditions?

**5.4.7 – Spring 2003 7B** Let  $f(z)$  be a function that is analytic in the unit disk  $\mathbb{D} = \{|z| < 1\}$ . Suppose that  $|f(z)| \leq 1$  in  $\mathbb{D}$ . Prove that if  $f(z)$  has at least two fixed points  $z_1$  and  $z_2$ , then  $f(z) = z$  for all  $z \in \mathbb{D}$ .

**5.3.8 – Spring 1995 18** Prove that there is no one-to-one conformal map of the punctured disk  $G = \{z \in \mathbb{C} : 0 < |z| < 1\}$  onto the annulus  $A = \{z \in \mathbb{C} : 1 < |z| < 2\}$ .

**5.5.3 – Fall 1999 4** Let the rational function  $f$  in the complex plane have no poles for

$\text{Im}(z) \geq 0$ . Prove that

$$\sup\{|f(z)| : \text{Im}(z) \geq 0\} = \sup\{|f(z)| : \text{Im}(z) = 0\}.$$

**5.5.8 – Spring 1997 4** Let  $f$  and  $g$  be two entire functions such that, for all  $z \in \mathbb{C}$ ,  $\text{Re}(f(z)) \leq k\text{Re}(g(z))$  for some real constant  $k$  (independent of  $z$ ). Show that there are constants  $a, b$  such that

$$f(z) = ag(z) + b.$$

**5.6.29 – Spring 1987 15** Prove or disprove: If the function  $f$  is analytic in the entire complex plane, and if  $f$  maps every unbounded sequence to an unbounded sequence, then  $f$  is a polynomial.