

NETWORK PARAMETERIZATIONS FOR THE GRASSMANNIAN

KELLI TALASKA AND LAUREN WILLIAMS

ABSTRACT. Deodhar introduced his decomposition of partial flag varieties as a tool for understanding Kazhdan-Lusztig polynomials. The Deodhar decomposition of the Grassmannian is also useful in the context of soliton solutions to the KP equation, as shown by Kodama and the second author. Deodhar components \mathcal{R}_D of the Grassmannian are in bijection with certain tableaux D called *Go-diagrams*, and each component is isomorphic to $(\mathbb{K}^*)^a \times (\mathbb{K})^b$ for some non-negative integers a and b .

Our main result is an explicit parameterization of each Deodhar component in the Grassmannian in terms of networks. More specifically, from a Go-diagram D we construct a weighted network N_D and its *weight matrix* W_D , whose entries enumerate directed paths in N_D . By letting the weights in the network vary over \mathbb{K} or \mathbb{K}^* as appropriate, one gets a parameterization of the Deodhar component \mathcal{R}_D . One application of such a parameterization is that one may immediately determine which Plücker coordinates are vanishing and nonvanishing, by using the Lindstrom-Gessel-Viennot Lemma. We also give a (minimal) characterization of each Deodhar component in terms of Plücker coordinates. A main tool for us is the work of Marsh and Rietsch [8] on Deodhar components in the flag variety.

CONTENTS

1. Introduction	1
2. Background on the Grassmannian	3
3. The main result: network parameterizations from Go-diagrams	6
4. The Deodhar decomposition of the Grassmannian	10
5. Formulas for entries of the MR-matrices	13
6. Proof of the main result	20
7. A characterization of Deodhar components in terms of Plücker coordinates	24
References	27

1. INTRODUCTION

There is a remarkable subset of the real Grassmannian $Gr_{k,n}(\mathbb{R})$ called its *totally non-negative part* $(Gr_{k,n})_{\geq 0}$ [7, 9], which may be defined as the subset of the real Grassmannian where all Plücker coordinates have the same sign. Postnikov showed that $(Gr_{k,n})_{\geq 0}$ has a decomposition into *positroid cells*, which are indexed by certain tableaux called \mathbb{J} -diagrams. He also gave explicit parameterizations of each cell. In particular, he showed that from each \mathbb{J} -diagram one can produce a planar *network*, and that one can write down a parameterization of the corresponding cell using the *weight matrix* of that network. This parameterization shows that the cell is isomorphic to $\mathbb{R}_{>0}^d$ for some d . Such a parameterization is convenient, because for example, one may read off formulas for Plücker coordinates from non-intersecting paths in the network, using the Lindstrom-Gessel-Viennot Lemma.

Date: February 17, 2013.

The first author was partially supported by NSF Grant DMS-1004532. The second author was partially supported by an NSF CAREER award and an Alfred Sloan Fellowship.

A natural question is whether these network parameterizations for positroid cells can be extended from $(Gr_{k,n})_{\geq 0}$ to the entire real Grassmannian $Gr_{k,n}(\mathbb{R})$. In this paper we give an affirmative answer to this question, by replacing the positroid cell decomposition with the Deodhar decomposition of the Grassmannian $Gr_{k,n}(\mathbb{K})$ (here \mathbb{K} is an arbitrary field).

The components of the Deodhar decomposition are not in general cells, but nevertheless have a simple topology: by [2, 3], each one is isomorphic to $(\mathbb{K}^*)^a \times (\mathbb{K})^b$. The relation of the Deodhar decomposition of $Gr_{k,n}(\mathbb{R})$ to Postnikov's cell decomposition of $(Gr_{k,n})_{\geq 0}$ is as follows: the intersection of a Deodhar component $\mathcal{R}_D \cong (\mathbb{R}^*)^a \times (\mathbb{R})^b$ with $(Gr_{k,n})_{\geq 0}$ is precisely one positroid cell isomorphic to $(\mathbb{R}_{>0})^a$ if $b = 0$, and is empty otherwise. In particular, when one intersects the Deodhar decomposition with $(Gr_{k,n})_{\geq 0}$, one obtains the positroid cell decomposition of $(Gr_{k,n})_{\geq 0}$. There is a related *positroid stratification* of the real Grassmannian, and each positroid stratum is a union of Deodhar components.

As for the combinatorics, components of the Deodhar decomposition are indexed by *distinguished subexpressions* [2, 3], or equivalently, by certain tableaux called *Go-diagrams* [6], which generalize \mathcal{J} -diagrams. In this paper we associate a network to each Go-diagram, and write down a parameterization of the corresponding Deodhar component using the weight matrix of that network. Our construction generalizes Postnikov's, but our networks are no longer planar in general.

Our main results can be summed up as follows. See Theorems 3.16 and 7.8 and the constructions preceding them for complete details.

Theorem. Let \mathbb{K} be an arbitrary field.

- Every point in $Gr_{k,n}(\mathbb{K})$ can be realized as the weight matrix of a unique network associated to a Go-diagram, and we can explicitly construct the corresponding network. The networks corresponding to points in the same Deodhar component have the same underlying graph, but different weights.
- Every Deodhar component may be characterized by the vanishing and nonvanishing of certain Plücker coordinates. Using this characterization, we can also explicitly construct the network associated to a point given either by a matrix representative or by a list of Plücker coordinates.

To illustrate the main results, we provide a small example here. More complicated examples may be seen throughout the rest of the paper.




FIGURE 1. The diagrams and networks associated to \mathcal{R}_{D_1} and \mathcal{R}_{D_2} in Example 1.1.

Example 1.1. Consider the Grassmannian $Gr_{2,4}$. The large Schubert cell in this Grassmannian can be characterized as

$$\Omega_\lambda = \{A \in Gr_{2,4} \mid \Delta_{1,2}(A) \neq 0\},$$

where Δ_J denotes the Plücker coordinate corresponding to the column set J in a matrix representative of a point in $Gr_{2,4}$. This Schubert cell contains multiple positroid strata, including $S_{\mathcal{I}}$, where \mathcal{I} is the Grassmann necklace $\mathcal{I} = (12, 23, 34, 14)$. This positroid stratum can also be characterized by the vanishing of certain Plücker coordinates:

$$S_{\mathcal{I}} = \{A \in Gr_{2,4} \mid \Delta_{1,2}(A) \neq 0, \Delta_{2,3}(A) \neq 0, \Delta_{3,4}(A) \neq 0, \Delta_{1,4}(A) \neq 0\}.$$

Figure 1 shows two Go-diagrams D_1 and D_2 and their associated networks. Note that the network on the right is not planar. The weight matrices associated to these diagrams are

$$\begin{pmatrix} 1 & 0 & -a_3 & -(a_3a_4 + a_3a_2) \\ 0 & 1 & a_1 & a_1a_2 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 & -a_3 & -a_3c_4 \\ 0 & 1 & 0 & a_2 \end{pmatrix}.$$

The positroid stratum $S_{\mathcal{I}}$ is the disjoint union of the two corresponding Deodhar components \mathcal{R}_{D_1} and \mathcal{R}_{D_2} , which can be characterized in terms of vanishing and nonvanishing of minors as:

$$\mathcal{R}_{D_1} = \{A \in S_{\mathcal{I}} \mid \Delta_{1,3} \neq 0\} \text{ and } \mathcal{R}_{D_2} = \{A \in S_{\mathcal{I}} \mid \Delta_{1,3} = 0\}.$$

Note that if one lets the a_i 's range over \mathbb{K}^* and lets c_4 range over \mathbb{K} , then we see that $\mathcal{R}_{D_1} \cong (\mathbb{K}^*)^4$ and $\mathcal{R}_{D_2} \cong (\mathbb{K}^*)^2 \times \mathbb{K}$.

There are several applications of our construction. First, as a special case of our theorem, one may parameterize all $k \times n$ matrices using networks. Second, by applying the Lindstrom-Gessel-Viennot Lemma to a given network, one may write down explicit formulas for Plücker coordinates in terms of collections of non-intersecting paths in the network. Third, building upon work of [6], we obtain (minimal) descriptions of Deodhar components in the Grassmannian, in terms of vanishing and nonvanishing of Plücker coordinates. It follows that each Deodhar component is a union of matroid strata.

Although less well known than the Schubert decomposition and matroid stratification, the Deodhar decomposition is very interesting in its own right. Deodhar's original motivation for introducing his decomposition was the desire to understand Kazhdan-Lusztig polynomials. In the flag variety, one may intersect two opposite Schubert cells, obtaining a Richardson variety, which Deodhar showed is a union of Deodhar components. Each Richardson variety $\mathcal{R}_{v,w}(q)$ may be defined over a finite field $\mathbb{K} = \mathbb{F}_q$, and in this case, the number of points determines the R -polynomials $R_{v,w}(q) = \#(\mathcal{R}_{v,w}(\mathbb{F}_q))$, introduced by Kazhdan and Lusztig [4] to give a recursive formula for the Kazhdan-Lusztig polynomials. Since each Deodhar component is isomorphic to $(\mathbb{F}_q^*)^a \times (\mathbb{F}_q)^b$ for some a and b , if one understands the decomposition of a Richardson variety into Deodhar components, then in principle one may compute the R -polynomials and hence Kazhdan-Lusztig polynomials.

Another reason for our interest in the Deodhar decomposition is its relation to soliton solutions of the KP equation. It is well-known that from each point A in the real Grassmannian, one may construct a soliton solution $u_A(x, y, t)$ of the KP equation. It was shown in recent work of Kodama and the second author [6] that when the time variable t tends to $-\infty$, the combinatorics of the solution $u_A(x, y, t)$ depends precisely on which Deodhar component A lies in.

The outline of this paper is as follows. In Section 2, we give some background on the Grassmannian and its decompositions, including the Schubert decomposition, the positroid stratification, and the matroid stratification. In Section 3, we present our main construction: we explain how to construct a network from each diagram, then use that network to write down a parameterization of a subset of the Grassmannian that we call a network component. In Section 4 we define Deodhar's decomposition of the flag variety, and its projection to the Grassmannian. We also describe parameterizations of Deodhar components in the flag variety which are due to Marsh and Rietsch [8]. In Sections 5 and 6 we prove that after a rational transformation of variables, our network parameterizations coincide with the projections of the Marsh-Rietsch parameterizations. Finally in Section 7 we give a characterization of Deodhar components in terms of the vanishing and nonvanishing of certain Plücker coordinates.

ACKNOWLEDGEMENTS: We are grateful to Sara Billey for numerous helpful comments on the first version of this paper. L.W. is also grateful to Yuji Kodama for their joint work on soliton solutions of the KP equation, which provided motivation for this project.

2. BACKGROUND ON THE GRASSMANNIAN

The *Grassmannian* $Gr_{k,n}$ is the space of all k -dimensional subspaces of an n -dimensional vector space \mathbb{K}^n . In this paper we will usually let \mathbb{K} be an arbitrary field, though we will often think of it as

\mathbb{R} or \mathbb{C} . An element of $Gr_{k,n}$ can be viewed as a full-rank $k \times n$ matrix modulo left multiplication by nonsingular $k \times k$ matrices. In other words, two $k \times n$ matrices represent the same point in $Gr_{k,n}$ if and only if they can be obtained from each other by row operations. Let $\binom{[n]}{k}$ be the set of all k -element subsets of $[n] := \{1, \dots, n\}$. For $I \in \binom{[n]}{k}$, let $\Delta_I(A)$ be the *Plücker coordinate*, that is, the maximal minor of the $k \times n$ matrix A located in the column set I . The map $A \mapsto (\Delta_I(A))$, where I ranges over $\binom{[n]}{k}$, induces the *Plücker embedding* $Gr_{k,n} \hookrightarrow \mathbb{K}\mathbb{P}^{\binom{[n]}{k}-1}$ into projective space.

We now describe several useful decompositions of the Grassmannian: the Schubert decomposition, the positroid stratification, and the matroid stratification. Note that the matroid stratification refines the positroid stratification, which refines the Schubert decomposition. The main subject of this paper is the *Deodhar decomposition* of the Grassmannian, which refines the positroid stratification, and is refined by the matroid stratification (as we prove in Corollary 7.9).

2.1. The Schubert decomposition of $Gr_{k,n}$. Throughout this paper, we identify partitions with their Young diagrams. Recall that the partitions λ contained in a $k \times (n-k)$ rectangle are in bijection with k -element subset $I \subset [n]$. The boundary of the Young diagram of such a partition λ forms a lattice path from the upper-right corner to the lower-left corner of the rectangle. Let us label the n steps in this path by the numbers $1, \dots, n$, and define $I = I(\lambda)$ as the set of labels on the k vertical steps in the path. Conversely, we let $\lambda(I)$ denote the partition corresponding to the subset I .

Definition 2.1. For each partition λ contained in a $k \times (n-k)$ rectangle, we define the *Schubert cell*

$$\Omega_\lambda = \{A \in Gr_{k,n} \mid I(\lambda) \text{ is the lexicographically minimal subset such that } \Delta_{I(\lambda)}(A) \neq 0\}.$$

As λ ranges over the partitions contained in a $k \times (n-k)$ rectangle, this gives the *Schubert decomposition* of the Grassmannian $Gr_{k,n}$, i.e.

$$Gr_{k,n} = \bigsqcup_{\lambda \subset (n-k)^k} \Omega_\lambda.$$

We now define the *shifted linear order* $<_i$ (for $i \in [n]$) to be the total order on $[n]$ defined by

$$i <_i i+1 <_i i+2 <_i \dots <_i n <_i 1 <_i \dots <_i i-1.$$

One can then define *cyclically shifted Schubert cells* as follows.

Definition 2.2. For each partition λ contained in a $k \times (n-k)$ rectangle, and each $i \in [n]$, we define the *cyclically shifted Schubert cell*

$$\Omega_\lambda^i = \{A \in Gr_{k,n} \mid I(\lambda) \text{ is the lexicographically minimal subset with respect to } <_i \text{ such that } \Delta_{I(\lambda)}(A) \neq 0\}.$$

2.2. The positroid stratification of $Gr_{k,n}$. The *positroid stratification* of the Grassmannian $Gr_{k,n}$ is obtained by taking the simultaneous refinement of the n Schubert decompositions with respect to the n shifted linear orders $<_i$. This stratification was first considered by Postnikov [9], who showed that the strata are conveniently described in terms of *Grassmann necklaces*, as well as *decorated permutations* and *J-diagrams*. Postnikov coined the terminology *positroid* because the intersection of the positroid stratification of the real Grassmannian with the *totally non-negative part of the Grassmannian* $(Gr_{k,n})_{\geq 0}$ gives a cell decomposition of $(Gr_{k,n})_{\geq 0}$ (whose cells are called *positroid cells*).

Definition 2.3. [9, Definition 16.1] A *Grassmann necklace* is a sequence $\mathcal{I} = (I_1, \dots, I_n)$ of subsets $I_r \subset [n]$ such that, for $i \in [n]$, if $i \in I_i$ then $I_{i+1} = (I_i \setminus \{i\}) \cup \{j\}$, for some $j \in [n]$ (j may coincide with i); and if $i \notin I_i$ then $I_{i+1} = I_i$. (Here indices i are taken modulo n .) In particular, we have $|I_1| = \dots = |I_n|$, which is equal to some $k \in [n]$. We then say that \mathcal{I} is a Grassmann necklace of *type* (k, n) .

Example 2.4. $\mathcal{I} = (1345, 3456, 3456, 4567, 4567, 1467, 1478, 1348)$ is an example of a Grassmann necklace of type $(4, 8)$.

Lemma 2.5. [9, Lemma 16.3] Given $A \in Gr_{k,n}$, let $\mathcal{I}(A) = (I_1, \dots, I_n)$ be the sequence of subsets in $[n]$ such that, for $i \in [n]$, I_i is the lexicographically minimal subset of $\binom{[n]}{k}$ with respect to the shifted linear order $<_i$ such that $\Delta_{I_i}(A) \neq 0$. Then $\mathcal{I}(A)$ is a Grassmann necklace of type (k, n) .

The *positroid stratification* of $Gr_{k,n}$ is defined as follows.

Definition 2.6. Let $\mathcal{I} = (I_1, \dots, I_n)$ be a Grassmann necklace of type (k, n) . The *positroid stratum* $S_{\mathcal{I}}$ is defined to be

$$S_{\mathcal{I}} = \{A \in Gr_{k,n} \mid \mathcal{I}(A) = \mathcal{I}\}.$$

Equivalently, each positroid stratum is an intersection of n cyclically shifted Schubert cells, that is,

$$S_{\mathcal{I}} = \bigcap_{i=1}^n \Omega_{\lambda(I_i)}^i.$$

Grassmann necklaces are in bijection with tableaux called *J-diagrams*.

Definition 2.7. [9, Definition 6.1] Fix k, n . A *J-diagram* $(\lambda, D)_{k,n}$ of type (k, n) is a partition λ contained in a $k \times (n - k)$ rectangle together with a filling $D : \lambda \rightarrow \{0, +\}$ of its boxes which has the *J-property*: there is no 0 which has a + above it and a + to its left.¹ (Here, “above” means above and in the same column, and “to its left” means to the left and in the same row.)

In Figure 2 we give an example of a J-diagram.

+	0	+	+
+	0	+	
0	0	+	
+	0		

FIGURE 2. A Le-diagram $L = (\lambda, D)_{k,n}$.

2.3. The matroid stratification of $Gr_{k,n}$.

Definition 2.8. A *matroid* of rank k on the set $[n]$ is a nonempty collection $\mathcal{M} \subset \binom{[n]}{k}$ of k -element subsets in $[n]$, called *bases* of \mathcal{M} , that satisfies the *exchange axiom*:

For any $I, J \in \mathcal{M}$ and $i \in I$ there exists $j \in J$ such that $(I \setminus \{i\}) \cup \{j\} \in \mathcal{M}$.

Given an element $A \in Gr_{k,n}$, there is an associated matroid \mathcal{M}_A whose bases are the k -subsets $I \subset [n]$ such that $\Delta_I(A) \neq 0$.

Definition 2.9. Let $\mathcal{M} \subset \binom{[n]}{k}$ be a matroid. The *matroid stratum* $S_{\mathcal{M}}$ is defined to be

$$S_{\mathcal{M}} = \{A \in Gr_{k,n} \mid \Delta_I(A) \neq 0 \text{ if and only if } I \in \mathcal{M}\}.$$

This gives a stratification of $Gr_{k,n}$ called the *matroid stratification*, or *Gelfand-Serganova stratification*.

Remark 2.10. Clearly the matroid stratification refines the positroid stratification, which in turn refines the Schubert decomposition.

¹This forbidden pattern is in the shape of a backwards L , and hence is denoted J and pronounced “Le.”

3. THE MAIN RESULT: NETWORK PARAMETERIZATIONS FROM GO-DIAGRAMS

In this section we define certain tableaux called *Go-diagrams*, then explain how to parameterize the Grassmannian using networks associated to Go-diagrams. First we will define more general tableaux called *diagrams*.

3.1. Diagrams and networks.

Definition 3.1. Let λ be a partition contained in a $k \times (n - k)$ rectangle. A *diagram* in λ is an arbitrary filling of the boxes of λ with pluses $+$, black stones \bullet , and white stones \circ .

To each diagram D we associate a network N_D as follows.

Definition 3.2. Let λ be a partition with ℓ boxes contained in a $k \times (n - k)$ rectangle, and let D be a diagram in λ . Label the boxes of λ from 1 to ℓ , starting from the rightmost box in the bottom row, then reading right to left across the bottom row, then right to left across the row above that, etc. The (*weighted*) *network* N_D associated to D is a directed graph obtained as follows:

- Associate an *internal vertex* to each $+$ and each \bullet ;
- After labeling the southeast border of the Young diagram with the numbers $1, 2, \dots, n$ (from northeast to southwest), associate a *boundary vertex* to each number;
- From each internal vertex, draw an edge right to the nearest $+$ -vertex or boundary vertex;
- From each internal vertex, draw an edge down to the nearest $+$ -vertex or boundary vertex;
- Direct all edges left and down. After doing so, k of the boundary vertices become *sources* and the remaining $n - k$ boundary vertices become *sinks*.
- If e is a horizontal edge whose left vertex is a $+$ -vertex (respectively a \bullet -vertex) in box b , assign e the weight a_b (respectively c_b). We think of a_b and c_b as indeterminates, but later they will be elements of \mathbb{K}^* and \mathbb{K} respectively.
- If e is a vertical edge, assign e the weight 1.

Note that in general such a directed graph is not planar, as two edges may cross over each other without meeting at a vertex. See Figure 3 for an example of a diagram and its associated network.




FIGURE 3. An example of a diagram and its associated network.

We now explain how to associate a *weight matrix* to such a network.

Definition 3.3. Let N_D be a network as in Definition 3.2. Let $I = \{i_1 < i_2 < \dots < i_k\} \subset [n]$ denote the sources. If P is a directed path in the network, let $w(P)$ denote the product of all weights along P .

If P is the empty path which starts and ends at the same boundary vertex, we let $w(P) = 1$. If r is a source and s is any boundary vertex, define

$$W_{rs} = \pm \sum_P w(P),$$

where the sum is over all paths P from r to s . The sign is chosen (uniquely) so that

$$\Delta_{I \setminus \{r\} \cup \{s\}}(W_D) = \sum_P w(P), \text{ where}$$

$$W_D = (W_{rs})$$

is the $k \times (n - k)$ weight matrix. We make the convention that the rows of W_D are indexed by the sources i_1, \dots, i_k from top to bottom, and its columns are indexed by $1, 2, \dots, n$ from left to right. An equivalent way to define the sign of W_{rs} is to let $q = |\{s + 1, s + 2, \dots, t - 1\} \cap I|$, i.e. the number of sources which are strictly between s and t . Then the sign of W_{rs} is $(-1)^q$.

Example 3.4. The weight matrix associated to the network in Figure 3 is

$$\begin{pmatrix} 1 & a_9 & 0 & 0 & a_9 a_{10} & 0 & -a_9 a_{10}(a_{11} + c_7) & -a_9 a_{10}(a_{11} a_{12} + a_{11} c_5 + a_8 + c_7 c_5) \\ 0 & 0 & 1 & 0 & -a_6 & 0 & a_6 c_7 & a_6 a_8 + a_6 c_7 c_5 \\ 0 & 0 & 0 & 1 & 0 & 0 & -a_4 & -a_4 c_5 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & a_2 \end{pmatrix}$$

3.2. Distinguished expressions. We now review the notion of distinguished subexpressions, as in [2] and [8]. This definition will be essential for defining Go-diagrams. We assume the reader is familiar with the (strong) Bruhat order $<$ on $W = \mathfrak{S}_n$, and the basics of reduced expressions, as in [1].

Let $\mathbf{w} := s_{i_1} \dots s_{i_m}$ be a reduced expression for $w \in W$. A *subexpression* \mathbf{v} of \mathbf{w} is a word obtained from the reduced expression \mathbf{w} by replacing some of the factors with 1. For example, consider a reduced expression in \mathfrak{S}_4 , say $s_3 s_2 s_1 s_3 s_2 s_3$. Then $s_3 s_2 1 s_3 s_2 1$ is a subexpression of $s_3 s_2 s_1 s_3 s_2 s_3$. Given a subexpression \mathbf{v} , we set $v_{(k)}$ to be the product of the leftmost k factors of \mathbf{v} , if $k \geq 1$, and $v_{(0)} = 1$.

Definition 3.5. [8, 2] Given a subexpression \mathbf{v} of a reduced expression $\mathbf{w} = s_{i_1} s_{i_2} \dots s_{i_m}$, we define

$$J_{\mathbf{v}}^{\circ} := \{k \in \{1, \dots, m\} \mid v_{(k-1)} < v_{(k)}\},$$

$$J_{\mathbf{v}}^{+} := \{k \in \{1, \dots, m\} \mid v_{(k-1)} = v_{(k)}\},$$

$$J_{\mathbf{v}}^{\bullet} := \{k \in \{1, \dots, m\} \mid v_{(k-1)} > v_{(k)}\}.$$

The expression \mathbf{v} is called *non-decreasing* if $v_{(j-1)} \leq v_{(j)}$ for all $j = 1, \dots, m$, e.g. $J_{\mathbf{v}}^{\bullet} = \emptyset$.

Definition 3.6 (Distinguished subexpressions). [2, Definition 2.3] A subexpression \mathbf{v} of \mathbf{w} is called *distinguished* if we have

$$(3.1) \quad v_{(j)} \leq v_{(j-1)} s_{i_j} \quad \text{for all } j \in \{1, \dots, m\}.$$

In other words, if right multiplication by s_{i_j} decreases the length of $v_{(j-1)}$, then in a distinguished subexpression we must have $v_{(j)} = v_{(j-1)} s_{i_j}$.

We write $\mathbf{v} < \mathbf{w}$ if \mathbf{v} is a distinguished subexpression of \mathbf{w} .

Definition 3.7 (Positive distinguished subexpressions). We call a subexpression \mathbf{v} of \mathbf{w} a *positive distinguished subexpression* (or a PDS for short) if

$$(3.2) \quad v_{(j-1)} < v_{(j-1)} s_{i_j} \quad \text{for all } j \in \{1, \dots, m\}.$$

In other words, it is distinguished and non-decreasing.

Lemma 3.8. [8] Given $v \leq w$ and a reduced expression \mathbf{w} for w , there is a unique PDS \mathbf{v}_+ for v in \mathbf{w} .

s_5	s_4	s_3	s_2	s_1
s_6	s_5	s_4	s_3	s_2
s_7	s_6	s_5	s_4	s_3

15	14	13	12	11
10	9	8	7	6
5	4	3	2	1

15	12	9	6	3
14	11	8	5	2
13	10	7	4	1

FIGURE 4. The labeling of a the boxes of a partition by simple generators s_i , and two reading orders.

3.3. Go-diagrams. In this section we explain how to index distinguished subexpressions by certain tableaux called *Go-diagrams*, which were introduced in [6]. Go-diagrams are fillings of Young diagrams by pluses $+$, *black stones* \bullet , and *white stones* \circ .²

Fix k and n . Let $W_k = \langle s_1, s_2, \dots, \hat{s}_{n-k}, \dots, s_{n-1} \rangle$ be a parabolic subgroup of $W = \mathfrak{S}_n$. Let W^k denote the set of minimal-length coset representatives of W/W_k . Recall that a *descent* of a permutation π is a position j such that $\pi(j) > \pi(j+1)$. Then W^k is the subset of permutations of \mathfrak{S}_n which have at most one descent; and that descent must be in position $n-k$.

It is well-known that elements w of W^k can be identified with partitions λ_w contained in a $k \times (n-k)$ rectangle: if $w = (w_1, \dots, w_n) \in W^k$ then $\{w_{n-k+1}, w_{n-k+2}, \dots, w_n\}$ is a subset of size k , which gives rise to a partition, as described at the beginning of Section 2.1. We refer to this partition as λ_w .

Moreover, it follows from [11] and [10] that the reduced expressions of $w \in W^k$ correspond to certain *reading orders* of the boxes of the partition λ_w . Specifically, let Q^k be the poset whose elements are the boxes of a $k \times (n-k)$ rectangle; if b_1 and b_2 are two adjacent boxes such that b_2 is immediately to the left or immediately above b_1 , we have a cover relation $b_1 < b_2$ in Q^k . The partial order on Q^k is the transitive closure of $<$. Now label the boxes of the rectangle with simple generators s_i as in the figure below. If b is a box of the rectangle, then let s_b denote its label by a simple generator. Let $w_0^k \in W^k$ denote the longest element in W^k . Then the set of reduced expressions of w_0^k can be obtained by choosing a linear extension of Q^k and writing down the corresponding word in the s_i 's. We call such a linear extension a *reading order*; two linear extensions are shown in the figure below. Additionally, given a partition λ_w contained in the $k \times (n-k)$ rectangle (chosen so that the upper-left corner of its Young diagram is aligned with the upper-left corner of the rectangle), and a linear extension of the sub-poset of Q^k comprised of the boxes of λ , the corresponding word in s_i 's is a reduced expression of the permutation $w \in W^k$. Moreover, all reduced expressions of w can be obtained by varying the linear extension.

Definition 3.9. [6, Section 4] Fix k and n . Let $w \in W^k$, let \mathbf{w} be a reduced expression for w , and let \mathbf{v} be a distinguished subexpression of \mathbf{w} . Then w and \mathbf{w} determine a partition λ_w contained in a $k \times (n-k)$ rectangle together with a reading order of its boxes. The *Go-diagram* associated to \mathbf{v} and \mathbf{w} is a filling of λ_w with pluses and black and white stones, such that: for each $k \in J_{\mathbf{v}}^{\circ}$ we place a white stone in the corresponding box; for each $k \in J_{\mathbf{v}}^{\bullet}$ we place a black stone in the corresponding box of λ_w ; and for each $k \in J_{\mathbf{v}}^{+}$ we place a plus in the corresponding box of λ_w .

Remark 3.10. By [6, Section 4], whether or not a filling of a partition λ_w is a Go-diagram does not depend on the choice of reading order of the boxes of λ_w .

Definition 3.11. We define the *standard reading order* of the boxes of a partition to be the reading order which starts at the rightmost box in the bottom row, then reads right to left across the bottom

²In KW2, we used a slightly different convention and used blank boxes in place of $+$'s.


row, then right to left across the row above that, then right to left across the row above that, etc. This reading order is illustrated at the right of the figure below.

By default, we will use the standard reading order in this paper.

Example 3.12. Let $k = 3$ and $n = 7$, and let $\lambda = (4, 3, 1)$. The standard reading order is shown at the right of the figure below.




Then the following diagrams are Go-diagrams of shape λ .



They correspond to the expressions $s_6s_3s_4s_5s_1s_2s_3s_4$, $s_61s_411s_2s_31$, and $1s_3s_41s_111s_4$. The first and second are positive distinguished subexpressions (PDS's), and the third one is a distinguished subexpression (but not a PDS).

Note that the following diagram of shape λ is not a Go-diagram. It corresponds to the word $11s_41s_1s_211$, which is not distinguished.



Remark 3.13. The Go-diagrams associated to PDS's are in bijection with \mathbb{J} -diagrams, see [6, Section 4]. Note that the Go-diagram associated to a PDS contains only pluses and white stones. This is precisely a \mathbb{J} -diagram.

If we choose a reading order of λ_w , then we will also associate to a Go-diagram of shape λ_w a *labeled Go-diagram*, as defined below. Equivalently, a labeled Go-diagram is associated to a pair (\mathbf{v}, \mathbf{w}) .

Definition 3.14. [6, Definition 4.15] Given a reading order of λ_w and a Go-diagram of shape λ_w , we obtain a *labeled Go-diagram* by replacing each \circ with a 1, replacing each box b containing a \bullet with a -1 and an m_i , and replacing each box b containing a $+$ by a p_i , where the subscript i corresponds to the label of b inherited from the reading order.

The labeled Go-diagrams corresponding to the examples above using the standard reading order are:

1	1	1	1
1	1	1	
1			

p_8	1	1	p_5
p_4	1	p_2	
1			

$\frac{-1}{m_8}$	p_7	p_6	1
p_4	1	1	
p_1			

3.4. The main result. To state the main result, we now consider Go-diagrams (not arbitrary diagrams), the corresponding networks (*Go-networks*), and the corresponding weight matrices.

Definition 3.15. Let D be a Go-diagram contained in a $k \times (n - k)$ rectangle. We define a subset \mathcal{R}_D of the Grassmannian $Gr_{k,n}$ by letting each variable a_i of the weight matrix (Definition 3.3) range over all nonzero elements \mathbb{K}^* , and letting each variable c_i of the weight matrix range over all elements \mathbb{K} . We call \mathcal{R}_D the *network component associated to D* .

Theorem 3.16. Let D be a Go-diagram contained in a $k \times (n - k)$ rectangle. Suppose that D has t pluses and u black stones. Then \mathcal{R}_D is isomorphic to $(\mathbb{K}^*)^t \times \mathbb{K}^u$. Furthermore, $Gr_{k,n}$ is the disjoint union of the network components \mathcal{R}_D , as D ranges over all Go-diagrams contained in a $k \times (n - k)$ rectangle. In other words, each point in the Grassmannian $Gr_{k,n}$ can be represented uniquely by a weighted network associated to a Go-diagram.

A more refined version of Theorem 3.16 is given in Corollary 6.9.

Corollary 3.17. Every matrix can be represented by a unique weighted network associated to a Go-diagram.

Proof. Suppose $M = (m_{i,j})$ is a $k \times (n - k)$ matrix. Let $A(M) = (a_{i,j})$ be the full rank $k \times n$ matrix with an identity submatrix in the first k columns and the remaining columns given by $a_{i,j+k} = (-1)^{i+1} m_{n+1-i,j}$. Then $A(M)$ represents an element in the Grassmannian $Gr_{k,n}$, so Theorem 3.16 applies. Moreover, the minors of M are in bijection with the $k \times k$ minors of $A(M)$, so if $A(M)$ is represented by the network N , we see that $m_{i,j}$ enumerates paths from the boundary source i to the boundary vertex j in N . \square

We will prove Theorem 3.16 by showing that each network component \mathcal{R}_D from a Go-diagram coincides with a (*projected*) *Deodhar component* $\mathcal{P}_{\mathbf{v},\mathbf{w}}$ in the Grassmannian. (Therefore we may refer to each \mathcal{R}_D as a *Deodhar component*.) More specifically, such Deodhar components have parameterizations due to Marsh and Rietsch [8], and we will show that after an invertible transformation of variables, our network parameterizations coincide with theirs.

4. THE DEODHAR DECOMPOSITION OF THE GRASSMANNIAN

In this section we review Deodhar's decomposition of the flag variety G/B [2], and the parameterizations of the components due to Marsh and Rietsch [8]. The Deodhar decomposition of the Grassmannian is obtained by projecting the Deodhar decomposition of G/B to the Grassmannian [3].

4.1. The flag variety. Let \mathbb{K} be a field, and let G denote the special linear group $SL_n = SL_n(\mathbb{K})$. Fix a maximal torus T , and opposite Borel subgroups B^+ and B^- , which consist of the diagonal, upper-triangular, and lower-triangular matrices, respectively. Let U^+ and U^- be the unipotent radicals of B^+ and B^- ; these are the subgroups of upper-triangular and lower-triangular matrices with 1's on the

which is

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_2 & -m_5 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & p_8 & 1 & 0 & 0 & 0 & 0 \\ 0 & -p_4 & -m_7 p_8 & -m_7 + p_{12} & 1 & 0 & 0 & 0 \\ 0 & 0 & -p_6 p_8 & -p_6 + p_{11} p_{12} & p_{11} & 1 & 0 & 0 \\ 0 & 0 & 0 & p_{10} p_{11} p_{12} & p_{10} p_{11} & p_{10} & 1 & 0 \\ 0 & 0 & 0 & p_9 p_{10} p_{11} p_{12} & p_9 p_{10} p_{11} & p_9 p_{10} & p_9 & 1 \end{pmatrix}.$$

The following result from [8] gives an explicit parametrization for the Deodhar component $\mathcal{R}_{\mathbf{v}, \mathbf{w}}$. We use Proposition 4.3 as the *definition* of $\mathcal{R}_{\mathbf{v}, \mathbf{w}}$.

Proposition 4.3. [8, Proposition 5.2] The map $(\mathbb{K}^*)^{|\mathcal{J}^\downarrow|} \times \mathbb{K}^{|\mathcal{J}^\circ|} \rightarrow G_{\mathbf{v}, \mathbf{w}}$ from Definition 4.1 is an isomorphism. The map $g \mapsto g \cdot B^+$ defines an isomorphism

$$(4.3) \quad G_{\mathbf{v}, \mathbf{w}} \xrightarrow{\sim} \mathcal{R}_{\mathbf{v}, \mathbf{w}}$$

between the subset $G_{\mathbf{v}, \mathbf{w}}$ of the group, and the Deodhar component $\mathcal{R}_{\mathbf{v}, \mathbf{w}}$ in G/B .

Suppose that for each $w \in W$ we choose a reduced expression \mathbf{w} for w . Then it follows from [2] and [8, Section 4.4] that

$$(4.4) \quad \mathcal{R}_{v, w} = \bigsqcup_{\mathbf{v} \prec \mathbf{w}} \mathcal{R}_{\mathbf{v}, \mathbf{w}} \quad \text{and} \quad G/B = \bigsqcup_{w \in W} \left(\bigsqcup_{\mathbf{v} \prec \mathbf{w}} \mathcal{R}_{\mathbf{v}, \mathbf{w}} \right),$$

where in the first sum \mathbf{v} ranges over all distinguished subexpressions for v in w , and in the second sum \mathbf{v} ranges over all distinguished subexpressions of \mathbf{w} . These two decompositions are called the *Deodhar decompositions* of $\mathcal{R}_{v, w}$ and G/B .

Remark 4.4. Although the Deodhar decomposition of $\mathcal{R}_{v, w}$ depends on the choice \mathbf{w} of reduced expression for w , its projection to the Grassmannian does not depend on \mathbf{w} [6, Proposition 4.16].

4.3. Projections of Deodhar components to the Grassmannian. Following [6], we now consider the projection of the Deodhar decomposition to the Grassmannian $Gr_{k, n}$ for $k < n$. Given $w = (w(1), w(2), \dots, w(n)) \in W^k$, we let $I(w) = \{w(n-k+1), w(n-k+2), \dots, w(n)\}$. The map I gives a bijection between W^k and k -element subsets of $[n]$.

Let $\pi_k : G/B \rightarrow Gr_{k, n}$ be the projection from the flag variety to the Grassmannian; this is an isomorphism on each $\mathcal{R}_{v, w}$. For each $w \in W^k$ and $v \leq w$, define $\mathcal{P}_{v, w} = \pi_k(\mathcal{R}_{v, w})$. Then by [7] we have a decomposition

$$(4.5) \quad Gr_{k, n} = \bigsqcup_{w \in W^k} \left(\bigsqcup_{v \leq w} \mathcal{P}_{v, w} \right).$$

For each reduced decomposition \mathbf{w} for $w \in W^k$, and each $\mathbf{v} \prec \mathbf{w}$, we define $\mathcal{P}_{\mathbf{v}, \mathbf{w}} = \pi_k(\mathcal{R}_{\mathbf{v}, \mathbf{w}})$. Now if for each $w \in W^k$ we choose a reduced decomposition \mathbf{w} , then we have

$$(4.6) \quad \mathcal{P}_{v, w} = \bigsqcup_{\mathbf{v} \prec \mathbf{w}} \mathcal{P}_{\mathbf{v}, \mathbf{w}} \quad \text{and} \quad Gr_{k, n} = \bigsqcup_{w \in W^k} \left(\bigsqcup_{\mathbf{v} \prec \mathbf{w}} \mathcal{P}_{\mathbf{v}, \mathbf{w}} \right),$$

where in the first sum \mathbf{v} ranges over all distinguished subexpressions for v in w , and in the second sum \mathbf{v} ranges over all distinguished subexpressions of \mathbf{w} .

Proposition 4.3 gives a concrete way to think about the projected Deodhar components $\mathcal{P}_{\mathbf{v}, \mathbf{w}}$. The projection $\pi_k : G/B \rightarrow Gr_{k,n}$ maps $g \cdot B^+ \in R_{\mathbf{v}, \mathbf{w}}$ (where $g \in G_{\mathbf{v}, \mathbf{w}}$) to the span of the leftmost k columns of g . More specifically, it maps

$$g = \begin{pmatrix} g_{n,n} & \cdots & g_{n,n-k+1} & \cdots & g_{n,1} \\ \vdots & & \vdots & & \vdots \\ g_{1,n} & \cdots & g_{1,n-k+1} & \cdots & g_{1,1} \end{pmatrix} \longrightarrow M = \begin{pmatrix} g_{1,n-k+1} & \cdots & g_{n,n-k+1} \\ \vdots & & \vdots \\ g_{1,n} & \cdots & g_{n,n} \end{pmatrix}$$

We call the resulting $k \times n$ matrix $M = (M_{st})$ the *MR-matrix*. To simplify the notation later, we will label its rows from top to bottom by i_1, i_2, \dots, i_k , where $\{i_1 < \dots < i_k\} = I(w)$.

Remark 4.5. Recall from Section 3.3 that in the Grassmannian setting (i.e. $W_k = \langle s_1, s_2, \dots, \hat{s}_{n-k}, \dots, s_{n-1} \rangle$ is a parabolic subgroup of $W = \mathfrak{S}_n$), the distinguished subexpressions of W^k are in bijection with Go-diagrams. Therefore each Go-diagram gives rise to an MR-matrix.

Example 4.6. We continue Example 4.2. Note that $w \in W^k$, where $k = 2$. Then the map $\pi_2 : G_{\mathbf{v}, \mathbf{w}} \rightarrow Gr_{2,5}$ is given by

$$g = \left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ p_2 & -m_5 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & p_8 & 1 & 0 & 0 & 0 & 0 \\ 0 & -p_4 & -m_7 p_8 & -m_7 + p_{12} & 1 & 0 & 0 & 0 \\ 0 & 0 & -p_6 p_8 & -p_6 + p_{11} p_{12} & p_{11} & 1 & 0 & 0 \\ 0 & 0 & 0 & p_{10} p_{11} p_{12} & p_{10} p_{11} & p_{10} & 1 & 0 \\ 0 & 0 & 0 & p_9 p_{10} p_{11} p_{12} & p_9 p_{10} p_{11} & p_9 p_{10} & p_9 & 1 \end{array} \right) \longrightarrow$$

$$M = \begin{pmatrix} p_9 p_{10} p_{11} p_{12} & p_{10} p_{11} p_{12} & -p_6 + p_{11} p_{12} & -m_7 + p_{12} & 1 & 0 & 0 & 0 \\ 0 & 0 & -p_6 p_8 & -m_7 p_8 & p_8 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -p_4 & 0 & -m_5 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & p_2 & 0 & 1 \end{pmatrix}$$

We label the rows of M from top to bottom by the index set $\{1, 3, 4, 6\}$, and the columns from left to right by the index set $\{1, 2, \dots, 8\}$, so e.g. $M_{34} = -m_7 p_8$.

The following lemma is a consequence of [6, Section 5.1] and in particular [6, Corollary 5.8].

Lemma 4.7. Let $M = M_D$ be the MR-matrix associated to the diagram D . The leftmost nonzero entry in row i_ℓ of M is in column i_ℓ . Furthermore, that entry is equal to $(-1)^b \prod p_i$, where b is the number of black stones in the row i_ℓ of D , and the product is over all boxes in the row i_ℓ of the labeled Go-diagram of D containing a p_i .

5. FORMULAS FOR ENTRIES OF THE MR-MATRICES

In this section we consider arbitrary diagrams (not necessarily Go-diagrams) contained in a $k \times (n-k)$ rectangle and the corresponding MR-matrices, obtained by multiplying factors $\dot{s}_i, y_i(p_j), x_i(m_j) \dot{s}_i^{-1}$ as specified by the filling of the diagram, and then projecting the resulting $n \times n$ matrix to a $k \times n$ matrix. We will give formulas for the entries of the MR-matrices in terms of *pseudopaths* in the corresponding network. For the purpose of giving this formula, we will replace weights a_i and c_j on the edges of the network by weights p_i and m_j .

Recall that if D is a diagram, its network N_D has three types of vertices: $+$ -vertices, \bullet -vertices, and boundary vertices. A *step* on a network is an edge between two vertices. Let W denote a single step west, S denote a single step south, and E denote either a single step east, or an *east-west combination step* consisting of a step east, followed by a step west ending at a \bullet . Let A^* indicate 0 or more instances of a step of type A .

Definition 5.1. A *pseudopath* \tilde{P} on a network is a path on the (undirected version of the) network such that:

- it starts and ends at two different boundary vertices, or else is the empty path from a boundary vertex to itself;
- it does not cross the same edge twice;
- its sequence of steps (for a nonempty path) has the form

$$WW^*S(EE^*S)^*E^*.$$

In particular, a pseudopath may not take two consecutive steps south.

Definition 5.2. The *weight* $w(\tilde{P})$ of a pseudopath \tilde{P} in a network is a Laurent monomial in p_i 's and m_j 's, which is obtained by multiplying the following terms:

- $\frac{1}{p_i}$ for every step west along an edge weighted p_i ;
- p_i for every step east along an edge weighted p_i which is preceded by a step east;
- m_j for every step west along an edge weighted m_j ;
- $(-1)^{b+w}$, where b (respectively w) is the number of black (resp. white) stones that the pseudopath skips over in the horizontal (resp. vertical) direction, when we superimpose the Go-diagram onto the network.

Example 5.3. In Figure 5, there are two pseudopaths from 1 to 4, with weights $\frac{1}{p_9 p_{10} p_{11}}$ and $\frac{-m_7}{p_9 p_{10} p_{11} p_{12}}$, and there is one pseudopath from 1 to 5, with weight $\frac{1}{p_9 p_{10} p_{11} p_{12}}$.




FIGURE 5. The two pseudopaths from 1 to 4 and the unique pseudopath from 1 to 5, indicated in bold. Note that the pseudopath in the middle figure contains an east-west combination step.

Definition 5.4. If M is an MR-matrix, we will let \tilde{M} denote the matrix obtained from M by rescaling rows so that the leftmost nonzero entry in each row is 1.

Definition 5.5. If D is a diagram contained in a $k \times (n - k)$ rectangle, then we let $i_1 < i_2 < \dots < i_k$ denote the labels of the sources in the corresponding network. If M and \tilde{M} are the corresponding $k \times n$ MR and rescaled MR-matrices associated to D , then we will index their rows by i_1, \dots, i_k from top to bottom, and their columns by $1, 2, \dots, n$ from left to right.

Theorem 5.6. Let D be a diagram contained in a $k \times (n - k)$ rectangle, and let $\widetilde{M} = (\widetilde{M}_{st})$ be the corresponding $k \times n$ rescaled MR-matrix. Then

$$\widetilde{M}_{st} = \sum_{\widetilde{P}} w(\widetilde{P}),$$

where the sum is over all pseudopaths from the source s to the boundary vertex t in the network .

The proof of Theorem 5.6 will follow from Theorem 5.10 and Lemma 5.11.

Example 5.7. The MR-matrix M from Example 4.6 corresponds to the network from Figure 5. The rows of M are indexed by 1, 3, 4, 6 from top to bottom. Note that after we rescale the rows of M , obtaining \widetilde{M} , we have $\widetilde{M}_{14} = \frac{1}{p_9 p_{10} p_{11}} - \frac{m_7}{p_9 p_{10} p_{11} p_{12}}$, and $\widetilde{M}_{15} = \frac{1}{p_9 p_{10} p_{11} p_{12}}$. This agrees with our pseudopath computation from Example 5.3.

Next we will give a formula for entries of MR-matrices, in terms of pseudopaths in *modified networks*.

Definition 5.8. Given a network N_D with k sources labeled i_1, \dots, i_k and n boundary vertices, we obtain from it a corresponding *modified network* N'_D , by:

- adding k new boundary vertices to the left of N_D , labeled i'_1, \dots, i'_k from top to bottom;
- adding a new horizontal edge which connects i'_j to the leftmost internal vertex of the network.

See Figure 6 for the modified network associated to the network from Figure 5.

Definition 5.9. A *pseudopath* P on a modified network is a path on the modified network which:

- starts at one of the boundary vertices labeled i'_1, \dots, i'_k , and ends at one of the boundary vertices labeled $1, 2, \dots, n$;
- takes a sequence of steps which has the form

$$(EE^*S)^*E^*.$$

The arrows in Figure 6 indicate the allowed directions in which a path may travel.

The *weight* of a pseudopath in a modified network is defined the same way as the weight of a pseudopath in a network (see Definition 5.2). Note that since a pseudopath in a modified network does not contain steps west along edges weighted p_i , its weight is a monomial (not a Laurent monomial).




FIGURE 6. Example of a modified network.

Theorem 5.10. Let D be a diagram contained in a $k \times (n - k)$ rectangle, and let $M = (M_{st})$ be the corresponding $k \times n$ MR-matrix. Then

$$M_{st} = \sum_P w(P),$$

where the sum is over all pseudopaths in the modified network from the boundary vertex s' to the boundary vertex t .

Lemma 5.11. Theorems 5.6 and 5.10 are equivalent.

Proof. There is an obvious bijection between pseudopaths in a network starting at boundary vertex s , and pseudopaths in the corresponding modified network starting at boundary vertex s' . The weights of the corresponding pseudopaths are the same except for a factor of $(-1)^b \prod p_i$, where b is the number of \bullet -vertices in row s of the network, and the p_i 's range over all edge weights in row s .

On the other hand, Lemma 4.7 implies that the leftmost nonzero entry of row s of the MR-matrix M is precisely the quantity $(-1)^b \prod p_i$ above. Therefore Theorems 5.6 and 5.10 are equivalent. \square

By Lemma 5.11, it suffices to prove just one of the two theorems. We will prove Theorem 5.10. Our strategy will be to interpret entries of the MR-matrix in terms of paths in a chip network, and then construct a weight-preserving bijection between these paths and between pseudopaths in the modified network.

Definition 5.12. A *chip* is one of the three configurations shown in Figure 7. We call the three configurations $y_i(p)$ - or y_i -chips, s_i -chips, and $x_i(m)$ - or x_i -chips, respectively.




FIGURE 7. The three types of chips.

Definition 5.13. A *chip network* is a concatenation of chips. Note that it has n boundary vertices at the left and n boundary vertices at the right. Let g be a product of factors of the form $y_i(p)$, \hat{s}_i , and $x_i(m)\hat{s}_i^{-1}$. We associate a chip network C_g to every such product g by concatenating the chips corresponding to the factors of g in the order given by the factorization.

Example 5.14. The chip network C_g associated to the product g from (4.2) is shown in Figure 8.




FIGURE 8. The chip network corresponding to the product

$$g = \dot{s}_6 y_7(p_2) \dot{s}_4 y_5(p_4) x_6(m_5) \dot{s}_6^{-1} y_3(p_6) x_4(m_7) \dot{s}_4^{-1} y_5(p_8) y_1(p_9) y_2(p_{10}) y_3(p_{11}) y_4(p_{12}).$$

Definition 5.15. A route Q in a chip network is a path in the network whose steps all travel east (or southeast or northeast for slanted edges). The weight $w(Q)$ of such a route is the product of all weights on its edges. To each chip network C we associate a weight matrix $x(C) = x_{ij}$ where $x_{ij} = \sum_Q w(Q)$ and the sum is over all routes from the boundary vertex i at the west to the boundary vertex j at the east.

It is simple to verify the following result. Recall our convention from Section 4.1 that the rows of g are labeled from bottom to top, and the columns of g are labeled from right to left.

Lemma 5.16. Let g be a product of factors of the form $y_i(p)$, \dot{s}_i , and $x_i(m)\dot{s}_i^{-1}$. Then the weight matrix $x(C_g)$ of the chip network C_g associated to g coincides with the matrix g .

We now prove Theorem 5.10.

Proof. Let D be a diagram contained in a $k \times (n - k)$ rectangle, and N'_D the corresponding modified network. Let $i'_1 < \dots < i'_k$ be the labels of the sources of N'_D . Let g be the product of factors of the form $y_i(p)$, \dot{s}_i , and $x_i(m)\dot{s}_i^{-1}$ which is encoded by D , and let M be the corresponding MR-matrix, whose rows are indexed from top to bottom by i_1, \dots, i_k . Recall that the projection from g to M switches rows and columns, and the columns labeled $n - k + 1, n - k + 2, \dots, n$ in g become rows labeled i_1, i_2, \dots, i_k in M . Therefore to prove Theorem 5.10, it suffices to prove that for all $1 \leq t \leq n$ and $1 \leq s \leq k$ we have

$$(5.1) \quad g_{t, s+(n-k)} = \sum_P w(P),$$

where the sum is over all pseudopaths P from i'_s to t in the modified network.

By Lemma 5.16, the matrix g coincides with the weight matrix $x(C_g) = (x_{st})$ of the chip network associated to g . Therefore by (5.1) it suffices to prove that for all $1 \leq t \leq n$ and $1 \leq s \leq k$ we have

$$(5.2) \quad x_{t, s+(n-k)} = \sum_P w(P),$$

where the sum is over all pseudopaths P from i'_s to t in the modified network.

Recall from Definition 5.15 that $x_{ij} = \sum_Q w(Q)$, where the sum is over all routes Q in the chip network from the boundary vertex i at the west to the boundary vertex j at the east. To prove (5.2), we will

give a weight-preserving bijection between pseudopaths P in the modified network from i'_s to t , and routes Q in the chip network from the boundary vertex t at the west to the boundary vertex $s + (n - k)$ at the east. More specifically, given a pseudopath P , we will examine its sequence of steps from source to sink, and explain how to build the corresponding route Q in the chip network. As illustrated in Figure 9, each step in a pseudopath corresponds to a portion of a route in a chip network. (Note that our bijection will build the route in the chip network from east to west, rather than west to east.)

Recall from Figure 4 that each modified network comes from a diagram, and that every box of a diagram is naturally associated with a simple generator s_i . Therefore every internal vertex in a modified network is naturally associated with a simple generator s_i for some i . We will call this the *position* of the vertex.

Let us consider the various kinds of steps in a pseudopath. Such steps naturally fall into one of the following categories (illustrated in Figure 9):

0. A single step east, which starts at a source, and ends at position s_i .
- I. A single step east, which is preceded by a south step, and followed by an east or south step. Such a step starts and ends at positions s_i and s_j (for $i > j$), and is labeled by some weight p . It may skip over some (positions corresponding to) white and black stones in the Go-diagram.
- II. An east-west combination step, which is preceded by a south step, and travels from position s_i to s_k to s_j (where $i > j > k$). The two components of such a step are labeled by some weights p and m , and may skip over some white and black stones.
- III. A single step east, which is preceded by an east step, and followed by an east or south step. Such a step starts and ends at positions s_i and s_j (for $i > j$), and is labeled by some weight p . It may skip over some white and black stones.
- IV. An east-west combination step, which is preceded by an east step, and travels from position s_i to s_k to s_j (where $i > j > k$). The two components of such a step are labeled by some weights p and m , and may skip over some white and black stones.
- V. A south step, which starts and ends at positions s_i and s_j (for $i < j$). Such a step may skip over some white and black stones.

The above steps in a pseudopath correspond to the following portion of a route in a chip network:

0. Start at the boundary vertex $i + 1$ at the east of the chip network.
- I. Start at level i , then travel west straight across the $y_i(p)$ chip. For each white or black stone (say in position s_ℓ) which lies in between positions s_i and s_j , travel northwest through the corresponding s_ℓ or x_ℓ -chip, ending at level $j + 1$.
- II. Start at level i , then travel west straight across the $y_i(p)$ chip. For each white or black stone (say in position s_ℓ) which lies in between positions s_i and s_j , travel northwest through the corresponding s_ℓ or x_ℓ -chip. Finally, travel along the -1 -edge and then the m -edge of the $x_j(m)$ chip, ending at level $j + 1$.
- III. Start at level $i + 1$, then travel northwest along the p -edge in the $y_i(p)$ chip. For each white or black stone (say in position s_ℓ) which lies in between positions s_i and s_j , travel northwest through the corresponding s_ℓ or x_ℓ -chip, ending at level $j + 1$.
- IV. Start at level $i + 1$, then travel northwest along the p -edge in the $y_i(p)$ chip. For each white or black stone (say in position s_ℓ) which lies in between positions s_i and s_j , travel northwest through the corresponding s_ℓ or x_ℓ -chip. Finally, travel along the -1 -edge and the m -edge of the $x_j(m)$ chip, ending at level $j + 1$.
- V. Start at level $i + 1$. For each white or black stone (say in position s_ℓ) which lies in between positions s_i and s_j , travel southwest through the corresponding s_ℓ or x_ℓ -chip, ending at level j .

It is a straightforward exercise to verify that this map is a bijection between pseudopaths P from i'_s to t in the modified network, and routes Q between the t vertex at the west and the $s + (n - k)$ vertex




FIGURE 9. Steps in pseudopaths and their corresponding fragments of the chip network.

at the east in the chip network. Moreover, the weights of P and Q are equal. See Figure 10 for example of entire pseudopaths and routes. \square




FIGURE 10. Pseudopaths in the modified network and their corresponding routes in the chip network.

6. PROOF OF THE MAIN RESULT

Let D be a diagram which contains t pluses and u black stones. In Section 6.1 we will define an isomorphism

$$\Psi = \Psi_D : (\mathbb{K}^*)^t \times \mathbb{K}^u \longrightarrow (\mathbb{K}^*)^t \times \mathbb{K}^u$$

which maps each parameter from the weight matrix of the network N_D to a Laurent monomial in the parameters used in the MR-matrix $M = M_D$. Then in Section 6.2, we will show that after applying Ψ , our network parameterization of the network component \mathcal{R}_D coincides with the corresponding MR-parameterization of the projected Deodhar component $\mathcal{P}_{\mathbf{v}, \mathbf{w}}$. Combining this fact with Proposition 4.3 yields Theorem 3.16.

6.1. A rational transformation of parameters.

Definition 6.1. Let D be a diagram, and let b_0 be a box of D containing a $+$ or \bullet . Let b_1 be the nearest box to the right of b_0 which contains a $+$ (if it exists). Let R_t be the set of boxes in the same row as b_0 which are to the right of b_0 and left of b_1 . Let R_ℓ be the set of boxes in the same column as b_0 and below b_0 . If b_1 exists, let R_r be the set of boxes in the same column as b_1 and below b_1 (otherwise $R_r = \emptyset$). (See Figure 11.) Let R_r^+ (resp. R_ℓ^+) be the set of boxes in R_r (resp. R_ℓ) containing a $+$. Let $R = R_r \cup R_\ell \cup R_t$ and let R^\bullet be the set of boxes in R containing a \bullet .

If b_0 contains a $+$, then define

$$\Psi(a_{b_0}) = \frac{(-1)^{|R^\bullet|} \prod_{b \in R_r^+} p_b}{p_{b_0} \prod_{b \in R_\ell^+} p_b}.$$

And if b_0 contains a \bullet , then define

$$\Psi(c_{b_0}) = \frac{m_{b_0}(-1)^{|R^\bullet|} \prod_{b \in R_r^+} p_b}{\prod_{b \in R_\ell^+} p_b}.$$

We also extend the definition of Ψ to all polynomials in the a_b 's and c_b 's by requiring it to be a ring homomorphism.




FIGURE 11. The definition of b_0 , b_1 , R_t , R_ℓ , and R_r .

Remark 6.2. Clearly $\Psi = \Psi_D$ is an isomorphism from $(\mathbb{K}^*)^t \times \mathbb{K}^u$ to itself.

Example 6.3. Consider the network from Figure 3 (shown again in Figure 12). Then we have $\Psi(a_2) = \frac{1}{p_2}$, $\Psi(a_4) = \frac{1}{p_4}$, $\Psi(c_5) = \frac{m_5}{p_2}$, $\Psi(a_6) = \frac{1}{p_6}$, $\Psi(c_7) = \frac{m_7}{p_4}$, $\Psi(a_8) = \frac{1}{p_2 p_8}$, $\Psi(a_9) = \frac{1}{p_9}$, $\Psi(a_{10}) = \frac{1}{p_6 p_{10}}$, $\Psi(a_{11}) = \frac{-p_6}{p_4 p_{11}}$, and $\Psi(a_{12}) = \frac{p_4}{p_2 p_8 p_{12}}$.

From Definition 6.1, one may easily deduce a formula for $\Psi(w(P))$, where P is a path in N_D . We will state this formula in terms of the Go-diagram. Note that one may identify a path P in N_D with a connected sequence \mathcal{P} of boxes in the Go-diagram, where any two adjacent boxes must share a side. We call a box b in \mathcal{P} a *corner box* if the path P turns from west to south, or from south to west, at the vertex associated to b . (Such a box b in D must contain a $+$ or \bullet .)

Proposition 6.4. Let P be a path in the network N_D , which we identify with a sequence \mathcal{P} of boxes in the Go-diagram D . Among the boxes in \mathcal{P} , let B_1 denote the subset containing a $+$; let B_2 denote the subset containing a \bullet which are corner boxes of \mathcal{P} ; and let B_3 denote the subset containing a \bullet which are not corner boxes of \mathcal{P} . Then


$$(6.1) \quad \Psi(w(P)) = \frac{(-1)^{|B_3|} \prod_{b \in B_2} m_b}{\prod_{b \in B_1} p_b}.$$

Proof. We leave this as an exercise for the reader. It is a simple consequence of Definition 6.1. \square

Example 6.5. Let P_1 and P_2 be the paths shown in Figure 12. Then $\Psi(w(P_1)) = \Psi(a_9 a_{10} a_{11} c_5) = \frac{-m_5}{p_2 p_4 p_9 p_{10} p_{11}}$ and $\Psi(w(P_2)) = \Psi(a_9 a_{10} c_5 c_7) = \frac{m_5 m_7}{p_2 p_4 p_6 p_9 p_{10}}$.

6.2. Applying row operations to the rescaled MR-matrix.

Theorem 6.6. Let D be a diagram, let $M = M_D$ be the corresponding MR-matrix, and let L be the matrix we obtain by putting M into reduced row-echelon form. Let $W_D = (W_{ij})$ be the weight matrix associated to D , and let $\Psi(W_D)$ be the matrix obtained from W_D by applying the rational map Ψ to each entry. Then $\Psi(W_D) = L$.

FIGURE 12. Two paths in N_D and their corresponding sequences of boxes in D .

Proof. To prove Theorem 6.6, we start by considering the rescaled MR-matrix \widetilde{M} . Its rows are indexed by the set i_1, \dots, i_k , the set of sources of the network N_D , and the leftmost nonzero entry in every row is a 1. Moreover, by Lemma 4.7, the 1 in row i_ℓ is located in column i_ℓ . The entries of the reduced row echelon matrix L obtained from \widetilde{M} are given by the formula

$$(6.2) \quad L_{st} = \widetilde{M}_{st} + \sum_{s < j_1 < \dots < j_r < t} (-1)^r \widetilde{M}_{sj_1} \widetilde{M}_{j_1 j_2} \dots \widetilde{M}_{j_r t},$$

where the sum ranges over all nonempty subsets $\{j_1, \dots, j_r\} \subset \{i_1, \dots, i_k\}$ of sources of the network between s and t .

By Theorem 5.6, the entry \widetilde{M}_{st} equals $\sum_{\widetilde{P}} w(\widetilde{P})$, where the sum is over all pseudopaths in the network N_D from the source s to the boundary vertex t . Therefore the right-hand side of (6.2) can be interpreted as a generating function for all concatenations of pseudopaths, where the first pseudopath starts at s , and the last pseudopath ends at t .

Let us identify a pseudopath with its collection of directed edges. Given a set of pseudopaths on N_D , we define its *signed union* to be the union of directed edges that one obtains by taking the multiset of all directed edges in the pseudopaths, and then cancelling pairs which traverse the same edge but in opposite directions. We define the *weight* of a set of pseudopaths to be the product of the weights of each pseudopath in the set.

Our goal is to show that after cancellation, the only terms which survive on the right-hand side of (6.2) correspond to concatenations of pseudopaths whose signed union forms a directed path (and not merely a pseudopath) from s to t in N_D . This will allow us to relate (6.2) to W_{st} , which is defined as a sum over all paths from s to t in N_D .

Definition 6.7. A *U-turn* pseudopath in a network is a pseudopath whose sequence of steps has the form $(WW^*S)E^*$.

First note that any path P in a network N_D has a unique decomposition as a signed union of U-turn pseudopaths. Moreover, the products of the weights of the pseudopaths is precisely the quantity on the right-hand side of (6.1). See Figure 13 for an example of the decomposition into U-turn pseudopaths.

This observation on the decomposition of paths may be generalized to pseudopaths. Consider a pseudopath \widetilde{P} which is not a path, and turns from south to east precisely q times (for $q \geq 1$). Then for each $0 \leq r \leq q$, there are $\binom{q}{r}$ decompositions of \widetilde{P} as a signed union of r pseudopaths. Moreover, each set of pseudopaths forming a decomposition of \widetilde{P} has the same weight. See Figure 14 for an example of all decompositions of a pseudopath as a signed union of pseudopaths. It is easy to check that each decomposition has the same weight.




FIGURE 13. A path in N_D and its decomposition into U-turn pseudopaths.




FIGURE 14. An example of the 2^q decompositions of a path in N_D into pseudopaths.

Note that since all decompositions of a pseudopath have the same weight, and because

$$\binom{q}{0} - \binom{q}{1} + \binom{q}{2} - \dots \pm \binom{q}{q} = 0 \text{ for } q \geq 1,$$

the net contribution of the corresponding concatenations of pseudopaths in (6.2) is 0.

More generally, a term on the right-hand side of (6.2) corresponds to a concatenation of pseudopaths (whose signed union may not be a pseudopath). However, just as before, one may decompose the signed union \tilde{P} in 2^q ways, where q is the number of times that \tilde{P} turns from south to east. And again, for $q \geq 1$, the net contribution of the corresponding concatenations of pseudopaths in (6.2) is 0. Therefore when one interprets the right-hand side of (6.2) as a sum over concatenations of pseudopaths, the only terms that are not cancelled are the terms corresponding to concatenations of pseudopaths whose signed union is a directed path.

Example 6.8. In Figure 15, the left and middle diagrams show a pseudopath from 1 to 5, and a concatenation of two pseudopaths from 1 to 5, whose weights cancel each other. The right diagram shows a path from 1 to 5, written as a signed union of two pseudopaths, which will not be cancelled by any other term.

It is a simple exercise to verify that the absolute value of the weight of such a concatenation is precisely the absolute value of the right-hand side of (6.1). We also need to verify that the signs agree. Once we do this, then since W_{st} is the sum of all weights of paths in N_D from s to t , $\Psi(W_{st})$ equals the corresponding term in the expression for L_{st} in (6.2), so the proof is done.

We now check that the signs agree. More specifically, consider a concatenation of $(r+1)$ pseudopaths whose signed union is a directed path P . Using Definition 5.2, the total sign associated to the concatenation of pseudopaths from (6.2) is $(-1)^{r+b+w}$, where b (respectively w) is the number of black (resp. white) stones that P skips over in the horizontal (resp. vertical) direction. (Note that for the purpose

of computing b and w , we can count stones skipped by P here, as opposed to the set of pseudopaths whose signed union is P .) Meanwhile, using Definition 3.3 and Proposition 6.4, the total sign associated to the directed path P in the expression $\Psi(W_{st})$ is $(-1)^{q+|B_3|}$, where q is the number of sources in the network which are strictly between s and t , and $|B_3|$ is the number of non-corner black stones in the boxes of the Go-diagram which P traces out. We need to show that $(-1)^{r+b+w} = (-1)^{q+|B_3|}$.

Note that $|B_3|$ is the number of black stones skipped either vertically or horizontally by P . Let b_v (resp. w_v) denote the number of black stones (resp. white stones) skipped vertically by P . Similarly, let b_h (resp. w_h) denote the number of black stones (resp. white stones) skipped horizontally by P . With this notation, we need to show that $(-1)^{b_h+w_h+r} = (-1)^{b_h+b_v+q}$, i.e. that $(-1)^{w_v+r} = (-1)^{b_v+q}$.

To prove this, we will show that $q - r = w_v + b_v$. Note that $q - r$ is the number of sources strictly between s and t which are *not* sources of any pseudopath in the pseudopath decomposition of P , i.e. which are not in the set $\{j_1, \dots, j_r\}$ from (6.2). Recall that the pseudopath decomposition of P is a U-turn pseudopath decomposition; therefore, a source between s and t lies in $\{j_1, \dots, j_r\}$ if and only if P has a vertical edge ending at a $+$ in this row. Otherwise P skips a black or white stone in this row. This proves that $q - r = w_v + b_v$, and hence completes the proof of Theorem 6.6. \square

We have now shown that after an invertible transformation of the parameters, our network parameterization of \mathcal{R}_D coincides with the corresponding MR-parameterization of the projected Deodhar component $\mathcal{P}_{\mathbf{v},\mathbf{w}}$. Combining this result with Proposition 4.3 yields Theorem 3.16.

Our proof yields the following statement.

Corollary 6.9. Let D be the Go-diagram associated to the distinguished subexpression \mathbf{v} of \mathbf{w} . Then $\mathcal{R}_D = \mathcal{P}_{\mathbf{v},\mathbf{w}}$ as subsets of $Gr_{k,n}$. Furthermore,

$$Gr_{k,n} = \bigsqcup_D \mathcal{R}_D,$$

where the union is over all Go-diagrams D contained in a $k \times (n - k)$ rectangle.




FIGURE 15. The three possible concatenations of pseudopaths from 1 to 5.

7. A CHARACTERIZATION OF DEODHAR COMPONENTS IN TERMS OF PLÜCKER COORDINATES

In this section we characterize Deodhar components in the Grassmannian by a list of vanishing and nonvanishing Plücker coordinates. Our main result in this section is Theorem 7.8. The proof uses results of Kodama and the second author from [6], which gave formulas for Plücker coordinates of Deodhar components.

7.1. Plücker coordinates of Deodhar components in terms of the MR parameters. Consider the Deodhar component $\mathcal{P}_{\mathbf{v},\mathbf{w}} \subset Gr_{k,n}$, where \mathbf{w} is a reduced expression for $w \in W^k$ and $\mathbf{v} \prec \mathbf{w}$. In this section we will review some formulas from [6] for the Plücker coordinates of the elements of $\mathcal{P}_{\mathbf{v},\mathbf{w}}$, in terms of the parameters which Marsh and Rietsch [8] used to define $G_{\mathbf{v},\mathbf{w}}$.

7.1.1. *Formulas for Plücker coordinates.*

Theorem 7.1. [6, Lemma 5.1 and Theorem 5.2] Let $\mathbf{w} = s_{i_1} \dots s_{i_m}$ be a reduced expression for $w \in W^k$ and $\mathbf{v} \prec \mathbf{w}$ be a distinguished subexpression for v . Let $A = \pi_k(g) \in \mathcal{P}_{\mathbf{v}, \mathbf{w}}$ for any $g \in G_{\mathbf{v}, \mathbf{w}}$. Then the lexicographically minimal and maximal nonzero Plücker coordinates of A are Δ_I and $\Delta_{I'}$, where $I = w\{n, n-1, \dots, n-k+1\}$ and $I' = v\{n, n-1, \dots, n-k+1\}$. If we write $g = g_1 \dots g_m$ as in Definition 4.1, then


$$(7.1) \quad \Delta_I(A) = (-1)^{|J_{\mathbf{v}}^{\bullet}|} \prod_{i \in J_{\mathbf{v}}^+} p_i \quad \text{and} \quad \Delta_{I'}(A) = 1.$$

Remark 7.2. If we write $I = \{i_1, \dots, i_k\}$, then $I' = vw^{-1}\{i_1, \dots, i_k\}$.


Definition 7.3. [6, Definition 5.4] Let $W = \mathfrak{S}_n$, let $\mathbf{w} = s_{i_1} \dots s_{i_m}$ be a reduced expression for $w \in W^k$ and choose $\mathbf{v} \prec \mathbf{w}$. This determines a Go-diagram D of shape $\lambda = \lambda_w$. Let b be any box of D . Note that the set of all boxes of D which are weakly southeast of b forms a Young diagram λ_b^{in} ; also the complement of λ_b^{in} in λ is a Young diagram which we call λ_b^{out} (see Example 7.4 below). By looking at the restriction of \mathbf{w} to the positions corresponding to boxes of λ_b^{in} , we obtained a reduced expression \mathbf{w}_b^{in} for some permutation w_b^{in} , together with a distinguished subexpression \mathbf{v}_b^{in} for some permutation v_b^{in} . Similarly, by using the positions corresponding to boxes of λ_b^{out} , we obtained $\mathbf{w}_b^{\text{out}}$, w_b^{out} , $\mathbf{v}_b^{\text{out}}$, and v_b^{out} . When the box b is understood, we will often omit the subscript b .

For any box b , note that it is always possible to choose a reading order of $\lambda = \lambda_w$ which orders all the boxes of λ^{out} after those of λ^{in} . We can then adjust \mathbf{w} accordingly; this does not affect whether the corresponding expression \mathbf{v} is distinguished. Having chosen such a reading order, we can then write $\mathbf{w} = \mathbf{w}^{\text{in}}\mathbf{w}^{\text{out}}$ and $\mathbf{v} = \mathbf{v}^{\text{in}}\mathbf{v}^{\text{out}}$. We then use g^{in} and g^{out} to denote the corresponding factors of $g \in G_{\mathbf{v}, \mathbf{w}}$. We define $J_{\mathbf{v}^{\text{out}}}^+$ to be the subset of $J_{\mathbf{v}}^+$ coming from the factors of \mathbf{v} contained in \mathbf{v}^{out} . Similarly, for $J_{\mathbf{v}^{\text{out}}}^{\circ}$ and $J_{\mathbf{v}^{\text{out}}}^{\bullet}$.

Example 7.4. Let $W = \mathfrak{S}_7$ and $\mathbf{w} = s_4s_5s_2s_3s_4s_6s_5s_1s_2s_3s_4$ be a reduced expression for $w \in W^3$. Let $\mathbf{v} = s_4s_511s_41s_5s_111s_4$ be a distinguished subexpression. So $w = (3, 5, 6, 7, 1, 2, 4)$ and $v = (2, 1, 3, 4, 6, 5, 7)$. We can represent this data by the poset λ_w and the corresponding Go-diagram:



Let b be the box of the Young diagram which is in the second row and the second column (counting from left to right). Then the diagram below shows: the boxes of λ^{in} and λ^{out} ; a reading order which puts the boxes of λ^{out} after those of λ^{in} ; and the corresponding labeled Go-diagram. Using this reading order, $\mathbf{w}^{\text{in}} = s_4s_5s_2s_3s_4$, $\mathbf{w}^{\text{out}} = s_6s_5s_1s_2s_3s_4$, $\mathbf{v}^{\text{in}} = s_4s_511s_4$, and $\mathbf{v}^{\text{out}} = 1s_5s_111s_4$.



Note that $J_{\mathbf{v}^{\text{out}}}^{\bullet} = \{7, 11\}$ and $J_{\mathbf{v}^{\text{out}}}^+ = \{6, 9, 10\}$. Then $g \in G_{\mathbf{v}, \mathbf{w}}$ has the form

$$g = g^{\text{in}} g^{\text{out}} = (\dot{s}_4 \dot{s}_5 y_2(p_3) y_3(p_4) \dot{s}_4) (y_6(p_6) x_5(m_7) \dot{s}_5^{-1} \dot{s}_1 y_2(p_9) y_3(p_{10}) x_4(m_{11}) s_4^{-1}).$$

When we project the resulting 7×7 matrix to its first three columns, we get the matrix

$$A = \begin{pmatrix} -p_9 p_{10} & -p_3 p_{10} & -p_{10} & -m_{11} & 0 & -1 & 0 \\ 0 & -p_3 p_4 & -p_4 & -m_7 & 1 & 0 & 0 \\ 0 & 0 & 0 & p_6 & 0 & 0 & 1 \end{pmatrix}$$

Theorem 7.5. [6, Theorem 5.6] Let $\mathbf{w} = s_{i_1} \dots s_{i_m}$ be a reduced expression for $w \in W^k$ and $\mathbf{v} \prec \mathbf{w}$, and let D be the corresponding Go-diagram. Choose any box b of D , and let $v^{\text{in}} = v_b^{\text{in}}$ and $w^{\text{in}} = w_b^{\text{in}}$, and $v^{\text{out}} = v_b^{\text{out}}$ and $w^{\text{out}} = w_b^{\text{out}}$. Let $A = \pi_k(g)$ for any $g \in G_{\mathbf{v}, \mathbf{w}}$, and let $I = w\{n, n-1, \dots, n-k+1\}$. If b contains a $+$, define $I_b = v^{\text{in}}(w^{\text{in}})^{-1} I \in \binom{[n]}{k}$. If b contains a white or black stone, define $I_b = v^{\text{in}} s_b (w^{\text{in}})^{-1} I \in \binom{[n]}{k}$. If we write $g = g_1 \dots g_m$ as in Definition 4.1, then

- (1) If b contains a $+$, then $\Delta_{I_b}(A) = (-1)^{|J_{\mathbf{v}^{\text{out}}}^{\bullet}|} \prod_{i \in J_{\mathbf{v}^{\text{out}}}^+} p_i$.
- (2) If b contains a white stone, then $\Delta_{I_b}(A) = 0$.
- (3) If b contains a black stone, then $\Delta_{I_b}(A) = (-1)^{|J_{\mathbf{v}^{\text{out}}}^{\bullet}|+1} m_b \prod_{i \in J_{\mathbf{v}^{\text{out}}}^+} p_i + \Delta_{I_b}(A_b)$, where m_b is the parameter corresponding to b , and A_b is the matrix A with $m_b = 0$.

Example 7.6. We continue Example 7.4. By Theorem 7.1, $I = w\{5, 6, 7\} = \{1, 2, 4\}$ and $I' = v\{5, 6, 7\} = \{5, 6, 7\}$, and the lexicographically minimal and maximal nonzero Plücker coordinates for A are $\Delta_I(A) = p_3 p_4 p_6 p_9 p_{10}$ and $\Delta_{I'}(A) = 1$; this can be verified for the matrix A above.

We now verify Theorem 7.5 for the box b labeled 5 in the reading order. Then $I_b = v^{\text{in}}(w^{\text{in}})^{-1} I = \{1, 4, 6\}$. Theorem 7.5 says that $\Delta_{I_b}(A) = 0$, since this box contains a white stone. The analogous computations for the boxes labeled 7, 6, 4, 3, 2, 1, respectively, yield $\Delta_{1,5,7} = -p_9 p_{10}$, $\Delta_{1,2,7} = p_3 p_4 p_9 p_{10}$, $\Delta_{1,4,5} = p_6 p_9 p_{10}$, $\Delta_{1,3,4} = p_4 p_6 p_9 p_{10}$, $\Delta_{1,2,4} = p_3 p_4 p_6 p_9 p_{10}$, and $\Delta_{1,2,4} = p_3 p_4 p_6 p_9 p_{10}$. These can be checked for the matrix A above.

Proposition 7.7. [6, Corollary 5.11] For any box b , the rescaled Plücker coordinate

$$\frac{\Delta_{I_b}(A)}{\prod_{i \in J_{\mathbf{v}^{\text{out}}}^+} p_i}$$

depends only on the parameters $p_{b'}$ and $m_{b'}$ which correspond to boxes b' weakly southeast of b in the Go-diagram.

7.2. The characterization of Deodhar components in terms of minors. Given a Go-diagram D of shape λ , contained in a $k \times (n-k)$ rectangle, let $I = I(\lambda)$. It is not hard to check that if D corresponds to the distinguished subexpression \mathbf{v} of the reduced expression \mathbf{w} , then $I = w\{n, n-1, \dots, n-k+1\}$.

Theorem 7.8. Let D be a Go-diagram of shape λ contained in a $k \times (n-k)$ rectangle. Let $A \in Gr_{k,n}$. Then A lies in the Deodhar component \mathcal{R}_D if and only if the following conditions are satisfied:

- (1) $\Delta_{I_b}(A) = 0$ for all boxes in D containing a white stone.
- (2) $\Delta_{I_b}(A) \neq 0$ for all boxes in D containing a $+$.
- (3) $\Delta_{I(\lambda)}(A) \neq 0$.
- (4) $\Delta_J(A) = 0$ for all k -subsets J which are lexicographically smaller than $I(\lambda)$.

Proof. Suppose that A lies in the Deodhar component \mathcal{R}_D . Then by Theorem 7.1, conditions (3) and (4) hold. And by Theorem 7.5, conditions (1) and (2) hold.

Now suppose that $A \in Gr_{k,n}$, and conditions (1)-(4) hold. We want to show that $A \in \mathcal{R}_D$. Since the Deodhar components partition $Gr_{k,n}$, it suffices to show that A cannot lie in $\mathcal{R}_{D'}$ for any other Go-diagram D' . For the sake of contradiction, assume that $A \in \mathcal{R}_{D'}$. Then by conditions (3) and (4)

above, and Theorem 7.1, it follows that D and D' must be Go-diagrams of the same shape. Therefore D and D' correspond to distinguished subexpressions \mathbf{v} and \mathbf{v}' of the same reduced expression \mathbf{w} .

Choose a reading order for the boxes of the Go-diagrams D and D' , and let b be the first box in that order where D and D' differ. Then without loss of generality, in D the box b must contain a $+$, and in D' the box b must contain a stone. (Because \mathbf{v} and \mathbf{v}' are distinguished subexpressions of the same reduced word \mathbf{w} , which agree in the first ℓ factors, and differ in the $(\ell + 1)$ st factor – therefore one of \mathbf{v} and \mathbf{v}' must use the $(\ell + 1)$ st simple generator s_b , and one must omit it.) In fact it follows from the definition of distinguished subexpression, and the fact that D corresponds to a distinguished subexpression, that the box b in D' must contain a white stone, not a black one. (When building a distinguished subexpression from left to right, if choosing the next simple generator s_b would decrease the length of the word so far, then one must choose s_b .)

Now note that the minor which Theorem 7.5 associates to box b in D is Δ_{I_b} , where $I_b = v^{\text{in}}(w^{\text{in}})^{-1}(I)$, and the minor which the theorem associates to box b in D' is $\Delta_{I'_b}$, where $I'_b = v'^{\text{in}}s_b(w^{\text{in}})^{-1}(I)$. But now note that $v'^{\text{in}}s_b = v^{\text{in}}$. Therefore $I_b = I'_b$. So then by Theorem 7.5, if $A \in \mathcal{R}_{D'}$, then $\Delta_{I_b}(A) = 0$, while by condition (2), $\Delta_{I_b}(A) \neq 0$. This is a contradiction. \square

Corollary 7.9. The Deodhar decomposition of the Grassmannian is a coarsening of the matroid stratification: in other words, each Deodhar component is a union of matroid strata.

Proof. Each matroid stratum is defined by specifying that certain Plücker coordinates are nonzero while the rest are zero. Therefore the corollary is an immediate consequence of Theorem 7.8. \square

Remark 7.10. Theorem 7.8 implicitly gives an algorithm for determining the Deodhar component and corresponding network of a point of the Grassmannian, given by a matrix representative or by a list of its Plücker coordinates. The steps are as follows.

- (1) Find the lexicographically minimal nonzero Plücker coordinate Δ_I . Then the Go-diagram has shape $\lambda(I)$. Fix a reading order for this shape.
- (2) We determine how to fill each box, working in the reading order, as follows. First check whether the box b is forced to contain a black stone. If so, proceed to the next box. If not, look at Δ_{I_b} . If this Plücker coordinate is zero, b must contain a white stone, and if it is nonzero, b must contain a $+$. Proceed to the next box. This process will completely determine the Go-diagram.
- (3) Given the Go-diagram, we know what the underlying graph of the network must be. To determine the weights on horizontal edges, work through them in the reading order again. The Plücker coordinate Δ_{I_b} will only use edge weights a_b (when b contains a $+$) or c_b (when b contains a black stone) and weights $a_{b'}$ and $c_{b'}$ corresponding to boxes b' which are earlier than b in the reading order. Thus, we may use the Lindström-Gessel-Viennot Lemma recursively to determine each weight a_b or c_b .

REFERENCES

- [1] A. Bjorner and F. Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, 231, Springer, New York, 2005.
- [2] V. Deodhar, On some geometric aspects of Bruhat orderings. I. A finer decomposition of Bruhat cells, *Invent. Math.* 79 (1985), no. 3, 499–511.
- [3] V. Deodhar, On some geometric aspects of Bruhat orderings II. The parabolic analogue of Kazhdan-Lusztig polynomials, *J. Alg.* 111 (1987), 483–506.
- [4] D. Kazhdan and G. Lusztig, Representations of Coxeter groups and Hecke algebras, *Invent. Math.* 53 (1979), no. 2, 165–184.
- [5] D. Kazhdan and G. Lusztig, *Schubert varieties and Poincaré duality*, Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), Proc. Sympos. Pure Math., XXXVI, Amer. Math. Soc., Providence, R.I., 1980, pp. 185–203. MR 84g:14054
- [6] Y. Kodama, L. Williams, The Deodhar decomposition of the Grassmannian and the regularity of KP solitons, arXiv:1204.6446.

- [7] G. Lusztig, Total positivity in partial flag manifolds, *Representation Theory*, 2 (1998) 70-78.
- [8] R. Marsh and K. Rietsch, Parametrizations of flag varieties, *Representation Theory*, 8 (2004).
- [9] A. Postnikov, Total positivity, Grassmannians, and networks, <http://front.math.ucdavis.edu/math.CO/0609764>.
- [10] R. Proctor, Bruhat lattices, plane partition generating functions, and minuscule representations. *European J. Combin.* 5 (1984), no. 4, 331–350.
- [11] J. Stembridge, On the Fully Commutative Elements of Coxeter Groups. *J. Algebraic Combinatorics* 5 (1996), 353–385.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720-3840
E-mail address: `talaska@math.berkeley.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720-3840
E-mail address: `williams@math.berkeley.edu`