

## Quiz 5: Differentiation Rules

Your name:

Discussions 201, 203 // 2018-10-12

**Problem 1** (5 points). Differentiate the function  $f(x) = \sqrt{e^{-x} + 2}$ .

*Solution:* Apply the power rule and the chain rule (twice):

$$\begin{aligned} f'(x) &= \frac{1}{2}(e^{-x} + 2)^{-1/2} \frac{d}{dx}(e^{-x} + 2) \\ &= \frac{e^{-x} \frac{d}{dx}(-x)}{2\sqrt{e^{-x} + 2}} \\ &= \frac{-1}{2e^x \sqrt{e^{-x} + 2}}. \end{aligned}$$

□

**Problem 2** (1 + 1 + 3 = 5 points). Consider the function  $y = x^{10} + \sin(x)$ . Find each of the following.

(a)  $\frac{dy}{dx} = 10x^9 + \cos(x)$

(b)  $\frac{d^2y}{dx^2} = 90x^8 - \sin(x)$

(c)  $\frac{d^{99}y}{dx^{99}} = -\cos(x)$

**Problem 3** (5 points). Find values of  $a$  and  $b$  that make the below function differentiable at  $t = 0$ .

$$f(t) = \begin{cases} te^t + 3 \tan t & \text{for } t \leq 0 \\ \frac{at + b}{t + 1} & \text{for } t > 0. \end{cases}$$

*Solution:* First, we need  $b = 0$  in order for the function to be continuous:

$$\begin{aligned} \lim_{t \rightarrow 0^+} f(t) &= b \\ f(0) &= 0e^0 + 3 \tan 0 = 0. \end{aligned}$$

We also need the derivatives of the two pieces to match up at  $t = 0$ .

$$\begin{aligned} \frac{d}{dt}(te^t + 3 \tan t) &= te^t + e^t + 3 \sec^2(t), \\ \frac{d}{dt}\left(\frac{at}{t+1}\right) &= \frac{a(t+1) - at}{(t+1)^2} \\ &= \frac{a}{(t+1)^2}. \end{aligned}$$

Equating these at  $t = 0$  gives the condition

$$\begin{aligned} 0e^0 + e^0 + 3 \sec^2(0) &= \frac{a}{1^2} \\ 4 &= a \end{aligned}$$

so  $a = 4$  and  $b = 0$ .

□