

CLASSICAL–QUANTUM CORRESPONDENCE IN LINDBLAD EVOLUTION

JEFFREY GALKOWSKI*, AND MACIEJ ZWORSKI

WITH AN APPENDIX BY ZHEN HUANG AND MACIEJ ZWORSKI

ABSTRACT. We show that for the Lindblad evolution defined using (at most) quadratically growing classical Hamiltonians and (at most) linearly growing classical jump functions (quantized into jump operators assumed to satisfy certain ellipticity conditions and modeling interaction with a larger system), the evolution of a quantum observable remains close to the classical Fokker–Planck evolution in the Hilbert–Schmidt norm for times vastly exceeding the Ehrenfest time (the limit of such agreement with no jump operators). The time scale is the same as in the recent papers by Hernández–Ranard–Riedel but the statement and methods are different. The appendix presents numerical experiments illustrating the classical/quantum correspondence in Lindblad evolution and comparing it to the mathematical results.

1. INTRODUCTION

In quantum mechanics a system is often described using a *density matrix*, that is a positive operator of trace one on a Hilbert space. In this paper the Hilbert space will be given by $L^2(\mathbb{R}^n)$ so that the density operator is then

$$Au(x) = \sum_j p_j \langle u, u_j \rangle u_j(x), \quad p_j \geq 0, \quad \sum_j p_j = 1, \quad \langle u_j, u_i \rangle = \delta_{ij}.$$

If the system evolves according to the Schrödinger equation $(ih\partial_t + P)v(t) = 0$, where P is a self-adjoint unbounded operator on $L^2(\mathbb{R}^n)$ then (note the sign convention) the density matrix evolves by the Schrödinger propagation of u_j 's. That gives the following equation:

$$\partial_t A(t) = \mathcal{L}_0 A(t), \quad \mathcal{L}_0 A := \frac{i}{h}[P, A], \quad A(t) = e^{t\mathcal{L}_0} A(0) = e^{itP/h} A(0) e^{-itP/h}. \quad (1.1)$$

This evolution clearly preserves density matrices. Gorini–Kossakowski–Sudarshan [GKS76] and Lindblad [Li76] generalized this by showing (in the setting of matrices and of bounded operators, respectively) that semigroups preserving the trace and

*Corresponding author: j.galkowski@ucl.ac.uk.

complete positivity¹ are generated by operators of the form

$$\mathcal{L}A := \frac{i}{\hbar}[P, A] + \frac{\gamma}{\hbar} \sum_{j=1}^J (L_j A L_j^* - \frac{1}{2}(L_j^* L_j A + A L_j^* L_j)), \quad \gamma \geq 0. \quad (1.2)$$

The corresponding evolution equation is called the Lindblad master equation or the GKLS equation and we refer to \mathcal{L} as the Lindbladian – see [ChPa17] for a history of this discovery and pointers to the literature. The operators L_j are called jump operators and they describe a dissipative (see (1.4) below) interaction of a system evolving according to (1.1) with a larger “open” system. (Hence the term “jump” as L_j describe the effect of moving to that larger system.) In general, J is allowed to be infinite, but in this paper we consider the case of $J < \infty$. The scaling γ/\hbar in front of the dissipative term corresponds to the $1/\hbar$ scaling in front $[P, A]$: the dissipative term can be written as a commutator – see (5.1). Then in general, in the notation of (2.1), $\frac{i}{\hbar}[a^w(x, \hbar D), b^w(x, \hbar D)] = \{a, b\}^w(x, \hbar D) + \mathcal{O}(\hbar)$. Hence having a factor of γ/\hbar is natural with γ measuring the strength of the interaction with the larger system.

1.1. Assumptions on P and L_j and Fokker–Planck evolution. In this paper we will consider (1.2) with P and L_j 's given by *pseudodifferential operators* (see (2.1) for the notation $a^w(x, \hbar D)$), that is semiclassical quantizations of classical observables, satisfying the following assumptions:

$$\begin{aligned} P &= p^w(x, \hbar D), \quad |\partial^\alpha p(x, \xi)| \leq C_\alpha, \quad |\alpha| \geq 2, \quad p = \bar{p}, \\ L_j &= \ell_j^w(x, \hbar D), \quad |\partial^\alpha \ell_j(x, \xi)| \leq C_\alpha, \quad |\alpha| \geq 1, \quad 1 \leq j \leq J. \end{aligned} \quad (1.3)$$

If in (1.2), $A = a^w(x, \hbar D)$, then the leading part of the semiclassical expansion of $\mathcal{L}A$ (see the derivation in §5) is given by the action, Qa , of the following Fokker–Planck operator

$$Q := H_p + \frac{\gamma}{2i} \sum_{j=1}^J (2\{\ell_j, \bar{\ell}_j\} - \ell_j H_{\bar{\ell}_j} + \bar{\ell}_j H_{\ell_j}) + \frac{\hbar\gamma}{4} \sum_{j=1}^J (H_{\bar{\ell}_j} H_{\ell_j} + H_{\ell_j} H_{\bar{\ell}_j}). \quad (1.4)$$

Here, $H_f := \sum_{j=1}^n \partial_{\xi_j} f \partial_{x_j} - \partial_{x_j} f \partial_{\xi_j}$ is the Hamiltonian vector field of $f = f(x, \xi)$, and $\{f, g\} := H_f g$ is the Poisson bracket. We note that H_p is anti-selfadjoint with respect to the standard measure on $\mathbb{R}^n \times \mathbb{R}^n$. Since

$$\frac{1}{2i} (2\{\ell_j, \bar{\ell}_j\} - \ell_j H_{\bar{\ell}_j} + \bar{\ell}_j H_{\ell_j}) =: \frac{1}{i} \{\ell_j, \bar{\ell}_j\} + B_j, \quad B_j^* = -B_j, \quad (1.5)$$

¹A linear map $T : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ ($\mathcal{B}(\mathcal{H})$ denotes the set of bounded operators on a Hilbert space \mathcal{H}) is completely positive if for every $k \in \mathbb{N}$, $\text{Id}_k \otimes T : \mathbb{M}(k, \mathbb{C}) \otimes \mathcal{B}(\mathcal{H}) \rightarrow \mathbb{M}(k, \mathbb{C}) \otimes \mathcal{B}(\mathcal{H})$ maps positive operators to positive operators. Here $\mathbb{M}(k, \mathbb{C})$ denotes the space of $k \times k$ matrices with complex entries.




FIGURE 1. With P as in (B.1) and L_j as in (B.2), we show contour plots of the evolutions of $a(t, x, \xi)$ under the Fokker–Planck equation (1.4) and the evolution of $b(t, x, \xi)$, where $A(t) = b^w(t, x, hD_x)$ with $A(t)$ evolving under the Lindblad equation (1.2): i.e. b is the full Weyl symbol of the operator $A(t)$. The initial data is given by a coherent state (B.7) with $h_0 = h = 2^{-4}$. The breakdown of the classical/quantum correspondence is clearly visible when $\gamma = 0$ (Schrödinger evolution (1.1)) while a very good agreement is seen for $\gamma = 1$. For an animated version, see https://math.berkeley.edu/~zworski/Lin_vs_FP.mp4 and for a description of the numerical schemes, Appendix B.

the self-adjoint contribution to the second term is given by the real valued function

$$\mu := \frac{1}{2i} \sum_{j=1}^J \{\ell_j, \bar{\ell}_j\}. \quad (1.6)$$

It is interpreted as friction. Finally, the last term in (1.4) is self-adjoint and non-negative. Assumptions (1.3) show that μ is bounded ($\mu \in S(1)$ in the notation of §2).

Example. Suppose $J = 2n$ and $\ell_j = x_j$, $\ell_{j+n} = \xi_j$ for $j \leq n$. Then

$$Q = H_p + \frac{1}{2}\gamma h(\Delta_x + \Delta_\xi). \quad (1.7)$$

When $\gamma = 0$ (that is, when we consider (1.1)) classical quantum correspondence in the evolution is described using *Egorov's theorem* – see [Zw12, Theorem 11.12, §11.5] and references given there. Here we present it slightly differently, using the Hilbert–Schmidt norm of the operator – see Theorem 3 for a general version. For the evolution (1.1) with $A(0) = (2\pi h)^{n/2} a_0^w(x, hD)$ where $a_0 \in C_c^\infty(\mathbb{R}^{2n})$ is h -independent (so that $\|A(0)\|_{\mathcal{L}_2} = \|a_0\|_{L^2}$) we have

$$\|A(t) - \text{Op}((\exp tH_p)^* a_0)\|_{\mathcal{L}_2} \leq C e^{3\Gamma t} h^2, \quad (1.8)$$

where $\|\bullet\|_{\mathcal{L}_2}$ denotes the Hilbert–Schmidt norm and

$$\Gamma := \sup_{|\alpha|=2} \sup_{\mathbb{R}^{2n}} |\partial^\alpha p(x, \xi)|. \quad (1.9)$$

For a more precise version of Γ , under additional assumptions on p , in terms of Lyapunov exponents of the flow of H_p see [OlBo23, Appendix C] and references given there. For a relation between (1.9) and the flow see Lemma 3.1.

The estimate (1.8) is not optimal, but as $\|a^w(x, hD)\|_{\mathcal{L}_2} = (2\pi h)^{-n/2} \|a\|_{L^2(\mathbb{R}^{2n})}$, (1.8) indicates the basic principle that the agreement with classical evolution breaks down at times proportional to $\log(1/h)$, the Ehrenfest time.

Motivated by recent papers [HRR25, HRR23] by Hernández–Ranard–Riedel we consider the question of an agreement with classical evolution for much longer times: the quantum evolution is given by $e^{t\mathcal{L}}$ where \mathcal{L} is the Lindblad operator (1.2) and the classical evolution by e^{tQ} , where Q is the Fokker–Planck operator (1.4). The results are shown in Theorem 1 for the special case of h -independent symbols, and in Theorem 4 for the more general situation of initial condition in exotic symbol classes. We show that agreement holds in Hilbert–Schmidt norms. The main advantage lies in an easy characterization of Hilbert–Schmidt pseudodifferential operators and in the simplicity of L^2 estimates for the Fokker–Planck evolution defined using (1.4).

Remark. As was shown by Davies [Da77], the operator of the form (1.2) generates a positivity preserving contraction on the Banach space of self-adjoint trace class operators provided that

$$Y := iP - \frac{1}{2} \sum_{j=1}^J L_j^* L_j$$

is the infinitesimal generator of a strongly continuous one parameter contraction semi-group on $L^2(\mathbb{R}^n)$. In our case, this follows from the Hille–Yosida theorem and Proposition A.2 (see the proof of Proposition 4.6 for a similar argument with \mathcal{L}_2 playing the role of L^2).

As in [HRR25, HRR23] we make a strong non-degeneracy assumption:

$$\mathbf{HH}^* \geq cI_{\mathbb{C}^{2n}}, \quad \mathbf{H} := [H_{\ell_1}, \dots, H_{\ell_J}, H_{\bar{\ell}_1}, \dots, H_{\bar{\ell}_J}] \in M_{2n \times 2J}(\mathbb{C}). \quad (1.10)$$

This cumbersome looking condition corresponds to ellipticity of the second order operator appearing in the classical Fokker–Planck equation (1.4) corresponding to (1.2) – see example (1.7) and Remark 5 after Theorem 1. We also need a more technical condition

$$|\partial^\alpha \operatorname{Im} \ell_j| |\ell_j| + |\operatorname{Im} \ell_j| |\partial^\alpha \ell_j| \leq C_\alpha, \quad |\alpha| \geq 2. \quad (1.11)$$

1.2. Lindblad propagation for h -independent observables. With this notation in place we have a special case of Theorem 4 in §6:

Theorem 1. *Suppose that \mathcal{L} is given by (1.2), assumptions (1.3), (1.10), and (1.11) hold and $h^{\frac{1}{3}} \leq \gamma \leq 1$. If $a_0 \in C_c^\infty(\mathbb{R}^{2n})$ is h -independent and $A(t)$ satisfies*

$$\partial_t A(t) = \mathcal{L}A(t), \quad A(0) = (2\pi h)^{n/2} a_0^w(x, hD), \quad \|A(0)\|_{\mathcal{L}_2} = \|a_0\|_{L^2(\mathbb{R}^{2n})},$$

then for all $\varepsilon > 0$ there is $C > 0$ such that,

$$\|A(t) - a(t)^w(x, hD)\|_{\mathcal{L}_2} \leq C e^{(M_0 + C_0 h)\gamma t} t h^{\frac{1}{2}} \gamma^{-\frac{3}{2}} (1 + t \gamma^{\frac{3}{2}} h^{\frac{1}{2}})^\varepsilon, \quad (1.12)$$

where

$$(\partial_t - Q)a(t) = 0, \quad a(0) = (2\pi h)^{n/2} a_0, \quad M_0 := \sup \mu.$$

When $\mu \equiv 0$ (see (1.6)) then (1.12) improves to

$$\|A(t) - a(t)^w(x, hD)\|_{\mathcal{L}_2} \leq C e^{Ch^2 \gamma t} t h^{\frac{1}{2}} \gamma^{-\frac{3}{2}}. \quad (1.13)$$

Remarks. 1. When $L_j = L_j^*$ then the estimates above are valid for $\gamma \leq h^{-1}$ with additional factors of $1 + \gamma$ present. This follows from the examination of the proof of Proposition 5.1. We do not stress this point as, following [HRR25, HRR23], we are interested in the improvement in the classical/quantum correspondence for weak couplings to the larger system.

2. We have not investigated Egorov’s theorem (see §3) for the Lindblad evolution and for $\gamma < h^{\frac{1}{3}}$ our estimates are worse than (1.8). The optimality of the $h^{\frac{1}{3}}$ threshold is not clear (see Appendix B) and hence a better estimate than (1.8) may be possible.

3. When in (1.3) p is quadratic and ℓ_j ’s are linear than the agreement of the two evolutions is exact. This corresponds to the same phenomenon in the case of Egorov’s theorem – see [Zw12, Theorem 11.9].




FIGURE 2. A comparison of classical evaluations with and without damping in the setting of Theorem 1 with P as in (B.1) and L_j as in (B.2). We choose a fixed a_0 given by (B.7) with $h_0 = 2^{-3}$ and take $h = 2^{-5}$. More quantitative results are presented in Appendix B but the striking difference between the evolutions is clearly visible.

4. When $p(x, \xi)$ is confining (for instance $p(x, \xi) \geq |x|^2 + |\xi|^2$ and subharmonic outside of a compact set) then Proposition 7.4 and (7.11) show that in Example (1.7) (and most likely in greater generality), $\|a^w(t)\|_{\mathcal{L}_2} \geq \|a^w(0)\|_{\mathcal{L}_2}/C$ for $t \leq h^{-\nu}$, $\nu > 0$. That means that the estimates (1.12) and (1.13) are meaningful for long times.
5. To see the reason for the powers of h , γ , and t in (1.12) consider the simplest case given in (1.7). The classical (Fokker–Planck) evolution is then

$$(\partial_t - H_p - \varepsilon^2 \Delta_{x,\xi})a(t) = 0, \quad \varepsilon := \sqrt{\gamma h/2}, \quad a(t) = a(t, x, \xi).$$

The solutions satisfy the following estimate (immediate if $H_p = 0$), see Proposition 5.1 (see (5.7)):

$$\sum_{|\alpha| \leq k} \|(\varepsilon \partial_{x,\xi})^\alpha a(t)\|_{L_{x,\xi}^2} \leq C \sum_{|\alpha| \leq k} \|(\varepsilon \partial_{x,\xi})^\alpha a(0)\|_{L_{x,\xi}^2}. \quad (1.14)$$

The key fact is that there is *no* dependence on t – that is not the case for the evolution by H_p alone, see (3.7). The composition formula for pseudodifferential operators in Lemma 2.2 shows that $\mathcal{L}[a(t)^w(x, hD)] \equiv (Qa(t))^w(x, hD)$ modulo terms quantizing functions bounded by the size of $h^2\partial^3 a(t)$. These can be estimated using (1.14) where in the case of (1.16) and for $|\alpha| = 3$,

$$h^2 \|\partial_{x,\xi}^\alpha a(t)\|_{L_{x,\xi}^2} \leq Ch^2 \varepsilon^{-3} \sum_{|\alpha| \leq 3} \|(\varepsilon \partial_{x,\xi})^\beta a(0)\|_{L_{x,\xi}^2} \leq (2\pi h)^{\frac{n}{2}} C \gamma^{-\frac{3}{2}} h^{\frac{1}{2}}. \quad (1.15)$$

To get (1.21) we write

$$A(t) - a(t)^w(x, hD) = \int_0^t e^{(t-s)\mathcal{L}} (\mathcal{L}a(s)^w(x, hD) - (Qa(s))^w(x, hD)) ds,$$

which together with (1.15) and the fact that $\|A(0)\|_{\mathcal{L}_2} = \|a_0\|_{L^2}$, gives (1.13). The extra growth in (1.12) results from friction which is absent in this example. We used here the fact that in the example $e^{t\mathcal{L}}$ is a contraction in the Hilbert–Schmidt norm – in general there could be exponential growth produced by the friction term; this is reflected by the exponential prefactor in (1.12).

6. The class of operators P satisfying (1.3) includes Schrödinger operator whose classical dynamics exhibits chaotic behaviour. In that case one expects optimality of $t \sim \log(1/h)$ limit for classical–quantum correspondence for (1.1). For instance we could take

$$p(x, \xi) = \xi_1^2 + \xi_2^2 + x_1^2 + x_2^2 + \lambda(x)(x_1^2 x_2 - \frac{1}{3} x_2^3),$$

where $\lambda \in C_c^\infty(\mathbb{R}^2; [0, 1])$ and $\lambda = 1$ near 0.

7. Compared to the models used in the physics literature – see Unruh–Zurek [UnZu89] for the pioneering discussion of the classical/quantum correspondence for open systems – the ellipticity hypothesis (1.10) made in [HRR25] and here is too strong. Rather than (1.7), one should consider $\ell_j = x_j$, $0 \leq j \leq J = n$ so that the Fokker–Planck operators is given by $Q = H_p + \frac{1}{2}\gamma h \Delta_\xi$. This would require more subtle subelliptic estimates (see Smith [Sm20] for a recent treatment with an asymptotic parameter) than (1.14). Some recent advances are described in §1.4. Gong–Brumer [GoBr99] showed numerically that for such operators with chaotic classical dynamics for p , the classical/quantum correspondence persists for long times.

1.3. Lindblad propagation for mixtures of Gaussian states. We now state a special case of our theorem where we consider mixtures of Gaussian states in a sense similar to that in [HRR25]. For that we define the standard coherent states:

$$\psi_{(x_0, \xi_0)} = (2\pi h)^{-\frac{n}{4}} e^{-(x-x_0)^2/2h} e^{i(x-x_0)\xi_0/h}, \quad \|\psi_{(x_0, \xi_0)}\|_{L^2(\mathbb{R}^n)} = 1.$$

The corresponding density operator is

$$\begin{aligned} A_{(x_0, \xi_0)} u &:= \psi_{(x_0, \xi_0)} \langle u, \psi_{(x_0, \xi_0)} \rangle, \quad A_{(x_0, \xi_0)} = a_0^w(x, hD), \\ a_{(x_0, \xi_0)}(x, \xi) &= 2^n \exp\left(-\frac{1}{h} \left((x - x_0)^2 + (\xi - \xi_0)^2 \right)\right). \end{aligned} \quad (1.16)$$

We note that in our result the Gaussian $(2\pi h)^{-\frac{n}{4}} e^{-(x-x_0)^2/2h}$ could be replaced by $\alpha h^{-n/4} \psi((x-x_0)/\sqrt{h})$ where $\psi \in \mathcal{S}(\mathbb{R}^{2n})$ and $\alpha = 1/\|\psi\|_{L^2}$.

For a probability measure λ_h on $\mathbb{R}^n \times \mathbb{R}^n$ we define the mixture of Gaussian states:

$$\begin{aligned} A_{\lambda_h} u &:= \int \psi_{(x_0, \xi_0)} \langle u, \psi_{(x_0, \xi_0)} \rangle d\lambda_h(x_0, \xi_0), \quad A_{\lambda_h} = a_{\lambda_h}^w(x, hD), \\ a_{\lambda_h}(x, \xi) &= 2^n \int \exp\left(-\frac{1}{h} \left((x - x_0)^2 + (\xi - \xi_0)^2 \right)\right) d\lambda_h(x_0, \xi_0). \end{aligned} \quad (1.17)$$

We note that $\|A_{\lambda_h}\|_{\mathcal{L}_1} = 1$. For the Hilbert Schmidt norm we calculate

$$(2\pi h)^{-n} 2^{2n} \int e^{-\frac{1}{h}((x-x_0)^2 + (x-y_0)^2 + (\xi-\xi_0)^2 + (\xi-\eta_0)^2)} dx d\xi = e^{-\frac{1}{h}((x_0-y_0)^2 + (\xi_0-\eta_0)^2)},$$

so that

$$\|A_{\lambda_h}\|_{\mathcal{L}_2}^2 = \iint \exp\left(-\frac{1}{h} \left((x_0 - y_0)^2 + (\xi_0 - \eta_0)^2 \right)\right) d\lambda_h(x_0, \xi_0) d\lambda_h(y_0, \eta_0).$$

If $\lambda_h = \mu(x, \xi) dx d\xi$, where μ is smooth and h independent we are close to the case considered in Theorem 1 and $\|A_{\lambda_h}\|_{\mathcal{L}_2} \sim h^{n/2}$.

As in (1.8) when $\gamma = 0$, we obtain a version of Egorov's Theorem: for the solution of (1.1) with $A(0) = a_{\lambda_h}^w(x, hD)$,

$$\|A(t) - ((\exp tH_p)^* a_{\lambda_h})^w(x, hD)\|_{\mathcal{L}_2} \leq C e^{3\Gamma t} h^{\frac{1}{2}} \|A(0)\|_{\mathcal{L}_2}. \quad (1.18)$$

On the other hand, for the Lindblad evolution, the quantum–classical agreement lasts substantially longer as can be seen from the following special case of Theorem 4

Theorem 2. *Suppose that \mathcal{L} is given by (1.2), assumptions (1.3), (1.10), and (1.11) hold and $\gamma < h^{-1}$. If, in the notation of (1.17), $A(t)$ satisfies*

$$\partial_t A(t) = \mathcal{L}A(t), \quad A(0) = A_{\lambda_h}, \quad (1.19)$$

then for all $\varepsilon > 0$ there is $C > 0$ such that

$$\|A(t) - a(t)^w(x, hD)\|_{\mathcal{L}_2} \leq C e^{(M_0 + Ch)\gamma t} t (1 + \gamma^{-\frac{3}{2}}) h^{\frac{1}{2}} (1 + t\gamma^{\frac{3}{2}} h^{\frac{1}{2}})^\varepsilon \|A_{\lambda_h}\|_{\mathcal{L}_2}, \quad (1.20)$$

where

$$(\partial_t - Q)a(t) = 0, \quad a(0) = a_{\lambda_h}, \quad M_0 := \sup \mu.$$

When $\mu \equiv 0$ (see (1.6)) then (1.20) improves to

$$\|A(t) - a(t)^w(x, hD)\|_{\mathcal{L}_2} \leq C e^{Ch^2\gamma t} t (1 + \gamma^{-\frac{3}{2}}) h^{\frac{1}{2}} \|A_{\lambda_h}\|_{\mathcal{L}_2}, \quad (1.21)$$




FIGURE 3. The analogue of Figure 2 with the initial condition given by a coherent state (B.7) with $h_0 = h = 2^{-5}$. Again the improvement in the agreement of quantum and classical evolutions is striking.

Remark. The time scales appearing in Theorems 1, 2 and 4 agree with the time scales in [HRR25], as long as $\gamma \leq 1$: Theorem 3.1 there gives the bound $C \max(1, \gamma^{-\frac{3}{2}}) h^{\frac{1}{2}} t$ for a two tier comparison of evolution of specially constructed Gaussian states. Under the assumptions in Theorem 1 it reads as

$$\begin{aligned} \|A(t) - \tilde{a}(t)^w(x, hD)\|_{\mathcal{L}_1} &\leq C \max(1, \gamma^{-\frac{3}{2}}) h^{\frac{1}{2}} t, \\ h^{-n} \|\tilde{a}(t) - a(t)\|_{L^1(\mathbb{R}^{2n})} &\leq C \max(1, \gamma^{-\frac{3}{2}}) h^{\frac{1}{2}} t, \end{aligned} \tag{1.22}$$

where $\|\bullet\|_{\mathcal{L}_1}$ is the trace class norm. Remarkably, since the semigroup $e^{t\mathcal{L}}$ is contracting on trace class operators, there is no exponential growth even when friction is positive. The estimate does *not* provide a bound on $A(t) - a(t)^w(x, hD)$ in any norm, but has the following natural consequence [HRR25, (1.7)]:

$$\begin{aligned} \text{tr}(A(t)b^w(x, hD)) - (2\pi h)^{-n} \int_{\mathbb{R}^{2n}} a(t, x, \xi) b(x, \xi) dx d\xi \\ = \mathcal{O}(t \max(1, \gamma^{-\frac{3}{2}}) h^{\frac{1}{2}}) (\|b^w(x, hD)\|_{L^2 \rightarrow L^2} + \|b\|_{L^\infty}). \end{aligned}$$

This is (typically) stronger than the corresponding consequence of (1.20):

$$\begin{aligned} \operatorname{tr}(A(t)b^w(x, hD)) &- (2\pi h)^{-n} \int_{\mathbb{R}^{2n}} a(t, x, \xi)b(x, \xi)dx d\xi \\ &= \mathcal{O}(e^{(M_0+Ch)\gamma t} t(1 + \gamma^{-\frac{3}{2}})h^{\frac{1}{2}}(1 + t\gamma^{\frac{3}{2}}h^{\frac{1}{2}})^{+0})(2\pi h)^{-n/2}\|b\|_{L^2}. \end{aligned}$$

(Here and later we use the notation $a \leq b^{+0}$ if for all $\varepsilon > 0$ there is $C_\varepsilon > 0$ such that $a \leq C_\varepsilon b^\varepsilon$.) We stress, however, that Theorem 4 below applies to very general initial states $A(0)$ of which Gaussian states or their mixtures are an example. In addition, at the cost of further terms in the expansion, it gives approximation of the Lindblad evolution modulo $\mathcal{O}(e^{(M_0+Ch)\gamma t}(th^{1/2}\gamma^{-\frac{3}{2}})^N(1 + t\gamma^{\frac{3}{2}}h^{\frac{1}{2}})^{+0})$ for any N . For instance, when

$$\gamma = h^\delta, \quad \delta < \frac{1}{3}, \quad t \leq h^{-\nu}, \quad \nu < \frac{1}{2} \min(2\delta, 1 - 3\delta), \quad (1.23)$$

this gives an expansion modulo $\mathcal{O}(h^\infty)$.

This paper is self-contained except for some basic facts about semiclassical quantization from [Zw12, Chapter 4]. It is organized as follows. In §2 we review the definition of pseudodifferential operators and symbol classes. We introduce a new L^2 -based symbol class which is natural for the study of Hilbert–Schmidt operators, and show the properties of the corresponding pseudodifferential calculus. In §3 we present a variant of Egorov’s theorem with Hilbert–Schmidt norm and in §4 we prove mapping properties of $e^{t\mathcal{L}}$. §5 is then devoted to estimates on the Fokker–Planck evolution. A general result about agreement of classical and quantum dynamics in Hilbert–Schmidt norm is proved in §6. In §7, we consider situations where we can effectively control the Hilbert–Schmidt norm of the Lindblad evolution from below. Finally, in the Appendix A, we review some properties of pseudodifferential operators with quadratic symbol growth and in Appendix B we describe some numerical experiments on the Lindblad evolution.

1.4. Recent developments. Since this paper was posted two interesting works on the classical/quantum correspondence appeared. In [Li24] Li provided estimates on the *trace class* norm of the difference of the Lindblad evolution $A(t)$ and $\operatorname{Op}(a(t))$ where $a(t)$ evolves according to the Fokker–Planck equation (1.4). That was done using L^1 based symbol classes and precise L^1 derivative estimates for the Fokker–Planck evolution. The jump operators were assumed self-adjoint, $L_j = L_j^*$ and γ was in the range $1 \leq \gamma \leq h^{-1}$. The model case is given by (1.7).

Smith [Sm25] used the results of [Sm20] and the general strategy of this paper to obtain the same Hilbert–Schmidt estimates as here but under hypoellipticity rather than ellipticity assumptions. The basic example (matching examples considered in the physics literature – see [UnZu89],[GoBr99] and references given there) is given by (1.7) but with $L_j = x_j$, $j = 1, \dots, n$. This corresponds to replacing $\Delta_x + \Delta_\xi$ by Δ_ξ in (1.7) and imposing certain natural conditions on p (roughly so that $H_p + \Delta_\xi$ satisfies Hörmander’s bracket conditions). At the moment in addition, [Sm25] requires

that $p(x, \xi) = |\xi|^2 + |x|^2$ (quadratic and elliptic) for $|x| + |\xi| > R$ and that $L_j = L_j^*$. Removing the various conditions in the two papers constitutes an exciting challenge.

Acknowledgements. We would like to thank Simon Becker for pointing out reference [Da77] which clarified the trace class properties of $e^{t\mathcal{L}}$ and Zhenhao Li for helpful comments on early versions of this paper. We are also grateful to Felipe Hernández, Daniel Ranard and Jess Riedel for an illuminating discussion of the results and methods of [HRR25]. We would also like to thank the two referees for their careful reading and constructive comments.

JG acknowledges support from EPSRC grants EP/V001760/1 and EP/V051636/1 as well as Leverhulme Trust grant RPG-2023-325 and MZ from the NSF grant DMS-1901462 and the Simons Foundation under a “Moiré Materials Magic” grant. MZ would like to thank Paul Brumer for introducing him, many years ago in the context of discussing [GoBr99], to the problem of classical/quantum correspondence for Lindbladians. MZ is grateful to University College London for the generous hospitality during the writing of this paper.

2. SYMBOL SPACES AND QUANTIZATION

The operators introduced in §1 are defined using *pseudodifferential operators* which are obtained by a Weyl quantization process: at first for $a \in \mathcal{S}(\mathbb{R}_x^n \times \mathbb{R}_\xi^n)$ (here \mathcal{S} denotes Schwartz functions, that is functions u for which $x^\alpha \partial^\beta u$ are bounded for all multiindices α and β ; \mathcal{S}' denotes its dual, the space of tempered distributions – see [Zw12, Chapter 3]) we define

$$\text{Op}(a)u = a^w(x, hD, h)u := \frac{1}{(2\pi h)^n} \int a\left(\frac{x+y}{2}, \xi\right) e^{\frac{i}{h}\langle x-y, \xi \rangle} u(y) dy d\xi. \quad (2.1)$$

The Hilbert–Schmidt norm has a clean expression in terms of the symbol a (PDE parlance for classical observables):

$$\|\text{Op}(a)\|_{\mathcal{L}_2}^2 = \text{tr} \text{Op}(a) \text{Op}(a)^* = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^{2n}} |a(x, \xi)|^2 dx d\xi.$$

This is in contrast with the trace class norm which does not have an easy characterization in terms of a and its estimates require L^1 norm of derivatives of a – see [DiSj99, Chapter 9].

In this paper we consider different classes of symbols for which (2.1) remains valid and has interesting composition properties (as an operator $\text{Op}(a) : \mathcal{S} \rightarrow \mathcal{S}'$ the operator (2.1) is well defined for $a \in \mathcal{S}'(\mathbb{R}^{2n})$ – see [Zw12, Theorem 4.2]). We first recall the standard symbol class: for $m : \mathbb{R}^{2n} \rightarrow [0, \infty)$ satisfying $m(z)/m(w) \leq C(1 + |z - w|)^N$,

$$a \in S_\rho(m) \implies |\partial_z^\alpha a(z, h)| \leq C_\alpha h^{-\rho|\alpha|} m(z), \quad z = (x, \xi) \in \mathbb{R}^{2n}. \quad (2.2)$$

When $\rho = 0$ we write $S(m)$ and when $m = 1$, S_ρ .

The next class corresponds to the conditions in (1.3): for smooth function on \mathbb{R}^{2n} ,

$$u(z, h) \in S_{(k)} \iff |\partial_z^\alpha u(z, h)| \leq C_\alpha, \quad |\alpha| \geq k, \quad (2.3)$$

with constants C_α independent of h . The seminorms are given by the best constants C_α .

In dealings with Hilbert–Schmidt operators it is natural to consider symbols whose bounds are defined using L^2 norms. For smooth functions on $C^\infty(\mathbb{R}^{2n})$ depending on parameters h we define, for $0 \leq \rho < 1$,

$$a \in S_\rho^{L^2} \iff h^{-\frac{n}{2}} \|\partial_z^\alpha a\|_{L^2(\mathbb{R}^{2n})} \leq C_\alpha h^{-\rho|\alpha|}. \quad (2.4)$$

with the obvious seminorms. We note that the Sobolev embedding theorem and an interpolation argument show that $|\partial_z^\alpha a| \leq C'_\alpha h^{-\rho(|\alpha|+n+\delta)+\frac{n}{2}}$, for any $\delta > 0$. Hence for $\rho = 0$ the L^2 based spaces are contained in $h^{\frac{n}{2}}S(1)$ defined above, and in general

$$S_\rho^{L^2} \subset h^{-\rho(n+)+\frac{n}{2}}S_\rho(1). \quad (2.5)$$

It will also be useful to consider mixed spaces obtained by taking tensor products:

$$c(z, w) \in S \otimes S_\rho^{L^2} \iff h^{-\frac{n}{2}} \|\sup_z \partial_z^\alpha \partial_w^\beta c(z, \bullet)\|_{L^2} \leq C_{\alpha\beta} h^{-\rho|\beta|}, \quad z, w \in \mathbb{R}^{2n}. \quad (2.6)$$

We stress that we always demand that $0 \leq \rho < 1$.

Remark Another choice of the norm could be given by $\sup_z \|\partial_z^\alpha \partial_w^\beta c(z, \bullet)\|_{L^2}$ and both agree on products. The choice in definition (2.6) is motivated by the fact that $\|f(w, w)\|_{L_w^2} \leq \|\sup_z |f(z, w)|\|_{L_w^2}$ which does not work for the other choice.

For the properties of operators which are quantizations of $a \in S_\rho(m)$ see [Zw12, Chapter 4]. The same methods apply to operators obtained from $a \in S_{(k)}$ and are reviewed in the appendix. In particular we obtain spectral properties of operators quantizing $S_{(2)}$. Since the properties of $S_\rho^{L^2}$ and $0 \leq \rho < 1$ are more unusual we present them in this section. We start with

Lemma 2.1. *Suppose that $Q : \mathbb{R}^{2n} \times \mathbb{R}^{2n} \rightarrow \mathbb{R}$ is a non-degenerate bilinear quadratic form. Then, using definition (2.6),*

$$e^{ihQ(D_z, D_w)} : S \otimes S_\rho^{L^2} \rightarrow S \otimes S_\rho^{L^2}, \quad (2.7)$$

is continuous and for every N

$$e^{ihQ(D_z, D_w)} a - e^{\frac{i\pi}{4} \operatorname{sgn} Q} \sum_{k=0}^{N-1} \left(\frac{h}{i}\right)^k \frac{1}{k!} Q(D_z, D_w)^k a(z, w) \in h^{N(1-\rho)} S \otimes S_\rho^{L^2}, \quad (2.8)$$

where $\operatorname{sgn} Q$ is the signature of Q considered as a quadratic form on $\mathbb{R}^{2n} \times \mathbb{R}^{2n}$.

Proof. We denote by B the symmetric matrix corresponding to our quadratic form: $Q(\zeta, \omega) = \frac{1}{2}\langle B(\zeta, \omega), (\zeta, \omega) \rangle$. For $a \in S(1) \otimes S_\rho^{L^2} \subset h^{-\rho(n+)+\frac{n}{2}}S(1) \otimes S_\rho(1)$, hence the expression

$$c(z, w) := e^{ihQ(D_z, D_w)}a(z, w)$$

makes sense as an element in \mathcal{S}' (to see this, we apply e.g. [Zw12, Theorem 4.17] with for each fixed value of h) and by [Zw12, Theorem 4.8], for $a \in \mathcal{S}$,

$$c(z, w) = \frac{|\det B|^{-\frac{1}{2}}}{(2\pi h)^{2n}} \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h}\varphi(z_1, z_2)} a(z + z_1, w + w_1) dz_1 dw_1,$$

where

$$\varphi(z_1, w_1) = -\frac{1}{2}\langle B^{-1}(z_1, w_1), (z_1, w_1) \rangle.$$

Since $a \in h^{-\rho(n+)+\frac{n}{2}}S(1) \otimes S_\rho(1)$, this integral can be understood in the sense of oscillatory integrals and defines an element of \mathcal{S}' – see [Zw12, §3.6]. Recall also that oscillatory integrals allow for integrations by parts.

Set $v_1 = h^{-\rho}w_1$, and $\chi \in C_c^\infty(\mathbb{R}^{2n} \times \mathbb{R}^{2n})$ with $\chi \equiv 1$ near 0 and $\text{supp } \chi \subset B(0, 1)$. Then using the fact that $w_1 \mapsto \varphi(z_1, w_1)$ is linear, we obtain

$$\begin{aligned} c(z, w) &= \frac{|\det B|^{-\frac{1}{2}}}{(2\pi h^{1-\rho})^{2n}} \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h^{1-\rho}}\varphi(z_1, v_1)} a(z + z_1, w + h^\rho v_1) dz_1 dv_1, \\ &= \frac{|\det B|^{-\frac{1}{2}}}{(2\pi h^{1-\rho})^{2n}} \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h^{1-\rho}}\varphi(z_1, v_1)} \chi(z_1, v_1) a(z + z_1, w + h^\rho v_1) dz_1 dv_1 \\ &\quad + \frac{|\det B|^{-\frac{1}{2}}}{(2\pi h^{1-\rho})^{2n}} \int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h^{1-\rho}}\varphi(z_1, v_1)} (1 - \chi(z_1, v_1)) a(z + z_1, w + h^\rho v_1) dz_1 dv_1 \\ &=: c_1(z, w) + c_2(z, w) \end{aligned}$$

We start by considering c_1 . In this case, the integrand is compactly supported and we may apply the method of stationary phase [Zw12, Theorem 3.16 and Theorem 3.17]. That gives

$$\begin{aligned} &\left| \partial_z^{\alpha_1} \partial_w^{\alpha_2} \left(c_1(z, w) - e^{\frac{i\pi}{4} \text{sgn } B} \sum_{k=0}^{N-1} \left(\frac{h^{1-\rho}}{i} \right)^k \frac{1}{k!} \left(Q(D_{z_1}, D_{v_1})^k a(z + z_1, w + h^\rho v_1) \Big|_{z_1=v_1=0} \right) \right) \right| \\ &\leq C_N h^{(1-\rho)N} \sum_{|\beta_1|+|\beta_2| \leq 2N+4n+1} h^{-\rho|\alpha_2|} \sup_{|(z_1, v_1)| < 1} |\partial_{z_1}^{\beta_1+\alpha_1} \partial_{v_1}^{\beta_2+\alpha_2} a(z + z_1, w + h^\rho v_1)| \\ &=: C_N h^{(1-\rho)N} \sum_{|\beta_1|+|\beta_2| \leq 2N+4n+1} R_{\alpha\beta}(z, w), \end{aligned}$$

with the estimates on the remainder provided by Sobolev's embedding:

$$|R_{\alpha\beta}(z, w)|^2 \leq h^{-2\rho|\alpha_2|} \sum_{\gamma \leq 2n+1} \|\partial_{z_1}^{\beta_1+\alpha_1+\gamma_1} \partial_{v_1}^{\beta_2+\alpha_2+\gamma_2} a(z + \cdot, w + h^\rho \cdot)\|_{L^2(B_{\mathbb{R}^{2n}}(0,1))}^2.$$

Hence, with $B := B_{\mathbb{R}^{4n}}(0, 1)$,

$$\begin{aligned} \int_{\mathbb{R}^{2n}} \sup_z |R_{\alpha\beta}(z, w)|^2 dw &\leq h^{-2\rho|\alpha_2|} \sum_{\gamma \leq 2n+1} \int_{\mathbb{R}^{2n}} \int_B \sup_z |\partial_{(z_1, v_1)}^{\alpha+\beta+\gamma} a(z + z_1, w + h^\rho v_1)|^2 dz_1 dv_1 dw \\ &\leq \int_B h^{2\rho(|\gamma_2|+|\beta_2|)} \left\| \sup_z |\partial_{(z, w)}^{\alpha+\beta+\gamma} a(z, \cdot)| \right\|_{L^2}^2 dz_1 dv_1 \\ &\leq C \sum_{\gamma \leq 2n+1} h^{2\rho(|\gamma_2|+|\beta_2|)} \left\| \sup_z |\partial_{(z, w)}^{\alpha+\beta+\gamma} a(z, \cdot)| \right\|_{L^2}^2. \end{aligned}$$

In particular, this implies that

$$c_1(z, w) - e^{\frac{i\pi}{4} \operatorname{sgn} B} \sum_{k=0}^{N-1} \left(\frac{h^{1-\rho}}{i} \right)^k \frac{1}{k!} \left(Q(D_{z_1}, D_{v_1})^k a(z + z_1, w + h^\rho v_1) \right) \Big|_{v_1=z_1=0}$$

is in $h^{(1-\rho)N} S \otimes S_\rho^{L^2}$.

We now consider the remaining term in c , c_2 , and note that on $\operatorname{supp}(1 - \chi)$, $|\partial_{(z_1, v_1)} \varphi(z_1, v_1)| \geq c \langle (z_1, v_1) \rangle$. Hence, integration by parts (justified by the definition of the oscillatory integral) yields, for $N > 2n + 1$,

$$\begin{aligned} &h^{(\rho-1)(2N-4n)} \left\| \sup_z \partial_{(z, w)}^\alpha c_2(z, \cdot) \right\|_{L^2}^2 \\ &\leq C_N \int \sup_z \left(\iint \sum_{|\beta_1|+|\beta_2| \leq N} \langle (z_1, v_1) \rangle^{-N} \left| \partial_{(z, w)}^\alpha \left(\partial_z^{\beta_1} (h^\rho \partial_w)^{\beta_2} a(z + z_1, w + h^\rho v_1) \right) \right| dz_1 dv_1 \right)^2 dw \\ &\leq C_N \iiint \sum_{|\beta_1|+|\beta_2| \leq N} \langle (z_1, v_1) \rangle^{-2N+2n+1} \sup_z \left| \partial_{(z, w)}^\alpha \partial_z^{\beta_1} (h^\rho \partial_w)^{\beta_2} a(z + z_1, w + h^\rho v_1) \right|^2 dz_1 dv_1 dw \\ &\leq C_N \iint \sum_{|\beta_1|+|\beta_2| \leq N} \langle (z_1, v_1) \rangle^{-2N+2n+1} \left\| \sup_z \partial_{(z, w)}^\alpha \partial_z^{\beta_1} (h^\rho \partial_w)^{\beta_2} a(z, \cdot) \right\|_{L^2}^2 dz_1 dv_1 \\ &\leq C_N \sum_{|\beta_1|+|\beta_2| \leq N} \left\| \sup_z \partial_{(z, w)}^\alpha \partial_z^{\beta_1} (h^\rho \partial_w)^{\beta_2} a(z, \cdot) \right\|_{L^2}^2. \end{aligned}$$

Hence, we have $c_2 \in h^{(N-2n)(1-\rho)} S \otimes S_\rho^{L^2}$ for arbitrary N and $c \in S \otimes S_\rho^{L^2}$. The argument also shows that the map from a to c is continuous, and (2.8) holds. \square

We can write the composition law for operators in $S_\rho^{L^2}$ with $S_{(k)}$.

Lemma 2.2. *Let $0 \leq \rho < 1$, $k \geq 0$, $a \in S_{(k)}$, $b \in S_\rho^{L^2}$. Then,*

$$\operatorname{Op}(a) \operatorname{Op}(b) = \operatorname{Op}(c),$$

where c has the following expansion: for $N \geq k$,

$$c(x, \xi) - \sum_{j=0}^{N-1} \frac{1}{j!} \left(\frac{h}{2i} \sigma(D_x, D_\xi, D_y, D_\eta)^j a(x, \xi) b(y, \eta) \right) \Big|_{\substack{y=x \\ \eta=\xi}} \in h^{N(1-\rho)} S_\rho^{L^2}. \quad (2.9)$$

Proof. Writing $z = (x, \xi)$, $w = (y, \eta)$, we have

$$\text{Op}(a) \text{Op}(b) = \text{Op}(c), \quad c(z) := e^{ihA(D_{z,w})} a(z) b(w)|_{z=w},$$

where $A(D_{z,w}) := -\frac{1}{2}\sigma(D_x, D_\xi, D_y, D_\eta)$. By Taylor's formula

$$c(z, h) = \sum_{\ell=0}^{N-1} \frac{1}{\ell!} (ihA(D))^\ell (a(z)b(w))|_{z=w} + R_N(z, h)$$

where

$$R_N(z, h) : \frac{1}{(N-1)!} \int_0^1 (1-t)^{N-1} e^{ihtA(D)} (ihA(D))^N ((a(z)b(w))|_{z,w}) dt.$$

For $N \geq k$,

$$A(D_{z,w})^N a(z)b(w) \in h^{-N\rho} S \otimes S_\rho^{L^2}.$$

Hence, Lemma 2.1 applies and $e^{ihtA(D)} : S \otimes S_\rho^{L^2} \rightarrow S \otimes S_\rho^{L^2}$ has uniform bounds in $t \in [0, 1]$. Now, for $e \in S \otimes S_\rho^{L^2}$, we have

$$\|\partial_w^\alpha e(w, w)\|_{L^2} \leq C \sum_{|\beta| \leq |\alpha|} \|\partial_{(z,w)}^\beta e(z, w)|_{w=z}\|_{L_w^2} \leq C \sum_{|\beta| \leq |\alpha|} \|\sup_z |\partial_{(z,w)}^\beta e(z, \cdot)|\|_{L^2}.$$

We conclude that $R_N \in h^{(1-\rho)N} S_\rho^{L^2}$ which is (2.9). \square

3. EGOROV'S THEOREM REVISITED

We give a variant of Egorov's theorem which is analogous to Theorems 1 and 4 and uses propagation of quantum observables in symbol classes $S_\rho^{L^2}$ introduced in §2. In fact, the proof of Theorem 4 follows the same strategy with improved estimates coming from diffusion estimates: Lemma 3.1 below (see also (3.7)) is replaced by Proposition 5.1.

We start with a lemma relating the constant Γ in (1.9) to the properties of the flow (see [Zw12, Lemma 11.11] for a slightly different version)

Lemma 3.1. *Let $\varphi_t := \exp tH_p$ where p satisfies (1.3). Then*

$$\sup_{(x,\xi) \in \mathbb{R}^{2n}} |\partial^\alpha \varphi_t(x, \xi)| \leq C_\alpha e^{\Gamma|\alpha|t}, \quad \alpha \in \mathbb{N}^{2n}, \quad |\alpha| > 0. \quad (3.1)$$

In the proof of Lemma 3.1 we use the following version of Grönwall's inequality:

Lemma 3.2. *Let $\Gamma \in \mathbb{R}$ and suppose that $u : \mathbb{R} \rightarrow \mathbb{R}$ is continuous and satisfies*

$$u(t) \leq v(t) + \Gamma \int_0^t u(s) ds. \quad (3.2)$$

Then,

$$u(t) \leq v(t) + \Gamma \int_0^t e^{\Gamma(t-s)} v(s) ds, \quad t \geq 0.$$

Proof. Define $w(t) := \int_0^t u(s)ds$. Then, w is continuously differentiable and satisfies

$$w'(t) \leq v(t) + \Gamma w(t), \quad w(0) = 0.$$

Hence, conjugating by $e^{-\Gamma t}$ and integrating gives

$$w(t) \leq \int_0^t e^{\Gamma(t-s)} v(s) ds,$$

which, after substitution in (3.2), finishes the proof. \square

Proof of Lemma 3.1. The proof of (3.1) is an induction on $|\alpha|$. The first step is the case of $|\alpha| = 1$. Since $(d/dt)\varphi_t = H_p(\varphi_t)$,

$$\frac{d}{dt} (\partial^\alpha \varphi_t) = \partial H_p(\varphi_t) \partial^\alpha \varphi_t, \quad \partial^\alpha \varphi(0) = \alpha. \quad (3.3)$$

Since the entries of the matrix ∂H_p are bounded by Γ , integration gives

$$\sup_{\mathbb{R}^{2n}} |\partial^\alpha \varphi_t|_{\ell^\infty} \leq 1 + \Gamma \int_0^t \sup_{\mathbb{R}^{2n}} |\partial^\alpha \varphi_s|_{\ell^\infty} ds.$$

Lemma 3.2 then gives (3.1) for $|\alpha| = 1$.

Now assume $|\alpha| = \ell$ and suppose the estimate (3.1) is valid for all multiindices β with $1 \leq |\beta| < \ell$. We differentiate (3.3), to find

$$\frac{d}{dt} (\partial^\alpha \varphi_t) = \partial H_p(\varphi_t) \partial^\alpha \varphi_t + g(t), \quad (3.4)$$

where $g(t)$ is a sum of terms having the form

$$g_{\alpha\beta} \circ \varphi_t \partial^{\beta_1} \varphi_t \cdots \partial^{\beta_k} \varphi_t, \quad g_{\alpha\beta} \in S(1),$$

for $\beta_1 + \cdots + \beta_k = \alpha$ and $0 < |\beta_j| < |\alpha| = \ell$ ($j = 1, \dots, k$). The induction hypothesis implies $\sup_{\mathbb{R}^{2n}} |g(t)|_{\ell^\infty} \leq C e^{\Gamma|\alpha||t|}$. Integrating as above, we obtain

$$\sup_{\mathbb{R}^{2n}} |\partial^\alpha \varphi_t|_{\ell^\infty} \leq C e^{\Gamma|\alpha|t} + \Gamma \int_0^t \sup_{\mathbb{R}^{2n}} |\partial^\alpha \varphi_s|_{\ell^\infty} ds.$$

and we can use Lemma 3.2 to obtain (3.1). \square

Theorem 3. *Suppose that \mathcal{L}_0 is given by (1.1) with P satisfying (1.3) and $0 \leq \rho < \frac{2}{3}$. If $A(t)$ satisfies (in the notation of §2)*

$$\partial_t A(t) = \mathcal{L}_0 A(t), \quad A(0) = \text{Op}(a_0), \quad a_0 \in S_\rho^{L^2},$$

Then, for every N there exist $C_N > 0$ and $a(t) \in S_{\rho(t)}^{L^2}$ such that for Γ given by (1.9) and

$$\rho(t) := \rho + \frac{\Gamma t}{|\log h|} \leq \frac{2}{3}, \quad (3.5)$$

$a(t) - (\exp tH_p)^ a_0 \in h^{2-3\rho} e^{3\Gamma t} S_{\rho(t)}^{L^2}$ and*

$$\|A(t) - \text{Op}(a(t))\|_{\mathcal{L}_2} \leq C_N e^{3N\Gamma t} h^{N(2-3\rho)}. \quad (3.6)$$

Proof. We define

$$U_0(t)b := (\exp tH_p)^*b, \quad \partial_t U_0(t) = H_p U_0(t), \quad U_0(0) = I,$$

and note that using the definition (3.5) and Lemma 3.1 we have

$$U(t-s) : S_{\rho(s)}^{L^2} \rightarrow S_{\rho(t)}^{L^2}, \quad (3.7)$$

To construct $a(t)$ we start with $a_0(t) := U_0(t)a_0$ so that $a_0(t) \in S_{\rho(t)}^{L^2}$. Set $A_0(t) := \text{Op}(a_0(t))$. Then, using Lemma 2.2 we obtain

$$\dot{A}_0(t) = \text{Op}(\dot{a}_0(t)) = \text{Op}(H_p a_0(t)) = \mathcal{L}_0 A_0(t) + \text{Op}(e_0(t)), \quad e_0(t) \in h^{(2-3\rho)} e^{3\Gamma t} S_{\rho(t)}^{L^2}.$$

Suppose now that we found

$$a_j(t) \in h^{(2-3\rho)j} e^{3j\Gamma t} S_{\rho(t)}^{L^2} \quad j = 0, \dots, N-1$$

such that, with $A_{N-1} := \sum_{j=0}^{N-1} \text{Op}(a_j(t))$, we have

$$\dot{A}_{N-1} = \mathcal{L}_0 A_{N-1}(t) + \text{Op}(e_N(t)), \quad e_N(t) \in h^{(2-3\rho)N} e^{3N\Gamma t} S_{\rho(t)}^{L^2}.$$

Using e_N we define

$$a_N(t) := - \int_0^t U_0(t-s) e_N(s) ds, \quad \partial_t a_N = H_p a_N - e_N, \quad a_N(0) = 0.$$

Then, using (3.7),

$$a_N(t) \in h^{(2-3\rho)N} e^{3N\Gamma t} S_{\rho(t)}^{L^2},$$

and hence, with $A_N(t) = A_{N-1}(t) + \text{Op}(a_N(t))$, we have

$$\begin{aligned} \dot{A}_N(t) &= \mathcal{L}_0 A_{N-1}(t) + \text{Op}(e_N(t)) + \text{Op}(\dot{a}_N(t)) \\ &= \mathcal{L}_0 A_{N-1}(t) + \text{Op}(H_p a_N(t)) \\ &= \mathcal{L}_0 A_N(t) + \text{Op}(e_{N+1}(t)), \quad e_{N+1}(t) \in h^{(2-3\rho)(N+1)} e^{3(N+1)\Gamma t} S_{\rho(t)}^{L^2}. \end{aligned}$$

Note that in the last line we used Lemma 2.2 to obtain the estimates on e_{N+1} . This gives $a = \sum_{j \leq N} a_j$.

To compare $A_N(t) := \text{Op}(a(t))$ to $A(t)$, we use the fact that $e^{t\mathcal{L}_0}$ preserves the Hilbert–Schmidt norm (see (1.1)):

$$\|A(t) - A_N(t)\|_{\mathcal{L}_2} \leq \int_0^t \left\| e^{(t-s)\mathcal{L}_0} \text{Op}(e_{N+1}(s)) \right\|_{\mathcal{L}_2} ds \leq C_N h^{(2-3\rho)(N+1)} e^{3(N+1)\Gamma t}.$$

This completes the proof \square

4. THE SEMIGROUP GENERATED BY THE LINDBLADIAN.

We prove here that the Lindblad evolution is well defined in the space of Hilbert–Schmidt operators. This is done under the assumption (1.3) alone.

To describe the action of \mathcal{L} on operators $\mathcal{S} \rightarrow \mathcal{S}'$, we identify such operators with their Schwartz kernels in $\mathbb{R}^n \times \mathbb{R}^n$ and consider

$$\mathcal{L}_1 : \mathcal{S}'(\mathbb{R}^n \times \mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n \times \mathbb{R}^n), \quad \mathcal{L}_0 : \mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n). \quad (4.1)$$

More precisely, for $K \in \mathcal{S}'(\mathbb{R}^n \times \mathbb{R}^n)$, and $\chi \in \mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n)$ we denote by $K(\chi)$ the distributional pairing, formally equal to $\int K(x, y)\chi(x, y)dx dy$. Then for $A, B : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ we define, $(A \otimes B)K \in \mathcal{S}'(\mathbb{R}^n \times \mathbb{R}^n)$ by

$$(A \otimes B)K(\varphi \otimes \psi) := K(A^t \varphi \otimes B \psi), \quad \varphi, \psi \in \mathcal{S}(\mathbb{R}^n), \quad (\varphi \otimes \psi)(x, y) := \varphi(x)\psi(y),$$

where A^t is the transpose of A : for $f, g \in \mathcal{S}(\mathbb{R}^n)$, $(Af)(g) = f(A^t g)$ (this also defines the action of A on \mathcal{S}'). We note that if we identify the Schwartz kernels with operators then $(A \times B)K = AKB$.

In this notation

$$\begin{aligned} \mathcal{L}_1 := & \frac{i}{h} (\text{Op}(p) \otimes I - I \otimes \text{Op}(p)) + \frac{\gamma}{h} \sum_j ((\text{Op}(l_j) \otimes I)(I \otimes \text{Op}(\bar{l}_j))) \\ & - \frac{1}{2} \sum_j (\text{Op}(\bar{l}_j) \text{Op}(l_j) \otimes I + I \otimes \text{Op}(\bar{l}_j) \text{Op}(l_j)), \end{aligned} \quad (4.2)$$

and $\mathcal{L}_0 := \mathcal{L}_1|_{\mathcal{S}(\mathbb{R}^{2n})}$.

The following lemma describes \mathcal{L}_1 in a way that allows an application of Proposition A.2, which in turn provides the definition of \mathcal{L} as an unbounded operator on \mathcal{L}_2 .

Lemma 4.1. *The operator $\mathcal{L}_1 : \mathcal{S}'(\mathbb{R}^{2n}) \rightarrow \mathcal{S}'(\mathbb{R}^{2n})$ defined by (4.2) is given by $\mathcal{L}_1 = \text{Op}(L)$, where where $L = L(x, \xi, y, \eta) \in C^\infty(\mathbb{R}^{4n})$ satisfies*

$$|\partial^\alpha L| \leq C_\alpha(1 + |x| + |\xi| + |y| + |\eta|), \quad |\alpha| \geq 1. \quad (4.3)$$

Moreover, identifying the Hilbert–Schmidt class $\mathcal{L}_2(L^2(\mathbb{R}^n))$ with $L^2(\mathbb{R}^n \times \mathbb{R}^n)$ using Schwartz kernels, the Lindbladian \mathcal{L} with the domain

$$\mathcal{D}(\mathcal{L}) := \{A \in \mathcal{L}_2(L^2(\mathbb{R}^n)) : \mathcal{L}_1 A \in \mathcal{L}_2(L^2(\mathbb{R}^n))\}, \quad (4.4)$$

satisfies

$$\mathcal{L} = \overline{\mathcal{L}_0}, \quad \mathcal{L}^* = \overline{\mathcal{L}_0^*},$$

where $\mathcal{L}_0^* : \mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n)$ is the formal adjoint of \mathcal{L}_0 .

Proof. Using coordinates $((x, \xi), (y, \eta)) \in \mathbb{R}^{2n} \times \mathbb{R}^{2n}$ and denoting $\text{Op}_{\mathbb{R}^{2n}}$ the Weyl quantization on \mathbb{R}^{2n} , the definitions above show that

$$\text{Op}(a) \otimes I = \text{Op}_{\mathbb{R}^{2n}}(a(x, \xi)), \quad I \otimes \text{Op}(a) = \text{Op}_{\mathbb{R}^{2n}}(\tilde{a}(y, \eta)), \quad \tilde{a} := e^{i\langle hD_y, D_\eta \rangle} a.$$

(See [Zw12, Theorem 4.13]: if $\text{Op}(a) = \text{Op}_1(a_1)$, then $\text{Op}(\tilde{a}) = \text{Op}(a)^t = \text{Op}_0(a_1)$.) Consequently $a \in C^\infty(\mathbb{R}^{2n})$ satisfies, $|\partial_{(x,\xi)}^\alpha a(x, \xi)| \leq C_\alpha(1 + |x| + |\xi|)$ for $|\alpha| \geq 1$, and, by [Zw12, Theorem 4.17], so does \tilde{a} .

Since $\ell_j \in S_{(1)}$, by Proposition A.1 we have $\text{Op}(\bar{\ell}_j) \text{Op}(\ell_j) = \text{Op}(c_j)$, for c_j satisfying $|\partial_{(x,\xi)}^\alpha c_j(x, \xi)| \leq C_\alpha(1 + |x| + |\xi|)$ for $|\alpha| \geq 1$. Together with the facts that $\ell_j \in S_{(1)}$ and $p \in S_{(2)}$, this implies $\mathcal{L}_1 = \text{Op}_{\mathbb{R}^{2n}}(L)$, where L satisfies (4.3). Thus we can apply Proposition A.2 and the lemma follows. \square

The next lemma describe the adjoint of \mathcal{L} :

Lemma 4.2. *The adjoint of the Lindblad operator \mathcal{L} , \mathcal{L}^* , is given by*

$$\mathcal{L}^*B = -\frac{i}{\hbar}[P, B] + \frac{\gamma}{\hbar} \sum_j L_j^* B L_j - \frac{1}{2}(L_j^* L_j B + B L_j^* L_j), \quad (4.5)$$

with domain

$$\mathcal{D}(\mathcal{L}^*) = \{A \in \mathcal{L}_2 : \mathcal{L}^*A \in \mathcal{L}_2\},$$

where for any $A \in \mathcal{L}_2$, \mathcal{L}^*A is defined as an operator $\mathcal{S}' \rightarrow \mathcal{S}'$.

Proof. By Proposition A.2 it is sufficient to compute the formal adjoint in the action on operators $\mathcal{S}' \rightarrow \mathcal{S}$. Observe that, using cyclicity of the trace, for $A, B : \mathcal{S}' \rightarrow \mathcal{S}$,

$$\begin{aligned} \left\langle \frac{i}{\hbar}[P, A], B \right\rangle_{\mathcal{L}_2} &= \text{tr} \left(\frac{i}{\hbar}[P, A] B^* \right) = \frac{i}{\hbar} \text{tr} \left((PA - AP) B^* \right) = \frac{i}{\hbar} \text{tr} \left(A[P, B]^* \right) \\ &= \text{tr} \left(A \left(-\frac{i}{\hbar}[P, B] \right)^* \right) = \langle A, -\frac{i}{\hbar}[P, B] \rangle_{\mathcal{L}_2}, \\ \langle L_j A L_j^*, B \rangle_{\mathcal{L}_2} &= \text{tr} \left(L_j A L_j^* B^* \right) = \text{tr} \left(A L_j^* B^* L_j \right) = \langle A, L_j^* B L_j \rangle_{\mathcal{L}_2}, \\ \langle L_j^* L_j A, B \rangle_{\mathcal{L}_2} &= \text{tr} \left(L_j^* L_j A B^* \right) = \text{tr} \left(A B^* L_j^* L_j \right) = \langle A, L_j^* L_j B \rangle_{\mathcal{L}_2}, \end{aligned}$$

and similarly for $\langle A L_j^* L_j, B \rangle_{\mathcal{L}_2}$. \square

We next record some properties of \mathcal{L} and its adjoint.

Lemma 4.3. *For $A : \mathcal{S}' \rightarrow \mathcal{S}$,*

$$2 \text{Re} \langle \mathcal{L}A, A \rangle_{\mathcal{L}_2} = -\frac{\gamma}{\hbar} \sum_j \| [L_j, A] \|_{\mathcal{L}_2}^2 + \frac{\gamma}{\hbar} \left\langle \sum_j [L_j, L_j^*] A^*, A^* \right\rangle_{\mathcal{L}_2}, \quad (4.6)$$

and

$$2 \text{Re} \langle \mathcal{L}^*A, A \rangle_{\mathcal{L}_2} = -\frac{\gamma}{\hbar} \sum_j \| [L_j^*, A] \|_{\mathcal{L}_2}^2 + \frac{\gamma}{\hbar} \left\langle \sum_j [L_j, L_j^*] A, A \right\rangle_{\mathcal{L}_2}. \quad (4.7)$$

Proof. First, observe that (1.2) and (4.5) show

$$(\mathcal{L}A)^* = \mathcal{L}A^*, \quad (\mathcal{L}^*A)^* = \mathcal{L}^*A^*.$$

Thus, we compute

$$\begin{aligned}
2 \operatorname{Re} \langle \mathcal{L}A, A \rangle_{\mathcal{L}_2} &= \operatorname{tr}((\mathcal{L}A)A^* + A(\mathcal{L}A^*)) \\
&= \operatorname{tr} \left(\frac{i}{\hbar} [P, A]A^* + A \frac{i}{\hbar} [P, A^*] + \frac{\gamma}{\hbar} \sum_j \left(L_j A L_j^* A^* + A L_j A^* L_j^* \right) \right. \\
&\quad \left. - \frac{\gamma}{2\hbar} \sum_j \left(L_j^* L_j A A^* + A L_j^* L_j A^* + A L_j^* L_j A^* + A A^* L_j^* L_j \right) \right) \\
&= \operatorname{tr} \left(\frac{i}{\hbar} [P, A]A^* + A \frac{i}{\hbar} [P, A^*] \right) \\
&\quad + \frac{\gamma}{\hbar} \sum_j \operatorname{tr} \left(L_j A L_j^* A^* + A L_j A^* L_j^* - L_j^* L_j A A^* - A L_j^* L_j A^* \right).
\end{aligned}$$

Now,

$$\operatorname{tr} \left([P, A]A^* + A[P, A^*] \right) = \operatorname{tr} \left(P A A^* - A A^* P \right) = 0,$$

and

$$\begin{aligned}
&\operatorname{tr} \left(L_j A L_j^* A^* + A L_j A^* L_j^* - L_j^* L_j A A^* - A L_j^* L_j A^* \right) \\
&= \operatorname{tr} \left(- [L_j, A]([L_j, A])^* + L_j A A^* L_j^* + A L_j L_j^* A^* - L_j^* L_j A A^* - A L_j^* L_j A^* \right) \\
&= \operatorname{tr} \left(- [L_j, A]([L_j, A])^* + [L_j, L_j^*] A^* A \right).
\end{aligned}$$

Hence, (4.6) follows.

The computation for (4.7) is similar. Since the commutator part of \mathcal{L}^* has the same form as that of \mathcal{L} , we only need to compute

$$\begin{aligned}
&\operatorname{tr} \left(L_j^* A L_j A^* + A L_j^* A^* L_j - L_j^* L_j A A^* - A L_j^* L_j A^* \right) \\
&= \operatorname{tr} \left(- [L_j^*, A]([L_j^*, A])^* + L_j^* A A^* L_j - L_j^* L_j A A^* \right) \\
&= \operatorname{tr} \left(- [L_j^*, A]([L_j^*, A])^* + [L_j, L_j^*] A A^* \right),
\end{aligned}$$

and (4.7) follows. \square

The next lemma will be used to control the second terms on the right hand sides of (4.6) and (4.7).

Lemma 4.4. *Let $C_0 \in \mathbb{R}$ and suppose that $E : \mathcal{S} \rightarrow L^2$ is a self-adjoint operator on $L^2(\mathbb{R}^n)$ satisfying $E \leq C_0$. Then, for $B : \mathcal{S}' \rightarrow \mathcal{S}$,*

$$\left\langle EB, B \right\rangle_{\mathcal{L}_2} \leq C_0 \|B\|_{\mathcal{L}_2}^2. \quad (4.8)$$

Proof. To see this, observe that exists an L^2 -orthonormal basis u_j and $\lambda_j \geq 0$,

$$BB^* = \sum_j \lambda_j u_j \otimes u_j, \quad (f \otimes g)(\varphi) := f\langle \varphi, g \rangle.$$

We also note that if $\lambda_j > 0$ then $u_j \in \mathcal{S} \subset \mathcal{D}(E)$. Then,

$$\begin{aligned} \langle EB, B \rangle_{\mathcal{L}_2} &= \text{tr}(EBB^*) = \sum_j \langle EBB^* u_j, u_j \rangle_{L^2} = \sum_j \lambda_j \langle E u_j, u_j \rangle_{L^2} \\ &\leq C_0 \sum_j \lambda_j = C_0 \|B\|_{\mathcal{L}_2}^2, \end{aligned}$$

which is (4.8). \square

Next, we provide an estimate

Lemma 4.5. *Suppose that, as a bounded self-adjoint operator on $L^2(\mathbb{R}^n)$ (see (1.6))*

$$\sum_j [L_j, L_j^*] \leq \frac{2Mh}{\gamma}. \quad (4.9)$$

Then, for $A : \mathcal{S}' \rightarrow \mathcal{S}$ and $\lambda > 0$,

$$\lambda \|A\|_{\mathcal{L}_2} \leq \|(\mathcal{L} - M - \lambda)A\|_{\mathcal{L}_2}, \quad \lambda \|A\|_{\mathcal{L}_2} \leq \|(\mathcal{L}^* - M - \lambda)A\|_{\mathcal{L}_2}, \quad (4.10)$$

Proof. Observe that by Lemma 4.3, and Lemma 4.4

$$\begin{aligned} 2 \text{Re} \langle (\mathcal{L} - M - \lambda)A, A \rangle_{\mathcal{L}_2} &\leq -2\lambda \|A\|_{\mathcal{L}_2} - 2M \|A\|_{\mathcal{L}_2} + \frac{\gamma}{h} \left\langle \sum_j [L_j, L_j^*] A^*, A^* \right\rangle_{\mathcal{L}_2} \\ &\leq -2\lambda \|A\|_{\mathcal{L}_2}. \end{aligned}$$

Hence,

$$2\lambda \|A\|_{\mathcal{L}_2} \leq |2 \text{Re} \langle (\mathcal{L} - M - \lambda)A, A \rangle_{\mathcal{L}_2}| \leq 2 \|(\mathcal{L} - M - \lambda)A\|_{\mathcal{L}_2} \|A\|_{\mathcal{L}_2},$$

from which the first estimate in (4.10) follows. The argument for the second estimate is identical. \square

Proposition 4.6. *Suppose that (4.9) holds. Then the operator \mathcal{L} with domain $\mathcal{D}(\mathcal{L}) := \{A \in \mathcal{L}_2 : \mathcal{L}A \in \mathcal{L}_2\}$ generates a strongly continuous semigroup*

$$e^{t\mathcal{L}} : \mathcal{L}_2 \rightarrow \mathcal{L}_2 \quad \text{and} \quad \|e^{t\mathcal{L}}\|_{\mathcal{L}_2 \rightarrow \mathcal{L}_2} \leq e^{Mt}, \quad t \geq 0.$$

Proof. By Proposition A.2, or rather its proof (see (A.4)), and Lemma 4.1, for $A \in \mathcal{D}(\mathcal{L})$ there exists a sequence of operators $A_n : \mathcal{S}' \rightarrow \mathcal{S}$ such that $A_n \xrightarrow{\mathcal{L}_2} A$ and $\mathcal{L}A_n \xrightarrow{\mathcal{L}_2} \mathcal{L}A$. Hence, for $A \in \mathcal{D}(\mathcal{L})$ and $\lambda > 0$, Lemma 4.5 gives

$$\begin{aligned} \lambda \|A\|_{\mathcal{L}_2} &= \lambda \lim_{n \rightarrow \infty} \|A_n\|_{\mathcal{L}_2} \\ &\leq \lim_{n \rightarrow \infty} \|(\mathcal{L} - M - \lambda)A_n\|_{\mathcal{L}_2} = \|(\mathcal{L} - M - \lambda)A\|_{\mathcal{L}_2}. \end{aligned}$$

Similarly, for $A \in \mathcal{D}(\mathcal{L}^*)$, we have $A_n : \mathcal{S}' \rightarrow \mathcal{S}$ such that $A_n \xrightarrow{\mathcal{L}_2} A$ and $\mathcal{L}^* A_n \xrightarrow{\mathcal{L}_2} \mathcal{L}^* A$. This implies that for $A \in \mathcal{D}(\mathcal{L}^*)$, and $\lambda > 0$

$$\lambda \|A\|_{\mathcal{L}_2} \leq \|(\mathcal{L}^* - M - \lambda)A\|_{\mathcal{L}_2}.$$

In particular, $(\mathcal{L} - M - \lambda)^{-1} : \mathcal{L}_2 \rightarrow \mathcal{D}(\mathcal{L})$ exists and satisfies,

$$\|(\mathcal{L} - M - \lambda)^{-1}\|_{\mathcal{L}_2 \rightarrow \mathcal{L}_2} \leq \lambda^{-1}, \quad \lambda > 0.$$

The Hille-Yosida theorem then implies that $\mathcal{L} - M$ generates a strongly continuous semigroup $e^{t(\mathcal{L}-M)}$ satisfying

$$\|e^{t(\mathcal{L}-M)}\|_{\mathcal{L}_2 \rightarrow \mathcal{L}_2} \leq 1,$$

from which the proposition follows. \square

We conclude this section by showing how condition (4.9) is related to a lower bound on the friction (1.6)

Lemma 4.7. *Let*

$$M_0 := \sup_{\mathbb{R}^{2n}} \mu, \quad \mu := \frac{1}{2i} \sum_{j=1}^J \{\ell_j, \bar{\ell}_j\}. \quad (4.11)$$

Then there is $C_0 > 0$ such that (4.9) holds with

$$M = \gamma M_0 + C_0 h \gamma, \quad (4.12)$$

for $0 < h < 1$. Furthermore, if $\mu \equiv 0$, then (4.9) holds with

$$M = C_0 h^2 \gamma \quad (4.13)$$

for $0 < h < 1$.

Proof. Since (1.3) shows that $\mu \in S(1)$, the first estimate is a straightforward application of sharp Gårding inequality for the class $S(1)$ – see [DiSj99, Theorem 7.1] or [Zw12, §4.7.2]. When $\mu \equiv 0$, we use that $[L_j, L_j^*] = \text{Op}(\frac{h}{2i}\{\ell_j, \bar{\ell}_j\} + h^3 e)$ for some $e \in S(1)$ and hence the second estimate follows. \square

5. THE CLASSICAL DYNAMICS

It will be convenient to rewrite the Lindbladian as

$$\mathcal{L}A = \frac{i}{h}[P, A] + \frac{\gamma}{2h} \sum_j ([L_j A, L_j^*] + [L_j, A L_j^*]). \quad (5.1)$$

Our first goal is to motivate the classical Fokker–Planck equation (1.4) from the evolution equation for \mathcal{L} .

Observe that for $0 \leq \rho < 1$, and $a \in S_\rho^{L^2}$,

$$L_j A = \text{Op}(\ell_j a + \frac{h}{2i}\{\ell_j, a\} + h^{2-2\rho} e_1), \quad A L_j^* = \text{Op}(a \bar{\ell}_j + \frac{h}{2i}\{a, \bar{\ell}_j\} + h^{2-2\rho} e_2),$$

with $e_j \in S_\rho^{L^2}$. Hence for $a \in S_\rho^{L^2}$

$$\begin{aligned} \mathcal{L}A &= \text{Op}(H_p a) + \frac{\gamma}{2i} \sum_j \text{Op}((2\{\ell_j, \bar{\ell}_j\}a - \ell_j H_{\bar{\ell}_j} a + \bar{\ell}_j H_{\ell_j} a)) \\ &\quad + \frac{h\gamma}{4} \sum_j \text{Op}(H_{\bar{\ell}_j} H_{\ell_j} a + H_{\ell_j} H_{\bar{\ell}_j} a) + h^{2-3\rho}(1 + \gamma) \text{Op}(e), \end{aligned} \quad (5.2)$$

with $e \in S_\rho^{L^2}$. Heuristic arguments in the physics literature – see [HRR25] and the discussion and references given there – suggest that the natural classical evolution should be given by the equation up to the diffusion term $\sum_j H_{\bar{\ell}_j} H_{\ell_j} + H_{\ell_j} H_{\bar{\ell}_j}$ which is a non-positive differential operator acting on the classical observable a (see (1.7) for a striking example). Hence as the generator of the classical flow (a form of Fokker–Planck operator) we take $Q \in \text{Diff}^2(\mathbb{R}^{2n})$ given by

$$Q := H_p + \frac{\gamma}{2i} \sum_j (2\{\ell_j, \bar{\ell}_j\} - \ell_j H_{\bar{\ell}_j} + \bar{\ell}_j H_{\ell_j}) + \frac{h\gamma}{4} \sum_j (H_{\bar{\ell}_j} H_{\ell_j} + H_{\ell_j} H_{\bar{\ell}_j}).$$

The key estimate for evolution by Q is given as follows. We need here the additional technical assumption (1.11). To state the next estimate we recall the definition of semiclassical Sobolev norms:

$$\|u\|_{H_\varepsilon^s}^2 := \int (1 + |\varepsilon\zeta|^2)^s |\widehat{u}(\zeta)|^2 d\zeta, \quad \widehat{u}(\zeta) := \int u(z) e^{-iz\zeta} dz. \quad (5.3)$$

Proposition 5.1. *Suppose that (1.11) holds, and $0 \leq \gamma \leq 1$. Let $U(t) : L^2(\mathbb{R}^{2n}) \rightarrow L^2(\mathbb{R}^{2n})$ be defined by*

$$(\partial_t - Q)U(t) = 0, \quad U(0) = \text{Id}. \quad (5.4)$$

Then, for all $s \geq 0$, there is $C > 0$ such that for all $t \geq 0$,

$$\|U(t)\|_{H_\varepsilon^s \rightarrow H_\varepsilon^s} \leq C e^{M_0 \gamma t} (1 + t\gamma\varepsilon)^{+0}, \quad \|U(t)\|_{L^2 \rightarrow L^2} \leq C e^{M_0 \gamma t}, \quad (5.5)$$

where M_0 is given in (4.11), the norms are defined in (5.3), and

$$\varepsilon := \sqrt{\gamma h}. \quad (5.6)$$

If, $\sum_j \{\bar{\ell}_j, \ell_j\} \equiv 0$, that is there is no friction (1.6), then

$$\|U(t)\|_{H_\varepsilon^s \rightarrow H_\varepsilon^s} \leq C. \quad (5.7)$$

Remark. The estimates (5.5) and (5.7) do not address the smoothing effect of the evolution by (5.4). Obtaining quantitative estimates seems to require stronger assumptions than (1.3) and we restrict ourselves to that case.

Proof. Recall from (1.4) and (1.5) that Q is given by $H_p + \gamma \sum_j B_j + \mu$ plus a second order divergence form operator and the first two terms are anti-selfadjoint. Hence, for $u \in H^2$,

$$\operatorname{Re}\langle Qu, u \rangle = \frac{\gamma}{2i} \sum_j \langle \{\ell_j, \bar{\ell}_j\} u, u \rangle - \frac{h\gamma}{4} \sum_j (\|H_{\ell_j} u\|_{L^2}^2 + \|H_{\bar{\ell}_j} u\|_{L^2}^2).$$

We start with an estimate on the solution, v , to

$$e^{-tM_0\gamma}(\partial_t - Q)(e^{tM_0\gamma}v(t)) = (\partial_t - Q + M_0\gamma)v(t) = f, \quad v(0) = v_0. \quad (5.8)$$

We have

$$\begin{aligned} \langle f, v \rangle &= \operatorname{Re}\langle (\partial_t - Q + M_0\gamma)v, v \rangle \\ &= \frac{1}{2}\partial_t \|v\|_{L^2}^2 + \gamma \langle (M_0 - \sum_j \frac{1}{2i}\{\ell_j, \bar{\ell}_j\})v, v \rangle + \frac{h\gamma}{4} \sum_j (\|H_{\ell_j} v\|_{L^2}^2 + \|H_{\bar{\ell}_j} v\|_{L^2}^2). \end{aligned}$$

Hence,

$$\partial_t \|v\|_{L^2}^2 + \frac{h\gamma}{2} \sum_j (\|H_{\ell_j} v\|_{L^2}^2 + \|H_{\bar{\ell}_j} v\|_{L^2}^2) \leq 2|\langle f, v \rangle|.$$

For $T > 0$ the ellipticity hypothesis (1.10) then gives

$$\begin{aligned} \|v(T)\|_{L^2}^2 + \gamma hc \int_0^T \|\nabla v\|_{L^2}^2 dt &\leq \|v(T)\|_{L^2}^2 + \frac{h\gamma}{2} \int_0^T \sum_j (\|H_{\ell_j} v\|_{L^2}^2 + \|H_{\bar{\ell}_j} v\|_{L^2}^2) dt \\ &\leq 2 \int_0^T \|f(t)\|_{L^2} \|v(t)\|_{L^2} dt + \|v_0\|_{L^2}^2. \end{aligned} \quad (5.9)$$

Now let u solve

$$(\partial_t - Q + M_0\gamma)u = 0, \quad u(0) = u_0.$$

Then, applying (5.9), we obtain

$$\|u(T)\|_{L^2}^2 + c \int_0^T \|\varepsilon \nabla u(t)\|_{L^2}^2 ds \leq \|u_0\|_{L^2}^2, \quad \varepsilon = \sqrt{\gamma h}. \quad (5.10)$$

To proceed by induction let us assume that for $k \geq 0$

$$\begin{aligned} \sum_{0 \leq |\alpha| \leq k} \|(\varepsilon \partial)^\alpha u(T)\|_{L^2}^2 + \int_0^T \sum_{1 \leq |\alpha| \leq k+1} \|(\varepsilon \partial)^\alpha u(t)\|_{L^2}^2 \\ \leq C \sum_{|\beta| \leq k} \|(\varepsilon \partial)^\beta u_0\|_{L^2}^2 + CT\gamma\varepsilon \|u_0\|_{L^2}^2. \end{aligned} \quad (5.11)$$

We set

$$\begin{aligned} Q_1 &:= \sum_j i\{\bar{\ell}_j, \ell_j\}, & Q_2 &:= \frac{1}{2i} \sum_j (-\ell_j H_{\bar{\ell}_j} + \bar{\ell}_j H_{\ell_j}), \\ Q_3 &:= \frac{1}{4} \sum_j (H_{\bar{\ell}_j} H_{\ell_j} + H_{\ell_j} H_{\bar{\ell}_j}). \end{aligned}$$

so that

$$\begin{aligned} (\partial_t - Q + M_0)\partial^\alpha u &= [H_p, \partial^\alpha]u + \gamma[Q_1, \partial^\alpha]u + \gamma[Q_2, \partial^\alpha]u + \gamma h[Q_3, \partial^\alpha]u, \\ \partial^\alpha(0) &= \partial^\alpha u_0. \end{aligned} \tag{5.12}$$

We have the following estimates on the commutators appearing on the right hand side:

$$\begin{aligned} \|[H_p, \partial^\alpha]u\|_{L^2} &\leq C \sum_{1 \leq |\beta| \leq |\alpha|} \|\partial^\beta u\|_{L^2}, & \|[Q_1, \partial^\alpha]u\|_{L^2} &\leq C \sum_{0 \leq |\beta| \leq |\alpha|-1} \|\partial^\beta u\|_{L^2}, \\ \|[Q_2, \partial^\alpha]u\|_{L^2} &\leq C \sum_{1 \leq |\beta| \leq |\alpha|} \|\partial^\beta u\|_{L^2}, & \|[Q_3, \partial^\alpha]u\|_{L^2} &\leq C \sum_{1 \leq |\beta| \leq |\alpha|+1} \|\partial^\beta u\|_{L^2}. \end{aligned} \tag{5.13}$$

It is important here that in the estimates *not* involving Q_1 , we have $|\beta| \geq 1$ on the right hand sides. To obtain the estimate on commutators with Q_2 , we use assumption (1.11).

Applying (5.9) to (5.12) and using (5.13) we obtain

$$\begin{aligned} &\sum_{|\alpha|=k+1} \|\partial^\alpha u(T)\|_{L^2}^2 + c\gamma h \int_0^T \sum_{|\alpha|=k+1} \|\nabla \partial^\alpha u\|_{L^2}^2 dt \\ &\leq C \int_0^T \sum_{|\alpha|=k+1} \left(\sum_{1 \leq |\beta| \leq k+1} \|\partial^\beta u\|_{L^2} + \gamma \|u\|_{L^2} + \gamma h \sum_{1 \leq |\beta'| \leq k+2} \|\partial^{\beta'} u\|_{L^2} \right) \|\partial^\alpha u\|_{L^2} dt \\ &\quad + \sum_{|\alpha|=k+1} \|\partial^\alpha u_0\|_{L^2}^2 \\ &\leq C \int_0^T \sum_{|\alpha|=k+1} \left(\sum_{1 \leq |\beta| \leq k+1} \|\partial^\beta u\|_{L^2} + \gamma h \sum_{|\beta'|=k+2} \|\partial^{\beta'} u\|_{L^2} \right) \|\partial^\alpha u\|_{L^2} dt \\ &\quad + \sum_{|\alpha|=k+1} \|\partial^\alpha u_0\|_{L^2}^2 + CT\gamma \|u_0\|_{L^2}^2. \end{aligned}$$

Young's inequality ($2ab \leq \delta^{-1}a^2 + \delta b^2$) allows us to move the highest order terms from the right hand side to the left hand side and that gives

$$\begin{aligned} &\sum_{|\alpha|=k+1} \|\partial^\alpha u(T)\|_{L^2}^2 + c\gamma h \int_0^T \sum_{|\alpha|=k+1} \|\nabla \partial^\alpha u\|_{L^2}^2 dt \\ &\leq C \int_0^T \sum_{1 \leq |\beta| \leq k+1} \|\partial^\beta u\|_{L^2}^2 dt + \sum_{|\alpha|=k+1} \|\partial^\alpha u_0\|_{L^2}^2 + CT\gamma \|u_0\|_{L^2}^2. \end{aligned}$$

We now use the inductive hypothesis (5.11) (with $\varepsilon = \sqrt{\gamma h} \leq 1$) to obtain

$$\begin{aligned}
& \sum_{|\alpha|=k+1} \|(\varepsilon\partial)^\alpha u(T)\|_{L^2}^2 + c \int_0^T \sum_{|\beta|=k+2} \|(\varepsilon\partial)^\beta u\|_{L^2}^2 dt \\
& \leq C \sum_{1 \leq |\beta| \leq k+1} \varepsilon^{2(k+1-|\beta|)} \int_0^T \|(\varepsilon\partial)^\beta u\|_{L^2}^2 dt + \sum_{|\alpha|=k+1} \|(\varepsilon\partial)^\alpha u_0\|_{L^2}^2 + CT\gamma\varepsilon^{2(k+1)} \|u_0\|_{L^2}^2. \\
& \leq C \left(\sum_{1 \leq |\beta| \leq k+1} \varepsilon^{2(k+1-|\beta|)} \left(\sum_{|\alpha| \leq |\beta|} \|(\varepsilon\partial)^\alpha u_0\|_{L^2}^2 + CT\gamma(\gamma h) \|u_0\|_{L^2}^2 \right) + \sum_{|\alpha|=k+1} \|(\varepsilon\partial)^\alpha u_0\|_{L^2}^2 \right. \\
& \quad \left. + CT\gamma\varepsilon^{2(k+1)} \|u_0\|_{L^2}^2 \right) \leq C \sum_{|\alpha| \leq k+1} \|(\varepsilon\partial)^\alpha u_0\|_{L^2}^2 + CT\gamma\varepsilon \|u_0\|_{L^2}^2.
\end{aligned}$$

Combined with the inductive hypothesis this shows that (5.11) holds with k replaced by $k+1$.

Returning to (5.8) we see that (5.11) gives

$$\|U(t)\|_{H_\varepsilon^s \rightarrow H_\varepsilon^s} \leq C e^{M_0 \gamma t} \sqrt{1 + t\gamma\varepsilon}, \quad \|U(t)\|_{L^2 \rightarrow L^2} \leq C e^{M_0 \gamma t},$$

for every s . Interpolation between these two estimates provides the improved version in (5.5).

When $\sum_j \{\bar{\ell}_j, \ell_j\} \equiv 0$ then we can take $M_0 \equiv 0$ and $Q_1 \equiv 0$ in the proof and that gives (5.7) (note that in this case Q_1 vanishes and hence the last term on the right hand side of (5.11) does not appear). \square

6. AGREEMENT OF QUANTUM AND CLASSICAL DYNAMICS

In this section we obtain an accurate approximation to the solution of the Lindblad master equation which is a far reaching strengthening of Theorem 1 in §1. We remark that Theorem 4 relies only on the estimates provided by Proposition 5.1 and the abstract machinery of §4.

Theorem 4. *Suppose that \mathcal{L} is given by (1.2), assumptions (1.3), (1.10), and (1.11) hold, $h^{2\rho-1} \leq \gamma \leq 1$ for some $0 \leq \rho \leq \frac{2}{3}$. There is $C_0 > 0$ such that if $A(t)$ satisfies (in the notation of §2)*

$$\partial_t A(t) = \mathcal{L}A(t), \quad A(0) = \text{Op}(a_0), \quad a_0 \in S_\rho^{L^2},$$

then, for every N there exist $C_N > 0$ and $a(t) \in S_\rho^{L^2}$ such that

$$\begin{aligned}
\|A(t) - \text{Op}(a(t))\|_{\mathcal{L}_2} & \leq C_N e^{(M_0 + C_0 h)\gamma t} (1 + \gamma^{\frac{3}{2}} h^{\frac{1}{2}} t)^{+0} t^{N+1} h^{(2-3\rho)(N+1)}, \\
a(t) - U(t)a_0 & \in e^{M_0 \gamma t} t h^{(2-3\rho)} (1 + t\gamma^{\frac{3}{2}} h^{\frac{1}{2}})^{+0} S_\rho^{L^2},
\end{aligned} \tag{6.1}$$

where $U(t)$ was defined by (5.4).

If $\sum_j \{\ell_j, \bar{\ell}_j\} \equiv 0$, that is there is no friction (1.6), then

$$\begin{aligned} \|A(t) - \text{Op}(a(t))\|_{\mathcal{L}_2} &\leq C_N e^{C_0 h^2 \gamma t} t^{N+1} h^{(2-3\rho)(N+1)}, \\ a(t) - U(t)a_0 &\in h^{(2-3\rho)} t S_\rho^{L^2}, \end{aligned} \quad (6.2)$$

Proof of Theorem 2 assuming Theorem 4. Let $a_0 \in S_{1/2}^{L^2}$. Then observe that by Proposition 4.6 and Lemma 4.7, together with the fact that for $a_0 \in S_{1/2}^{L^2}$, $\|\text{Op}(a_0)\|_{\mathcal{L}_2} \leq C$,

$$\|A(t)\|_{\mathcal{L}_2} \leq C e^{(M_0 + C_0 h) \gamma t}.$$

Next, using Proposition 5.1

$$\|\text{Op}(U(t)a_0)\|_{\mathcal{L}_2} \leq C e^{M_0 \gamma t}.$$

Therefore, since our estimates are trivially valid when $t(1 + \gamma^{-\frac{3}{2}})h^{\frac{1}{2}} \geq 1$, we may assume without loss of generality that $t(1 + \gamma^{-\frac{3}{2}})h^{\frac{1}{2}} \leq 1$.

We can assume that $\gamma = h^{2\rho-1}$ for some $\rho \geq \frac{1}{2}$, so that the estimate (6.1) reads

$$\begin{aligned} \|A(t) - \text{Op}(a(t))\|_{\mathcal{L}_2} &\leq C_N e^{(M_0 + C_0 h) \gamma t} (t h^{\frac{1}{2}} \gamma^{-\frac{3}{2}})^{N+1} (1 + t \gamma^{\frac{3}{2}} h^{\frac{1}{2}})^{+0} \\ a(t) - U(t)a_{\lambda_h} &\in e^{(M_0 + C_0 h) \gamma t} (t h^{\frac{1}{2}} \gamma^{-\frac{3}{2}}) (1 + t \gamma^{\frac{3}{2}} h^{\frac{1}{2}})^{+0} S_\rho^{L^2}. \end{aligned} \quad (6.3)$$

Hence, since $t \gamma^{-\frac{3}{2}} h^{\frac{1}{2}} \leq 1$, the estimate (1.20) follows in this case. \square

Proof of Theorem 4. Define $a_0(t) := U(t)a_0$, with U given in (5.4). Then, recalling that $\varepsilon = \sqrt{\gamma h/2}$, $h^\rho \leq \varepsilon \leq h^{\frac{1}{2}}$, (5.5) gives

$$a_0(t) \in e^{M_0 \gamma t} (1 + t \gamma^{\frac{3}{2}} h^{\frac{1}{2}})^{+0} S_\rho^{L^2}, \quad \text{uniformly in } t \geq 0.$$

Set $A_0(t) := \text{Op}(a_0(t))$. Then, using Lemma 2.2 as in the derivation of (5.2), we obtain

$$\dot{A}_0(t) = \text{Op}(\dot{a}_0(t)) = \text{Op}(Qa_0(t)) = \mathcal{L}A_0(t) + \text{Op}(e_1(t)),$$

where

$$e_1(t) \in h^{2-3\rho} e^{M_0 \gamma t} (1 + t \gamma^{\frac{3}{2}} h^{\frac{1}{2}})^{+0} S_\rho^{L^2}.$$

Suppose, by induction that we have found

$$a_j(t) \in e^{M_0 \gamma t} t^j h^{(2-3\rho)j} (1 + t \gamma^{\frac{3}{2}} h^{\frac{1}{2}})^{+0} S_\rho^{L^2}, \quad j = 0, \dots, N-1$$

such that, with $A_{N-1} := \sum_{j=0}^{N-1} \text{Op}(a_j(t))$, we have

$$\dot{A}_{N-1} = \mathcal{L}A_{N-1}(t) + \text{Op}(e_N(t)),$$

with

$$e_N(t) \in e^{M_0 \gamma t} t^{N-1} (1 + t \gamma^{\frac{3}{2}} h^{\frac{1}{2}})^{+0} S_\rho^{L^2}.$$

Using e_N we define

$$a_N(t) = - \int_0^t U(t-s) e_N(s) ds,$$

so that

$$a_N(t) \in e^{M_0\gamma t} t^N h^{(2-3\rho)N} (1 + \gamma^{\frac{3}{2}} h^{\frac{1}{2}} t)^{+0} S_\rho^{L^2},$$

and hence, with $A_N(t) = A_{N-1}(t) + \text{Op}(a_N(t))$, we have

$$\begin{aligned} \dot{A}_N(t) &= \mathcal{L}A_{N-1}(t) + \text{Op}(e_N(t)) + \text{Op}(\dot{a}_N(t)) \\ &= \mathcal{L}A_{N-1}(t) + \text{Op}(Qa_N(t)) \\ &= \mathcal{L}A_N(t) + \text{Op}(e_{N+1}(t)), \end{aligned}$$

with

$$e_{N+1} \in e^{M_0\gamma t} t^N h^{(2-3\rho)(N+1)} (1 + \gamma^{\frac{3}{2}} h^{\frac{1}{2}} t)^{+0} S_\rho^{L^2}.$$

Note that in the last line we used Lemma 2.2 to obtain the estimates on e_{N+1} . This gives $a = \sum_{j \leq N} a_j$.

We next use Proposition 4.6 and Lemma 4.7 to compare $A(t)$ and $A_N(t)$:

$$\begin{aligned} \|A(t) - A_N(t)\|_{\mathcal{L}_2} &\leq \int_0^t \|e^{(t-s)\mathcal{L}} \text{Op}(e_{N+1}(s))\|_{\mathcal{L}_2} ds \\ &\leq C_N e^{(M_0+C_0h)\gamma t} t^{N+1} h^{(2-3\rho)(N+1)} (1 + \gamma^{\frac{3}{2}} h^{\frac{1}{2}} t)^{+0}. \end{aligned}$$

The stronger version under the assumption that $\sum_j \{\ell_j, \bar{\ell}_j\} = 0$ follows from the stronger estimates in (4.13) and (5.7). \square

7. BOUNDS ON THE HILBERT–SCHMIDT NORM OF LINDBLAD EVOLUTION

In this section we use Theorem 4 to give lower bounds on the Hilbert–Schmidt norm of the Lindblad evolution in the case of Example (1.7). We will consider two special cases: quadratic Hamiltonians and confining Hamiltonians.

7.1. Quadratic Hamiltonians. We first show that when p is quadratic and the initial condition is Gaussian, it is possible to solve (1.4) exactly. For the purposes of this section, we let B be a real, symmetric, matrix and suppose that

$$p(x, \xi) := \frac{1}{2} \langle B\rho, \rho \rangle, \quad \rho := \begin{pmatrix} x \\ \xi \end{pmatrix}. \quad (7.1)$$

We also use the notation $\Omega := \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$ for the standard symplectic form.

Lemma 7.1. *Let p as in (7.1), A_0 be a real, symmetric, positive definite matrix. $\rho_0 \in \mathbb{R}^{2n}$, and u solve*

$$(\partial_t - H_p - \frac{1}{2}\gamma h \Delta_{x,\xi})u = 0, \quad u(0) = \exp\left(-\frac{1}{2h} \langle A_0(\rho - \rho_0, \rho - \rho_0) \rangle\right). \quad (7.2)$$

Then,

$$u(t) = e^{f(t)} \exp\left(-\frac{1}{2h} \langle A(t)(\rho - \rho_0(t), \rho - \rho_0(t)) \rangle\right),$$

where $A(0) = A_0$, $\rho_0(0) = \rho_0$, $f(0) = 0$, and

$$\begin{aligned} \dot{\rho}_0(t) &= -\Omega B \rho_0 \\ \dot{A}(t) &= (A + A^t)\Omega B - \frac{\gamma}{4}(A + A^t)^2 \quad \dot{f}(t) = -\frac{\gamma}{4} \operatorname{tr}(A(t) + A^t(t)). \end{aligned} \quad (7.3)$$

Proof. We compute

$$\begin{aligned} \partial_t u &= u \left(\dot{f} - \frac{1}{2h} \langle \dot{A}(\rho - \rho_0), \rho - \rho_0 \rangle + \frac{1}{2h} \langle A \dot{\rho}_0, \rho - \rho_0 \rangle + \frac{1}{2h} \langle A(\rho - \rho_0), \dot{\rho}_0 \rangle \right), \\ &= u \left(\dot{f} - \frac{1}{2h} \langle \dot{A}(\rho - \rho_0), \rho - \rho_0 \rangle + \frac{1}{2h} \langle (A + A^t) \dot{\rho}_0, \rho - \rho_0 \rangle \right), \\ H_p u &= u \left(-\frac{1}{2h} \langle A \Omega B \rho, (\rho - \rho_0) \rangle - \frac{1}{2h} \langle A(\rho - \rho_0), \Omega B \rho \rangle \right) \\ &= u \left(-\frac{1}{2h} \langle (A + A^t) \Omega B \rho_0, \rho - \rho_0 \rangle - \frac{1}{2h} \langle (A + A^t) \Omega B (\rho - \rho_0), \rho - \rho_0 \rangle \right) \\ \gamma h \Delta u &= u \gamma h \left(\frac{1}{4h^2} \langle (A + A^t)(\rho - \rho_0), (A + A^t)(\rho - \rho_0) \rangle - \frac{1}{2h} 2 \operatorname{tr}(A + A^t) \right) \\ &= u \gamma h \left(\frac{1}{4h^2} \langle (A + A^t)^2 (\rho - \rho_0), \rho - \rho_0 \rangle - \frac{1}{2h} \operatorname{tr}(A + A^t) \right). \end{aligned}$$

Then, using that u satisfies (7.2), and equating terms by homogeneity in $\rho - \rho_0$, we obtain (7.3). \square

Remark. As an easy corollary of Lemma 7.1, we see that if $A_0 = 2I$, $B = 0$, then

$$u(t) = \frac{1}{(1 + 2\gamma t)^n} e^{-\frac{1}{h(1+2\gamma t)} \langle \rho - \rho_0, \rho - \rho_0 \rangle}, \quad \|u(t)\|_{L^2} = \left(\frac{\pi h}{2(1 + 2\gamma t)} \right)^{\frac{n}{2}}.$$

When $p = 0$, the Lindblad evolution is exactly given by the Fokker–Planck evolution, and thus the solution $A(t)$ to (1.19), satisfies

$$\|A(t)\|_{\mathcal{L}_2} = \left(\frac{1}{1 + 2\gamma t} \right)^{\frac{n}{2}}.$$

7.2. Confining Hamiltonians. We next consider the case where the Hamiltonian p is confining. We assume in this subsection that there are $c, m > 0$ and $M \in \mathbb{R}$ such that

$$\begin{aligned} \Delta p &\geq 0, \quad p \geq c|\nabla p|^2, \quad \text{on } |p| \geq M, \\ |p| &\geq c\langle(x, \xi)\rangle^m - 1/c, \quad (x, \xi) \in \mathbb{R}^{2n}. \end{aligned} \quad (7.4)$$

We show in Proposition 7.4 that under this assumption, for sufficiently dispersed initial data, the Hilbert–Schmidt norm of the Lindblad evolution is bounded from below for long times.

We will use a maximum principle type argument to show that, in the presence of a confining Hamiltonian, the Fokker–Planck evolution remains well confined in L^1 for long times. We start by constructing an effective barrier with which to apply the maximum principle.




FIGURE 4. The $\|u(t)\|_{L^2}/\|u(0)\|_{L^2}$ for the solution to (7.2) with $A = 2I$ for various choices of B in (7.1).

Lemma 7.2. *Suppose that (7.4) holds. Then for any $f \in C^\infty(\mathbb{R})$, such that $\text{supp } f' \subset (M, \infty)$, $f, f'' \geq 0$, $f'' \in C_c^\infty$, there is $C > 0$ such that, defining*

$$g(t) := \frac{g(0)}{C\gamma h t g(0) + 1}, \quad v(t, x, \xi) := \exp(-g(t)f(p(x, \xi))), \quad (7.5)$$

we have

$$(\partial_t - H_p - \frac{1}{2}\gamma h\Delta)v \geq 0, \quad t \geq 0. \quad (7.6)$$

Proof. We calculate

$$\begin{aligned} v_t - \gamma h\Delta v &= [-g'(t)f(p) + \gamma h((-g^2[f'(p)]^2 + gf''(p))|\nabla p|^2 + gf'(p)\Delta p)]v \\ &\geq [-g'(t)f(p) + \gamma h(-g^2[f'(p)]^2|\nabla p|^2)]v \\ &\geq [-g'(t)f(p) + C\gamma h(-g^2[f'(p)]^2p)]v. \end{aligned} \quad (7.7)$$

Since $f \geq 0$ and $|f''(p)| \leq C$,

$$|f'(p)| \leq C\sqrt{f(p)}.$$

Hence, using that $f'' \in C_c^\infty$, we have that there is p_0 such that $f'(p) = L$ for $p \geq p_0$ large enough and $f'(p) = 0$ for $p < M$,

$$f(p) \geq \max(c[f'(p)]^2, L(p - p_0) + f(p_0)) \geq c[f'(p)]^2p.$$

Thus, for $C > 0$ large enough, and g given in (7.5) (so that $-g'(t) - C\gamma hg^2 = 0$) the last inequality in (7.7) gives (7.6). \square

In the next lemma, we show that, given some apriori assumptions on the Fokker-Planck solution, we are able to confine the majority of its L^1 mass to a bounded set. As a consequence, we obtain that the L^2 norm cannot decay for long times.

Lemma 7.3. *Suppose that (7.4) holds. Then, $\forall R_0, c_1 > 0 \exists R_1 > 0 \forall \delta, N > 0 \exists C_{N,\delta}$ such that if*

$$0 \leq a_0(x, \xi) \leq e^{-c_1\langle(x,\xi)\rangle^2/h} \|a_0\|_{L^\infty}, \quad |(x, \xi)| \geq R_0, \quad (7.8)$$

then for any solution $a(t) \in L^\infty(\{t > 0\} \times \mathbb{R}_{(x,\xi)}^{2n})$ to

$$(\partial_t - H_p - \frac{1}{2}\gamma h\Delta)a = 0, \quad a(0) = a_0,$$

we have

$$\|a(t)\|_{L^1(\mathbb{R}^d \setminus B(0, R_1))} = C_{N,\delta} h^N \|a_0\|_{L^\infty} \quad 0 \leq t \leq h^{-1+\delta} \gamma^{-1}. \quad (7.9)$$

In particular,

$$\|a_0\|_{L^1} - C_{N,\delta} h^N \|a_0\|_{L^\infty} \leq C \|a(t)\|_{L^2}, \quad (7.10)$$

Proof. We assume without loss of generality that $0 \leq a_0 \leq 1$. Let $M_0 \geq 0$ such that

$$p \geq M_0 \implies c\langle(x, \xi)\rangle^m / 2 \leq p \leq C_1\langle(x, \xi)\rangle^2 \quad \text{and} \quad |(x, \xi)| \geq R_0.$$

Then, let $\psi \in C_c^\infty((M_0, M_0+1); [0, \infty))$ with $\int \psi = 1$ and define $f(x) := \int_0^x \int_0^s \psi(t) dt ds$ so that $f''(x) = \psi(x)$, $\text{supp } f \subset (M_0, \infty)$ with $f, f'' \geq 0$. Let $R_1 \geq 0$ such that $p \geq M_0 + 1$ on $|(x, \xi)| \geq R_1$.

Since $f(p(x, \xi)) = 0$ on $|(x, \xi)| \leq R_0$, $f(p) \leq Cp \leq C\langle(x, \xi)\rangle^2$, and a_0 satisfies (7.8) there exists $c_0 > 0$ such that

$$\exp(-c_0 f(p(x, \xi))/h) \geq a_0.$$

We now apply Lemma 7.2 with $g(0) = c_0/h$: for v in (7.5)

$$(\partial_t - H_p - \frac{1}{2}\gamma h\Delta)(v - a) \geq 0, \quad t \geq 0.$$

The maximum principle [Co80, Theorem 1] then shows that $0 \leq a \leq v$ and consequently, using that $f(p) \geq cp \geq c\langle(x, \xi)\rangle^m$ on $\mathbb{R}^{2n} \setminus B(0, R_1)$,

$$\begin{aligned} \|a(t)\|_{L^1(\mathbb{R}^{2n} \setminus B(0, R_1))} &\leq \|v(t)\|_{L^1(\mathbb{R}^{2n} \setminus B(0, R_1))} \\ &\leq \int_{\mathbb{R}^{2n} \setminus B(0, R_1)} e^{-c_0\langle(x, \xi)\rangle^m / (C\gamma t + h)} dx d\xi \leq C e^{-c_0 / (C\gamma t + h)}, \end{aligned}$$

from which (7.9) follows.

To obtain (7.10), observe that

$$\partial_t \int a dx d\xi = \int (H_p + \frac{1}{2}\gamma h\Delta) a dx d\xi = 0.$$

Hence,

$$\begin{aligned} \|a_0\|_{L^1} = \|a(t)\|_{L^1} &\leq \|a(t)\|_{L^1(B(0, R_1))} + C_{N, \delta} h^N \\ &\leq C R_1^n \|a(t)\|_{L^2(B(0, R_1))} + C_{N, \delta} h^N \\ &\leq C R_1^n \|a(t)\|_{L^2} + C_{N, \delta} h^N. \end{aligned}$$

□

Finally, we show that the Hilbert–Schmidt norm of the Lindblad evolution with a confining Hamiltonian can, in many cases, be effectively controlled from below.

Proposition 7.4. *Suppose that \mathcal{L} is given by (1.2), assumptions (1.3), and (7.4) hold, that ℓ_j 's are as in (1.7). Let $0 \leq \rho \leq \frac{2}{3}$ and assume $h^{2\rho-1} \leq \gamma \leq 1$. There is $C_0 > 0$ such that for $a_0 \in S_\rho^{L^2}$ satisfying (7.8), for any $N > 0$ and $\delta > 0$, there is $C > 0$ such that*

$$(2\pi h)^{\frac{n}{2}} \operatorname{tr} A(0) - C(e^{C_0 h^2 \gamma t} t h^{(2-3\rho)} + h^N) \|a_0\|_{L^\infty} \leq C \|A(t)\|_{\mathcal{L}_2}$$

for $0 \leq t \leq h^{-1+\delta} \gamma^{-1}$. Here

$$\partial_t A(t) = \mathcal{L}A(t), \quad A(0) = \operatorname{Op}(a_0).$$

Remark. When $a_0 \in C_c^\infty(\mathbb{R}^{2n})$ is independent of h (as in Theorem 1) notice that $\|a_0\|_{L^1} \geq C \|a_0\|_{L^2}$, where the constant depends on the support and hence

$$\|A(0)\|_{\mathcal{L}_2} = (2\pi h)^{-\frac{n}{2}} \|a_0\|_{L^2} \leq C (2\pi h)^{-\frac{n}{2}} \|a_0\|_{L^1} = C (2\pi h)^{\frac{n}{2}} \operatorname{tr} A(0). \quad (7.11)$$

Proof. By Theorem 4,

$$\|A(t) - \operatorname{Op}(a(t))\|_{\mathcal{L}_2} \leq C e^{C_0 h^2 \gamma t} (1 + \gamma) t h^{(2-3\rho)} \quad (7.12)$$

where $a(t) = U(t)a_0$. In particular, $a(t)$ satisfies

$$(\partial_t - H_p - \frac{1}{2}\gamma h \Delta)a = 0, \quad a(0) = 0,$$

and $a(t) \in S_\rho^{L^2}$. Since $a(t) \in S_\rho^{L^2}$, uniformly in $t > 0$, by the Sobolev embedding, $a(t) \in L^\infty(\{t > 0\} \times \mathbb{R}_{(x,\xi)}^{2n})$ and hence applying Lemma 7.3 then yields

$$\begin{aligned} (2\pi h)^n \operatorname{tr} A(0) - Ch^N \|a_0\|_{L^\infty} &= \|a_0\|_{L^1} - Ch^N \|a_0\|_{L^\infty} \\ &\leq C \|a(t)\|_{L^2} = C(2\pi h)^{\frac{n}{2}} \|\operatorname{Op}(a(t))\|_{\mathcal{L}_2}. \end{aligned} \quad (7.13)$$

The Proposition now follows from combining (7.12) and (7.13). \square

APPENDIX A. OPERATORS WITH QUADRATIC SYMBOL GROWTH

We start with the composition formula of operators quantizing symbols in $S_{(k)}$ where that space was defined in (2.3).

Proposition A.1. *Suppose that $a_j \in S_{(k_j)}$, $j = 1, 2$. Then $\operatorname{Op}(a_1)\operatorname{Op}(a_2) = \operatorname{Op}(b)$, where for any $N \geq \max(k_1, k_2)$,*

$$b(x, \xi, h) - \sum_{\ell=0}^{N-1} \frac{1}{\ell!} \left(\frac{h}{2i} \sigma(D_x, D_\xi, D_y, D_\eta) \right)^\ell a_1(x, \xi) a_2(y, \eta)|_{x=y, \xi=\eta} \in h^N S_0, \quad (\text{A.1})$$

where σ is the standard symplectic form on \mathbb{R}^{2n} .

Remark. Note that b in the statement of the proposition is not necessarily in an $S_{(k)}$ class since they are not closed under multiplication.

Proof. We observe that $S_{(k)} \subset S(m_k)$, $m_k(x, \xi) = (1 + |x| + |\xi|)^k$. Hence [Zw12, Theorem 4.18] applies and, writing $z = (x, \xi)$, $w = (y, \eta)$,

$$b(z, h) = \exp(ihA(D))(a_1(x)a_2(w))|_{z=w}, \quad A(D_{z,w}) = -\frac{1}{2}\sigma(D_x, D_\xi, D_y, D_\eta).$$

By Taylor's formula,

$$b(z, h) = \sum_{\ell=0}^{N-1} \frac{1}{\ell!} (ihA(D))^\ell (a_1(z)a_2(w))|_{z=w} + R_N(z, h)$$

where

$$R_N(z, h) : \frac{1}{(N-1)!} (1-t)^{N-1} e^{ithA(D)} (ihA(D))^N ((a_1(z)a_2(w))|_{z,w}.$$

For $N \geq \max(k_1, k_2)$, $A(D)^N a_1(z)a_2(w) \in S_0(\mathbb{R}_{z,w}^{4n})$ and since $e^{ihtA(D)} : S_0(\mathbb{R}^{4n}) \rightarrow S(\mathbb{R}^{4n})$ (with uniform bounds for $0 \leq t \leq 1$ – see [Zw12, Theorem 4.17]) we conclude that $R_N \in h^N S_0(\mathbb{R}^{2n})$ which is (A.1). \square

We now present a general spectral result following the proof of a special case in [Hö95] (see the example in [Zw12, §C.2.2]):

Proposition A.2. *Suppose that $p(x, \xi) \in C^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ satisfies*

$$|\partial^\alpha p(x, \xi)| \leq C_\alpha(1 + |x| + |\xi|), \quad |\alpha| \geq 1, \quad (\text{A.2})$$

and define

$$\begin{aligned} N_p u &= p^w(x, D)u, \quad \mathcal{D}(N_p) := \mathcal{S}(\mathbb{R}^d), \\ M_p u &= p^w(x, D)u, \quad \mathcal{D}(M_p) := \{u \in L^2(\mathbb{R}^d) : p^w(x, D)u \in L^2(\mathbb{R}^d)\}, \end{aligned}$$

where in the case of $u \in L^2(\mathbb{R}^d) \subset \mathcal{S}'(\mathbb{R}^d)$ we consider $p^w(x, D)u \in \mathcal{S}'(\mathbb{R}^d)$. Then M_p is closed and

$$M_p = \overline{N_p}, \quad M_p^* = M_{\bar{p}}. \quad (\text{A.3})$$

Proof. We recall that $p^w(x, D) : \mathcal{S}' \rightarrow \mathcal{S}'$ is continuous and hence, if $u_j \rightarrow u$ and $p^w(x, D)u_j \rightarrow v$ in L^2 , then $u_j \rightarrow u$ in \mathcal{S}' . Consequently, $v = p^w(x, D)u \in L^2$, $u \in \mathcal{D}(M_p)$ and $M_p u = v$. This shows that M_p is closed.

To show that M_p is the closure of N_p we have to show that for any $u \in \mathcal{D}(M_p)$ there exists a family $u_\varepsilon \in \mathcal{S}$ such that $u_\varepsilon \rightarrow u$ and

$$p^w(x, D)u_\varepsilon \rightarrow p^w(x, D)u \quad \text{in } L^2 \text{ as } \varepsilon \rightarrow 0. \quad (\text{A.4})$$

To construct u_ε we take $\chi \in C_c^\infty(\mathbb{R}^{2d})$ equal to one in $B_{\mathbb{R}^{2n}}(0, 1)$, and put

$$u_\varepsilon := \chi_\varepsilon^w(x, D)u \in \mathcal{S}, \quad \chi_\varepsilon(x, \xi) := \chi(\varepsilon x, \varepsilon \xi), \quad u_\varepsilon \rightarrow u \quad \text{in } L^2.$$

Then $p^w u_\varepsilon = \chi_\varepsilon^w p^w u_\varepsilon + [p^w, \chi_\varepsilon^w]u_\varepsilon$ and as $\chi_\varepsilon^w p^w u \rightarrow p^w$ in L^2 , we need to show that

$$[p^w(x, D), \chi_\varepsilon^w(x, D)]u \rightarrow 0 \quad \text{in } L^2, \quad \varepsilon \rightarrow 0. \quad (\text{A.5})$$

To see this we note that [Zw12, Theorem 4.18] and the two term Taylor expansion of $e^{iA(D)}$ give

$$[p^w(x, D), \chi_\varepsilon(x, D)] = a_\varepsilon^w(x, D), \quad a_\varepsilon(x, \xi) = i\{\chi_\varepsilon, p\}(x, \xi) + b_\varepsilon(x, \xi), \quad (\text{A.6})$$

where

$$b_\varepsilon(x, \xi) := \int_0^1 (1-t) (e^{itA(D)}(iA(D))^2 (p(x, \xi)\chi_\varepsilon(y, \eta) - p(y, \eta)\chi_\varepsilon(x, \xi)))|_{x=y, \xi=\eta} dt,$$

and $A(D) := \sigma(D_x, D_\xi; D_y, D_\eta)$. In view of (A.2),

$$\{\chi_\varepsilon, p\}(x, \xi) = \varepsilon \sum_{j=1}^n (\partial_{x_j} p(x, \xi)(\partial_{\xi_j} \chi)(\varepsilon x, \varepsilon \xi) - \partial_{\xi_j} p(x, \xi)(\partial_{x_j} \chi)(\varepsilon x, \varepsilon \xi))$$

is bounded in $S(1)$, uniformly as $\varepsilon \rightarrow 0$.

To obtain estimates on b_ε we observe that, for some $c_{\alpha\beta} \in \mathbb{C}$,

$$\begin{aligned} & (iA(D))^2 (p(x, \xi)\chi_\varepsilon(y, \eta) - p(y, \eta)\chi_\varepsilon(x, \xi)) \\ &= \sum_{|\alpha|+|\beta|=2} c_{\alpha\beta} \varepsilon^2 (\partial^\alpha p(x, \xi) \partial^\beta \chi(\varepsilon y, \varepsilon \eta) - \partial^\alpha \chi(\varepsilon x, \varepsilon \xi) \partial^\beta p(y, \eta)) \in \varepsilon(S(m) + S(1/m)), \end{aligned}$$

where the order function is given by $m(x, \xi, y, \eta) := \langle x, \xi \rangle \langle y, \eta \rangle^{-1}$. The inclusion follows from the fact $\varepsilon \leq C \langle \varepsilon(x, \xi) \rangle^{-1}$ for $(x, \xi) \in \text{supp } \chi$ and from the assumption (A.2). By [Zw12, Theorem 4.17], the operators $e^{ithA(D)} : S(m^{\pm 1}) \rightarrow S(m^{\pm 1})$ are bounded uniformly in t . Since $m|_{x=y, \xi=\eta} = 1$, $b_\varepsilon \in \varepsilon S(1)$, and [Zw12, Theorem 4.23] gives, uniformly as $\varepsilon \rightarrow 0$,

$$\|b_\varepsilon^w(x, D)\|_{L^2 \rightarrow L^2} \leq C\varepsilon. \quad (\text{A.7})$$

We now choose $\psi \in C_c^\infty(\mathbb{R}^n)$ supported in $B_{\mathbb{R}^{2n}}(0, 1)$, equal to one near 0, and put $\psi_\varepsilon(x, \xi) = \psi(\varepsilon x, \varepsilon \xi)$. Then $\{\chi_\varepsilon, p\}(x, \xi) \psi_\varepsilon(x, \xi) \equiv 0$, and [Zw12, Theorems 4.18 and 4.23] imply

$$\|\{\chi_\varepsilon, p\}^w(x, D) \psi_\varepsilon^w(x, D)\|_{L^2 \rightarrow L^2} \leq C\varepsilon.$$

This and (A.7) give

$$\begin{aligned} [p^w(x, D), \chi_\varepsilon^w(x, D)] &= \{\chi_\varepsilon, p\}^w(x, D)(1 - \psi_\varepsilon^w(x, D)) + \{\chi_\varepsilon, p\}^w(x, D) \psi_\varepsilon^w(x, D) + b_\varepsilon^w(x, D) \\ &= \{\chi_\varepsilon, p\}^w(x, D)(1 - \psi_\varepsilon^w(x, D)) + \mathcal{O}(\varepsilon)_{L^2 \rightarrow L^2}. \end{aligned}$$

Since $\psi_\varepsilon^w(x, D)u \rightarrow u$ in L^2 and $\{\chi_\varepsilon, p\} \in S(1)$ (hence by [Zw12, Theorem 4.23] $\|\{\chi_\varepsilon, p\}^w(x, D)\|_{L^2 \rightarrow L^2}$ is uniformly bounded), this and (A.7) give (A.5).

It remains to show the last assertion in (A.3). For that we recall that $v \in \mathcal{D}(M_p^*)$ if and only if there exists $C = C(v)$ such that for all $u \in \mathcal{D}(M_p)$

$$\langle M_p u, v \rangle \leq C \|u\|_{L^2}. \quad (\text{A.8})$$

For $u \in \mathcal{S} \subset \mathcal{D}(M_p)$ we have $\langle M_p u, v \rangle = \langle u, \bar{p}^w(x, D)v \rangle$, where $\bar{p}^w(x, D)v \in \mathcal{S}'$ and (A.8) implies that $\bar{p}^w(x, D)v \in L^2$. Hence $M_p^* \subset M_{\bar{p}}$. Since M_p^* is closed, $N_{\bar{p}} \subset N_p^* = \bar{N}_p^* = M_p^*$. It follows that $M_{\bar{p}} = \bar{N}_{\bar{p}} \subset M_p^*$ and that $M_p^* = M_{\bar{p}}$. \square

APPENDIX B. BY ZHEN HUANG AND MACIEJ ZWORSKI

We describe the results of numerical experiments illustrating the difference between the Schrödinger (1.1) and Lindblad (1.2) evolutions, and their relation to the corresponding classical evolutions given by the Hamiltonian flow and the Fokker–Planck equation (1.4), respectively. In particular, for both the Hilbert–Schmidt and trace class norms, we compare the results to the bounds (1.13), (1.21), and the corresponding bounds in [HRR25, Theorem 1.1].

If the Hamiltonian acts on $L^2(\mathbb{R}^n)$ then the computations for the Lindblad evolution and the Fokker–Planck equation have to be performed in \mathbb{R}^{2n} , which is a dramatic increase of dimension. This forces us, at this early stage, to restrict our attention to $n = 1$ which does not allow for chaotic behaviour. However, see [HRR25, §1.3] for pointers to the physics literature where time dependent chaotic systems were considered.

We choose a one dimensional model in which a hyperbolic point occurs in the classical dynamics and hence we observe a $\log 1/h$ Ehrenfest time: the classical/quantum




FIGURE 5. Comparison of Lindblad and Fokker–Planck evolutions for P as in (B.1) and L_j as in (B.2) at time $T = 2$ and for initial data given by a coherent state (B.7) with $h_0 = h$, $x_0 = -1$, $\xi_0 = 0$, and $\gamma = h^\delta$, $\delta = 0, \frac{1}{12}, \frac{1}{6}, \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{5}{6}, 1$. The figure illustrates Theorem 2 and the estimates (1.22) from [HRR25], with the dashed lines given by $\log(1/h) \rightarrow \log h^{\frac{3\delta-1}{2}}$, which corresponds to the exponents in those estimates. The change of behaviour at $\delta = \frac{1}{3}$ (the norms of the error decrease with h for $\delta \leq \frac{1}{3}$) suggests that the bounds in (1.13) and (1.22) are accurate but we do not have conclusive evidence, see Figure 7. We also show the behaviour of the Hilbert–Schmidt norm which is consistent with the results of §7 (see the Remark after Theorem 2).

correspondence breaks down at that time – see (1.18) and Figures 2,3. The model is




FIGURE 6. An analogue of Figure 5 for the initial data given by a_0 with $h_0 = 2^{-3}$, $x_0 = -0.8$, $\xi_0 = 0$. The agreement of the evolutions seems dramatically better than the estimates given in Theorem 1, especially in view of the fact that those estimates are only non-trivial for $\gamma = h^\delta$ and $\delta \leq \frac{1}{3}$. The behaviour of the Hilbert Schmidt norm is consistent with the results of §7 (see Remark 2 after Theorem 1).

given by the Schrödinger operator with a double well potential:

$$\begin{aligned}
 P &= (hD_x)^2 + V(x), \quad D_x = (1/i)\partial_x, \\
 V(x) &= \left(x^2 - \frac{1}{4}\right)^2.
 \end{aligned}
 \tag{B.1}$$

We choose the jump operators, L_j , so that the Fokker-Planck equation (1.4) takes the simple form (1.7):

$$L_1 = x, \quad L_2 = hD_x, \quad L_j^* = L_j.
 \tag{B.2}$$

The specific choice of constants in V appearing in (B.1) is dictating by numerical considerations: we need our dynamics to be confined to a box of size $[2\pi, 2\pi]$.

We write the Lindblad evolution equation (1.2) as a differential equation for the Schwartz kernel of A , $A(x, y)$, $x, y \in \mathbb{R}$:

$$\begin{aligned} \partial_t A(t) &= \mathcal{L}A(t), \quad A(0) = A_0, \quad A = A(x, y), \\ (\mathcal{L}A)(x, y) &= \frac{i}{h} \left(-(h\partial_x)^2 + V(x) + (h\partial_y)^2 - V(y) \right) A \\ &\quad - \frac{\gamma}{2h} \left((x-y)^2 - (h\partial_x + h\partial_y)^2 \right) A. \end{aligned} \tag{B.3}$$

The Fokker-Planck equation (1.4) is an evolution equation for functions in phase space, that is, functions of (x, ξ) :

$$\begin{aligned} \partial_t a(t) &= Qa(t), \quad a(0) = a_0, \quad a(t) = a(t, x, \xi), \\ Q &= 2\xi\partial_x - V'(x)\partial_\xi + \frac{1}{2}\gamma h \left(\partial_x^2 + \partial_\xi^2 \right). \end{aligned} \tag{B.4}$$

B.1. Numerical schemes. We use the following numerical methods to solve (B.4). The operator Q is split into two parts, $Q = Q_1 + Q_2$, where

$$Q_1 := 2\xi\partial_x + \frac{1}{2}\gamma h\partial_x^2, \quad Q_2 := -V'(x)\partial_\xi + \frac{1}{2}\gamma h\partial_\xi^2. \tag{B.5}$$

On the Fourier transform side in the x variable (with the Fourier transform variable denoted by k_x), the operator Q_1 acts by multiplication:

$$\widehat{Q_1 a_1}(k_x, \xi) := \widehat{Q_1 a_1}(k_x, \xi) = \left(2i\xi k_x - \frac{1}{2}\gamma h k_x^2 \right) \widehat{a_1}(k_x, \xi).$$

and thus the corresponding evolution operator could also be written explicitly:

$$e^{t\widehat{Q_1}} \widehat{a_1}(k_x, \xi) = e^{t(2i\xi k_x - \frac{1}{2}\gamma h k_x^2)} \widehat{a_1}(k_x, \xi).$$

Similarly, on the Fourier transform side in the ξ variable (with the Fourier variable denoted by k_ξ), the operator Q_2 acts diagonally:

$$\widehat{Q_2 a_2}(x, k_\xi) := \widehat{Q_2 a_2}(x, k_\xi) = \left(-iV'(x)k_\xi - \frac{1}{2}\gamma h k_\xi^2 \right) \widehat{a_2}(x, k_\xi).$$

and the corresponding evolution operator is

$$e^{t\widehat{Q_2}} \widehat{a_2}(x, k_\xi) = e^{t(-iV'(x)k_\xi - \frac{1}{2}\gamma h k_\xi^2)} \widehat{a_2}(x, k_\xi).$$

This motivates the following operator-splitting scheme for time-stepping, known as the (second-order) Lie-Suzuki-Trotter formula:

$$\begin{aligned} e^{n\Delta t Q} &= \left(e^{\Delta t Q} \right)^n = \left(e^{\frac{\Delta t}{2} Q_1} e^{\Delta t Q_2} e^{\frac{\Delta t}{2} Q_1} \right)^n + \mathcal{O}((\Delta t)^2) \\ &= e^{\frac{\Delta t}{2} Q_1} \left(e^{\Delta t Q_2} e^{\Delta t Q_1} \right)^{n-1} e^{\Delta t Q_2} e^{\frac{\Delta t}{2} Q_1} + \mathcal{O}((\Delta t)^2). \end{aligned}$$

Numerically, the Fourier transform in 1D is handled by the 1D FFT function.




FIGURE 7. For the initial data in Figure 5 (a coherent state with an h dependent symbol) we take $\gamma = h^\delta$ with $\delta = \frac{1}{3}, \frac{1}{4}, \frac{1}{6}, \frac{1}{12}, 0$ and time equal to $T = h^{-\frac{1-3\delta}{2}}$ which is the limit of the validity of estimates in Theorem 2 and in (1.22) (from [HRR25]). The results confirm their validity but do not indicate optimality in our model.

To solve the Lindblad equation (B.3), we first change variable and define

$$B(w, z) := A(w + z, w - z).$$

The Lindbladian then becomes

$$\mathcal{L} = \frac{i}{h} (-h^2 \partial_z \partial_w + V(w + z) - V(w - z)) - \frac{\gamma}{2h} (4z^2 - h^2 \partial_w^2). \quad (\text{B.6})$$

As in (B.5) we write $\mathcal{L} = \mathcal{L}_1 + \mathcal{L}_2$,

$$\mathcal{L}_1 := \frac{i}{h} (V(w + z) - V(w - z)) - \frac{2\gamma}{h} z^2, \quad \mathcal{L}_2 := -ih \partial_z \partial_w + \frac{\gamma h}{2} \partial_w^2.$$




FIGURE 8. The analogue of Figure 7 with the initial data from Figure 6. It confirms the validity of the estimates in Theorem 1 but suggests that in the model we considered they might not be optimal.

The operator \mathcal{L}_2 acts by multiplication on the Fourier transform side:

$$\mathcal{L}_2 \widehat{B}(k_w, k_z) = \left(i\hbar k_w k_z - \frac{\gamma \hbar}{2} k_w^2 \right) \widehat{B}(k_w, k_z),$$

while \mathcal{L}_1 is already a multiplication operator in the physical space. We again proceed using the Lie–Suzuki–Trotter formula. The MATLAB codes are included in §B.3.

B.2. Numerical experiments. The numerical results are shown in figures in §1 and in this Appendix and we refer to their captions for the interpretation of the results.

When comparing the quantum (Lindblad) and classical (Fokker–Planck) evolution in Hilbert–Schmidt norms we take

$$A(0) = a(x, \hbar D, \hbar), \quad a(x, \xi, \hbar) = \sqrt{2\pi\hbar} a_0(x, \xi),$$

where

$$a_0(x, \xi) = \sqrt{\frac{2}{\pi h_0}} e^{-\frac{(x-x_0)^2}{h_0} - \frac{(\xi-\xi_0)^2}{h_0}}, \quad \|a_0\|_{L^2} = 1. \quad (\text{B.7})$$

The case of h_0 fixed but small (so that a_0 is numerically compactly supported) corresponds to the assumptions in Theorem 1 and $h_0 = h$ to those in Theorem 2, and in [HRR25]. When we compare trace class norms we normalize the initial data so that $\|A(0)\|_{\mathcal{L}_1} = 1$.

The initial value for Lindblad equation (B.6) (for the rotated Schwartz kernel B) is

$$B(w, z) = \frac{2}{\sqrt{\pi h}} e^{-\frac{(w-x_0)^2}{h_0}} e^{-\frac{h_0}{h^2} z^2} e^{i\frac{2\xi_0 z}{h}}. \quad (\text{B.8})$$

We recall the general relation between $a(x, \xi)$ and $B(w, z)$:

$$B(x, y) = \sqrt{\frac{2}{\pi h}} \int a(x, \xi) e^{i\frac{2y\xi}{h}} d\xi, \quad a(x, \xi) = \sqrt{\frac{1}{2\pi h}} \int B(x, y) e^{-i\frac{2\xi y}{h}} dy. \quad (\text{B.9})$$

B.3. Matlab codes. The first code solves the Fokker–Planck equation as described in §B.1:

```

1 function u = fp(dV, h, gamma_x, gamma_p, u, Nt, dt, N)
2 % solving Fokker-Planck equation on [-a,a] * [-a,a] using N * N grid points,
3 % with initial data u, time step dt, for Nt time steps,
4 % with force dV(x), semiclassical parameter h,
5 % and diffusion coefficients gamma_x and gamma_p.
6
7 % Setup grids
8 a = pi; dx = 2*a/N; dp = 2*a/N;
9 x = -a:dx:a-dx; p = -a:dp:a-dp; [XX, PP] = ndgrid(x, p);
10 K = [0:N/2-1 0 -N/2+1:-1]*(pi/a);
11 [Kx, Kp] = ndgrid(K, K);
12
13 % construct exp(Q1*dt/2) and exp(Q2*dt)
14 Q1 = 2*PP.*(1j*Kx) - gamma_x*h/2.0 * (Kx.^2);
15 Q2 = -dV(XX).*(1j*Kp) - gamma_p*h/2.0 * (Kp.^2);
16 expQ1_2 = exp(Q1*dt/2); expQ1 = exp(Q1*dt); expQ2 = exp(Q2*dt);
17 for n = 0 : Nt-1
18     % evolve Q1 for half a time step if t = 0
19     if n == 0
20         u1 = fft(u);
21         u1 = expQ1_2.*u1;
22         u = ifft(u1);
23     end

```

```

24
25 % evolve Q2 for a full time step
26 u1 = transpose(fft(transpose(u)));
27 u1 = expQ2.*u1;
28 u = transpose(ifft(transpose(u1)));
29
30 % evolve Q1 for a full time step
31 u1 = fft(u);
32 if n < Nt-1
33     u1 = expQ1.*u1;
34 else
35     u1 = expQ1_2.*u1; % evolve Q1 for a half time step if t = T
36 end
37 u = ifft(u1);
38 end

```

The next code gives the solution to the Lindblad equation (B.3): in the notation of (B.6) we obtain B (denoted by ρ in the code) with the initial data given by ρ . An additional feature is allowing for different γ 's in front of Δ_x and Δ_ξ in (1.7), γ_x and γ_p respectively. In our experiment we take $\gamma_x = \gamma_p$.

```

1 function rho = lindblad(V, h, gamma_x, gamma_p, rho, Nt, dt, N)
2 % solving Lindblad equation on [-a,a] * [-a,a] using N * N grid points,
3 % with initial data u, time step dt, for Nt time steps,
4 % with force V(x), semiclassical parameter h,
5 % and diffusion coefficients gamma_x and gamma_p.
6
7 % Setup grids
8 a = pi; dw = 2*a/N; dz = 2*a/N;
9 w = -a:dw:a-dw; z = -a:dz:a-dz; [ww, zz] = ndgrid(w, z);
10 K = [0:N/2-1 0 -N/2+1:-1]*(pi/a);
11 [Kw, Kz] = ndgrid(K, K);
12
13 % construct exp(Q1*dt/2) and exp(Q2*dt)
14 L1 = 1i*(V(ww+zz)-V(ww-zz))/h - 2*gamma_p*(zz.^2)/h;
15 L2 = 1i*(Kw.*Kz)*h-h*gamma_x*(Kw.^2)/2;
16 expL1dt_2 = exp(L1*dt/2.0); expL1dt = exp(L1*dt); expL2dt = exp(L2*dt);
17
18 for n = 0 : Nt-1
19     % evolve L1 for half a time step if t = 0
20     if n == 0

```

```

21     rho_tmp = expL1dt_2.*rho;
22 end
23
24 % evolve L2 for a full time step
25 rho_hat = ifft2(rho_tmp);
26 rho_hat = expL2dt.*rho_hat;
27 rho_tmp = fft2(rho_hat);
28
29 % evolve Q1 for a full time step
30 if n < Nt-1
31     rho_tmp = expL1dt.*rho_tmp;
32 else
33     % evolve Q1 for a half time step if t = T
34     rho = expL1dt_2.*rho_tmp;
35 end
36 end

```

The next matlab codes implements the transformations between $a(x, \xi)$ and $B(w, z)$ given in (B.9) via equidistance quadrature:

```

1 function rho = a2B(h, p, u)
2 % transform a(x,p) to B(w,z), h is the semiclassical parameter,
3 % p is the grid points as defined before, i.e. p = -a:dp:a-dp;
4 % u is function a(x,p) evaluated on the N*N mesh,
5 % as defined in previous codes.
6
7 Kernel_ifft = exp(1j*2*p'*p/h)*dp/(sqrt(2));
8 rho = u* Kernel_ifft;
9
10 end
11
12 function u = B2a(h, z, rho)
13 % transform B(w,z) to a(x,p), h is the semiclassical parameter,
14 % z is the grid points as defined before, i.e. z = -a:dz:a-dz;
15 % rho is function B(w,z) evaluated on the N*N mesh,
16 % as defined in previous codes.
17
18 Kernel_fft = exp(-1j*2*y'*y/h)*dy*sqrt(2)/(pi*h);
19 u = rho* Kernel_fft;
20
21 end

```

Finally we give the code for evaluating the Hilbert-Schmidt norm and the trace norm of data given as $B(w, z)$. Since $B(w, z) = A(w + z, w - z)$, therefore $\|A\|_{\text{HS}} = \frac{1}{\sqrt{2}}\|B\|_{\text{HS}}$. To calculate the trace norm, we first reconstruct $A(x, y)$ on a $2N \times 2N$ grid from $B(w, z)$ on a $N \times N$ grid by rotation and interpolation, and then calculate the trace norm (also known as the nuclear norm in linear algebra) of this $2N \times 2N$ matrix using singular value decomposition (SVD).

```

1  function [HS_norm, Tr_norm] = norm_calc(h, rho, a, N)
2  % calculate the Hilbert-Schmidt norm and the trace norm of B(w,z)
3  % h is the semiclassical parameter, rho is the function B(w,z)
4  % evaluated on the N*N mesh,
5  % a is the computation region,
6  % i.e. the calculation is done on [-a,a]*[-a,a],
7  % N is the number of grid points in each direction.
8
9  err_HS = sqrt((sum(abs(rho(:)).^2))*((2*a/N)^2) /2);
10
11 rho_rot = rho_rotate(rho);
12 err_trace = (sum(svd(rho_rot)))*(2*a/(2*N));
13 end
14
15 function rho_rotate = rho_rotate(rho)
16 % by rotation and interpolation, reconstruct A(x,y) from B(w,z):
17 N = size(rho,1); rho_reflect = rho(:, N:-1:1);
18 rho_rotate = zeros(2*N);
19 for i = 1:2*N-1
20     rho_diagonal = diag(rho_reflect, N-i);
21     L = length(rho_diagonal);
22     rho_fit = rho_diagonal;
23     if L>1
24         rho_fit = spline(1:L,rho_diagonal,1:0.5:L) ;
25     end
26     idy = (1:(2*L-1))-L + N+1;
27     rho_rotate(i,idy) = rho_fit;
28 end
29 end

```

REFERENCES

- [ChPa17] D. Chruściński and S. Pascazio, *A brief history of the GKLS equation*, Open Systems & Information Dynamics **24**(2017), 1740001.

- [Co80] C. Cosner, *Asymptotic behavior of solutions of second order parabolic partial differential equations with unbounded coefficients*, Journal of Differential Equations, **35**(1980), 407–428.
- [Da77] E.B. Davies, *Quantum dynamical semigroups and the neutron diffusion equation*, Reports on Mathematical Physics **11**(1977), 169–188.
- [DiSj99] M. Dimassi and J. Sjöstrand, *Spectral Asymptotics in the Semi-Classical Limit*, Cambridge U Press, 1999.
- [GoBr99] J. Gong and P. Brumer, *Decoherence and correspondence in conservative chaotic dynamics*, Phys. Rev. E **60**(1999), 1643–1647.
- [GKS76] V. Gorini, A. Kossakowski and E.C.G. Sudarshan, *Completely positive dynamical semigroups of N -level systems*, J. Math. Phys. **17**(1976), 821–825.
- [HRR25] F. Hernández, D. Ranard, C.J. Riedel, *Classical correspondence beyond the Ehrenfest time for open quantum systems with general Lindbladians*. Commun. Math. Phys. 406, **4** (2025).
- [HRR23] F. Hernández, D. Ranard, C.J. Riedel, *Decoherence ensures classicality beyond the Ehrenfest time as $\hbar \rightarrow 0$* , [arXiv:2306.13717](#)
- [Hö95] L. Hörmander, *Symplectic classification of quadratic forms and general Mehler formulas*, Math. Z. **219**(1995), 413–449.
- [Li24] Z. Li, *Long time quantum-classical correspondence for open systems in trace norm*. [arXiv:2408.16953](#).
- [Li76] G. Lindblad, *On the generators of quantum dynamical semigroups*, Commun. Math. Phys. **48**(1976), 119–130.
- [OlBo23] I. Oltman and Y. Borns-Weil, *Particle trajectories for quantum maps*, Ann. H. Poincaré (2023) published online, [arXiv:2210.03224](#).
- [Sm20] H.F. Smith, *Parametrix for a semiclassical subelliptic operator*, Anal.&PDE **7**(2020), 2375–2398.
- [Sm25] H.F. Smith, *Uniform subelliptic estimates for degenerating Fokker–Planck equations*, in preparation.
- [UnZu89] W.G. Unruh and W.H. Zurek, *Reduction of a wave packet in quantum orangeian motion*, Phys. Rev. D **40**(1989), 1071–1094.
- [Zw12] M. Zworski, *Semiclassical Analysis*, Graduate Studies in Mathematics **138**, AMS, 2012.

Email address: j.galkowski@ucl.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, WC1H 0AY, UK

Email address: zworski@math.berkeley.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA

Email address: hertz@math.berkeley.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA.